



Cite this: *Org. Biomol. Chem.*, 2015, **13**, 2026

A coumarin based gold(i)-alkynyl complex: a new class of supramolecular hydrogelators†

Artur J. Moro,^a Bertrand Rome,^{a,b} Elisabet Aguiló,^c Julià Arcau,^c Rakesh Puttreddy,^d Kari Rissanen,^d João Carlos Lima*^a and Laura Rodríguez*^c

A phosphine-gold(i)-alkynyl-coumarin complex, [Au(7-(prop-2-ine-1-yloxy)-1-benzopyran-2-one)-(DAPTA)] (**1**), was synthesized and the formation of long luminescent fibers in solution was characterized via fluorescence microscopy and dynamic light scattering. The fibers presented strong blue and green luminescence, suggesting that the gold(i) in the complex increased intersystem crossing due to the heavy atom effect, resulting in a significant increase in triplet emission. The X-ray structure of the fibers indicates that both aurophilic, π - π interactions and hydrogen bonding contribute to their formation in aqueous solvents.

Received 29th September 2014,
Accepted 5th December 2014

DOI: 10.1039/c4ob02077d

www.rsc.org/obc

Introduction

Supramolecular gels¹ have become a topic of increasing interest in recent years. Compared to traditional polymers, in which the individual monomers are linked irreversibly by covalent bonds, supramolecular linkages are reversible, which can result in new dynamic structures with interesting and new properties completely different from those observed on the individual molecules.²

In particular, self-assembled supramolecular hydrogels have attracted much attention during the past few decades because of promising high-technology applications in very diverse fields such as biomedicine, nanoelectronics and catalysis.^{3–9} These colloidal materials can be formed either by

low- or high-molecular-weight hydrogelators by means of non-covalent interactions (*e.g.*, π - π stacking, van der Waals forces, hydrogen bonds, and hydrophobic interactions).^{10,11} Among these materials, metallo-gels^{12–14} represent an important subclass where at least one metallic element has been incorporated into the gel network, but they have been a subject of study only in the last few years. The presence of metal ions should be highlighted since they may also modify the physical properties of the gels.¹⁵

In contrast to the development of coordination polymeric gelators, the use of discrete small-molecule metal complexes such as metallo-gelators has been relatively scarce and it was only recently that a number of reports on the gelation properties of metal complexes, such as Au(i), Au(III) and Pt(II), have appeared.^{12,15–18} Interestingly, this process becomes unpredictable in some cases.

The formation of the supramolecular gels and metallo-gels depends on the concentration of the monomer and on the strength of the non-covalent interactions responsible for holding the monomers together.^{2,19} In general, hydrogen bonds are among the most often used non-covalent interactions in the formation of supramolecular gels and polymers. Nevertheless, in the particular case of organometallic complexes like gold(i) derivatives, the possible establishment of Au(i)⋯Au(i) (aurophilic) bonds is also observed to be an important driving force responsible for the aggregation.

In fact, the formation of aurophilic interactions in the crystal structures of gold compounds, as well as in solution, has fascinated chemists for decades. The aggregation of gold compounds in this fashion arises from relativistic effects that are at a maximum for gold compared to other heavy elements.^{20–22} One reason for this interest is that gold compounds exhibit interesting luminescence properties, in both

^aREQUIMTE, Dep. de Química, Universidade Nova de Lisboa, Monte de Caparica, Portugal. E-mail: lima@fct.unl.pt; Fax: +351 212948550; Tel: +351 212948300 (Ext. 10923)

^bEcole polytechnique de Bruxelles, Université libre de Bruxelles, 50 Av F.D Roosevelt, 1050 Bruxelles, Belgium

^cDepartament de Química Inorgànica, Universitat de Barcelona, c/Martí i Franquès, 1-11, 08028 Barcelona, Spain. E-mail: laura.rodriguez@qi.ub.es; Fax: +34 934907725; Tel: +34 934039130

^dDepartment of Chemistry, Nanoscience Center, P.O. Box 35, 40014 Jyväskylä, Finland

† Electronic supplementary information (ESI) available: Hydrogel formation with complex **1** checked by the inverted tube method (Fig. S1); absorption spectra of **1** between 5 and 100 μ M in water. Inset: Plot of the absorption at 400 nm vs. concentration (Fig. S2); DLS auto-correlation data (left) and the resulting size distributions (right) of aqueous solutions of **1** at different concentrations in water (Fig. S3); schematic representation of (a) the star-shaped 1-D polymeric structure of **1** and (b) the observed π - π interactions (Fig. S4); 3-D crystal packing of **1** (Fig. S5). X-ray crystallographic data for **1** (Table S1). CCDC 1026387. For ESI and crystallographic data in CIF or other electronic format see DOI: 10.1039/c4ob02077d






Chart 1

the solid-state and in solution that could be modulated by the presence of these Au...Au interactions.^{22–27}

Aurophilic interactions can also lead to fascinating supra-molecular architectures^{21,22,28–32} and are known to present stabilization energies to their crystal structures in the range of 7–12 kcal mol⁻¹.^{33,34} This energy range should be stressed since it is comparable to that normally associated with conventional hydrogen bonding.

Among gold(i) supramolecular structures, gold(i) alkynyl derivatives are specially denoted due to the preference of gold(i) for a linear coordination geometry, together with the linearity of an acetylide unit and its π -unsaturated nature. These structural characteristics have made the alkynyl gold(i) complexes attractive building blocks for organometallic oligomeric and polymeric materials which may possess unique properties in different fields.²² Related to all of this, we have recently described the formation of luminescent hydrogelators with two gold(i) alkynyl derivatives with a very simple chemical structure ([Au(4-pyridylethynyl)(phosph)] {phosph = PTA (1,3,5-triaza-7-phosphaadamantane);¹⁷ DAPTA = 3,7-diacetyl-1,3,7-triaza-5-phosphabicyclo[3.3.1]nonane}).¹⁸ In this work, the synthesis of a new hydrogelator that contains a propargyloxycoumarin group instead of pyridylethynyl is described (Chart 1). In order to try to understand better the gelation process with these discrete molecules, we report on the analysis of the aggregation process by different spectroscopical and microscopic techniques and the results are compared with the X-ray crystal diffraction data retrieved for this complex.

Materials and methods

Reagents

The synthesis of [Au{7-(prop-2-inoxy)-1-benzopyran-2-one}(DAPTA)] (**1**) is described in previous reports.³⁵

Physical measurements

Absorption spectra were recorded on a Varian Cary 5000 UV-Vis spectrophotometer in a 3 mL (10 × 10 mm wide) quartz cell. Fluorescence measurements were performed on a Horiba-Jobin-Yvon SPEX Fluorolog 3.22 in a 3 mL (10 × 10 mm wide) quartz cell. Emission quantum yields were measured employing 7-methoxy-4-methylcoumarin as a reference ($\phi_F = 0.124$, 25 °C, methanol).

Dynamic light scattering experiments were performed on a Horiba Scientific Nanoparticle Analyzer SZ-100 in 3 mL (10 × 10 mm wide) quartz cells. The samples were filtered

before analysis using a 0.46 μ m polystyrene membrane disc filter.

Fluorescence microscopy was recorded on an Axioplan 2ie Zeiss imaging microscope equipped with a NikonDXM1200F digital camera. Excitation light for fluorescence imaging was selected using filters in the range of 300–400 and 450–490 nm respectively.

Preparation of the gel

Solid **1** was dissolved in water in different concentration ranges (0.01–0.02% by weight). The sample was maintained under sonication for *ca.* 20 minutes and left at room temperature at least for 4 hours.

X-ray crystal structure determination

Data for **1** were collected at 123 K on an Agilent SuperNova diffractometer with an Atlas detector using mirror-monochromated Cu-K α radiation ($\lambda = 1.54184$ Å). The CRYCALISPRO³⁶ program was used for the data collection and processing of complex **1**. The intensities were corrected for absorption using the multi-scan absorption correction method.³⁶ The structure of **1** was solved by direct methods with the SHELXS solution program.³⁷ The structure was refined by full-matrix least-squares calculations based on F^2 using SHELXL-2013³⁸ integrated into the WINGX³⁹ program package. Hydrogen atoms were included in calculated positions as riding atoms, with SHELXL-2013³⁸ defaults. X-ray crystallographic data for **1** are given in Table S1 (see ESI†).

CCDC-1026387 contains the supplementary crystallographic data for this paper.

Results and discussion

The synthesis and characterization of **1** was recently reported by us, and its photophysical properties were reported in methanol.³⁵ Herein we have described the observed formation of hydrogels (see Fig. S1†) and compared it with the previous data obtained with pyridylethynyl derivatives.^{17,18} In the case of **1**, apart from the Au(i) atoms, the structure of the complex possesses a phosphine and a coumarin unit at both ends of the molecule. While the former moiety provides an increase in hydrophilicity, the latter moiety presents strong absorption and emission in the UV region of the spectrum. Thus, we measured the optical properties of the coumarin as a means to study the formation of aggregates that ultimately lead to new luminescent soft materials such as hydrogels.

Absorption/fluorescence spectroscopy

The effect of concentration on the absorption and emission of complex **1** was primarily studied in methanol (Fig. 1) where aggregates are not expected to be formed.

As can be seen in Fig. 1a and its inset, no changes were observed both in the shape and in the values for absorbance maximum of the spectra up to 24 μ M, with a calculated molar absorptivity (ϵ) of 15 259 M⁻¹ cm⁻¹ at 320 nm. Although the



was obtained in the following sections). The hydrophobic parts are then surrounded by a shell of water molecules organized in such a way to shade it from the solvent. For aggregation to happen, this shell must be displaced momentarily, which creates a rise in entropy and thus changes the initial formula towards this modified thermodynamic equilibrium:

$$\Delta G_{\text{aggregation}} = \Delta H - T (\Delta S_{\text{aggregation}} + \Delta S_{\text{solvent}})$$

If the solvent contribution (positive) is bigger than the aggregation contribution (negative) to entropy, then increasing the temperature will lead to a faster aggregation by increasing the driving force.

Dynamic light scattering

In order to obtain more insight into the size of the aggregates, screening tests at different concentrations were performed using dynamic light scattering. The overall results show that aggregates can be found down to the micromolar concentration range in water (Fig. S3[†]). The size distribution of the aggregates allows the identification of a single population with a well-defined mean diameter (ranging from 70 to 140 nm, depending on the concentration). However, in all cases, a high polydispersity index is recorded, which is a strong indication of the presence of larger aggregates. Nevertheless, at the lowest concentrations (50 μM or lower) and after filtration with a 0.46 μm

filter, we can obtain good mono-exponential fittings of the auto-correlation curves, with low polydispersity index (PDI) values, suggesting the presence of spherical nanostructures.

The influence of temperature on the formation/destruction of aggregates was studied by performing temperature ramps at 38 μM in water as a means to complement the results of absorption experiments (Fig. 3).

As one can observe, a mono-exponential fitting of the DLS correlation function is achieved, with a monodisperse population of aggregates of about 90 nm, typical for micellar-like aggregates that could be created from hydrophobic interactions, as mentioned in the previous section. Raising the temperature to about 40 $^{\circ}\text{C}$ leads to a bi-exponential function due to the appearance of a long decay signal, a clear indication of the formation of bigger structures.

This fact can be explained based on the molecular structure of complex 1 if we consider that linear Au(I)-alkynyl complexes are known to form intermolecular auophilic bonds and produce gels.^{17,18} If this is the case, the formation of linear aggregates/fibers may explain the high PDI values, since the DLS models for the determination of hydrodynamics assume a spherical shape for the analyzed samples.

Upon further increasing the temperature, we can no longer observe the initial population of smaller aggregates (in the nm range), although one cannot exclude their presence since the




Fig. 3 Evolution of DLS scans upon increasing the temperature from 25 $^{\circ}\text{C}$ to 85 $^{\circ}\text{C}$ and cooling back to 25 $^{\circ}\text{C}$.



light scattering phenomenon is intrinsically linked to the size in a non-linear way, *i.e.*, bigger particles/structures scatter more light than smaller ones. Finally, by decreasing the temperature, the initial signal for small particles is partially recovered. These results are consistent with the aforementioned hypothesis for temperature-favoured aggregation.

Fluorescence microscopy

Images of the aggregates at different concentrations of **1** were obtained upon drying the solutions and analyzing the samples under the microscope. Fig. 4 shows bright field images of dried solutions of **1**.

Detailed observation of pictures from Fig. 4 reveals the evolution of aggregation upon increasing the concentrations of **1**. Indeed, at low concentration, only small spherical aggregates are observed which is consistent with the results from DLS, even though the size scales are quite different, since the solutions for DLS were previously filtered with a 460 nm filter and particles smaller than 460 nm are not seen under the microscope. A ten-fold more concentrated sample of **1** (image b) already reveals the presence of agglomerates of particles in a linear fashion, while at 1 mM a highly reticulated network of fibers can be clearly identified in the picture which accounts for the observed high polydispersity of the samples in DLS measurements.

Pictures were also taken under UV irradiation using two different bandpass filters. A detailed analysis of these pictures shows that, while green emission is present throughout the whole reticulated network of fibers (Fig. 5b), blue emission can only be detected at the extremities (Fig. 5a).

A possible explanation for this behavior is the heavy-atom effect induced by the gold atoms, *i.e.*, in the bulk of the aggregate the fibers are highly crosslinked. This causes the gold atoms and coumarins to be closer to each other than in the case of isolated linear fibers that are linked by aurophilic interactions.

Indeed, since it has been shown that this kind of compound shows a linear architecture L1–Au–L2 and that it aggregates by Au...Au interactions,²² the coumarin and gold atoms can never be closer than the intramolecular distance, which in the case of **1** consists of four chemical bonds due to the methoxy-alkynyl spacer. However, once this “dense” aggregation phase is reached, gold and coumarin units can be, on average, at a shorter distance than the intramolecular distance, thus allowing for a more efficient intersystem crossing to take place and resulting in coumarin phosphorescence emission. Previous data obtained with ([Au(4-pyridylethynyl)(phosph)]{phosph = PTA (1,3,5-triaza-7-phosphaadamantane)¹⁷ and DAPTA = 3,7-diacetyl-1,3,7-triaza-5-phosphabicyclo[3.3.1]nonane})¹⁸ complexes have been a good reference for the analysis of these gels. It must be highlighted that the comparison




Fig. 4 Bright field microscopic images of dried solutions of **1** at (a) 3.8 μM , (b) 38 μM and (c) 1000 μM .




Fig. 5 Fluorescence microscopy images of dried solutions of **1** at 1 mM using a bandpass filter of (a) 300–400 nm and (b) 450–490 nm.






Fig. 7 1-D polymeric structure of complex 1, hydrogen atoms and solvent molecules are omitted for clarity.




Fig. 8 Space fill representation of the 3-D crystal packing of complex 1 viewed down the *a*-axis displaying channels for a MOF-like motif. Hydrogen atoms and dichloromethane molecules are omitted for clarity.

favoured in water, being the driving force of the formation of gels. On the other hand, theoretical calculations carried out with similar aurophilic hydrogelators indicate a clear involvement of Au(I)⋯Au(I) interactions in the formation of these larger aggregates.³⁴

Conclusions

The design and characterization of an alkynyl gold(I) complex with a coumarin fluorophore as a new hydrogelator for the synthesis of luminescent soft materials has been performed.

Absorption and DLS studies in aqueous media suggested the formation of aggregates at low micromolar concentrations. The same techniques indicated that aggregation was favoured with increasing the temperature.

Fluorescence microscopy revealed the presence of luminescent fiber-like structures at millimolar concentrations. While strong blue emission (corresponding to the fluorescence of the coumarin) was only observed in isolated fibers, green emission (derived from coumarin phosphorescence) was observed throughout the bulk of the reticulated network of fibers. This

effect is accounted for by the spatial proximity of coumarins and gold(I) atoms, which leads to an increase in intersystem crossing and, consequently, quenching of fluorescence and enhancement of triplet emission (T_1-S_0 radiative pathway).

X-ray crystallography confirmed the existence of π - π interactions, hydrogen bondings as well as aurophilic interactions, resulting in highly ordered linear assemblies.

Acknowledgements

The support and sponsorship provided by COST Action CM1005 is acknowledged. Authors are also grateful to the Ministerio de Ciencia e Innovación of Spain (project CTQ2012-31335), Fundação para a Ciência e Tecnologia of Portugal (PTDC/QUI-QUI/112597/2009; PEST-C/EQB/LA0006/2011) and Academy of Finland (K.R., grant no. 263256 and 265328). A.M. thanks FCT for a post-doctoral grant (SFRH/BPD/69210/2010).

References

- 1 S. Dong, B. Zheng, D. Xu, X. Yan, M. Zhang and F. Huang, *Adv. Mater.*, 2012, **24**, 3191.
- 2 M. Fleischer and C. Schmuck, *Chem. Commun.*, 2014, **50**, 10464.
- 3 S. Banerjee, R. K. Das and U. Maitra, *J. Mater. Chem.*, 2009, **19**, 6649.
- 4 A. R. Hirst, B. Escuder, J. F. Miravet and D. K. Smith, *Angew. Chem., Int. Ed.*, 2008, **47**, 8002.
- 5 N. M. Sangeetha and U. Maitra, *Chem. Soc. Rev.*, 2005, **34**, 821.
- 6 T. Vermonden, R. Censi and W. E. Hennink, *Chem. Rev.*, 2012, **112**, 2853.
- 7 E. A. Appel, J. del Barrio, X. J. Loh and O. A. Scherman, *Chem. Soc. Rev.*, 2012, **41**, 6195.
- 8 D. D. Díaz, D. Kühbeck and R. J. Koopmans, *Chem. Soc. Rev.*, 2011, **40**, 427.
- 9 S. Saha, J. Bachl, T. Kundu, D. Díaz Díaz and R. Banerjee, *Chem. Commun.*, 2014, **50**, 7032.



