# Chemical Science



## **EDGE ARTICLE**

View Article Online
View Journal | View Issue



Cite this: Chem. Sci., 2016, 7, 1786

Received 7th November 2015 Accepted 14th December 2015

DOI: 10.1039/c5sc04237b

www.rsc.org/chemicalscience

## Aminofluorination: transition-metal-free N-F bond insertion into diazocarbonyl compounds†

Gui Chen,<sup>a</sup> Jinshuai Song,<sup>b</sup> Yinghua Yu,<sup>a</sup> Xuesong Luo,<sup>a</sup> Chunsen Li\*<sup>b</sup> and Xueliang Huang\*<sup>a</sup>


Gem-aminofluorination of diazocarbonyl compounds has been achieved for the first time. This reaction proceeds under mild conditions and does not require any transition-metal promoter or catalyst. Treatment of diazoesters with *N*-fluorobenzenesulfonimide (NFSI), which serves as both a fluorine and nitrogen source, results in the facile construction of C–N and C–F bonds, providing aminofluorination products in moderate to excellent yields. Kinetic studies and DFT calculations have provided valuable insight into the potential mechanism for this novel N–F bond insertion.

#### Introduction

Diazoacetate derivatives play an important role in the synthetic community. Over the past century, significant advances have been made towards the generation of carbenoid intermediates triggered by transition-metals. The resulting reactive species can undergo valuable transformations, such as three-membered ring (cyclopropane, cyclopropene) formation, X–H bond insertion (X = C, N, O, S, etc.) and ylide generation (Scheme 1a–c). van Vranken, Barluenga, Wang, and others have developed palladium- or copper-catalysed multiple component reactions of diazo compounds, which allow the installation of two separated functional moieties on the carbenic carbon via


a single operation (Scheme 1d).<sup>2</sup> Despite these important advances, the direct introduction of two functional groups to the same carbon center, namely gem-difunctionalization of donor/acceptor (D/A) carbenes, is still far from well developed.<sup>3,4</sup> It is of note that these processes at least involve one C–C bond formation.<sup>5</sup> Thus the studies on the gem-difunctionalization of D/A carbenes, which involves two distinct carbon heteroatom bond formations, would greatly enhance the synthetic applications of diazo compounds.

Single substitution of hydrogen with fluorine may alter the chemical and physical properties of a potential drug candidate by blocking undesired metabolism at a specific site. In contrast to the relatively large number of reports on the catalytic insertion of N–H bonds to α-diazocarbonyl compounds, a simple yet appealing concept for the transition-metal catalysed N–F bond insertion has not been realized thus far. *N*-Fluorobenzenesulfonimide (NFSI) is inexpensive and shelf stable, and is often employed as a mild electrophilic fluorinating or aminating reagent. Recently, Liu and Zhang demonstrated that NFSI could serve as both an amino and fluorine source for the transition-metal-catalysed aminofluorination of alkenes (Scheme 2a). Inspired by these seminal works, we envisioned that NFSI might be an ideal candidate for the transition-metal



Scheme 1 Representative reaction modes of a transition-metal carbenoid.

<sup>†</sup> Electronic supplementary information (ESI) available: Experimental procedures, and data for new compounds. CCDC 1417241. For ESI and crystallographic data in CIF or other electronic format see DOI: 10.1039/c5sc04237b



Scheme 2 Aminofluorination catalysed by transition-metals.

<sup>&</sup>quot;Key Laboratory of Coal to Ethylene Glycol and Its Related Technology, Fujian Institute of Research on the Structure of Matter, Chinese Academy of Sciences, Yangqiao west road 155", Fuzhou, Fujian 350002, China. E-mail: huangxl@fjirsm.ac.cn; Web: http://www.fjirsm.cas.cn/yjxt/yyhxyjzs/hxlktz/

bState Key Laboratory of Structural Chemistry, Fujian Institute of Research on the Structure of Matter, Chinese Academy of Sciences, Yangqiao west road 155#, Fuzhou, Fujian 350002, China. E-mail: chunsen.li@fjirsm.ac.cn

catalysed N–F bond insertion into D/A carbenes (Scheme 2b). Although procedures for the amination<sup>7,12</sup> or fluorination<sup>13,14</sup> of diazo compounds are known, to our knowledge, direct aminofluorination of diazo compounds remains unexplored. Herein, we present our primary results on gem-aminofluorination of diazocarbonyl compounds under mild conditions. Kinetic studies and DFT calculations shed light on the mechanism of the current N–F bond insertion.

#### Results and discussion

#### Optimization studies and substrate scope

We initiated our studies using the reaction of ethyl diazophenyl acetate 1a (ref. 15) with NFSI using CuBr as a precatalyst and bathocuproine (2,9-dimethyl-4,7-diphenyl-1,10-phenanthroline; BC) as a ligand. 11 Gratifyingly, in the presence of 5 mol% CuBr and 6 mol% BC, ethyl diazophenyl acetate 1a was completely consumed after stirring in a reaction medium of 1,2dichloroethane (DCE) at 60 °C for 21 h, and the desired product 2a was obtained in a 79% NMR yield (Table 1, entry 1). To our surprise, the reaction proceeded equally well in the absence of both CuBr and BC, giving 2a in 89% yield (Table 1, entry 2). The reaction could complete in a comparably short time when the reaction was carried out at an elevated temperature (Table 1, entries 2-4 vs. entry 5). A brief examination of the solvent effects revealed that DCE was still the best choice (Table 1, entries 6-14). Notably, the reaction can also be performed in water, giving 2a in a moderate yield, which indicates an environmentally

Table 1 Optimization of the reaction conditions

Ph CO <sub>2</sub> Et	+	PhO <sub>2</sub> S <sub>N</sub> SO <sub>2</sub> Ph	conditions	(PhO <sub>2</sub> S) <sub>2</sub> N F Ph CO <sub>2</sub> Et
<b>1a</b> , 0.15 mmol		NFSI, 0.18 mmol		2a

Entry <sup>a</sup>	Solvent	Temp (°C)	Time (h)	Yield <sup>b</sup> (%)
1 <sup>c</sup>	DCE	60	21	79
2	DCE	60	21	89
3	DCE	80	7	77
4	DCE	100	4	73
5	DCE	RT	24	17
6	$CH_2Cl_2$	60	24	81
7	$CHCl_3$	60	24	70
8	Toluene	60	24	58
9	MeCN	60	24	69
10	THF	60	24	47
11	Dioxane	60	24	22
12	DMF	60	24	0
13	MeOH	60	24	0
14	$H_2O$	60	12	62
$15^d$	DCE	60	24	82
$16^e$	DCE	60	48	97 <sup>f</sup>

 $^a$  All reactions were carried out under an atmosphere of argon in 0.15 mmol scale, [1a] = 0.10 M, [NFSI] = 0.12 M.  $^b$  Yields were determined by  $^{19}$ F NMR, using 1-bromo-4-fluorobenzene as the internal standard.  $^c$  In the presence of 5 mol% CuCl and 6 mol% BC.  $^d$  Under air.  $^e$  The reaction was carried out in 0.3 mmol scale, [1a] = 0.15 M, [NFSI] = 0.10 M.  $^f$  Isolated yield.

benign perspective (Table 1, entry 15). Due to the similar polarity between 2a and NFSI, a slight excess of 1a (1.5 eq.) was necessary to make sure that NFSI reached full conversion. In this case, 2a was isolated in a nearly quantitative yield (Table 1, entry 16).

With the optimized reaction conditions in hand (Table 1, entry 16), the generality and limitations of this metal-free N–F bond insertion were investigated. The results are summarized in Table 2. Most of the  $\alpha$ -diazoacetates reacted with NFSI smoothly to give the corresponding products in moderate to excellent yields. When changing the ester group (R²) from methyl to isobutyl, or benzyl, no significant decrease of the product yields was observed (Table 2, **2a–d**). When the diazo

Table 2 Geminal aminofluorination of various diazocompounds<sup>a</sup>

<sup>a</sup> All reactions were carried out in 0.30 mmol scale under argon at 60  $^{\circ}$ C, [1a] = 0.15 M, [NFSI] = 0.10 M, isolated yield. <sup>b</sup> Reaction was carried out at 80  $^{\circ}$ C.

Chemical Science Edge Article

compound contained a chiral auxiliary, an ester derived from (+)-menthol, a pair of diastereoisomers 2e were obtained in a ratio of 1:1. Interestingly, diazo compounds derived from cyclic esters and amides were proven to be viable substrates. N-F bond insertion of 1f and 1g gave the corresponding products 2f and 2g in 97% and 83% yields, respectively. The structure of 2f was confirmed by X-ray crystallographic analysis.16 The effects of substituents on the phenyl ring were also examined. Both electron-withdrawing and electron-donating groups on the aromatic ring of 1 were tolerated under the reaction conditions. The reaction of diazo acetates bearing mild electron-withdrawing or electron-donating groups (fluoro, chloro, bromo and methyl) at the para position of the phenyl ring gave the corresponding products in excellent yields (Table 2, 2h, 2i, 2m and 2o). Incorporation of one chloro group to the meta position had no obvious impact on the yields of the products (Table 2, 2i and 2k). Interestingly, the yield of 2l bearing two meta chloro substituents was decreased to 52%. While the diazo acetate bearing an ortho substituted group was not compatible to the current conditions (Table 2 and 2p), probably due to the steric hindrance effect. Similarly, strong electron-donating or -withdrawing substituents on the aromatic ring were amenable for the current aminofluorination, albeit giving moderate yields (Table 2, 2n, 2r-t). Of note, vinyl and alkyl moieties remained intact (Table 2, 2v and 2w), indicating that the involvement of a free carbene intermediate is less likely (vide infra). Notably, the reaction could be scaled up to gram scale without sacrificing the yield of 2a (5 mmol scale, 2.39 g 2a was obtained with a 97% yield). It is worthwhile to mention that the current conditions are not applicable to alkyl or heteroaryl acetate derived diazo compounds.17

#### Mechanistic studies

Having uncovered an efficient method for N–F bond insertion, we sought to gain more insight into the reaction mechanism. Thus a series of additional experiments were subsequently carried out. Performing the reaction under irradiation by UV light, 2a was obtained in a much lower yield (Scheme 3a). Furthermore, evolution of  $N_2$  was significantly slow in the absence of NFSI under otherwise identical conditions (Fig. S1†). These experiments also suggest that a pathway via a free carbene intermediate is unfavourable. Addition of the radical

Scheme 3 Preliminary mechanistic studies.

scavenger 2,6-di-tert-butyl-4-methylphenol (BHT) had no obvious influence on the reaction efficiency (Scheme 3b), which indicates that a mechanism that involves free radical species is also less likely. Additionally, considering the diazo carbon atom is mildly nucleophilic and might be trapped by the electrophilic fluorine atom, a subsequent displacement of dinitrogen by the imide moiety of NFSI would give 2a. With this consideration in mind, the following control experiment was carried out immediately. The combination of selectfluor, a reactive electrophilic fluorinating reagent, with tetrapropyl ammonium benzenesulfonimide did afford 2a in 8% yield (Scheme 3c).

Kinetic studies for the reaction of  ${\bf 1a}$  with NFSI were further performed to get a deeper understanding of the reaction mechanism. The reaction was found to be first-order in both  ${\bf 1a}$  and NFSI (Fig. S2–S5†). The activation parameters  $\Delta H^{\ddagger}=17.1$  kcal mol $^{-1}$  and  $\Delta S^{\ddagger}=-13.0$  cal mol $^{-1}$  K $^{-1}$  were obtained from Eyring plots by varying the temperature from 313 to 353 K (Fig. 1A, top). The negative  $\Delta S^{\ddagger}$  value may suggest the generation of a bimolecular transition state involving NFSI and  ${\bf 1a}$ . Similar kinetic behaviours were observed for the reactions of various para-substituted diazo phenyl acetates  ${\bf 1}$  with NFSI. A fairly linear Hammett correlation between  $\log(k_{\rm X}/k_{\rm H})$  and  $\sigma^+$  was obtained with a reaction constant of  $\rho=-0.81$  (Fig. 1B, bottom). The small negative  $\rho$  value suggests that the transition





Fig. 1 (A) (top) Plot of In(initial rate/T) vs. 1/T for the reaction between 1a and NFSI in DCE, [1a] = 0.15 M, [NFSI] = 0.1 M, slope =  $-8.61 \times 10^3$ , y-intercept =  $1.27 \times 10$ ,  $r^2 = 0.996$ . (B) (bottom) Hammett plot of  $\log(k_{\rm X}/k_{\rm H})$  vs.  $\sigma^+$  for the reaction of NFSI with para substituted diazo phenyl acetates 1 in DCE at 50 °C, [1] = 0.15 M, [NFSI] = 0.1 M, slope = -0.81, y-intercept =  $0.4 \times 10^{-2}$ ,  $r^2 = 0.995$ .

Edge Article Chemical Science



Scheme 4 Mechanistic rationale supported by DFT calculations.

state is weakly polarized with a positive charge at the reaction center. 18,19

DFT calculations were carried out to provide more details on the potential mechanism (Scheme 4).20 The reaction of 1a with NFSI encounters an activation enthalpy of 16.2 kcal mol<sup>-1</sup> with a slight loss of entropy ( $\Delta S^{\ddagger} = -9.0$  cal mol<sup>-1</sup> K<sup>-1</sup>), which matches well with the experimental values (vide supra). According to the downhill energy profile, the following steps are rather facile. The weakly polarized TS further collapses to an ion pair Int1 with an energy change of -41.8 kcal mol<sup>-1</sup>. Releasing N<sub>2</sub> from Int1 has been proven to be an exothermic process. Finally, an S<sub>N</sub>1 like reaction takes place to give 2a, with a total enthalpy change of -84.8 kcal mol<sup>-1</sup>. It is of note, that the formation of a procarbonium ion through the reaction of diazo carbon with Cl<sup>+</sup>, Br<sup>+</sup> and I<sup>+</sup> has been reported recently.<sup>21</sup> However, in our case, the replacement of NFSI by NXS (X = Cl,Br, I) results in no formation of the aminohalogenated product. It is well known that diazo compounds are not stable in the presence of a strong acid. Indeed, addition of strong Brønsted acids (CCl<sub>3</sub>CO<sub>2</sub>H or CF<sub>3</sub>CO<sub>2</sub>H) led to the rapid decomposition of 1a. However, the addition of acetic acid had no influence on the current N-F bond insertion.22,23

## Conclusions

In conclusion, we have developed an unprecedented N–F bond insertion into diazocarbonyl compounds. The current method represents a facile approach to construct C–N and C–F bonds on the same carbon without any transition-metal as a catalyst or promoter. Mechanistic studies, including kinetic experiments and DFT calculations, revealed that a reaction sequence of electrophilic activation of 1 by NFSI, followed by an  $S_N1$  like displacement of dinitrogen by benzenesulfonimide is preferred. Further study on the asymmetric variant as well as the difunctionalization of the metal carbene is ongoing in our laboratory.

## Acknowledgements

This work is supported by the National Natural Science Foundation of China (Grant No. 21402197, 21502190) and Fujian Institute of Research on the Structure of Matter, Chinese Academy of Sciences.

### Notes and references

- 1 For selected reviews, see: (a) M. P. Doyle, Chem. Rev., 1986, 86, 919; (b) T. Ye and M. A. McKervey, Chem. Rev., 1994, 94, 1091; (c) A. Padwa and D. J. Austin, Angew. Chem., Int. Ed., 1994, 33, 1797; (d) A. Padwa and M. D. Weingarten, Chem. Rev., 1996, 96, 223; (e) M. P. Doyle and D. C. Forbes, Chem. Rev., 1998, 98, 911; (f) M. P. Dovle, M. A. McKervey and T. Ye, Modern Catalytic Methods for Organic Synthesis with Diazo Compounds: From Cyclopropanes to Ylides, John Wiley & Sons, Inc., New York, NY, 1998; (g) H. M. L. Davies and R. E. J. Beckwith, Chem. Rev., 2003, 103, 2861; (h) Z. Zhang and J. Wang, Tetrahedron, 2008, 64, 6577; (i) Y. Zhang and J. Wang, Chem. Commun., 2009, 45, 5350; (j) M. P. Doyle, R. Duffy, M. Ratnikov and L. Zhou, Chem. Rev., 2010, 110, 704; (k) H. M. L. Davies and Y. Lian, Acc. Chem. Res., 2012, 45, 923; (l) S.-F. Zhu and Q.-L. Zhou, Acc. Chem. Res., 2012, 45, 1365; (m) X. Guo and W. Hu, Acc. Chem. Res., 2013, 46, 2427; (n) A. Ford, H. Miel, A. Ring, C. N. Slattery, A. R. Maguire and M. A. McKervey, Chem. Rev., 2015, 115, 9981.
- (a) K. L. Greenman, D. S. Carter and D. L. van Vranken, *Tetrahedron*, 2001, 57, 5219; For selected reviews, see: (b) J. Barluenga and C. Valdés, *Angew. Chem., Int. Ed.*, 2011, 50, 7486; (c) Q. Xiao, Y. Zhang and J. Wang, *Acc. Chem. Res.*, 2013, 46, 236; (d) Y. Zhang and J. Wang, *Eur. J. Org. Chem.*, 2011, 1015; (e) Y. Xia, Y. Zhang and J. Wang, *ACS Catal.*, 2013, 3, 2586.
- For recent selected examples, see: (a) W.-W. Chan, S.-F. Lo,
   Z. Zhao and W.-Y. Yu, J. Am. Chem. Soc., 2012, 134, 13565;
   (b) T. K. Hyster, K. E. Ruhl and T. Rovis, J. Am. Chem. Soc., 2013, 135, 5364;
   (c) Y. Xia, Z. Liu, Z. Liu, R. Ge, F. Ye,
   M. Hossain, Y. Zhang and J. Wang, J. Am. Chem. Soc., 2014, 136, 3013;
   (d) A. Yada, S. Fujita and M. Murakami, J. Am. Chem. Soc., 2014, 136, 7217;
   (e) F.-N. Ng, Y.-F. Lau, Z. Zhou and W.-Y. Yu, Org. Lett., 2015, 17, 1676.
- 4 For selected examples on Lewis acid promoted insertion to carbonyl compounds, see: (a) T. Hashimoto, Y. Naganawa and K. Maruoka, J. Am. Chem. Soc., 2009, 131, 6614; (b) D. C. Moebius and J. S. Kingsbury, J. Am. Chem. Soc., 2009, 131, 878; (c) W. Li, X. Liu, F. Tan, X. Hao, J. Zheng, L. Lin and X. Feng, Angew. Chem., Int. Ed., 2013, 52, 10883; W. Li, F. Tan, X. Hao, G. Wang, Y. Tang, X. Liu, L. Lin and X. Feng, Angew. Chem., Int. Ed., 2015, 54, 1608.
- 5 (a) R. Kudirka and D. L. van Vranken, J. Org. Chem., 2008, 73, 3585; (b) R. Kudirka, S. K. J. Devine, C. S. Adams and D. L. van Vranken, Angew. Chem., Int. Ed., 2009, 48, 3677; (c) P.-X. Zhou, Z.-Z. Zhou, Z.-S. Chen, Y.-Y. Ye, L.-B. Zhao, Y.-F. Yang, X.-F. Xia, J.-Y. Luo and Y.-M. Liang, Chem. Commun., 2013, 49, 561. For a recent example on Pd(0)-catalysed carbene insertion of Sn-Sn and Si-Si bonds, see: (d) Z. Liu, H. Tan, T. Fu, Y. Xia, D. Qiu, Y. Zhang and J. Wang, J. Am. Chem. Soc., 2015, 137, 12800.
- 6 W. K. Hagmann, J. Med. Chem., 2008, 51, 4359.
- 7 For recent selected examples, see: (a) S. Bachmann, D. Fielenbach and K. A. Jørgensen, *Org. Biomol. Chem.*,

2004, 2, 3044; (b) B. Liu, S.-F. Zhu, W. Zhang, C. Chen and Q.-L. Zhou, J. Am. Chem. Soc., 2007, 129, 5834; (c) E. C. Lee and G. C. Fu, J. Am. Chem. Soc., 2007, 129, 12066; (d) Z. Hou, J. Wang, P. He, J. Wang, B. Qin, X. Liu, L. Lin and X. Feng, Angew. Chem., Int. Ed., 2010, 49, 4763; (e) B. Xu, S.-F. Zhu, X.-L. Xie, J.-J. Shen and Q.-L. Zhou, Angew. Chem., Int. Ed., 2011, 50, 11483; (f) S.-F. Zhu, B. Xu, G.-P. Wang and Q.-L. Zhou, J. Am. Chem. Soc., 2012, 134, 436; (g) Y. Zhu, X. Liu, S. Dong, Y. Zhou, W. Li, L. Lin and X. Feng, Angew. Chem., Int. Ed., 2014, 53, 1636.

- 8 (a) J.-A. Ma and D. Cahard, Chem. Rev., 2004, 104, 6119; (b)
  J.-A. Ma and D. Cahard, Chem. Rev., 2008, 108, PR1; (c)
  T. Liang, C. Neumann and T. Ritter, Angew. Chem., Int. Ed., 2013, 52, 8214; (d) X. Yang, T. Wu, R. J. Phipps and F. D. Toste, Chem. Rev., 2015, 115, 826; (e)
  P. A. Champagne, J. Desroches, J. D. Hamel, M. Vandamme and J. F. Paquin, Chem. Rev., 2015, 115, 9073; (f) P. Chen and G. Liu, Eur. J. Org. Chem., 2015, 4295.
- 9 Y. Li and Q. Zhang, Synthesis, 2015, 159.

**Chemical Science** 

- 10 S. Qiu, T. Xu, J. Zhou, Y. Guo and G. Liu, J. Am. Chem. Soc., 2010, 132, 2856.
- 11 H. Zhang, Y. Song, J. Zhao, J. Zhang and Q. Zhang, Angew. Chem., Int. Ed., 2014, 53, 11079.
- 12 For selected examples on metal-free N-H insertion of diazoesters, see: (a) S. S. So and A. E. Mattson, J. Am. Chem. Soc., 2012, 134, 8798; (b) S. R. Hansen, J. E. Spangler, J. H. Hansen and H. M. L. Davies, Org. Lett., 2012, 14, 4626; (c) D. M. Nickerson, V. V. Angeles, T. J. Auvil, S. S. So and A. E. Mattson, Chem. Commun., 2013, 49, 4289.
- 13 (a) G. A. Olah and J. Welch, Synthesis, 1974, 896; (b) R. Pasceri, H. E. Bartrum, C. J. Hayes and C. J. Moody,

- Chem. Commun., 2012, 48, 12077; (c) A. K. Yadav, V. P. Srivastava and L. D. S. Yadav, Chem. Commun., 2013, 49, 2154; (d) J. Tao, R. Tran and G. K. Murphy, J. Am. Chem. Soc., 2013, 135, 16312.
- 14 It should be mentioned that slightly before the submission of this manuscript, an elegant work on fluorohalogenation of diazos from Katzenellenbogen and co-workers was published online, see: N. Yasui, C. G. Mayne and J. A. Katzenellenbogen, *Org. Lett.*, 2015, 17, 5540.
- 15 (a) G. Maas, Angew. Chem., Int. Ed., 2009, 48, 8186; (b) F. Ye,
  C. Wang, Y. Zhang and J. Wang, Angew. Chem., Int. Ed., 2014,
  53, 11625.
- 16 ESI†
- 17 See ESI† for the details.
- 18 J. Wang, F. Liang and B. Chen, J. Org. Chem., 1998, 63, 8589.
- 19 W.-L. Man, J. Xie, P.-K. Lo, W. W.-Y. Lam, S.-M. Yiu, K.-C. Lau and T.-C. Lau, *Angew. Chem., Int. Ed.*, 2014, 53, 8463.
- 20 See ESI† for the details on DFT calculation.
- 21 H. Mao, Z. Tang, H. Hu, Y. Cheng, W.-H. Zheng and C. Zhu, Chem. Commun., 2014, 50, 9773.
- 22 Details on the reactions in presence of Brønsted acids, see:

23 H. Peng, Z. Yuan, H.-Y. Wang, Y.-L. Guo and G. Liu, *Chem. Sci.*, 2013, 4, 3172.