



Cite this: *Phys. Chem. Chem. Phys.*,
2022, 24, 29381

High-dimensional neural network potentials for accurate vibrational frequencies: the formic acid dimer benchmark†

Dilshana Shanavas Rasheeda, *^a Alberto Martín Santa Daría, ^b
Benjamin Schröder, ^c Edit Mátyus ^b and Jörg Behler ^{‡a}

In recent years, machine learning potentials (MLP) for atomistic simulations have attracted a lot of attention in chemistry and materials science. Many new approaches have been developed with the primary aim to transfer the accuracy of electronic structure calculations to large condensed systems containing thousands of atoms. In spite of these advances, the reliability of modern MLPs in reproducing the subtle details of the multi-dimensional potential-energy surface is still difficult to assess for such systems. On the other hand, moderately sized systems enabling the application of tools for thorough and systematic quality-control are nowadays rarely investigated. In this work we use benchmark-quality harmonic and anharmonic vibrational frequencies as a sensitive probe for the validation of high-dimensional neural network potentials. For the case of the formic acid dimer, a frequently studied model system for which stringent spectroscopic data became recently available, we show that high-quality frequencies can be obtained from state-of-the-art calculations in excellent agreement with coupled cluster theory and experimental data.

Received 23rd August 2022,
Accepted 18th November 2022

DOI: 10.1039/d2cp03893e

rsc.li/pccp

1 Introduction

Due to its unique capabilities to process and analyze large amounts of data, machine learning has nowadays found numerous applications in chemistry and related fields.^{1–5} Starting with the work of Doren and coworkers in 1995,⁶ a prominent example is the representation of the atomic interactions with quantum mechanical accuracy by learning the potential energy surface (PES) from a set of known training points computed using accurate electronic structure methods. The resulting machine learning potentials (MLP) have rapidly evolved in the past two decades.^{7–12} While first-generation MLPs have been restricted to small molecular systems,^{13,14} second-generation MLPs like high-dimensional neural network

potentials (HDNNPs),¹⁵ Gaussian approximation potentials,¹⁶ spectral neighbor analysis potentials,¹⁷ moment tensor potentials,¹⁸ atomic cluster expansion¹⁹ and many others have paved the way to large-scale simulations of condensed systems, from water and aqueous solutions to bulk materials and interfaces. Long-range electrostatic interactions,^{20–22} non-local charge transfer,^{23–27} and magnetism^{28,29} can also be explicitly considered in modern MLPs. Neural networks have also been used to study reactive molecular systems in the gas phase.^{30–32}

Due to their very flexible but simple functional form MLPs offer advantages like a very high accuracy of about 1 meV per atom in reproducing known reference total energies, an efficiency close to empirical force fields, and an unbiased description of many types of atomic interactions – from covalent bonds *via* dispersion interactions to metallic and ionic bonding. Still, although some recent MLPs include physical terms like electrostatics, in general the high flexibility of MLPs is curse and blessing at the same time, since the functional forms employed in MLPs, like neural networks or kernel methods, are not guaranteed to yield the correct physical shape of the PES. Therefore, large training sets are needed to ensure that a reliable representation of the PES is obtained in the training process. Moreover, a careful validation is required, since energy and force predictions can be inaccurate when extrapolating to atomic configurations that are too different from those in the training sets.

^a Universität Göttingen, Institut für Physikalische Chemie, Theoretische Chemie, Tammannstraße 6, 37077 Göttingen, Germany.

E-mail: dilshana.rasheeda@chemie.uni-goettingen.de

^b ELTE, Eötvös Loránd University, Institute of Chemistry, Pázmány Péter sétány 1/A, 1117 Budapest, Hungary

^c Universität Göttingen, Institut für Physikalische Chemie, Tammannstraße 6, 37077 Göttingen, Germany

† Electronic supplementary information (ESI) available. See DOI: <https://doi.org/10.1039/d2cp03893e>

‡ Present address: Lehrstuhl für Theoretische Chemie II, Ruhr-Universität Bochum, 44780 Bochum, Germany, and Atomistic Simulations, Research Center Chemical Sciences and Sustainability, Research Alliance Ruhr.



These limited extrapolation capabilities of MLPs and the need for a dense enough sampling of the reference data set raise the question how modern MLPs can be validated. While for early MLPs constructed for small molecular systems a systematic, grid-based mapping of the underlying PES has been possible allowing for rigorous quality control, the situation is different for second-generation MLPs designed for very large systems. Here, the total energy is typically constructed as a sum of environment-dependent atomic energies, resulting in a linear scaling of the computational effort with system size enabling simulations of thousands of atoms. However, since the atomic energies, which are not quantum mechanical observables, can be considered as mathematical auxiliary quantities, error compensation might occur reducing the transferability of the potentials. Further, typically investigated quantities like the root mean squared errors (RMSE) of energies and forces are of limited use, since they can only be computed for the available reference structures, and as averaged properties they may not allow to assess the quality of all the fine details of the PES.

Here, we propose to employ the harmonic and anharmonic vibrational frequencies as a sensitive probe to validate the quality of MLPs pinpointing on close-to equilibrium structures, which are particularly important for spectroscopic applications. While far-from equilibrium behaviour is in principle equally interesting also for spectroscopy, any PES not being able to describe the equilibrium properties correctly could not be trusted for spectroscopic applications. This tool can thus be viewed as complementary to other procedures like active learning that are covering a wide range of non-equilibrium structures.^{33–35}

Vibrational frequencies have been studied since the advent of MLPs, mainly for small molecular systems, with great success,^{36,37} and very accurate approaches for constructing MLPs suitable for this purpose have been derived.^{38–40} Still, most of these methods exhibit an unfavorable scaling with system size. Second-generation potentials allow to address larger systems, and some vibrational studies for larger molecules,⁴¹ clusters,⁴² and condensed systems^{43–46} have been reported. However, MLPs suitable for condensed systems have been typically constructed relying on density functional theory (DFT), which, although offering a good compromise between accuracy and efficiency for many systems, does not provide spectroscopic-quality vibrational frequencies. Moreover, studying complex condensed systems does not allow to disentangle all the subtle atomic interactions, which would be required for a thorough validation of the PES.

In the present work, we fill this gap by investigating in detail the accuracy that can be achieved by HDNNPs as a typical example for second-generation MLPs. We explore the limits of this method, which has originally been designed for dealing with very large numbers of atoms, by carefully training a coupled-cluster-quality⁴⁷ PES for a moderately sized system. This does not only allow us to compute harmonic frequencies but also to assess the quality of highly accurate anharmonic frequencies using state-of-the-art methods under well-controlled conditions.

As a model system we have chosen the formic acid dimer (FAD), a doubly hydrogen-bonded complex, which in recent years has become a benchmark system for the development of molecular PESs and the calculation of accurate vibrational spectra thanks to the increasing body of gas-phase spectroscopic data.^{48–54} It is the smallest hydrogen bonded complex with double proton transfer and as such it is a system of high interest for spectroscopy and theoretical dynamic studies due to the possible delocalization of the nuclei over the two wells. This double proton transfer is challenging since it cannot be described by standard near-equilibrium tools based on normal coordinates and perturbation theory. Even for this seemingly small ten-atom system and its 24-dimensional PES the determination of accurate anharmonic frequencies is computationally demanding and thus hardly accessible by a direct application of wave function electronic structure methods. For this reason the standard approach is the intermediate construction of an analytic PES for the determination of vibrational frequencies.

Hence, in general, theoretical studies on vibrational frequencies have three limiting factors: the electronic structure method, its representation by a multi-dimensional PES function, and the vibrational treatment. Regarding the first two aspects, two *ab initio*-based analytic PESs have been proposed for FAD in the literature. In 2016, Qu and Bowman developed the first full-dimensional potential energy surface⁵⁵ (henceforth labelled as QB16) for FAD by fitting permutationally invariant polynomials, a very accurate method providing PESs of a quality similar to modern MLPs,⁵⁶ to 13 475 *ab initio* energy points computed at the CCSD(T)-F12/haTZ level of electronic structure theory. Later, they carried out vibrational configuration interaction computations in normal coordinates^{57–59} on this surface. Recently, two of us used the QB16 PES and tested, using a reduced dimensionality model, the utility of normal coordinates or a possible efficiency gain of using curvilinear (normal) coordinates⁶⁰ in the GENIUSH program.⁶¹ During this work, several fundamental, combination, and overtone frequencies in the fingerprint range were obtained in an excellent agreement with experiment.⁵³ However, two (totally symmetric) fundamental frequencies were obtained strongly blueshifted in comparison with the harmonic frequencies of the QB16 PES⁵⁵ and in comparison with the experimental value.⁵³ Based on these observations and due to some ‘artificial’ features of the QB16 PES that made it necessary to restrict the quadrature grid used for the vibrational computations, it was concluded that further work on improving the FAD PES is required.

In 2022, Käser and Meuwly reported another full-dimensional PES (Phys_{TL}) for FAD⁶² generated by the message passing neural network PhysNet.²² It is based on 26 000 MP2/aug-cc-pVTZ single point energies, which have then been transfer-learned employing 866 CCSD(T)/aug-cc-pVTZ energies to obtain an approximately coupled cluster-quality PES that has also been used in the computation of harmonic and anharmonic vibrations.

Regarding experiment, FAD is a very well-studied system (see, e.g. ref. 53 and 54 and references therein), and a wealth of data is available for the validation of theoretical frequencies.



In early work, thermal gas phase spectroscopy of FAD has been employed,^{63,64} while more recently jet-cooled infrared and Raman spectra of FAD in the monomer finger print region up to 1500 cm⁻¹ have been studied in ref. 54. Over the past one and a half decades, all intermolecular vibrational fundamentals and several combination and overtone bands of FAD have been determined in the gas phase with an experimental uncertainty of 1 cm⁻¹.^{48-52,65-69} The experimental results including a critical evaluation of theoretical work, which appear to be still lagging behind experiment, have been recently reviewed in ref. 53.

The aim of the present work is to develop a robust and full-dimensional high-quality HDNNP for FAD and to benchmark the obtained frequencies using the best available theoretical and experimental data. By robustness, we mean that the 24-dimensional (24D) hypersurface provides a faithful representation (possibly without ‘holes’ or other unphysical features) of the *ab initio* electronic energy over the relevant quantum dynamical range, and this robustness persists irrespective of the actual choice of (normal or curvilinear) internal coordinates. By accuracy, in this context, we mean that the fitted hypersurface reproduces *ab initio* points sufficiently closely. ‘Sufficient’ is determined in relation with prospective (ro)vibrational computations which are to be compared with (gas-phase) experimental infrared and Raman spectroscopy data.⁵³

Next to the RMSE, which is the most common quality measure of PESs, we make use of additional quantities in this work that allow further refinement of the potential energy surface for spectroscopic applications. For this purpose, first, we define an accuracy goal for the harmonic frequencies that are expected to be reproduced by the PES to within 10 cm⁻¹ with respect to the *ab initio* harmonic frequency values computed at the same level of electronic structure theory. Furthermore, to have a relatively compact assessment of the mode coupling representation in the PES, we test the second-order vibrational perturbation theory (VPT2) frequencies of the PES again compared to the direct *ab initio* values as well as experimental results. Finally, although a semi-rigid description involving relatively small amplitude vibrations about an equilibrium structure of the PES appears to be a good starting point for FAD, the concerted proton tunneling of the double hydrogen bond qualifies this complex for the family of systems with multiple (two) large-amplitude motions. ‘High’-dimensional systems with multiple large-amplitude motions, *i.e.*, motions in which nuclei are delocalized over multiple PES wells, are common in molecular systems and cannot be efficiently described by using perturbative methods developed about equilibrium structure properties (underlying the normal coordinate concept). An efficient quantum dynamics description of these types of systems is currently an active and challenging field for methodological developments.⁷⁰⁻⁷⁶ These developments can be tested and validated with respect to precise spectroscopy data, assuming that a faithful and accurate PES representation for the molecular system is available.

After giving a brief summary of the employed methods in Section 2 and the computational details in Section 3, the results are presented in Section 4. First, we assess the quality of the

HDNNP, which is obtained by iteratively increasing the amount of reference data, until a converged potential is obtained. This PES is then characterized by its harmonic frequencies, which are compared to coupled cluster data. Finally, we report anharmonic frequencies obtained from VPT2 and reduced-dimensionality variational calculations, which allow direct validation of the MLP using reference *ab initio* calculations and accurate experimental data.

2 Methods

2.1 High-dimensional neural network potentials

High-dimensional neural network potentials have been introduced in 2007 by Behler and Parrinello¹⁵ as the first type of MLP applicable to large condensed systems containing thousands of atoms. This is achieved by representing the potential energy E of the system as a sum of atomic energies E_i , which depend on the local chemical environment up to a cutoff radius,

$$E = \sum_{i=1}^{N_{\text{atom}}} E_i. \quad (1)$$

Each of these atomic energies is the output of an individual atomic feed-forward neural network describing the functional relation between the respective energy contribution and the local atomic structure. The weight parameters of these neural networks are determined in an iterative training process using known energies (and often also forces) obtained from reference electronic structure calculations. The weight parameters and architectures of all atomic neural networks of a given chemical element are constrained to be the same making the potential transferable to different system sizes.

To ensure the mandatory translational, rotational and permutational invariances of a HDNNP based PES, the local atomic environments are typically characterized by vectors of atom-centered symmetry functions (ACSF)⁷⁷ as geometric descriptors, which meet these requirements by construction and provide local structural fingerprints. The inclusion of a cutoff function ensures that the ACSFs smoothly decay to zero in value and slope at the cutoff radius, which is commonly chosen between 5 and 10 Å.

Several extensions to second-generation HDNNPs have been introduced in recent years, like the consideration of long-range electrostatic interactions based on environment-dependent charges,^{20,42} non-local charge transfer,²⁴ and magnetic interactions.²⁸ More details about HDNNPs and their properties can be found in several recent reviews.⁷⁸⁻⁸⁰ In the present work a second-generation HDNNP relying on “short-range” atomic energies only is employed, since for a comparably small benchmark system like the FAD all interactions can be fully described as a function of the local chemical environments provided that a sufficiently large cutoff radius is chosen.

2.2 Frequency calculations

2.2.1 Harmonic frequencies. The determination of harmonic frequencies ω is a routine task in theoretical spectroscopic investigations.⁸¹ Here, we construct the Cartesian second



derivative matrix (Hessian) by means of finite differences based on the HDNNP. Mass weighting and diagonalization then yields the desired harmonic vibrational frequencies. These are then compared directly to the corresponding results of reference *ab initio* calculations.

2.2.2 Second-order vibrational perturbation theory. The anharmonic fundamental transition frequency ν_i within VPT2 is given by⁸²

$$\nu_i = \omega_i + 2x_{ii} + \frac{1}{2} \sum_{j \neq i} x_{ij}, \quad (2)$$

where ω_i is the harmonic vibrational wavenumber and the x_{ij} are the anharmonicity constants that account for anharmonicity in the vibrational mode i as well as the coupling to other modes j . Eqn (2) is based on a quartic force field (QFF), *i.e.*, a potential energy expression in terms of dimensionless normal coordinates $\mathbf{q} = \{q_i\}$ given by a Taylor expansion up to fourth order,⁸²

$$E(\mathbf{q}) = \frac{1}{2} \sum_i \omega_i q_i^2 + \frac{1}{6} \sum_{ijk} \phi_{ijk} q_i q_j q_k + \frac{1}{24} \sum_{ijkl} \phi_{ijkl} q_i q_j q_k q_l. \quad (3)$$

In eqn (3) the so-called cubic and quartic force constants are denoted by ϕ_{ijk} and ϕ_{ijkl} , respectively. A contact transformation of the (ro)vibrational Hamiltonian^{83–85} up to second order then yields formulae for the x_{ij} in terms of molecular parameters,

$$x_{ii} = \frac{1}{4} \phi_{iii} - \frac{1}{16} \sum_j \phi_{ijj}^2 \frac{8\omega_i^2 - 3\omega_j^2}{\omega_j(4\omega_i^2 - \omega_j^2)} \quad (4)$$

and

$$x_{ij} = \frac{1}{4} \phi_{ijj} - \frac{1}{4} \sum_k \phi_{iik} \phi_{kjj} \frac{1}{\omega_k} - \frac{1}{2} \sum_k \phi_{ijk} \frac{\omega_k^2 - \omega_i^2 - \omega_j^2}{\Delta_{ijk}} + \sum_{\alpha} B_c^{\alpha} \zeta_{ij}^{\alpha} \left(\frac{\omega_i}{\omega_j} + \frac{\omega_j}{\omega_i} \right), \quad (5)$$

where the resonance denominator Δ_{ijk} is given by

$$\Delta_{ijk} = (\omega_i + \omega_j + \omega_k)(\omega_i + \omega_j - \omega_k)(\omega_i - \omega_j + \omega_k)(\omega_i - \omega_j - \omega_k). \quad (6)$$

The last term in eqn (5) depends on the three equilibrium rotational constants B_c^{α} and the Coriolis coupling constants ζ_{ij}^{α} .

A closer look at eqn (4) and (5) shows that they contain differences of harmonic frequencies in denominators which may lead to so-called Fermi-Resonances. Two cases need to be accounted for within VPT2: $\omega_i \approx 2\omega_j$ and $\omega_i \approx \omega_j + \omega_k$. In fact, such vibrational resonances have been well documented experimentally for FAD (see ref. 54 and references therein) and therefore can be expected to interfere in the present VPT2 treatment. In order to allow a comparison between VPT2 results and experiment a special treatment is required which is sometimes referred to as GVPT2.⁸⁶ First, a change in the contact transformation removes the resonance denominators from the x_{ij} to yield so-called *deperturbed* anharmonicity constants x_{ij}^* .⁸⁷ In a second step the resonating vibrational states are treated using perturbation theory for (near) degenerate states⁸⁸ where

the coupling matrix element depends on a cubic force constant, *i.e.* ϕ_{ijj} or ϕ_{ijk} .

2.2.3 Reduced-dimensionality variational vibrational computations. A complete quantum dynamical characterization of a molecular system can be obtained by the variational solution of the (ro)vibrational Schrödinger equation including the multi-dimensional PES as an ‘effective’ interaction acting among the nuclei. The vibrational Hamiltonian, as a sum of the PES, E , and the kinetic energy operator written in general coordinates is

$$\hat{H} = \frac{1}{2} \sum_{k=1}^D \sum_{l=1}^D \tilde{g}^{-1/4} \hat{p}_k G_{kl} \tilde{g}^{1/2} \hat{p}_l \tilde{g}^{-1/4} + E \quad (7)$$

where $\hat{p}_k = -i\partial/\partial q_k$ ($k = 1, \dots, D \leq 3N_{\text{atom}} - 6$), and the q_k are vibrational coordinates whose definitions (and choice of the body-fixed frame) are encoded in the mass-weighted metric tensor, $\mathbf{g} \in \mathbb{R}^{D \times D}$, with $\tilde{g} = \det \mathbf{g}$ and $\mathbf{G} = \mathbf{g}^{-1}$. Rigorous geometrical constraints ($D < 3N_{\text{atom}} - 6$) of selected internal coordinates can be introduced in the quantum mechanical vibrational model by ‘deleting’ the relevant rows and columns of the \mathbf{g} matrix. A numerical kinetic energy operator (KEO) approach based on this formalism has been implemented in the GENIUSH computer program,⁶¹ *i.e.*, \mathbf{g} , \tilde{g} , and \mathbf{G} are evaluated at grid points. The original implementation relied on the discrete variable representation (DVR)⁸⁹ of the vibrational Hamiltonian, but more recently, this numerical KEO approach has been used with finite basis representation (FBR) and the Smolyak scheme,^{70,71,90} which opens the route towards higher-dimensional computations. For FAD, the lowest-energy vibrational frequencies from the fingerprint range have been converged for a series of reduced-dimensionality vibrational models defined in ref. 60.

The first applications with the new HDNNP based PES developed in the present work use an 8-dimensional vibrational model termed 8D(\mathcal{Jt}). This approach includes the six intermolecular modes (\mathcal{J}) and the *cis-trans* torsional degree of freedom (t) of both monomers (see Fig. 1), which was found to perform reasonably well in ref. 60.




Fig. 1 Formic acid dimer in its equilibrium structure. The intermolecular ($R, \theta, \phi, \alpha, \beta, \gamma$) and the two *cis-trans* torsional ($\tau^{(1)}$ and $\tau^{(2)}$) coordinates used in the 8-dimensional variational vibrational computations are also shown.






Fig. 2 Energy difference $\Delta E = E_{CC} - E_{PES}$ as a function of the reference energy E_{CC} relative to the energy of the global minimum for the QB16 PES,⁵⁵ HDNNP1, HDNNP2, and HDNNP3 (=FAD-HDNNP). The root-mean-squared errors (RMSE) for the HDNNPs are provided in Table 1.

RMSE of only about 0.35 meV per atom (28 cm^{-1}) for HDNNP3 in the relevant energy range up to $0.1 E_h$ (ca. 21950 cm^{-1}).

4.2 Vibrational frequencies

4.2.1 Harmonic frequencies. As an important test for spectroscopic applications, next we compare the harmonic frequencies of the PESs with the respective *ab initio* values corresponding to the same level of electronic structure theory. As described in the introduction we consider an agreement within $\pm 10 \text{ cm}^{-1}$ as a requirement for this purpose.

Fig. 3 shows the harmonic frequency errors obtained for the three main HDNNPs of the PES refinement procedure. Similar to the energy RMSEs we find a continuous improvement starting from HDNNP1 showing some rather large deviations up to 52 cm^{-1} , which are decreasing to at most 27 cm^{-1} for HDNNP2, finally reaching a very high quality in case of HDNNP3 with the largest deviation in the PES *vs.* *ab initio* harmonic frequencies being less than 7 cm^{-1} . Overall, the frequency RMSEs of HDNNP1, HDNNP2, and HDNNP3 are about 27 cm^{-1} , 9 cm^{-1} , and 4 cm^{-1} , respectively. Consequently, we choose HDNNP3 as the production-quality PES for vibrational calculations, which will be called simply 'FAD-HDNNP' in the following.

The lower part of Fig. 3 provides a comparison of FAD-HDNNP harmonic frequencies with the other two published FAD PESs.^{55,62} Concerning the accuracy of the PES at the harmonic level, FAD-HDNNP seems to outperform the earlier QB16⁵⁵ and Phys_{TL}⁶² potentials, which exhibit RMSEs of 8 cm^{-1} and 14 cm^{-1} with maximum deviations of 28 and 30 cm^{-1} ,




Fig. 3 Deviations of the harmonic vibrational frequencies ω_i with respect to the reference *ab initio* results (tight settings). Top panel depicts the deviations $\Delta\omega_i = \omega_i - \omega_{ref,i}$ for different HDNNP versions during the potential refinement with HDNNP3 corresponding to the final FAD-HDNNP. The bottom panel compares these differences for harmonic frequencies calculated from the QB16 PES by Qu and Bowman,⁵⁵ the harmonic frequencies published by Käser and Meuwly⁶² based on the transfer-learned Phys_{TL}, and the final FAD-HDNNP (= HDNNP3) results. The reference *ab initio* results have been obtained at the level of electronic structure theory that was used for the development of the respective PES, i.e. fc-CCSD(T)/aug-cc-pVTZ for Phys_{TL} (see ref. 62 for details) and fc-CCSD(T)-F12a/haTZ for QB16 as well as FAD-HDNNP. Root mean squared errors (RMSE) are provided in the legends.

respectively, from the *ab initio* harmonic frequency values of the corresponding level of electronic structure theory.

The numerical values of the harmonic frequencies for the newly developed FAD-HDNNP (and also for the HDNNP1 and HDNNP2 development stages) are collected in Table 2 together with the *ab initio* frequencies. Interestingly, the numerical values of the *ab initio* frequencies at the chosen level of theory depend notably on the precise details of the MOLPRO computation setup as shown in the last two columns of the table. The determined 4 cm^{-1} RMSE with a maximum deviation of 7 cm^{-1} of FAD-HDNNP appears to have reached the current accuracy limit of the available electronic structure methodology employing commonly used default settings, as even the MOLPRO frequencies exhibit changes up to 13 cm^{-1} in exceptional cases like ω_{15} when using tight settings. Still, even in this case the RMSE of FAD-HDNNP with respect to the tight MOLPRO data is very similar to the RMSE with respect to the default settings with only a marginally increased maximum deviation of 8 cm^{-1} in case of ω_{12} and ω_{13} .

The equilibrium structure of FAD obtained from the reference *ab initio* calculations can be compared to the minima of



Table 2 Comparison of the harmonic frequencies ω_i (in cm^{-1}) for the three HDNNPs and the respective reference *ab initio* results obtained at the fc-CCSD(T)-F12a/haTZ level of theory. The latter are computed using either the standard convergence criteria (termed default) or stricter convergence criteria (termed tight) with respect to the electronic energy and geometry optimization as well as smaller step sizes in the numerical gradient and Hessian (*cf.* Section 3)

Mode	Sym.	HDNNP1	HDNNP2	FAD-HDNNP HDNNP3	<i>Ab initio</i>	
					Default	Tight
ω_1	A_g	3218	3212	3209	3203	3207
ω_2	A_g	3060	3077	3103	3105	3103
ω_3	A_g	1708	1723	1722	1717	1718
ω_4	A_g	1443	1485	1486	1484	1482
ω_5	A_g	1376	1417	1410	1413	1411
ω_6	A_g	1254	1249	1257	1257	1255
ω_7	A_g	689	681	688	688	686
ω_8	A_g	216	215	214	211	210
ω_9	A_g	164	173	167	171	170
ω_{10}	B_g	1073	1083	1083	1085	1083
ω_{11}	B_g	980	960	957	960	955
ω_{12}	B_g	257	241	257	258	249
ω_{13}	A_u	1128	1106	1109	1102	1101
ω_{14}	A_u	937	983	979	986	985
ω_{15}	A_u	213	161	180	186	173
ω_{16}	A_u	68	71	71	76	68
ω_{17}	B_u	3324	3323	3312	3305	3309
ω_{18}	B_u	3108	3085	3100	3101	3099
ω_{19}	B_u	1773	1783	1784	1782	1782
ω_{20}	B_u	1411	1459	1459	1456	1453
ω_{21}	B_u	1358	1414	1410	1405	1407
ω_{22}	B_u	1208	1247	1260	1260	1260
ω_{23}	B_u	704	718	712	716	715
ω_{24}	B_u	281	275	275	278	276

the HDNNPs. This is done in Table 3 and additional results obtained with the QB16 PES⁵⁵ are provided for comparison. Due to symmetry only 9 internal coordinates are required to characterize FAD. Similar to the harmonic frequencies, the agreement of the HDNNP geometrical parameters with the reference *ab initio* results improves with the refinement procedure. The largest difference is observed for the $r(\text{O} \cdots \text{O})$ distance which at the same time appears to be the most sensitive coordinate exemplified by a comparably large variation of 0.001 Å between the default and tight reference *ab initio*

Table 3 Geometrical parameters of the formic acid dimer minimum structure optimized at the reference *ab initio* level of theory and determined for different PESs. Bond lengths are provided in Ångströms and angles in degrees. HDNNP3 corresponds to the final FAD-HDNNP for spectroscopic use

Parameter	<i>Ab initio</i>					
	Default	Tight	QB16 ^a	HDNNP1	HDNNP2	HDNNP3
$r(\text{O}-\text{H})$	0.9934	0.9932	0.9927	0.9945	0.9925	0.9936
$r(\text{C}-\text{H})$	1.0929	1.0930	1.0929	1.0927	1.0937	1.0927
$r(\text{C}-\text{O})$	1.3113	1.3114	1.3116	1.3104	1.3121	1.3112
$r(\text{O} \cdots \text{O})$	2.6748	2.6758	2.6778	2.6729	2.6791	2.6709
$r(\text{C}=\text{O})$	1.2177	1.2176	1.2174	1.2192	1.2172	1.2178
$\angle \text{O}=\text{C}-\text{O}$	126.14	126.14	126.15	126.26	126.13	126.13
$\angle \text{O}=\text{C}-\text{H}$	122.02	122.03	122.05	122.04	122.15	121.99
$\angle \text{C}-\text{O}-\text{H}$	109.77	109.76	109.73	109.42	109.93	109.79
$\angle \text{O}-\text{H} \cdots \text{O}$	178.93	178.93	178.95	179.51	179.01	178.86

^a Qu and Bowman.⁵⁵

geometries. Nevertheless, the agreement of HDNNP 3 (= FAD-HDNNP) with the (tight) reference *ab initio* geometry is good with an RMSE of 0.002 Å for the bond distances and 0.04° for the angular coordinates.

Finally, we investigated the performance of the HDNNPs in describing the geometry and barrier height of the double proton transfer transition state. The details are provided in the ESI† and results are compared to new *ab initio* calculations. Although not the main focus of the present investigation, we find good agreement between FAD-HDNNP and the reference results with an RMSE of 0.002 Å for Cartesian coordinates and a deviation of only 12 cm^{-1} in the barrier height.

4.2.2 VPT2 frequencies. Since the harmonic frequencies are well described by the FAD-HDNNP, it is now appropriate to investigate the representation of mode couplings as a next step. A compact measure for the mode couplings near the equilibrium structure can be obtained from a comparison of the VPT2 frequencies computed using the PES with gas-phase experimental results^{49,51,54,64,103} as well as direct *ab initio* results which are still feasible for FAD, although the computations are very demanding taking several weeks. The corresponding anharmonic fundamental transition frequencies are provided in Table 4. The normal coordinate QFF parameters obtained from FAD-HDNNP and those calculated *ab initio* are compiled in the ESI,† and can be used for further assessment of the representation of the mode couplings. This allows a comparison which is completely unaffected by details of the VPT2 resonance treatment.

A direct comparison between VPT2-based results and experiment for the high-frequency hydrogen-stretching vibrations is problematic. This is on the one hand due to the fact that these fundamentals are in an energy range of already high state density which leads to substantial anharmonic couplings/resonances beyond what can be reliably treated using VPT2.⁵⁷ On the other hand experimental results for these fundamentals also carry a significant uncertainty which may exceed 10 cm^{-1} . Therefore these frequencies are not well suited for benchmarking purposes and we will below focus on the fundamentals < 1800 cm^{-1} . Nevertheless, the hydrogen stretching fundamental frequencies obtained with FAD-HDNNP agree well with the reference *ab initio* VPT2 results. A maximum deviation of 11 cm^{-1} is found for ν_1 which probably is due to the resonance effects mentioned above that render these frequencies highly sensitive to small details of the PES representation.

For the lower frequency modes FAD-HDNNP reproduces the reference *ab initio* results with an RMSE of 5 cm^{-1} where the maximum deviation (for resonance free fundamentals) of 23 cm^{-1} is observed for ν_{11} . Considering the deviation of the corresponding harmonic frequencies is only 2 cm^{-1} (*cf.* Table 2) this indicates a problem in correctly describing the anharmonicity in the symmetric OH out-of-plane bend. Overall the agreement between FAD-HDNNP and the experimental results is good with an RMSE of 9 cm^{-1} . The RMSE of the reference results is slightly better with 6 cm^{-1} but still the ν_{11} mode shows a large deviation of 12 cm^{-1} . Nevertheless, these results are well within the typical range of errors that are to be



Table 4 Comparison of the VPT2 fundamental frequencies (in cm^{-1}) obtained from the reference *ab initio* calculations and the FAD-HDNNP PES with experimental data^{49,51,54,64,103}

Mode	Sym.	<i>Ab initio</i> ^a	FAD-HDNNP	Exp.
ν_1	A_g	2909	2920	—
ν_2	A_g	2942	2948	—
ν_3	A_g	1672 ^b	1677 ^c	1664 ^d
ν_4	A_g	1431	1433	1430
ν_5	A_g	1375	1375	1375
ν_6	A_g	1225	1229	1224
ν_7	A_g	679	682	681
ν_8	A_g	194	197	194
ν_9	A_g	157	164	161
ν_{10}	B_g	1061	1058	1058
ν_{11}	B_g	923	934	911
ν_{12}	B_g	241	247	242
ν_{13}	A_u	1072	1074	1069
ν_{14}	A_u	959 ^e	964 ^f	939 ^g
ν_{15}	A_u	162	166	168
ν_{16}	A_u	67	68	69
ν_{17}	B_u	3044	3041	—
ν_{18}	B_u	2935	2941	—
ν_{19}	B_u	1741	1745	1741
ν_{20}	B_u	1406	1416	1407
ν_{21}	B_u	1372	1375	1372
ν_{22}	B_u	1234 ^h	1233 ⁱ	1234 ^j
ν_{23}	B_u	704	706	708
ν_{24}	B_u	262	264	264

^a CCSD(T)-F12a/hATZ using tight settings. ^b Fermi-resonance coupled with $\nu_4 + \nu_8$ at 1619 cm^{-1} . ^c Fermi-resonance coupled with $\nu_4 + \nu_8$ at 1625 cm^{-1} . ^d Experimental bands at 1664 and 1668 cm^{-1} . ^e Fermi-resonance coupled with $\nu_{12} + \nu_{23}$ at 938 cm^{-1} . ^f Fermi-resonance coupled with $\nu_{12} + \nu_{23}$ at 941 cm^{-1} . ^g Experimental bands at 939 and 953 cm^{-1} . ^h Fermi-resonance coupled with $\nu_{10} + \nu_{15}$ at 1218 cm^{-1} . ⁱ Fermi-resonance coupled with $\nu_{10} + \nu_{15}$ at 1219 cm^{-1} . ^j Experimental bands at 1220 , 1225 , and 1234 cm^{-1} .

expected for the underlying level of *ab initio* theory. To improve on this would require the inclusion of high-level corrections such as core-valence correlation and higher-order correlation beyond CCSD(T). While such composite schemes have been shown to provide high-quality potentials for small molecules,^{104–107} their computational cost is prohibitive for FAD.

For ν_{14} we observe a large deviation of 25 cm^{-1} when comparing the FAD-HDNNP and the experimental frequency of 964 and 939 cm^{-1} , respectively. In agreement with experimental results^{54,108} we find this mode to be in Fermi-resonance with $\nu_{12} + \nu_{23}$ at 941 cm^{-1} to be compared to the experimental value of 961 cm^{-1} . A close look at these numbers may indicate a possible misassignment. However, upon inspection of the VPT2 results we find that these vibrational states are in an almost perfect resonance (56:44 mixing) with deperturbed energies of 952 and 954 cm^{-1} for $(\nu_{12} + \nu_{23})^*$ and ν_{14}^* , respectively. As such this resonance is very sensitive to the level of theory and small changes in the PES can easily reverse the state ordering. Note that this situation is also present in the reference *ab initio* results and therefore appears to be due to the convergence of the PES with respect to the employed electronic structure level.

Finally, our calculations indicate a Fermi-resonance between ν_{22} and $\nu_{10} + \nu_{15}$ with anharmonic transition frequencies of 1234 and 1219 cm^{-1} , respectively. Nejad *et al.* have discussed

this resonance in detail,⁵⁴ proposing a new assignment for the transitions in the triplet of bands centered at 1220 , 1225 , and 1234 cm^{-1} , *i.e.*, a resonance triad assigned to $\nu_{10} + \nu_{15}$, $\nu_9 + \nu_{11} + \nu_{15}$ and ν_{22} . In contrast, we find no involvement of $\nu_9 + \nu_{11} + \nu_{15}$ in the VPT2 results for ν_{22} based on FAD-HDNNP. This (preliminary) result is also obtained from the *ab initio* reference calculations based on the resonance detection criterium for the ν_{10} fundamental. However, the situation is complicated by the fact that the ν_{11} shows the largest error (for fundamentals not affected by resonances) with respect to experiment of more than 20 cm^{-1} in case of FAD-HDNNP which in consequence shifts the $\nu_9 + \nu_{11} + \nu_{15}$ band by a similar amount and thus it can not interact with $\nu_{10} + \nu_{15}$ through the rather small coupling matrix element of only a few cm^{-1} . In contrast, for the reference *ab initio* calculations the VPT2 frequencies 1219 , 1226 , and 1233 cm^{-1} for $\nu_{10} + \nu_{15}$, $\nu_9 + \nu_{11} + \nu_{15}$ and ν_{22} , respectively, agree nicely with the experimental results. Upon enforcing the coupling between $\nu_{10} + \nu_{15}$ and $\nu_9 + \nu_{11} + \nu_{15}$ the previous frequencies change only slightly to 1218 , 1227 , and 1234 cm^{-1} but with a complicated mixing of the harmonic basis functions in agreement with the results of Nejad *et al.*⁵⁴ Clearly, a correct description of this intricate resonance triad both for the frequencies of the involved bands as well as the intensity pattern governed by the mixing ratios will be an excellent benchmark test for a theoretical spectroscopic description of FAD.

4.2.3 Reduced-dimensionality variational vibrational frequencies. Finally, to perform the 8D intermolecular-plus-torsion vibrational computations, we first determined a potential-optimized DVR for every vibrational degree of freedom following the procedure described in ref. 60. All remaining, *i.e.*, constrained, degrees of freedom have been fixed at the values of the FAD-HDNNP equilibrium geometry (*cf.* Table 3).

The one-dimensional cuts of the three investigated HDNNPs and the QB16 potential along the intermolecular coordinates plotted in Fig. 4 show that the newly developed FAD-HDNNP behaves well along all the intramolecular degrees of freedom consisting of the distance between the monomers and relative orientation (*cf.* Fig. 1). In contrast to the QB16 potential, which shows a low-energy oscillation for $\phi \approx 150^\circ$, no artificial cutoff of the primitive grid intervals has been found to be necessary for running the calculations.

The obtained vibrational frequencies converged for the 8D($\mathcal{J}\mathcal{T}$) intermolecular-plus-torsional representation are collected in Table 5. Overall the agreement for both QB16 and FAD-HDNNP with experiment is reasonably good with deviations of only a few cm^{-1} for most frequencies. However, similarly to previous results discussed in ref. 60, we can still observe a non-negligible blue shift in the problematic fundamental frequencies ν_8 and ν_9 . With the present, extensively tested PESs, we can identify two possible origins and corresponding solutions for this shift. First, either it is a consequence of the constrained coordinates resulting in a too steep potential energy well at the equilibrium cut, which may be overcome by relaxing the constraint coordinates. Or, second, the number of the active vibrational degrees of freedom should be increased





Fig. 4 1-dimensional cuts of the three investigated HDNNPs and the QB16 potential along the intermolecular coordinates (cf. Fig. 1) of the formic acid dimer used in the reduced-dimensionality variational computations.

Table 5 Vibrational energies referenced to the zero-point vibrational energy (in cm^{-1}) obtained with the 8D(\mathcal{J} t) intermolecular-torsional model in the GENIUSH program using the QB16 PES and FAD-HDNNP. The inactive degrees of freedom are fixed at their respective equilibrium values

Assignment ⁶⁰	$\tilde{\nu}_{\text{QB16}}^{60}$	$\tilde{\nu}_{\text{FAD-HDNNP}}$	$\tilde{\nu}_{\text{expt}}^{53}$
ν_{16}	70	70	69.2
$2\nu_{16}$	141	140	139
ν_{15}	162	171	168.5
ν_9/ν_8	191	190	161
ν_8/ν_9	208	210	194
$3\nu_{16}$	211	210	
$\nu_{15} + \nu_{16}$	232	240	
ν_{12}	239	243	242
ν_{24}	253	253	264
$\nu_9 + \nu_{16}$	262	260	
$\nu_8 + \nu_{16}$	277	279	
$4\nu_{16}$	281	280	
$\nu_{15} + 2\nu_{16}$	303	309	
$\nu_{12} + \nu_{16}$	310	311	311
$\nu_{24} + \nu_{16}$	323	322	
$2\nu_{15}$	324	330	336
$\nu_9 + 2\nu_{16}$	332	340	
$\nu_8 + 2\nu_{16}$	347	348	

in the GENIUSH program. However, in the latter case the computational cost would increase rapidly in DVR, and for this reason, it will be necessary to use the more efficient FBR-Smoljak representation.^{70,71,90}

In comparison with VPT2 (cf. Section 4.2.2), it is interesting to note that VPT2 gets all fundamental frequencies (including ν_8 and ν_9) mostly correct (cf. Table 4). In Fig. 5 of ref. 53, Nejad and Suhm contrasted the good performance of VPT2 ($\pm 5 \text{ cm}^{-1}$ with respect to experiment) against the very large deviations (15–40 cm^{-1}) of the fundamentally more complete VCI computations.¹⁰⁹ Their observations motivated further theoretical and

computational work, including an inquiry about the efficiency of normal coordinates for the intermolecular (low-frequency) vibrations for this system.⁶⁰

5 Summary & conclusions

In this work we have examined the accuracy of machine-learned potential energy surfaces using the prototypical case of high-dimensional neural network potentials. Like many other modern machine learning potentials, HDNNPs have been primarily developed to transfer the accuracy of electronic structure methods to very large systems containing thousands of atoms, with the aim to perform large-scale molecular dynamics simulations. Consequently, the validation of the resulting multidimensional PESs is a difficult task. In this benchmark study, we have selected a system of moderate size, the formic acid dimer, for which vibrational frequencies at the CCSD(T) level of theory are accessible as a probe for assessing the quality of the PES along with a wealth of experimental data.

For the development of the full-dimensional HDNNP we have pursued a three-step procedure based on increasing reference data sets in the training process. First, a generally faithful representation of the PES without artifacts like artificial “holes” of overall good quality is generated. Second, this surface is then iteratively refined by adding further points to ensure that the harmonic frequencies of the PES reproduce within $\pm 10 \text{ cm}^{-1}$ the harmonic frequencies of the CCSD(T) level of theory, which serves as our reference. Third, the representation of (lower-order) couplings is investigated by computing the VPT2 frequencies and the corresponding normal-coordinate quartic force constants in comparison with experimental data and reference *ab initio* results.

This carefully validated full-dimensional FAD-HDNNP surface has been used in (pilot) 8-dimensional, curvilinear, variational computations focusing on the low-energy intermolecular range. Further progress in the variational vibrational methodology is required for reaching a higher-energy spectral range including a higher number of active vibrational degrees of freedom.

The finally obtained FAD-HDNNP potential energy surface shows a very high quality, which, in combination with the wealth of available, high-quality experimental data,^{49,53,63–67,110} we expect to be very useful for future developments in quantum dynamics and spectroscopic applications, which rely on robust and accurate PESs.

Data availability

Data for this paper, including the newly calculated *ab initio* data set used in the fitting of fit FAD-HDNNP, the necessary RuNner input files for FAD-HDNNP, the *ab initio* points employed in the reference QFF generation *via* numerical differentiation, and the obtained QFF parameters (equilibrium geometry, normal coordinates and force constants) for both the *ab initio* reference as well as FAD-HDNNP are available at GRO.data



(10.25625/ZDGKYA).¹¹¹ Additionally, the ESI[†] provides details on the RuNNer settings and the ACSFs used during the HDNNP development, the transition state structure determined from different PESs and *ab initio* calculations, and details on the QFF parameters.

Conflicts of interest

There are no conflicts to declare.

Acknowledgements

This project has been funded by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) – 389479699/GRK2455. EM is grateful for a DFG Mercator Fellowship granted in the framework of this project. AMSD and EM are grateful for the financial support of the Swiss National Science Foundation (PROMYS Grant, No. IZ11Z0_166525). We thank Chen Qu and Joel Bowman for sharing with us their formic acid dimer PES and for providing their *ab initio* data set. Computing time has been available through the DFG project INST186/1294-1 FUGG (Project No. 405832858). Discussions with Ricardo Mata, Rainer Ostwald, Arman Nejad, Martin Suhm and Guntram Rauhut are gratefully acknowledged.

Notes and references

- 1 K. T. Butler, D. W. Davies, H. Cartwright, O. Isayev and A. Walsh, *Nature*, 2018, **559**, 547–555.
- 2 A. C. Mater and M. L. Coote, *J. Chem. Inf. Model.*, 2019, **59**, 2545–2559.
- 3 J. Gasteiger and J. Zupan, *Angew. Chem., Int. Ed. Engl.*, 1993, **32**, 503–527.
- 4 B. Sanchez-Lengeling and A. Aspuru-Guzik, *Science*, 2018, **361**, 360–365.
- 5 M. Meuwly, *Chem. Rev.*, 2021, **121**, 10218–10239.
- 6 T. B. Blank, S. D. Brown, A. W. Calhoun and D. J. Doren, *J. Chem. Phys.*, 1995, **103**, 4129–4137.
- 7 J. Behler, *J. Chem. Phys.*, 2016, **145**, 170901.
- 8 V. L. Deringer, M. A. Caro and G. Csányi, *Adv. Mater.*, 2019, **31**, 1902765.
- 9 F. Noé, A. Tkatchenko, K.-R. Müller and C. Clementi, *Ann. Rev. Phys. Chem.*, 2020, **71**, 361–390.
- 10 O. T. Unke, S. Chmiela, H. E. Saucedo, M. Gastegger, I. Poltavsky, K. T. Schütt, A. Tkatchenko and K.-R. Müller, *Chem. Rev.*, 2021, **121**, 10142–10186.
- 11 P. Friederich, F. Häse, J. Proppe and A. Aspuru-Guzik, *Nat. Mater.*, 2021, **20**, 750–761.
- 12 J. Behler and G. Csányi, *Eur. Phys. J. B*, 2021, **94**, 142.
- 13 C. M. Handley and P. L. A. Popelier, *J. Phys. Chem. A*, 2010, **114**, 3371–3383.
- 14 J. Behler, *Phys. Chem. Chem. Phys.*, 2011, **13**, 17930–17955.
- 15 J. Behler and M. Parrinello, *Phys. Rev. Lett.*, 2007, **98**, 146401.
- 16 A. P. Bartók, M. C. Payne, R. Kondor and G. Csányi, *Phys. Rev. Lett.*, 2010, **104**, 136403.
- 17 A. P. Thompson, L. P. Swiler, C. R. Trott, S. M. Foiles and G. J. Tucker, *J. Comput. Phys.*, 2015, **285**, 316–330.
- 18 A. V. Shapeev, *Multiscale Model. Simul.*, 2016, **14**, 1153–1173.
- 19 R. Drautz, *Phys. Rev. B*, 2019, **99**, 014104.
- 20 N. Artrith, T. Morawietz and J. Behler, *Phys. Rev. B: Condens. Matter Mater. Phys.*, 2011, **83**, 153101.
- 21 K. Yao, J. E. Herr, D. W. Toth, R. Mckintyre and J. Parkhill, *Chem. Sci.*, 2018, **9**, 2261–2269.
- 22 O. T. Unke and M. Meuwly, *J. Chem. Theory Comput.*, 2019, **15**, 3678–3693.
- 23 S. A. Ghasemi, A. Hofstetter, S. Saha and S. Goedecker, *Phys. Rev. B: Condens. Matter Mater. Phys.*, 2015, **92**, 045131.
- 24 T. W. Ko, J. A. Finkler, S. Goedecker and J. Behler, *Nat. Commun.*, 2021, **12**, 398.
- 25 X. Xie, K. A. Persson and D. W. Small, *J. Chem. Theory Comput.*, 2020, **16**, 4256–4270.
- 26 L. Jacobson, J. Stevenson, F. Ramezanghorbani, D. Ghoreishi, K. Leswing, E. Harder and R. Abel, *J. Chem. Theory Comput.*, 2022, **18**, 2354–2366.
- 27 T. W. Ko, J. A. Finkler, S. Goedecker and J. Behler, *Acc. Chem. Res.*, 2021, **54**, 808–817.
- 28 M. Eckhoff and J. Behler, *npj Comput. Mater.*, 2021, **7**, 170.
- 29 I. Novikov, B. Grabowski, F. Körmann and A. Shapeev, *npj Comput. Mater.*, 2022, **8**, 13.
- 30 B. Jiang, J. Li and H. Guo, *Int. Rev. Phys. Chem.*, 2016, **35**, 479–506.
- 31 J. Li, K. Song and J. Behler, *Phys. Chem. Chem. Phys.*, 2019, **21**, 9672–9682.
- 32 D. Lu, J. Behler and J. Li, *J. Phys. Chem. A*, 2020, **124**, 5737–5745.
- 33 N. Artrith and J. Behler, *Phys. Rev. B: Condens. Matter Mater. Phys.*, 2012, **85**, 045439.
- 34 E. V. Podryabinkin and A. V. Shapeev, *Comput. Mater. Sci.*, 2017, **140**, 171–180.
- 35 J. S. Smith, B. Nebgen, N. Lubbers, O. Isayev and A. E. Roitberg, *J. Chem. Phys.*, 2018, **148**, 241733.
- 36 F. V. Prudente, P. H. Acioli and J. J. Soares Neto, *J. Chem. Phys.*, 1998, **109**, 8801–8808.
- 37 A. C. P. Bittencourt, F. V. Prudente and J. D. M. Vianna, *Chem. Phys.*, 2004, **297**, 153–161.
- 38 S. Manzhos and T. Carrington Jr., *J. Chem. Phys.*, 2007, **127**, 014103.
- 39 M. Malshe, R. Narulkar, L. M. Raff, M. Hagan, S. Bukkapatnam, P. M. Agrawal and R. Komanduri, *J. Chem. Phys.*, 2009, **130**, 184102.
- 40 T. Carrington Jr., *J. Chem. Phys.*, 2017, **146**, 120902.
- 41 M. Gastegger, J. Behler and P. Marquetand, *Chem. Sci.*, 2017, **8**, 6924–6935.
- 42 T. Morawietz, V. Sharma and J. Behler, *J. Chem. Phys.*, 2012, **136**, 064103.
- 43 V. Quaranta, M. Hellström, J. Behler, J. Kullgren, P. Mitev and K. Hermansson, *J. Chem. Phys.*, 2018, **148**, 241720.
- 44 T. Morawietz, O. Marsalek, S. R. Pattenaude, L. M. Streaker, D. Ben-Amotz and T. E. Markland, *J. Phys. Chem. Lett.*, 2018, **9**, 851–857.



- 45 S. Shepherd, J. Lan, D. M. Wilkins and V. Kapil, *J. Phys. Chem. Lett.*, 2021, **12**, 9108–9114.
- 46 G. M. Sommers, M. F. Calegari Andrade, L. Zhang, H. Wang and R. Car, *Phys. Chem. Chem. Phys.*, 2020, **22**, 10592–10602.
- 47 R. J. Bartlett and M. Musial, *Rev. Mod. Phys.*, 2007, **79**, 291–352.
- 48 M. Herman, R. Georges, M. Hepp and D. Hurtmans, *Int. Rev. Phys. Chem.*, 2000, **19**, 277–325.
- 49 R. Georges, M. Freytes, D. Hurtmans, I. Kleiner, J. Vander Auwera and M. Herman, *Chem. Phys.*, 2004, **305**, 187–196.
- 50 P. Zielke and M. A. Suhm, *Phys. Chem. Chem. Phys.*, 2007, **9**, 4528.
- 51 Z. Xue and M. A. Suhm, *J. Chem. Phys.*, 2009, **131**, 054301.
- 52 F. Kollipost, R. W. Larsen, A. V. Domanskaya, M. Nörenberg and M. A. Suhm, *J. Chem. Phys.*, 2012, **136**, 151101.
- 53 A. Nejad and M. A. Suhm, *J. Indian Inst. Sci.*, 2020, **100**, 1–15.
- 54 A. Nejad, K. A. E. Meyer, F. Kollipost, Z. Xue and M. A. Suhm, *J. Chem. Phys.*, 2021, **155**, 224301.
- 55 C. Qu and J. M. Bowman, *Phys. Chem. Chem. Phys.*, 2016, **18**, 24835–24840.
- 56 B. J. Braams and J. M. Bowman, *Int. Rev. Phys. Chem.*, 2009, **28**, 577–606.
- 57 C. Qu and J. M. Bowman, *J. Phys. Chem. Lett.*, 2018, **9**, 2604–2610.
- 58 C. Qu and J. M. Bowman, *J. Chem. Phys.*, 2018, **148**, 241713.
- 59 C. Qu and J. M. Bowman, *Faraday Discuss.*, 2018, **212**, 33–49.
- 60 A. Martín Santa Daría, G. Avila and E. Mátyus, *Phys. Chem. Chem. Phys.*, 2021, **23**, 6526–6535.
- 61 E. Mátyus, G. Czako and A. G. Császár, *J. Chem. Phys.*, 2009, **130**, 134112.
- 62 S. Käser and M. Meuwly, *Phys. Chem. Chem. Phys.*, 2022, **24**, 5269–5281.
- 63 R. C. Millikan and K. S. Pitzer, *J. Am. Chem. Soc.*, 1958, **80**, 3515–3521.
- 64 J. E. Bertie and K. H. Michaelian, *J. Chem. Phys.*, 1982, **76**, 886.
- 65 V. V. Matyilitsky, C. Riehn, M. F. Gelin and B. Brutschy, *J. Chem. Phys.*, 2003, **119**, 10553–10562.
- 66 F. Ito and T. Nakanaga, *Chem. Phys.*, 2002, **277**, 163–169.
- 67 J. E. Bertie, K. H. Michaelian, H. H. Eysel and D. Hager, *J. Chem. Phys.*, 1986, **85**, 4779–4789.
- 68 O. Birer and M. Havenith, *Annu. Rev. Phys. Chem.*, 2009, **60**, 263.
- 69 M. Ortlieb and M. Havenith, *J. Phys. Chem. A*, 2007, **111**, 7355.
- 70 G. Avila and E. Mátyus, *J. Chem. Phys.*, 2019, **150**, 174107.
- 71 G. Avila and E. Mátyus, *J. Chem. Phys.*, 2019, **151**, 154301.
- 72 G. Avila, D. Papp, G. Czako and E. Mátyus, *Phys. Chem. Chem. Phys.*, 2020, **22**, 2792–2802.
- 73 X.-G. Wang and T. Carrington, *J. Chem. Phys.*, 2020, **152**, 204311.
- 74 P. M. Felker, D. Lauvergnat, Y. Scribano, D. M. Benoit and Z. Bačić, *J. Chem. Phys.*, 2019, **151**, 124311.
- 75 Y. Liu, J. Li, P. M. Felker and Z. Bačić, *Phys. Chem. Chem. Phys.*, 2021, **23**, 7101–7114.
- 76 A. Chen, D. M. Benoit, Y. Scribano, A. Nauts and D. Lauvergnat, *J. Chem. Theory Comput.*, 2022, **18**, 4366–4372.
- 77 J. Behler, *J. Chem. Phys.*, 2011, **134**, 074106.
- 78 J. Behler, *Int. J. Quantum Chem.*, 2015, **115**, 1032–1050.
- 79 J. Behler, *Angew. Chem., Int. Ed.*, 2017, **56**, 12828–12840.
- 80 J. Behler, *Chem. Rev.*, 2021, **121**, 10037–10072.
- 81 E. B. Wilson, J. C. Decius and P. C. Cross, *Molecular Vibrations: The Theory of Infrared and Raman Vibrational Spectra*, Dover Publications, 1980.
- 82 D. A. Clabo, W. D. Allen, R. B. Remington, Y. Yamaguchi and H. F. Schaefer, *Chem. Phys.*, 1988, **123**, 187–239.
- 83 G. Amat, H. H. Nielsen and G. Tarrago, *Rotation-Vibration of Polyatomic Molecules*, Dekker, New York, 1971.
- 84 D. Papoušek and M. R. Aliev, *Molecular Vibrational-rotational Spectra: Theory and Applications of High Resolution Infrared, Microwave and Raman Spectroscopy of Polyatomic Molecules*, Elsevier Science Ltd, 1982.
- 85 M. R. Aliev and J. K. G. Watson, in *Molecular Spectroscopy: Modern Research*, ed. K. N. Rao, Academic Press, 1985, pp. 1–67.
- 86 M. Piccardo, J. Bloino and V. Barone, *Int. J. Quantum Chem.*, 2015, **115**, 948–982.
- 87 H. H. Nielsen, in *Handbuch der Physik*, ed. S. Flügge, Springer, Berlin, 1959, vol. 37, part I, pp. 173–313.
- 88 S. Califano, *Vibrational States*, John Wiley & Sons, Inc., London, 1976.
- 89 J. C. Light and T. Carrington Jr., *Discrete-Variable Representations and their Utilization*, John Wiley & Sons, Ltd, 2000, pp. 263–310.
- 90 A. Martín Santa Daría, G. Avila and E. Mátyus, *J. Mol. Spectrosc.*, 2022, **385**, 111617.
- 91 H.-J. Werner, P. J. Knowles, G. Knizia, F. R. Manby, M. Schütz, P. Celani, W. Györfy, D. Kats, T. Korona, R. Lindh, A. Mitrushenkov, G. Rauhut, K. R. Shamasundar, T. B. Adler, R. D. Amos, S. J. Bennie, A. Bernhardsson, A. Berning, D. L. Cooper, M. J. O. Deegan, A. J. Dobbyn, F. Eckert, E. Goll, C. Hampel, A. Hesselmann, G. Hetzer, T. Hrenar, G. Jansen, C. Köppl, S. J. R. Lee, Y. Liu, A. W. Lloyd, Q. Ma, R. A. Mata, A. J. May, S. J. McNicholas, W. Meyer, T. F. Miller III, M. E. Mura, A. Nicklass, D. P. O'Neill, P. Palmieri, D. Peng, K. Pflüger, R. Pitzer, M. Reiher, T. Shiozaki, H. Stoll, A. J. Stone, R. Tarroni, T. Thorsteinsson, M. Wang and M. Welborn, *MOLPRO, version, a package of ab initio programs*, see <https://www.molpro.net>.
- 92 T. B. Adler, G. Knizia and H.-J. Werner, *J. Chem. Phys.*, 2007, **127**, 221106.
- 93 G. Knizia, T. B. Adler and H.-J. Werner, *J. Chem. Phys.*, 2009, **130**, 054104.
- 94 H.-J. Werner, G. Knizia, T. B. Adler and O. Marchetti, *Z. Phys. Chem.*, 2010, **224**, 493–511.
- 95 R. A. Kendall, T. H. Dunning and R. J. Harrison, *J. Chem. Phys.*, 1992, **96**, 6796–6806.



- 96 T. Dunning, *J. Chem. Phys.*, 1989, **90**, 1007–1023.
- 97 F. Weigend, *Phys. Chem. Chem. Phys.*, 2002, **4**, 4285–4291.
- 98 B. Ziegler and G. Rauhut, *J. Chem. Phys.*, 2016, **144**, 114114.
- 99 R. Ramakrishnan and G. Rauhut, *J. Chem. Phys.*, 2015, **142**, 154118.
- 100 T. B. Blank and S. D. Brown, *J. Chemometrics*, 1994, **8**, 391–407.
- 101 A. Singraber, J. Behler and C. Dellago, *J. Chem. Theory Comput.*, 2019, **15**, 1827–1840.
- 102 S. Plimpton, *J. Comput. Phys.*, 1995, **117**, 1.
- 103 F. Ito and T. Nakanaga, *Chem. Phys.*, 2002, **277**, 163.
- 104 O. L. Polyansky, A. G. Császár, S. V. Shirin, N. F. Zobov, P. Barletta, J. Tennyson, D. W. Schwenke and P. J. Knowles, *Science*, 2003, **299**, 539–542.
- 105 D. Feller, K. A. Peterson and D. A. Dixon, *Mol. Phys.*, 2012, **110**, 2381–2399.
- 106 M. B. Gardner, B. R. Westbrook, R. C. Fortenberry and T. J. Lee, *Spectrochim. Acta, Part A*, 2021, **248**, 119184.
- 107 B. Schröder and P. Sebald, *J. Mol. Spectrosc.*, 2022, **386**, 111628.
- 108 K. A. E. Meyer and M. A. Suhm, *J. Chem. Phys.*, 2018, **149**, 104307.
- 109 C. Qu and J. M. Bowman, *Phys. Chem. Chem. Phys.*, 2019, **21**, 3397–3413.
- 110 M. Gantenberg, M. Halupka and W. Sander, *Chem. – Eur. J.*, 2000, **6**, 1865–1869.
- 111 D. S. Rasheeda, A. M. Santa Daría, B. Schröder, E. Mátyus and J. Behler, Replication Data for: High-dimensional neural network potentials for accurate vibrational frequencies: The formic acid dimer benchmark, *GRO.data*, V1, 2022, DOI: [10.25625/ZDGKYA](https://doi.org/10.25625/ZDGKYA).

