

Cite this: *RSC Med. Chem.*, 2022, **13**, 72

Metabolic labelling of cancer cells with glycodendrimers stimulate immune-mediated cytotoxicity†

David Goyard, Peremobowei Iyanu Diriwari and Nathalie Berthet *

The recruitment of antibody naturally present in human blood stream at the surface of cancer cells have been proved a promising immunotherapeutic strategy to fight cancer. Antibody recruiting molecules (ARMs) combining tumor and antibody binding modules have been developed for this purpose, however the formation of the interacting complex with both antibody and cell is difficult to optimize to stimulate immune-mediated cytotoxicity. To circumvent this limitation, we report herein a more direct approach combining cell metabolism of azido-sugar and bio-orthogonal click chemistry to conjugate at the cell glycolyx structurally well-defined glycodendrimers as antibody binding module (ABM). We demonstrate that this strategy allows not only the recruitment of natural antibody at the surface of isolated cells or solid tumor models but also activate a cytotoxic response with human serum as unique source of immune effectors.

Received 5th August 2021,
Accepted 3rd November 2021

DOI: 10.1039/d1md00262g

rsc.li/medchem

Introduction

The exploitation of endogenous antibodies naturally present in the blood stream of all individuals has recently emerged as an alternative immuno-strategy to fight cancer.^{1–3} To this aim, synthetic antibody recruiting molecules (ARMs) combining two binding modules, one for tumor cell (TBM) and the other for the antibody recruitment (ABM, typically dinitrophenol, α Gal or Rha) have been demonstrated to successfully trigger immune cytotoxicity against cancer cells by CDC or ADCC mechanisms.^{4–7} From the first generation of ARMAs to the more sophisticated antibody recruiting polymers (ARPs)⁸ or glycodendrimers (ARGs),⁸ significant advances have been made in the understanding of functional and structural requirements to improve immunological effects. If the multivalent presentation of ABM was clearly demonstrated as a key element to recruit endogenous antibodies,^{10,11} the major shortcoming of this approach concerns the TBM which has to ensure the binding of the cell surface without promoting internalisation to maintain the ABM exposure and accessibility at the cell surface. By doing so, the recruiting molecule can promote the formation of a reversible ternary interacting complex with antibodies and cancer cell. When suitable conditions are used to control this complex

equilibrium process,¹² the immune-mediated cytotoxic effect can be activated against the cancer cell line.^{13,14} To avoid the utilization of TBM and the problems associated with interacting systems involving three partners, the insertion of dinitrophenol or carbohydrate haptens in the cell membrane either with a lipid anchor or by covalent conjugation was proved to be a valuable alternative as a simplified antibody recruiting system.^{8,15–19}


In this regard, the use of well-known substrates for cell surface engineering, such as the tetraacetyl-*N*-azidoacetyl-mannosamine (Ac₄ManNAz), represents a powerful and reliable method to modify the glycolyx with unnatural recognition moieties. Once internalized, the intracellular metabolism of Ac₄ManAz into azido sialic acid indeed leads to azido group expression onto extracellular glycans that can be engaged in bio-orthogonal reaction to further decorate the cell membrane by copper-free strain-promoted azide-alkyne cycloaddition (SPAAC).^{20–25} Several groups used similar approach to conjugate diverse haptens on the membrane of different cells to promote immune-mediated cytotoxicity.^{15,16,26} In this study, we reasoned that the conjugation of clustered ABM at the surface of cancer cells would represent a robust antibody recruiting approach. We recently identified an ARG composed of a tetravalent cluster of cRGD as TBM and an hexadecavalent dendrimer of Rha as ABM to redirect natural antibodies against cancer cells expressing $\alpha_v\beta_3$ integrins.¹¹ This compound has been shown to stimulate immune-mediated cytotoxicity against this cell line. Among the variety of the tested compounds, we demonstrated that ARGs presenting the high rhamnose

Univ. Grenoble Alpes, CNRS, DCM UMR 5250, F-38000 Grenoble, France.

E-mail: nathalie.berthet@univ-grenoble-alpes.fr

† Electronic supplementary information (ESI) available. See DOI: 10.1039/d1md00262g






Scheme 1 Strategy combining glycometabolism and bio-orthogonal click chemistry to label cells with clustered rhamnose antigen and activate immune response against cancer cells.

valency with a precise spatial orientation were able to recruit antibodies from human serum at the surface of cancer cells and to subsequently activate an immune response.⁹ Herein, we exploited a metabolic labelling strategy to attach multivalent ABMs at the cell surface. To this aim, Ac₄ManNAz was first delivered to BT-549 for the surface tagging with azido group (Scheme 1, step 1). The coupling with dibenzocyclooctyne (DBCO)-bearing ABMs presenting one, four or sixteen rhamnoses to the cell membrane *via* copper-free chemistry was performed in the second step (Scheme 1, step 2). Finally, we evaluated the recruitment of natural antibodies present in human serum (Scheme 1, step 3) and the subsequent stimulation of the immune response (Scheme 1, step 4).

Results and discussion

ABMs were synthesized as previously reported.¹¹ Briefly, mono- **1** and tetraazido **2** cyclodecapeptides were functionalized *via* CuAAC using propargylated α -L-Rha to



Scheme 2 Synthesis of DBCO-ABM conjugates. Reagents and conditions: i) pentynoic acid NHS ester, DIPEA, DMF, 2 h, r.t.; ii) propargyl α -L-rhamnopyranoside or **2**, CuSO₄·5H₂O, THPTA, sodium ascorbate, DMF/PBS (p.H. 7.5) (1:1), 2h., r.t.; iii) **5**, PyBOP, DIPEA, DMF 4 h, r.t.

afford glycopeptides **3** and **4** respectively. Subsequent amide coupling between carboxylic acid-bearing DBCO derivative **5** and the free lysine side chain of compounds **3** and **4** yielded mono- and tetravalent compounds **DBCO-ABM1** and **DBCO-ABM4** respectively. Tetravalent glycocluster **4** was functionalized with pentynoic acid and subjected to another CuAAC reaction with scaffold **2**. The resulting hexadecaivalent glycodendrimer **6** was finally conjugated with DBCO derivative **5** to afford **DBCO-ABM16**. The synthesis route is depicted in Scheme 2. Final compounds were characterized by HRMS, ¹H NMR and analytical RP-HPLC before biological studies (see ESI†).

We first determined the optimal concentration of Ac₄ManNAz for metabolic labelling with azido groups of the targeted cells. For that, the triple negative breast cancer cell line BT-549 was cultivated with various concentrations of Ac₄ManNAz for 24 h and the conversion in azido sialic acid was followed by treatment with the commercial DBCO-PEG4-Fluor 545. The analyse by flow cytometry and confocal microscopy of the cell fluorescence enabled to determine optimal cell surface labelling for a concentration in Ac₄ManNAz of 50 μ M. Negligible fluorescence was observed for cells untreated with Ac₄ManNAz confirming the absence of non-specific DBCO-dye binding to the cells (Fig. S1, ESI†).

We next investigated the coupling of DBCO-conjugates (**DBCO-ABM16**, Fig. 1 and **DBCO-ABM1-4**, Fig. S2 ESI†) by SPAAC to azido-bearing cells. Extracellular Rha exposure was revealed using anti-rhamnose IgM naturally present in human serum (HS). Azido tagged BT-549 cells were treated with or without DBCO-ABM (concentrations from 0.1 to 10 μ M) or DBCO-PEG 5 used as negative control. The SPAAC coupling of the DBCO-conjugates was evaluated by flow cytometry and confocal microscopy after successive incubations with HS and AlexaFluor488- anti-human IgM secondary antibody. A dose-dependent effect with a maximum fluorescence intensity at 10 μ M was observed with all DBCO-ABMs (Fig. 1a, Fig. S2, ESI†) demonstrating the efficiency of the bio-orthogonal SPAAC coupling. The dose-response curve of the fluorescence of treated cells with various concentrations of **DBCO-ABM16** (Fig. S3, ESI†) clearly showed a plateau at 10 μ M. This result was confirmed by confocal microscopy of cells tagged with **ABM16** (Fig. 1b) which revealed an intense fluorescence homogenously distributed around the cell surface at 5 and 10 μ M of DBCO-conjugate, while no significant fluorescence was detected at lower concentration. Of note, no fluorescence was observed for azido-bearing cells or those treated with 10 μ M of DBCO-PEG 5, thus indicating the absence of non-specific binding of anti-Rha antibodies.

The influence of the Rha density on the recruitment of serum antibodies was next evaluated. Cells pre-incubated with or without Ac₄ManNAz were treated with 10 μ M of mono, tetra- and hexadecaivalent DBCO-ABMs. The anti-Rha IgM recruitment was revealed by immuno-fluorescence as described earlier and analysed by flow cytometry and






Fig. 1 DBCO-ABM16 coupling to azido groups exposed at the cell surface and anti-Rha recruitment. BT-549 cells were incubated with Ac_4ManNAz for azido labelling before treatment with DBCO-ABM16 or DBCO-PEG 5 (10–0.1 μM). Azido labelled cells untreated with DBCO-conjugate were used as control. SPAAC coupling of the DBCO-conjugates to the cell surface was revealed using anti-Rha IgM naturally present in HS and a fluorescent secondary antibody. Cell fluorescence was analysed by flow cytometry (a. data are presented as mean \pm SD for $n = 3$ measurements) and confocal microscopy (b. scale bar: 10 μm).

confocal microscopy. As shown by the histogram overlays reported in Fig. 2a, antibodies present in HS are efficiently recruited by all modified cell surface with DBCO-derivatives with a significantly higher efficiency for cells displaying the higher Rha density at their surface. Confocal microscopy confirmed this result since the highest fluorescence and homogenous dye labelling was observed with the **ABM16** (Fig. 2b).

Interestingly, mono- and tetravalent **ABM1** and **ABM4** covalently anchored to the cell membrane were able to recruit antibodies present in HS while ARGs previously reported with low Rha density failed.¹¹ These results demonstrate the interest of the metabolic labelling strategy to recruit antibody onto cell surface. In addition, flow cytometry analysis and confocal microscopy experiment with cells untreated with Ac_4ManNAz or DBCO conjugates revealed negligible fluorescent intensity, thus suggesting the binding specificity of anti-Rha antibodies (Fig. 2b).

Antibody recruitment being dependent on the presence and the persistence of ABMs on the cell surface, we next studied the stability of the cell labelling with ABM. Cells pre-treated with or without Ac_4ManNAz were cultured for 1–8 h with 10 μM of DBCO-ABM16 prior to being incubated with HS and the secondary antibody. The cell fluorescence was then analysed by flow cytometry. As shown in Fig. 3




Fig. 2 a. Flow cytometry analysis of anti-Rha recruitment by labelled BT-549 cells with mono-, tetra- and hexadecavalent-ABMs. Azido cells were treated with 10 μM of DBCO-conjugates before incubation with HS. The antibody recruitment was revealed with AF488 conjugated secondary antibody. For both cases, non-labelled cells were used as control. b. Corresponding confocal microscopy images (scale bar: 10 μm).

persistent cell labelling with **ABM16** is observed after 8 hours for cells pre-treated with Ac_4ManNAz and an increase of the fluorescent intensity is also measured with increasing incubation time. These results demonstrate the stability of cell labelling with ABM over the time. However a fluorescent




Fig. 3 Stability test of the ABM16 displayed on the cell surface. BT-549 cells metabolically labelled with Ac_4ManNAz or unlabelled cells (used as negative control) were incubated for different time (0, 1, 5 and 8 h) with DBCO-ABM16 (10 μM , final concentration) prior being treated successively with HS and the AF488 conjugated secondary antibody. The fluorescence of the cells was analysed by flow cytometry.



decorate the cell glycocalyx with antibody recruiting modules. We have clearly observed that such structural modification has the ability to redirect natural antibodies present in human serum, not only against BT-549 cell line but also against 3D-spheroid model of solid tumor. More importantly, the covalent grafting of chemically well-defined ABM with high Rha density at the cell surface was found to stimulate immune-mediated cytotoxicity of BT-549. By avoiding the use of tumor binding module which significantly complicates the utilization of bimodal antibody recruiting molecules, we believe that the present strategy combining delivery of azido sugar to cancer cell and bio-orthogonal ligation of synthetic antibody recruiting modules is highly promising in the context of cancer immunotherapy. The exploitation of this approach with azido monosaccharide embedded with targeting agent and cleavable linker is however required for a selective delivery in cancer cells. We are currently investigating strategies based on the use of receptor-mediated endocytosis or enzyme activatable azido sugar by leveraging dysregulated mechanisms leading to enzyme-overexpression in cancer cells. The results will be reported in due course.

Experimental

Materials

Tetraacetylated *N*-azidoacetyl-*D*-mannosamine (Ac₄ManNAz), Alexa-Fluor 488-coupled anti-human IgM were purchased from Fisher Scientific. All other chemical reagents were purchased from Aldrich (Saint-Quentin Fallavier, France) or Acros (Noisy-Le-Grand, France) and were used without further purification. All protected amino acids and Fmoc-Gly-Sasrin@resin was obtained from Advanced ChemTech Europe (Brussels, Belgium), BachemBiochimie SARL (Voisins-Les-Bretonneux, France) and FranceBiochem S.A. (Meudon, France). Human serum (HS) were obtained from a healthy human male donor (Etablissement Français du Sang, EFS Grenoble). Cell Cytotoxicity assay kit abcam 112 118 was from Abcam. All experiments were performed in accordance with the Guidelines of the “Etablissement Français du Sang (EFS) Auvergne-Rhône Alpes 2017-2958”, and Experiments were approved by the ethics committee at the EFS AURA 21-033. Informed consents were obtained from human participants of this study. For peptides and glycopeptides, analytical RP-HPLC was performed on a Waters alliance 2695 separation module, equipped with a Waters 2489 UV/visible detector. Analyses were carried out at 1.23 mL min⁻¹ (Waters XBridge Shield RP18 3.5 μM, C18, 100 × 4.6 mm) with UV monitoring at 214 nm and 250 nm using a linear A–B gradient (buffer A: 0.09% CF₃CO₂H in water; buffer B: 0.09%CF₃CO₂H in 90% acetonitrile). Preparative HPLC was performed on waters equipment consisting of a Waters 2545 controller and a Waters 2487 dual absorbance detector. Purifications were carried out at 22 mL min⁻¹ (VP 250 × 21 mm nucleosil100-7 C18) with UV monitoring at 214 nm and 250 nm using a linear A–B gradient. ¹H NMR spectra were recorded on

BrukerAvanceIII 500 MHz spectrometers and chemical shifts (δ) were reported in parts per million (ppm). Spectra were referenced to the residual proton solvent peaks relative to the signal of D₂O (4.79 ppm). All mass spectrometry characterizations were performed at mass spectrometry facility, PCN-ICMG, Grenoble. ESI⁺ high resolution mass spectra of peptides and glycopeptides were measured on a LTQ Orbitrap XL spectrometer from Thermo Scientific.

Synthetic procedures and characterization

General procedure for the amide coupling of DBCO derivative 5. A solution of **5** (2 eq.), PyBOP (2 eq.) and DIPEA (4 eq.) in dry DMF (0.5 mL) was stirred at room temperature for 1 hour then a solution of rhamnosylated glycocluster or glycodendrimer (1 eq.) in dry DMF (0.5 mL) was added. The mixture was stirred at r.t. for 2 hours, diluted in H₂O/TFA (99.9: 0.1) and purified by semi preparative RP-HPLC. Fractions containing the product were collected and lyophilized to obtain the desired compound as a white fluffy solid.

Compound DBCO-ARM1. Prepared according to the general procedure from **3** (8.5 mg, 7.6 μmol). The crude mixture was purified (5–100% B in 20 min) to afford the title compound (9.6 mg, 5.7 μmol, 75%). HRMS (ESI⁺) *m/z*: calcd for C₇₉H₁₁₆N₁₆O₂₅ [M + 2H]²⁺: 844.4143, found 844.4176; RP-HPLC: *R*_t = 5.79 min (C18, λ = 214 nm, 5–100% B in 15 min).

Compound DBCO-ARM4. Prepared according to the general procedure from **4** (11.8 mg, 6.1 μmol). The crude mixture was purified (5–100% B in 20 min) to afford the title compound (12.3 mg, 4.9 μmol, 81%). HRMS (ESI⁺) *m/z* calcd for C₁₁₅H₁₇₃N₂₅O₃₇ [M + 2H]²⁺: 1248.1207, found 1248.1255; RP-HPLC: *R*_t = 6.72 min (C18, λ = 214 nm, 5–100% B in 15 min).

Compound DBCO-ARM16. Prepared according to the general procedure from **6** (20.0 mg, 2.2 μmol). The crude mixture was purified (0–80% B in 20 min) to afford the title compound (15.1 mg, 1.5 μmol, 71%). HRMS (ESI⁺) *m/z* calcd for C₄₃₁H₆₆₉N₁₁₇O₁₄₁ [M + 6H]⁶⁺: 1623.3124, found 1623.3195; RP-HPLC: *R*_t = 7.59 min (C18, λ = 214 nm, 0–80% B in 15 min).

Instrumentation

Flow cytometry analysis was performed on a BD LSR FORTRESSA Flow cytometer (laser excitation at λ = 488 nm, emission bypass filter at 525/50).

Confocal microscopy images of 2D and 3D-cultures were taken with a TCS SP8 CSU Leica (HC PL APO CS2, 40×/1.30 OIL, laser excitation at λ = 488 nm and fluorescence emission collected between λ = 495 and 545 nm). Images were processed using ImageJ software.

The solution-absorbance in microplates was read on a POLARstar Omega plate reader (BMG labtech).

Cell culture method

BT-549 cells were purchased from ATCC and were cultured in RPMI 1640 medium supplemented with 10% fetal bovine serum, 100 μg mL⁻¹ streptomycin and 100 U mL⁻¹ penicillin (all from Sigma-Aldrich). Cells were maintained in an



Acknowledgements

This work was supported by CNRS, Université Grenoble Alpes, ICMG FR 2607, the French ANR project Glyco@Alps (ANR-15-IDEX-02), Labex ARCANE and CBH-EUR-GS (ANR-17-EURE-0003). We acknowledge the European Research Council Consolidator Grant “LEGO” (647938). We thank Cecile Cottet (LBFA, Grenoble) for confocal microscopy experiments, the data analysis and for giving access to flow cytometer.

References

- 1 K. Sasaki, M. Harada, Y. Miyashita, H. Tagawa, A. Kishimura, T. Mori and Y. Katayama, *Chem. Sci.*, 2020, **11**, 3208.
- 2 B. Lake, N. Serniuck, E. Kapcan, A. Wang and A. F. Rullo, *ACS Chem. Biol.*, 2020, **15**, 1089.
- 3 H. Hong, C. Li, L. Gong, J. Wang, D. Li, J. Shi, Z. Zhou, Z. Huang and Z. Wu, *Chem. Sci.*, 2021, **12**, 4623.
- 4 P. J. McEnaney, C. G. Parker, A. X. Zhang and D. A. Spiegel, *ACS Chem. Biol.*, 2012, **7**, 1139.
- 5 P. J. McEnaney, C. G. Parker and A. X. Zhang, in *Annu. Rep. Med. Chem.*, 2017, p. 481.
- 6 R. T. C. Sheridan, J. Hudon, J. A. Hank, P. M. Sonde and L. L. Kiessling, *ChemBioChem*, 2014, **15**, 1393–1398.
- 7 S. Achilli, N. Berthet and O. Renaudet, *RSC Chem. Biol.*, 2021, **2**, 713.
- 8 A. Uvyn, R. De Coen, M. Gruijs, C. W. Tuk, J. De Vrieze, M. van Egmond and B. G. De Geest, *Angew. Chem.*, 2019, **131**, 13122.
- 9 B. Todaro, S. Achilli, B. Liet, E. Laigre, C. Tiertant, D. Goyard, N. Berthet and O. Renaudet, *Biomater. Sci.*, 2021, **9**, 4076.
- 10 A. Uvyn and B. G. De Geest, *ChemBioChem*, 2020, **21**, 3036.
- 11 B. Liet, E. Laigre, D. Goyard, B. Todaro, C. Tiertant, D. Boturyn, N. Berthet and O. Renaudet, *Chem. – Eur. J.*, 2019, **25**, 15508.
- 12 E. F. Douglass, C. J. Miller, G. Sparer, H. Shapiro and D. A. Spiegel, *J. Am. Chem. Soc.*, 2013, **135**, 6092.
- 13 C. B. Calson, P. Mowery, R. M. Owen, E. C. Dykhuizen and L. L. Kiessling, *ACS Chem. Biol.*, 2007, **2**, 119.
- 14 A. F. Rullo, K. J. Fitzgerald, V. Muthusamy, M. Liu, C. Yuan, M. Huang, M. Kim, A. E. Cho and D. A. Spiegel, *Angew. Chem., Int. Ed.*, 2016, **55**, 3642.
- 15 A. Uvyn, R. De Coen, O. De Wever, K. Deswarte, B. N. Lambrecht and B. G. De Geest, *Chem. Commun.*, 2019, **55**, 10952.
- 16 S. Li, B. Yu, J. Wang, Y. Zheng, H. Zhang, M. J. Walker, Z. Yuan, H. Zhu, J. Zhang, P. G. Wang and B. Wang, *ACS Chem. Biol.*, 2018, **13**, 1686.
- 17 C. E. Jakobsche, C. G. Parker, R. N. Tao, M. D. Kolesnikova, E. F. Douglass and D. A. Spiegel, *ACS Chem. Biol.*, 2013, **8**, 2404.
- 18 B. Lin, X. Wu, H. Zhao, Y. Tian, J. Han, J. Liu and S. Han, *Chem. Sci.*, 2016, **7**, 3737.
- 19 X. Li, X. Rao, L. Cai, X. Liu, H. Wang, W. Wu, C. Zhu, M. Chen, P. G. Wang and W. Yi, *ACS Chem. Biol.*, 2016, **11**, 1205.
- 20 N. J. Agard, J. A. Prescher and C. R. Bertozzi, *J. Am. Chem. Soc.*, 2004, **126**, 15046.
- 21 E. M. Sletten and C. R. Bertozzi, *Acc. Chem. Res.*, 2011, **44**, 666–676.
- 22 J. C. Jewett and C. R. Bertozzi, *Chem. Soc. Rev.*, 2010, **39**, 1272–1279.
- 23 Y. Takayama, K. Kusamori and M. Nishikawa, *Molecules*, 2019, **24**, 172.
- 24 A. Mongis, F. Piller and V. Piller, *Bioconjugate Chem.*, 2017, **28**, 1151.
- 25 H. Wang and D. J. Mooney, *Nat. Chem.*, 2020, **12**, 1102.
- 26 S. H. Park, H. Jung, H. Lee, T. M. Kim, J. W. Cho, W. D. Jang, J. Y. Hyun and I. Shin, *Chem. Commun.*, 2020, **56**, 10650.
- 27 X. Li, X. Xu, X. Rao, Y. Tian and W. Yi, *Carbohydr. Res.*, 2017, **452**, 25.
- 28 C. J. Lovitt, T. B. Shelper and V. M. Avery, *Biology*, 2014, **3**, 345.
- 29 D. Anton, H. Burckel, E. Josset and G. Noel, *Int. J. Mol. Sci.*, 2015, **16**, 5517.

