

Cite this: *RSC Med. Chem.*, 2022, **13**, 72

Metabolic labelling of cancer cells with glycodendrimers stimulate immune-mediated cytotoxicity†

David Goyard, Peremobowei Iyanu Diriwari and Nathalie Berthet *

The recruitment of antibody naturally present in human blood stream at the surface of cancer cells have been proved a promising immunotherapeutic strategy to fight cancer. Antibody recruiting molecules (ARMs) combining tumor and antibody binding modules have been developed for this purpose, however the formation of the interacting complex with both antibody and cell is difficult to optimize to stimulate immune-mediated cytotoxicity. To circumvent this limitation, we report herein a more direct approach combining cell metabolism of azido-sugar and bio-orthogonal click chemistry to conjugate at the cell glycolyx structurally well-defined glycodendrimers as antibody binding module (ABM). We demonstrate that this strategy allows not only the recruitment of natural antibody at the surface of isolated cells or solid tumor models but also activate a cytotoxic response with human serum as unique source of immune effectors.

Received 5th August 2021,
Accepted 3rd November 2021

DOI: 10.1039/d1md00262g

rsc.li/medchem

Introduction

The exploitation of endogenous antibodies naturally present in the blood stream of all individuals has recently emerged as an alternative immuno-strategy to fight cancer.^{1–3} To this aim, synthetic antibody recruiting molecules (ARMs) combining two binding modules, one for tumor cell (TBM) and the other for the antibody recruitment (ABM, typically dinitrophenol, α Gal or Rha) have been demonstrated to successfully trigger immune cytotoxicity against cancer cells by CDC or ADCC mechanisms.^{4–7} From the first generation of ARM to the more sophisticated antibody recruiting polymers (ARPs)⁸ or glycodendrimers (ARGs),⁸ significant advances have been made in the understanding of functional and structural requirements to improve immunological effects. If the multivalent presentation of ABM was clearly demonstrated as a key element to recruit endogenous antibodies,^{10,11} the major shortcoming of this approach concerns the TBM which has to ensure the binding of the cell surface without promoting internalisation to maintain the ABM exposure and accessibility at the cell surface. By doing so, the recruiting molecule can promote the formation of a reversible ternary interacting complex with antibodies and cancer cell. When suitable conditions are used to control this complex

equilibrium process,¹² the immune-mediated cytotoxic effect can be activated against the cancer cell line.^{13,14} To avoid the utilization of TBM and the problems associated with interacting systems involving three partners, the insertion of dinitrophenol or carbohydrate haptens in the cell membrane either with a lipid anchor or by covalent conjugation was proved to be a valuable alternative as a simplified antibody recruiting system.^{8,15–19}

In this regard, the use of well-known substrates for cell surface engineering, such as the tetraacetyl-*N*-azidoacetyl-mannosamine (Ac₄ManNAz), represents a powerful and reliable method to modify the glycolyx with unnatural recognition moieties. Once internalized, the intracellular metabolism of Ac₄ManAz into azido sialic acid indeed leads to azido group expression onto extracellular glycans that can be engaged in bio-orthogonal reaction to further decorate the cell membrane by copper-free strain-promoted azide-alkyne cycloaddition (SPAAC).^{20–25} Several groups used similar approach to conjugate diverse haptens on the membrane of different cells to promote immune-mediated cytotoxicity.^{15,16,26} In this study, we reasoned that the conjugation of clustered ABM at the surface of cancer cells would represent a robust antibody recruiting approach. We recently identified an ARG composed of a tetravalent cluster of cRGD as TBM and an hexadecavalent dendrimer of Rha as ABM to redirect natural antibodies against cancer cells expressing $\alpha_v\beta_3$ integrins.¹¹ This compound has been shown to stimulate immune-mediated cytotoxicity against this cell line. Among the variety of the tested compounds, we demonstrated that ARGs presenting the high rhamnose

Univ. Grenoble Alpes, CNRS, DCM UMR 5250, F-38000 Grenoble, France.

E-mail: nathalie.berthet@univ-grenoble-alpes.fr

† Electronic supplementary information (ESI) available. See DOI: 10.1039/d1md00262g






Scheme 1 Strategy combining glycometabolism and bio-orthogonal click chemistry to label cells with clustered rhamnose antigen and activate immune response against cancer cells.

valency with a precise spatial orientation were able to recruit antibodies from human serum at the surface of cancer cells and to subsequently activate an immune response.⁹ Herein, we exploited a metabolic labelling strategy to attach multivalent ABMs at the cell surface. To this aim, Ac₄ManNAz was first delivered to BT-549 for the surface tagging with azido group (Scheme 1, step 1). The coupling with dibenzocyclooctyne (DBCO)-bearing ABMs presenting one, four or sixteen rhamnoses to the cell membrane *via* copper-free chemistry was performed in the second step (Scheme 1, step 2). Finally, we evaluated the recruitment of natural antibodies present in human serum (Scheme 1, step 3) and the subsequent stimulation of the immune response (Scheme 1, step 4).

Results and discussion

ABMs were synthesized as previously reported.¹¹ Briefly, mono- **1** and tetraazido **2** cyclodecapeptides were functionalized *via* CuAAC using propargylated α -L-Rha to



Scheme 2 Synthesis of DBCO-ABM conjugates. Reagents and conditions: i) pentynoic acid NHS ester, DIPEA, DMF, 2 h, r.t.; ii) propargyl α -L-rhamnopyranoside or **2**, CuSo₄·5H₂O, THPTA, sodium ascorbate, DMF/PBS (p.H. 7.5) (1:1), 2h., r.t.; iii) **5**, PyBOP, DIPEA, DMF 4 h, r.t.

afford glycopeptides **3** and **4** respectively. Subsequent amide coupling between carboxylic acid-bearing DBCO derivative **5** and the free lysine side chain of compounds **3** and **4** yielded mono- and tetravalent compounds **DBCO-ABM1** and **DBCO-ABM4** respectively. Tetravalent glycocluster **4** was functionalized with pentynoic acid and subjected to another CuAAC reaction with scaffold **2**. The resulting hexadecaivalent glycodendrimer **6** was finally conjugated with DBCO derivative **5** to afford **DBCO-ABM16**. The synthesis route is depicted in Scheme 2. Final compounds were characterized by HRMS, ¹H NMR and analytical RP-HPLC before biological studies (see ESI†).

We first determined the optimal concentration of Ac₄ManNAz for metabolic labelling with azido groups of the targeted cells. For that, the triple negative breast cancer cell line BT-549 was cultivated with various concentrations of Ac₄ManNAz for 24 h and the conversion in azido sialic acid was followed by treatment with the commercial DBCO-PEG4-Fluor 545. The analyse by flow cytometry and confocal microscopy of the cell fluorescence enabled to determine optimal cell surface labelling for a concentration in Ac₄ManNAz of 50 μ M. Negligible fluorescence was observed for cells untreated with Ac₄ManNAz confirming the absence of non-specific DBCO-dye binding to the cells (Fig. S1, ESI†).

We next investigated the coupling of DBCO-conjugates (**DBCO-ABM16**, Fig. 1 and **DBCO-ABM1-4**, Fig. S2 ESI†) by SPAAC to azido-bearing cells. Extracellular Rha exposure was revealed using anti-rhamnose IgM naturally present in human serum (HS). Azido tagged BT-549 cells were treated with or without DBCO-ABM (concentrations from 0.1 to 10 μ M) or DBCO-PEG 5 used as negative control. The SPAAC coupling of the DBCO-conjugates was evaluated by flow cytometry and confocal microscopy after successive incubations with HS and AlexaFluor488- anti-human IgM secondary antibody. A dose-dependent effect with a maximum fluorescence intensity at 10 μ M was observed with all DBCO-ABMs (Fig. 1a, Fig. S2, ESI†) demonstrating the efficiency of the bio-orthogonal SPAAC coupling. The dose-response curve of the fluorescence of treated cells with various concentrations of **DBCO-ABM16** (Fig. S3, ESI†) clearly showed a plateau at 10 μ M. This result was confirmed by confocal microscopy of cells tagged with **ABM16** (Fig. 1b) which revealed an intense fluorescence homogenously distributed around the cell surface at 5 and 10 μ M of DBCO-conjugate, while no significant fluorescence was detected at lower concentration. Of note, no fluorescence was observed for azido-bearing cells or those treated with 10 μ M of DBCO-PEG 5, thus indicating the absence of non-specific binding of anti-Rha antibodies.

The influence of the Rha density on the recruitment of serum antibodies was next evaluated. Cells pre-incubated with or without Ac₄ManNAz were treated with 10 μ M of mono, tetra- and hexadecaivalent DBCO-ABMs. The anti-Rha IgM recruitment was revealed by immuno-fluorescence as described earlier and analysed by flow cytometry and






Fig. 1 DBCO-ABM16 coupling to azido groups exposed at the cell surface and anti-Rha recruitment. BT-549 cells were incubated with Ac₄ManNAz for azido labelling before treatment with DBCO-ABM16 or DBCO-PEG 5 (10–0.1 μM). Azido labelled cells untreated with DBCO-conjugate were used as control. SPAAC coupling of the DBCO-conjugates to the cell surface was revealed using anti-Rha IgM naturally present in HS and a fluorescent secondary antibody. Cell fluorescence was analysed by flow cytometry (a. data are presented as mean ± SD for *n* = 3 measurements) and confocal microscopy (b. scale bar: 10 μm).

confocal microscopy. As shown by the histogram overlays reported in Fig. 2a, antibodies present in HS are efficiently recruited by all modified cell surface with DBCO-derivatives with a significantly higher efficiency for cells displaying the higher Rha density at their surface. Confocal microscopy confirmed this result since the highest fluorescence and homogenous dye labelling was observed with the **ABM16** (Fig. 2b).

Interestingly, mono- and tetravalent **ABM1** and **ABM4** covalently anchored to the cell membrane were able to recruit antibodies present in HS while ARGs previously reported with low Rha density failed.¹¹ These results demonstrate the interest of the metabolic labelling strategy to recruit antibody onto cell surface. In addition, flow cytometry analysis and confocal microscopy experiment with cells untreated with Ac₄ManNAz or DBCO conjugates revealed negligible fluorescent intensity, thus suggesting the binding specificity of anti-Rha antibodies (Fig. 2b).

Antibody recruitment being dependent on the presence and the persistence of ABMs on the cell surface, we next studied the stability of the cell labelling with ABM. Cells pre-treated with or without Ac₄ManNAz were cultured for 1–8 h with 10 μM of DBCO-ABM16 prior to being incubated with HS and the secondary antibody. The cell fluorescence was then analysed by flow cytometry. As shown in Fig. 3




Fig. 2 a. Flow cytometry analysis of anti-Rha recruitment by labelled BT-549 cells with mono-, tetra- and hexadecavalent-ABMs. Azido cells were treated with 10 μM of DBCO-conjugates before incubation with HS. The antibody recruitment was revealed with AF488 conjugated secondary antibody. For both cases, non-labelled cells were used as control. b. Corresponding confocal microscopy images (scale bar: 10 μm).

persistent cell labelling with **ABM16** is observed after 8 hours for cells pre-treated with Ac₄ManNAz and an increase of the fluorescent intensity is also measured with increasing incubation time. These results demonstrate the stability of cell labelling with ABM over the time. However a fluorescent




Fig. 3 Stability test of the ABM16 displayed on the cell surface. BT-549 cells metabolically labelled with Ac₄ManNAz or unlabelled cells (used as negative control) were incubated for different time (0, 1, 5 and 8 h) with DBCO-ABM16 (10 μM, final concentration) prior being treated successively with HS and the AF488 conjugated secondary antibody. The fluorescence of the cells was analysed by flow cytometry.






Fig. 4 *In vitro* human blood compatibility of DBCO-ABMs (1, 4 and 16). hRBCs were incubated with DBCO-ABMs at 100 μ M (final concentration). Red blood cells treatment with Triton 1% or phosphate-buffered saline (PBS) was used as positive and negative controls respectively. a. Morphology of treated hRBCs was observed under a light microscopy. b. Hemolysis ratio (%) of hRBCs after incubation with DBCO-ABM conjugates.

signal similar to the autofluorescence of the cells was observed for non azido-labelled cells incubated with the DNCO conjugate even after 8 hours showing one more time the specific ABM labelling for cells pre-treated with $Ac_4ManNAz$.

Since DBCO-ABM conjugates were attached onto the cells for antibodies recruitment in the human serum, we also evaluated their blood compatibility in an *in vitro* assay. This was performed by determining the effect of DBCO-ABMs (1, 4 and 16) on human red blood cells (hRBC). Like saline treatment, incubating hRBC with all DBCO-ABMs in saline did not cause detectable changes on morphology of hRBC under light microscopy even at a concentration 10 times higher than which was showing cytotoxic effect (100 μ M) (Fig. 4a). The hemolysis ratios for each DBCO-ABM treatments was calculated. Less than 2% of hemolysis was obtained for all conjugates, indicating a good blood compatibility of the DBCO conjugates (Fig. 4b).

We next evaluated the ability of the recruited antibodies to promote immune response leading to cancer cell destruction. Azido tagged cells treated with or without DBCO-ABMs were incubated with human serum as unique source of antibodies and protein complement and the cell viability assay which measures the complement-dependent cytotoxicity (CDC) was examined as reported previously. The cytotoxicity effect was found to increase with the ABM valency (Fig. 5), the lowest effect being observed for cells modified with the monovalent ABM1 (20%) and the highest with the hexadecaivalent ABM16 (30%). This result is in good correlation with the antibody recruitment observed by flow cytometry. By comparison with previous studies reported by other groups,^{16,27} the conjugation of ABM16 promote similar cytotoxic effect at much lower concentration, thus suggesting the importance of a dense presentation in hapten to the cell surface.

In order to evaluate the ability of ABMs to recruit antibody in solid tumor context, we cultured 3D-spheroid which are commonly used as *in vitro* model of such tumor.^{28,29} Spheroids grown 4 days were cultured with or without $Ac_4ManNAz$ (untreated spheroids were used as control) prior to be treated with DBCO-ABM16 and HS (as 2D culture). As described before,



Fig. 5 Cytotoxicity of BT-549 cells induced by ABMs displayed to the cell surface following the metabolic labelling process in the presence of HS (average value of three independent experiments). Azido cells untreated with DBCO-ABMs exposed to HS were used as control. Data are presented as mean \pm SD ($n = 3$).




Fig. 6 Confocal microscopy images (maximum intensity projections) of BT-549 human breast cancer spheroids, cultured (a.) or without (b.) $Ac_4ManNAz$ and treated with 10 μ M of DBCO-ABM16, HS and AF488-secondary antibodies. Scale bar is 50 μ m. (c.) Average AF488 fluorescence intensity of BT-549 spheroids following the same treatment as described in (a.) and (b.).

the recruitment of endogenous anti-Rha was revealed with AF488-conjugated anti-human secondary antibody.

Confocal microscopy images (Fig. 6) showed high fluorescence for spheroids cultured with $Ac_4ManNAz$ and treated with DBCO-ABM16 compared to the control spheroids. The measure of the ratio of the Fluorescence intensity reported to the spheroid surface revealed a fluorescence intensity 2 times higher for cells incubated with $Ac_4ManNAz$ compared to the control spheroids. These results clearly indicate the effectiveness of the spheroid labelling and the accessibility of Rha antigens for IgM recruitment at the cell surface to induce a targeted immune response against BT-549 tumor cells.

Conclusions

In summary, we have demonstrated the robustness of cell glycometabolism and bio-orthogonal click chemistry to



incubator at 37 °C in a 5% CO₂ humidified atmosphere and allowed for downstream experiment until 90% confluency.

Determination of the optimal concentration of Ac₄ManN₃ for *in vitro* metabolic cell labelling

BT-549 cells were seeded at a density of about 150 000 cells per well in 6-well plates and cultured for 24 h in the medium described in the cell culture method section. Various dilutions of Ac₄ManNAz were prepared and added directly to the culture medium to get final concentrations from 1 μM to 50 μM and the culture was continued for 48 h. The culture medium was then removed and cells were washed 3 times with PBS prior to be incubated with 50 μM DBCO-PEG₄-Fluor 545 diluted in HBSS for 1 h at 37 °C (50 mM stock solution in DMSO was prepared). Control cells without azido sugar received the same amount of DMSO. Cells were washed again 3 times with PBS then were detached by treatment with 0.5 mL per well of trypsin-EDTA for 5 min at 37 °C. Afterwards, the harvest cells were centrifuged, washed with 1 mL of PBS and resuspended in 500 μL of PBS for analysis by flow cytometry.

In vitro coupling of DBCO-ARMs to azido labeled cells and antibody recruitment analysis

BT-549 cells (150 000 cells per well in 6-well plates) were cultured for 24 h prior to be incubated with 50 μM Ac₄ManNAz as described above. After culture medium removing, cells were washed 3 times with PBS then the copper-free click chemistry of azido-labeled cells with various concentrations of each DBCO-ABMs in HBSS (0 μM, 0.1 μM, 0.5 μM, 5 μM and 10 μM) was performed for 1 h at 37 °C. Cells were then washed twice with PBS and were detached with trypsin-EDTA as described previously. Washed cells were then incubated with 50% human serum in PBS-1% BSA for 1 h at 37 °C. Following 1 wash with PBS-1% BSA, cells were incubated with 1:400 AF488-conjugated goat anti-human antibody for 1 h at 37 °C. Finally, the cells were washed and resuspended in 500 μL PBS and analyzed for FITC intensity by flow cytometry and confocal microscopy.

Complement-mediated cytotoxicity

BT-549 cells were plated onto the 96-well plate at the concentration of 20 000 cells per well and left to adhere for 24 h prior to be incubated with 50 μM Ac₄ManNAz for 48 h at 37 °C, 5% CO₂. Plates were then washed twice with PBS and click chemistry with 10 μM DBCO-ABMs in HBSS was performed for 1 h at 37 °C. Solution was then removed and cells were incubated with 100 μL of HS 50% in HBSS for 2 h at 37 °C. Subsequently, 20% (v/v) abcam reagent was added and the plates were further incubated for 3 h at 37 °C, 5% CO₂. The plates were then read at OD₅₇₀ to OD₆₀₅ on microplate reader. The percentage of cell cytotoxicity was calculated using the following formula:

$$\% \text{cytotoxicity} = [1 - (R_{\text{sample}} - R_0)/(R_{\text{ctrl}} - R_0)] \times 100$$

R_{sample} was the absorbance ratio of OD₅₇₀/OD₆₀₅ for the cells labelled with ARMs and incubated with HS.

R_{ctrl} was the absorbance ratio of OD₅₇₀/OD₆₀₅ for unlabelled cells incubated with HS.

R_0 was the averaged background (non-cell control) absorbance ratio OD₅₇₀/OD₆₀₅.

BT-549 tumor spheroids

Spheroids were generated by plating BT-549 cells at 5000 cells per well into ultralow adherence-96-well plates (Corning). Spheroids grew in complete medium as in 2D-cultures in the final volume of 200 μL. After 72 h, spheroids were treated with 50 μM Ac₄ManNAz and 10 μM DBCO-ABMs in the same manner as in 2D-cultures. Spheroids were imaged using confocal microscope (TCS SP8 CSU, Leica, laser excitation at $\lambda = 448$ nm and fluorescence emission collected between $\lambda = 495$ and 545 nm) equipped with a 40× objective and analysis was performed with the ImageJ software.

Human red blood cell hemolysis assay

All experiments were performed in accordance with the Guidelines of the “Etablissement Français du Sang (EFS) Auvergne-Rhône Alpes 2017-2958”, and Experiments were approved by the ethics committee at the EFS AURA 21-033. Informed consents were obtained from human participants of this study. Human red blood (hRBC) were obtained by centrifugation of 2 mL of whole blood collected in heparin tubes, from healthy donors (EFS Grenoble) at 800 g for 10 min at room temperature. After removal of the plasma, hRBC were further purified by washing three times with PBS and resuspended in PBS at a final concentration of 2%. The hRBC suspensions (180 μL) were incubated with 20 μL of DBCO-ABM 1, 4 or 16 (1 mM in PBS) while parallel assays were performed using PBS as negative control and a 1% (w/v) solution of Triton X-100 as positive control. After incubation at 37 °C in 5% CO₂ for 2 h, each sample was centrifuged for 10 min at 800 g and 100 μL of the supernatant was transferred to a 96-well microplate. Absorbance at 540 nm was measured using a microplate reader and the results were expressed as a percentage of hemoglobin released relative to the positive control. All treatments were performed in triplicates and the hemolysis ratio (%) was calculated using equation:

$$\begin{aligned} \text{Hemolysis ratio}(\%) \\ = \frac{A(\text{sample}) - A(\text{negative control})}{A(\text{positive control}) - A(\text{negative control})} \times 100 \end{aligned}$$

The integrity of the precipitated hRBC was checked by morphological observations under light microscopy.

Conflicts of interest

There are no conflicts to declare.



Acknowledgements

This work was supported by CNRS, Université Grenoble Alpes, ICMG FR 2607, the French ANR project Glyco@Alps (ANR-15-IDEX-02), Labex ARCANE and CBH-EUR-GS (ANR-17-EURE-0003). We acknowledge the European Research Council Consolidator Grant “LEGO” (647938). We thank Cecile Cottet (LBFA, Grenoble) for confocal microscopy experiments, the data analysis and for giving access to flow cytometer.

References

- 1 K. Sasaki, M. Harada, Y. Miyashita, H. Tagawa, A. Kishimura, T. Mori and Y. Katayama, *Chem. Sci.*, 2020, **11**, 3208.
- 2 B. Lake, N. Serniuck, E. Kapcan, A. Wang and A. F. Rullo, *ACS Chem. Biol.*, 2020, **15**, 1089.
- 3 H. Hong, C. Li, L. Gong, J. Wang, D. Li, J. Shi, Z. Zhou, Z. Huang and Z. Wu, *Chem. Sci.*, 2021, **12**, 4623.
- 4 P. J. McEnaney, C. G. Parker, A. X. Zhang and D. A. Spiegel, *ACS Chem. Biol.*, 2012, **7**, 1139.
- 5 P. J. McEnaney, C. G. Parker and A. X. Zhang, in *Annu. Rep. Med. Chem.*, 2017, p. 481.
- 6 R. T. C. Sheridan, J. Hudon, J. A. Hank, P. M. Sonde and L. L. Kiessling, *ChemBioChem*, 2014, **15**, 1393–1398.
- 7 S. Achilli, N. Berthet and O. Renaudet, *RSC Chem. Biol.*, 2021, **2**, 713.
- 8 A. Uvyn, R. De Coen, M. Gruijs, C. W. Tuk, J. De Vrieze, M. van Egmond and B. G. De Geest, *Angew. Chem.*, 2019, **131**, 13122.
- 9 B. Todaro, S. Achilli, B. Liet, E. Laigre, C. Tiertant, D. Goyard, N. Berthet and O. Renaudet, *Biomater. Sci.*, 2021, **9**, 4076.
- 10 A. Uvyn and B. G. De Geest, *ChemBioChem*, 2020, **21**, 3036.
- 11 B. Liet, E. Laigre, D. Goyard, B. Todaro, C. Tiertant, D. Boturyn, N. Berthet and O. Renaudet, *Chem. – Eur. J.*, 2019, **25**, 15508.
- 12 E. F. Douglass, C. J. Miller, G. Sparer, H. Shapiro and D. A. Spiegel, *J. Am. Chem. Soc.*, 2013, **135**, 6092.
- 13 C. B. Calson, P. Mowery, R. M. Owen, E. C. Dykhuizen and L. L. Kiessling, *ACS Chem. Biol.*, 2007, **2**, 119.
- 14 A. F. Rullo, K. J. Fitzgerald, V. Muthusamy, M. Liu, C. Yuan, M. Huang, M. Kim, A. E. Cho and D. A. Spiegel, *Angew. Chem., Int. Ed.*, 2016, **55**, 3642.
- 15 A. Uvyn, R. De Coen, O. De Wever, K. Deswarte, B. N. Lambrecht and B. G. De Geest, *Chem. Commun.*, 2019, **55**, 10952.
- 16 S. Li, B. Yu, J. Wang, Y. Zheng, H. Zhang, M. J. Walker, Z. Yuan, H. Zhu, J. Zhang, P. G. Wang and B. Wang, *ACS Chem. Biol.*, 2018, **13**, 1686.
- 17 C. E. Jakobsche, C. G. Parker, R. N. Tao, M. D. Kolesnikova, E. F. Douglass and D. A. Spiegel, *ACS Chem. Biol.*, 2013, **8**, 2404.
- 18 B. Lin, X. Wu, H. Zhao, Y. Tian, J. Han, J. Liu and S. Han, *Chem. Sci.*, 2016, **7**, 3737.
- 19 X. Li, X. Rao, L. Cai, X. Liu, H. Wang, W. Wu, C. Zhu, M. Chen, P. G. Wang and W. Yi, *ACS Chem. Biol.*, 2016, **11**, 1205.
- 20 N. J. Agard, J. A. Prescher and C. R. Bertozzi, *J. Am. Chem. Soc.*, 2004, **126**, 15046.
- 21 E. M. Sletten and C. R. Bertozzi, *Acc. Chem. Res.*, 2011, **44**, 666–676.
- 22 J. C. Jewett and C. R. Bertozzi, *Chem. Soc. Rev.*, 2010, **39**, 1272–1279.
- 23 Y. Takayama, K. Kusamori and M. Nishikawa, *Molecules*, 2019, **24**, 172.
- 24 A. Mongis, F. Piller and V. Piller, *Bioconjugate Chem.*, 2017, **28**, 1151.
- 25 H. Wang and D. J. Mooney, *Nat. Chem.*, 2020, **12**, 1102.
- 26 S. H. Park, H. Jung, H. Lee, T. M. Kim, J. W. Cho, W. D. Jang, J. Y. Hyun and I. Shin, *Chem. Commun.*, 2020, **56**, 10650.
- 27 X. Li, X. Xu, X. Rao, Y. Tian and W. Yi, *Carbohydr. Res.*, 2017, **452**, 25.
- 28 C. J. Lovitt, T. B. Shelper and V. M. Avery, *Biology*, 2014, **3**, 345.
- 29 D. Anton, H. Burckel, E. Josset and G. Noel, *Int. J. Mol. Sci.*, 2015, **16**, 5517.

