



Showcasing research from Professor Yun-Hong Zhang's laboratory, School of Chemistry and Chemical Engineering, Beijing Institute of Technology, Beijing, China.

Directly measuring Fe(III)-catalyzed SO₂ oxidation rate in single optically levitated droplets

We directly measure the kinetics of Fe(III)-catalyzed SO₂ oxidation at the surface of a single droplet trapped by optical tweezers. The Fe(III)-catalyzed SO₂ reactive uptake coefficient is 2 or 3 orders higher than that of the auto-oxidation without TMI. The reaction rate decreases by up to a decade when the Fe(III)/S(vi) coexisting time increases, owing to a complexation between Fe(III) and S(vi).

As featured in:




See Pai Liu, Yun-Hong Zhang *et al.*, *Environ. Sci.: Atmos.*, 2023, 3, 298.



Cite this: *Environ. Sci.: Atmos.*, 2023, 3, 298

Directly measuring Fe(III)-catalyzed SO₂ oxidation rate in single optically levitated droplets†

Xue Cao, Zhe Chen, Yu-Xin Liu, Xin-Bo Jing, Lin-Fang Li, Pai Liu* and Yun-Hong Zhang *

Sulfate aerosols are produced in China's winter haze at an unresolved rapid rate. Such fast kinetics may arise from a heterogeneous SO₂ conversion in urban aerosols, which differs significantly from the aqueous S(IV) oxidation in bulk solutions. Given the uniqueness of aerosols as a multiphase reactor, it is preferable to measure the heterogeneous SO₂ conversion rate *in situ*, ideally in levitated microdroplets. Here, we directly measure the Fe(III)-catalyzed SO₂ conversion in single microdroplets trapped and levitated with a gradient-force aerosol optical tweezer. The sulfate formation rate was inferred from the droplet's growth rate driven by the heterogeneous reaction. Our results show that the Fe(III)-catalyzed SO₂ conversion in aerosols is 2 to 3 decades faster than that determined in bulk solutions. The SO₂ reactive uptake coefficient at pH ~5.0 and 298 K is on the order of 10⁻⁴ to 10⁻³. The reaction rate scales with droplet surface area, indicating that the major reaction location is the air–water interface. This interfacial reaction is further corroborated by a positive kinetic salt effect, a trait of the interaction between ions and the neutral molecules, such as Fe(III) ions and SO₂ molecules. The reaction rate decreases by up to a decade, as the Fe(III)/S(VI) coexisting time increases, possibly owing to a complexation between Fe(III) and S(VI) ions.

Received 22nd September 2022
Accepted 8th December 2022

DOI: 10.1039/d2ea00125j

rsc.li/esatmospheres

Environmental significance

Sulfate is a major secondary inorganic component of atmospheric particulate matter, and its production is closely related to urban haze formation. Sulfate can be produced rapidly *via* a heterogeneous process, in which SO₂ molecules are directly converted at the aerosol surface, catalyzed by the transition metal ions. In this study, we directly measure the kinetics of Fe(III)-catalyzed SO₂ oxidation at the aerosol surface. The kinetic effects of SO₂ concentration, total Fe(III) molar fraction, ambient humidity, and hours of S(VI)/Fe(III) coexistence are investigated. We emphasize that the SO₂ conversion rate in incipient aerosols and in aged ones can differ by nearly a decade, which should be incorporated in the air-quality models.

1 Introduction

The rapid formation of sulfate aerosols is the chemical driver of the severe winter haze events in China, exerting tremendous negative impacts on social economy and public health.^{1–4} During the past decade, SO₂ emission in China's urban area has been effectively controlled: The annual average SO₂ concentration decreased from 9.1 ppb in 2013 to 1.7 ppb in 2018.⁵ Yet, during the same period, sulfate aerosol concentration in polluted air only decreased moderately, from 19.2 to 12.1 μg m⁻³.⁶ And severe air pollution events still occurred in winter seasons.^{4,7,8} To further contain secondary sulfate pollutants, we need to thoroughly understand the mechanisms and kinetics of SO₂ conversion in polluted air.

Per the traditional view, SO₂ in the gas phase is oxidized by OH radicals⁹ and stable Criegee intermediates.¹⁰ Aqueous reaction pathways include the oxidation of S(IV) by O₃, O₂ (catalyzed by transition metal ions, TMI), and H₂O₂ dissolved in cloud and fog droplets.^{11,12} But these mechanisms cannot explain the rapid sulfate PM_{2.5} formation in the north China plain (NCP). The sulfate PM_{2.5} concentration simulated with air quality models¹³ were 3–4 times lower than that measured in the atmosphere. This gap has motivated many research efforts to discover the hitherto unidentified sulfate sources in polluted environments.^{11,14–22} New aqueous SO₂ conversion mechanisms have been identified, and the kinetics of existing mechanisms have been updated.

The significance of aqueous SO₂ conversion, however, was questioned in recent kinetic studies.^{1,23} The concern partly arises from a reaction space constraint.^{1,23} The volume of aerosol water in polluted air is several decades smaller than that of clouds, and such small aerosol water volume precludes large-scale sulfate formation *via* aqueous routes.¹ Instead, interfacial reactions – a direct conversion of SO₂ molecules at the

Institute of Chemical Physics, School of Chemistry and Chemical Engineering, Beijing Institute of Technology, 100081, Beijing, China. E-mail: pliu@bit.edu.cn; yhz@bit.edu.cn

† Electronic supplementary information (ESI) available. See DOI: <https://doi.org/10.1039/d2ea00125j>



undergoing reaction. These parameters are tabulated in Table S1.† Note that the concentration of reactant gases was estimated with their dilution ratio. This technique is the same as that in our previous publication.²⁶

2.4 Measuring reaction rate

When RH is fixed, the sulfate solute molarity, $[S_{(VI)}]$, is also a constant, owing to an equilibrium of water partition. At such an equilibrium condition, an increasing sulfate mass inside the droplet will cause an increasing droplet volume. In other words, converting SO_2 to sulfate inside a droplet will induce a hygroscopic growth of the droplet. Such reaction-induced droplet growth facilitates us to measure the reaction rate precisely, if we can measure the droplet size precisely, for example:

$$R_{S(VI)} = \frac{dS(VI)}{dt} = \frac{[S(VI)] \times 4\pi r_0^2 dr}{dt} \quad (3)$$

Here, $R_{S(VI)}$ is the reaction rate in mol s^{-1} ; r is droplet radius, subscript zero indicates the initial state, and t is time. The droplet size during the reaction, $r(t)$, was inferred from the whispering gallery mode wavelength λ_{WGM} per the Mie scattering calculation²⁸ established by Preston and Reid (refer to ESI Fig. S1† for typical droplet growth data). It is also implicitly assumed in eqn (3) that the curvature effect of the droplet surface is insignificant. Such an assumption requires that the increase in droplet radius during the reaction is negligible when compared with the initial radius. In the experiments, we guaranteed that the increase in r is always less than 5% of r_0 . The reactant SO_2 concentration ranged between 25 and 10^3 ppb. Each kinetic measurement was conducted at fixed RH conditions (60, 70, or 80%). The O_2 in the background air served as the oxidizer.

3 Results and discussion

3.1 Influence of SO_2 concentration

We first discuss the kinetic effect of SO_2 concentration. In these experiments, the droplet pH was maintained at ~ 5.0 ; the ambient RH, at $\sim 60\%$; $Fe(III)$ molar fraction, 0.01%. SO_2 concentration varied between 25 and 10^3 ppb. Fig. 1 plots the reaction rate $R_{S(VI)}$ as a function of SO_2 concentration. Note that the unit of $R_{S(VI)}$ here is mole of sulfate produced in a unit time per a single droplet. The trends in Fig. 1 show that the sulfate formation rate correlates positively with SO_2 concentration. Quantitatively, we show that a linear fitting (red line in Fig. 1) describes the data best, indicating that the reaction rate is first order in SO_2 . This observation agrees with our knowledge of the $Fe(III)$ -catalyzed SO_2 conversion.³⁰

3.2 The location of heterogeneous reaction

The size dependence of reaction rate provides us insight on the location of heterogeneous reactions.²⁶ For example, if SO_2 are first dissolved in aerosol water and the $S(IV)$ are oxidized homogeneously therein, the reaction rate will be proportional to droplet volume, $R_{S(VI)} \propto r_0^3$; if the SO_2 are directly converted at the droplet surface, the reaction rate will be proportional to



Fig. 1 The formation rate of sulfate inside a single levitated droplet, $R_{S(VI)}$, as a function of SO_2 concentration. The droplets were seeded with AS and IS mixture. The molar fraction of $Fe(III)$ is 0.01%. Ambient RH was about 60%. The droplet pH was buffered at ca. 5.0 by dissolving an 8 ppm NH_3 into droplet water. The solid line represents a linear fitting on the dataset. Error bars arose from the uncertainty in determining dr/dt from the $r(t)$ data. The values of error stemmed from the 95% confidence interval of dr/dt values.

droplet surface area, $R_{S(VI)} \propto r_0^2$. We evaluated these scaling laws by using the droplet growth rate, dr/dt , which was measured directly during reactions. Eqn (3) indicates that $dr/dt \propto R_{S(VI)} \times r_0^{-2}$ holds for short-range droplet growth driven by sulfate formation. Then, the kinetic scaling laws can be reduced to $dr/dt \propto r_0$ for aqueous reaction, and to $dr/dt = \text{constant}$ for the interfacial reaction.

Fig. 2 shows the dr/dt of droplets within two size bins: the smaller droplets around 4.5 μm radii, and the larger ones around 7.0 μm radii. The pH of these droplets was maintained at ~ 5.0 ; the SO_2 concentration, at 25 ppb; $Fe(III)$ molar fraction, at 0.01%; RH, at $\sim 60\%$. Inside these two size bins, we repeatedly



Fig. 2 Droplet growth rate during reaction, dr/dt , as a function of droplet initial radius, r_0 . The ambient RH conditions were 60%; the SO_2 concentration, at 25 ppb; the droplet pH, at ~ 5.0 . Gray circles represent the original dr/dt data; Black circles represent the mean values of dr/dt and r_0 for each cluster of data. Error bars represent on standard deviation.



measured dr/dt more than ten times (see the gray circles). Then we calculated the mean values for dr/dt and r_0 , as well as their one standard deviation (black circles and error bars). Despite the fluctuation, the mean dr/dt values for droplets inside each size bin are clearly independent of the corresponding mean r_0 values. (The mean dr/dt was $0.0131 \pm 0.002 \text{ nm s}^{-1}$ and $0.0128 \pm 0.002 \text{ nm s}^{-1}$, respectively, when the mean r_0 was $4.525 \pm 0.308 \text{ }\mu\text{m}$ and $6.936 \pm 0.207 \text{ }\mu\text{m}$.) With this constant dr/dt , we infer that the reaction occurs primarily at the droplet surface. Were aqueous reactions the case, dr/dt would increase linearly with r_0 , and the dr/dt data at $\sim 7 \text{ }\mu\text{m}$ would instead be clustered around 0.02 nm s^{-1} .

In the following, we normalize the sulfate formation rate with SO_2 concentration and droplet surface area, per the expression of reactive uptake coefficient Γ_{SO_2} :²⁶

$$\Gamma_{\text{SO}_2} = \frac{dS(\text{VI})/dt}{Z} \quad (4.1)$$

and

$$Z = \frac{1}{4} \sqrt{\frac{8RT}{\pi M_{\text{SO}_2}}} A [\text{SO}_{2(\text{g})}]. \quad (4.2)$$

Here, Z is the collision efficiency between SO_2 molecules and droplet (unit, mol s^{-1}); the parameters R , T , M_{SO_2} , A , and $[\text{SO}_{2(\text{g})}]$ are gas constant ($\text{J mol}^{-1} \text{K}^{-1}$), temperature (K^{-1}), SO_2 molecular weight (kg mol^{-1}), and droplet initial surface area (m^2), and SO_2 gas concentration ($\text{mol m}_{\text{air}}^{-3}$), respectively.

3.3 Influence of metal concentration

Not all $\text{Fe}(\text{III})$ in aerosol water are soluble. The soluble $\text{Fe}(\text{III})$ ions include FeOH^{2+} , $\text{Fe}(\text{OH})_2^+$, and $\text{Fe}_2(\text{OH})_2^{4+}$; their concentration in dilute solution might be estimated per the precipitation equilibrium relationship. But it remains unclear whether such a relationship still provides accurate predictions in aerosol water at the non-ideal, high-ionic-strength condition. Here, we empirically measured the reaction rate at changing concentrations of total $\text{Fe}(\text{III})$, including both soluble and insoluble $\text{Fe}(\text{III})$. In these experiments, droplet pH was maintained at ~ 5 ; ambient RH, at $\sim 60\%$. The molar fraction of total $\text{Fe}(\text{III})$ in droplet solute varied between 0.0001% and 0.1% (correspondingly, the total $\text{Fe}(\text{III})$ concentration varied between $\sim 6.2 \text{ }\mu\text{M}$ and $\sim 6.2 \text{ mM}$.)

The results are shown in Fig. 3. Although the total $\text{Fe}(\text{III})$ varied across three orders of magnitude, the reaction rate at $\text{pH} \approx 5$ remains a constant. This constant reaction rate indicates that most of the $\text{Fe}(\text{III})$ in droplets was insoluble. Such an observation of excessive total $\text{Fe}(\text{III})$ agrees with traditional kinetic studies. For example, Martin *et al.*³¹ found that, at $\text{pH} \approx 5$, the first-order rate coefficient of $\text{Fe}(\text{III})$ -catalyzed oxidation of aqueous $\text{S}(\text{IV})$ is also independent of $\text{Fe}(\text{III})$ concentration. In polluted air, the $\text{Fe}(\text{III})$ concentration is about $18 \text{ ng m}_{\text{air}}^{-3}$, and the aerosol water content is about $300 \text{ }\mu\text{g m}_{\text{air}}^{-3}$.²⁰ These conditions suggest that the total $\text{Fe}(\text{III})$ molarity in aerosol water is $\sim 1 \text{ mM}$ (dotted line in Fig. 3), far beyond the solubility limit. Therefore, one may assume that the reaction kinetics at fixed pH conditions is zeroth order in the total airborne $\text{Fe}(\text{III})$.




Fig. 3 Reaction rate is independent of the total $\text{Fe}(\text{III})$ concentration in microdroplets. The gray circle represents the reactive uptake coefficient Γ_{SO_2} data obtained from each experiment run. The black circle represents the mean Γ_{SO_2} values of the data in each bin, and the error bar represents one standard deviation. The vertical dotted line indicates the typical value of total $\text{Fe}(\text{III})$ molarity in aerosols at haze conditions. The droplet pH was maintained at ~ 5 ; the ambient RH, at $\sim 60\%$.

3.4 Influences of ambient humidity

We next investigate the influence of ambient humidity on reaction rate. In these experiments, the SO_2 was maintained at 25 ppb; $\text{Fe}(\text{III})$ molar fraction, at 0.01%. Ambient RH varied among 60, 70, and 80%. The results are plotted in Fig. 4. As RH decreases from 80 to 60%, the mean reaction rate increases by a factor of ~ 2 . The faster reaction at lower RH conditions may arise from the kinetic salt effect. When RH decreases, droplet water becomes more concentrated, and droplet ionic strength I



Fig. 4 The reaction rate decreases as ambient relative humidity (RH) increases. The gray circle represents the reactive uptake coefficient Γ_{SO_2} data obtained from each experiment run. The black circle represents the mean Γ_{SO_2} values of the data in each bin, and the error bar represents one standard deviation. The droplet pH was maintained at ~ 5 ; $\text{Fe}(\text{III})$ molar fraction, 0.01%.



increases. (For example, ESI Fig. S2† shows that the I of ammonium sulfate droplet increases from ~ 17 to 35 mol, as the ambient RH decreases from 80 to 60%.) Reactions are accelerated at the high I condition if the rate-limiting step of the reaction involves an interaction between charge neutral molecules and ions^{1,26,32} here, the neutral SO_2 molecules and the Fe(III) ions. This positive kinetic salt effect also indicates that SO_2 is directly converted at the droplet surface, without an *a priori* dissolution and dissociation.¹ For example, SO_2 and O_2 may react with Fe(III) at droplet surface, producing Fe(II) and SO_5^- radicals. Instead, if the SO_2 were first dissolved and dissociated into S(IV) ions, then the rate-limiting step would involve an interaction between S(IV) ions and Fe(III) ions. Such ion-ion reaction would be inhibited when I is large.¹⁷ One may refer to Angle *et al.*'s recent kinetic study¹⁷ for such a slower TMI-catalyzed S(IV) oxidation at higher I conditions.

The faster reaction rate at lower RH conditions is unlikely due to the enrichment effect (*i.e.*, droplet solvent evaporates, the aqueous reactant is concentrated, and the reaction is accelerated.) We find that this explanation is not very compelling, because when the SO_2 molecules are directly converted at the aerosol surface, their availability is unaffected by the enrichment of the aqueous phase. The Fe(III) catalysts do exist in the aqueous phase. But, when the total Fe(III) concentration is much greater than the solubility limit, the actual concentration of soluble Fe(III) is unlikely to be affected by the enrichment of droplet water either.

3.5 Influence of $\text{Fe(III)}/\text{S(VI)}$ coexisting time

The aqueous Fe(III) -catalyzed S(IV) conversion is self-inhibiting because the reaction product S(VI) is an effective complexation agent to the Fe(III) catalysts.^{17,24} Martin and Hill²⁴ reported that the Fe(III) -catalyzed reaction in bulk solution decelerates significantly when the S(VI) concentration increases from a μM to a mM level. Here, we investigate the influence of $\text{Fe(III)}/\text{S(VI)}$ coexisting time on the heterogeneous reaction rate. In this part of experiment, we aged the AS/IS mixture solutions for 12, 36, or 48 hours before nebulizing them into aerosols. The reaction rate in the droplets made from aged solutions was then compared with that in the droplets made from the fresh solution. The SO_2 was maintained at 25 ppb; Fe(III) molar fraction, at 0.01%. Ambient RH, at 60%. Fig. 5 plots the Γ_{SO_2} as a function of the $\text{Fe(III)}/\text{S(VI)}$ coexisting time. The reaction rate decelerates by almost a decade, as the $\text{Fe(III)}/\text{S(VI)}$ coexisting time increased from 0 to 48 hours. This observation indicates that the $\text{Fe(III)}/\text{S(VI)}$ complexation is also a significant factor affecting the heterogeneous SO_2 conversion. The Fe(III) -catalyzed reaction may be a decade slower in real-world sulfate aerosols that have gone through an elongated aging process.

3.6 Comparing with other reaction mechanisms

Table 1 compares the kinetic data obtained in the present study with those reported in ref. 1, 26 and 31. Per our present study, the Fe(III) -catalyzed SO_2 oxidation occurs primarily at aerosol surface. At room temperature and a pH ~ 5 condition, such a reaction exhibits Γ_{SO_2} on the order of 10^{-4} . This reaction rate



Fig. 5 Reaction rate decreases as the $\text{Fe(III)}/\text{S(VI)}$ coexisting time increases. The coexisting time refers to how long the AS/IS mixture solution was aged before being nebulized into aerosols. The gray circle represents the reactive uptake coefficient Γ_{SO_2} data obtained from each experiment run. The black circle represents the mean Γ_{SO_2} values of the data in each bin, and the error bar represents one standard deviation. The droplet pH was maintained at ~ 5 ; the ambient RH, at $\sim 60\%$; Fe(III) molar fraction, at 0.01%.

is 2–3 decades faster than what the solution chemistry predicts. For example, Martin *et al.*³¹ reported that the Fe(III) -catalyzed S(IV) oxidation in bulk solution at pH 5.0 has a first-order rate coefficient of $\sim 10^{-3} \text{ s}^{-1}$. This rate coefficient corresponds to a Γ_{SO_2} of $\sim 1.7 \times 10^{-6}$ for the heterogeneous reaction inside the droplets with a $\sim 10 \mu\text{m}$ radius (refer to Table 1 footnotes for details of calculation).

The much faster reaction rate observed in our present study is not an experimental artefact. Previously, Chen *et al.*²⁶ used the AOT to measure the uncatalyzed SO_2 oxidation rate in levitated droplets. In that study, we conducted the experiment by using the same AOT apparatus and the same ammonium sulfate chemicals, but we did not intentionally add any Fe(III) to the droplets. At such conditions, the reaction at pH ~ 5.0 exhibited a Γ_{SO_2} on the orders of 10^{-7} to 10^{-6} . This comparison indicates that the 2-to-3-decade faster reaction observed here is due to our intentional addition of Fe(III) to droplet water. In other words, if such faster kinetics were a systematic artefact of the AOT apparatus, Chen *et al.* would also observe a Γ_{SO_2} on the order of 10^{-4} for the uncatalyzed reactions.

One should also note that the Γ_{SO_2} on the order of 10^{-4} to 10^{-3} is even greater than what is needed to explain the missing sulfate in haze events. Such an observation indicates that the Fe(III) -catalyzed heterogeneous SO_2 oxidation in the atmosphere is not as fast as that in the laboratory aerosols, particularly those freshly-made ones. As we discussed earlier, the catalytic capability of Fe(III) is impaired if Fe(III) and S(VI) coexists for a long time. On the other hand, recent studies proposed that a strong electric field exists at the air–water interface, triggering a spontaneous conversion of some OH^- ions at droplet surface into OH radicals and free electron.^{33,34} Such unique environment at air–water interface may also alter the kinetics of redox reactions.



Table 1 Kinetics of heterogeneous SO₂ oxidation at room temperature

References	Catalysts	Reaction location	Conditions	Uptake coef.
The present study	Fe(III)	Aerosol surface	pH 5; 60% RH; incipient aerosols pH 5; 80% RH; incipient aerosols pH 5; 60% RH; aged aerosols	$9.9(\pm 1.7) \times 10^{-4}$ $5.3(\pm 1.2) \times 10^{-4}$ $1.4(\pm 0.2) \times 10^{-4}$
Martin <i>et al.</i> (1991)	Fe(III)	Bulk solution ^a	pH 5; $r = 10 \mu\text{m}$ (assumed)	$\sim 1.7 \times 10^{-6}$
Chen <i>et al.</i> (2022)	Uncatalyzed	Aerosol surface ^b	pH 3.5–5.5; 60% RH pH 3.5–5.5; 80% RH	$\sim 4.9 \times 10^{-7}$ $\sim 8.1 \times 10^{-8}$
		Aerosol interior	pH 5; 60% RH; $r = 10 \mu\text{m}$ (assumed) pH 5; 80% RH; $r = 10 \mu\text{m}$ (assumed)	$\sim 6.6 \times 10^{-6}$ $\sim 5.5 \times 10^{-7}$
Wang <i>et al.</i> (2021)	Mn(II)	Aerosol surface ^c	pH 4–5; >80% RH; 0.81% Mn(II) pH 4–5; >80% RH; 5.54% Mn(II)	$\sim 1.6 \times 10^{-4}$ $\sim 7.1 \times 10^{-4}$

^a Martin *et al.*³¹ measured the rate of Fe(III)-catalyzed S(IV) oxidation in bulk solutions. They found that the first order rate coefficient k of this reaction is *ca.* 10^{-3} s^{-1} at pH 5.0 condition. One may estimate the corresponding uptake coefficient Γ_{SO_2} by using the eqn (4.1) in this paper. Specifically, rate per droplet is $\text{dS(IV)}/\text{dt} = (4/3)\pi r^3 k[\text{S(IV)}]$; the sulfite molarity $[\text{S(IV)}] = H_{\text{SO}_2} P_{\text{SO}_2} (1 + K_{1s}[\text{H}^+]^{-1} + K_{1s}K_{2s}[\text{H}^+]^{-2})$, where H_{SO_2} and P_{SO_2} respectively denote the Henry's law constant of SO₂ and SO₂ gas partial pressure. The K_{1s} , and K_{2s} respectively denote the first and second dissociation equilibrium constants of sulfite hydrate. Droplet radius r was assumed to be 10 μm matching our experimental condition. The P_{SO_2} is trivial and can take any arbitrary value. ^b Chen *et al.*²⁶ measured the uncatalyzed SO₂ oxidation rate in levitated aerosol microdroplets. The Γ_{SO_2} was calculated with eqn (5) and (8) in their paper. ^c Wang *et al.*¹ measured the Mn(II)-catalyzed SO₂ oxidation rate by using a smoke chamber. The Γ_{SO_2} was calculated with the dataset shown in the Fig. 1b in their paper.

4 Conclusion

In this work, we directly measured Fe(III)-catalyzed SO₂ conversion inside optically levitated microdroplets. We show that aerosol optical tweezers – coupled with cavity-enhanced Raman spectroscopy – can be utilized to determine the rate of heterogeneous reactions between SO₂ gas and microdroplets. Our findings show that the rate of Fe(III)-catalyzed SO₂ oxidation is first order in SO₂ concentration. During the heterogeneous reaction, the droplet growth rate is independent of the droplet initial radius, indicating that the sulfate production rate (per droplet) is proportional to the droplet surface area. Such a surface scaling law in turn suggests that the heterogeneous reaction occurs primarily at the air–water interface. The reaction rate is faster at lower RH conditions, possibly owing to a positive kinetic salt effect for ion–molecule reactions. The reaction rate at pH ~ 5 remains constant despite the total Fe(III) molar fraction varies across three decades, indicating that most Fe(III) is insoluble. This excessive Fe(III) condition also holds valid for real-world aerosols in polluted air. The reaction rate decelerates nearly one decade as the Fe(III)/S(VI) coexists for two days, indicating that the contribution of the Fe(III)-catalyzed reaction to sulfate may significantly diminish in aged aerosols.

Author contributions

P. L. and Y. Z. designed the research; X. C. and Z. C. performed the experiments; P. L. performed the theoretical calculations; P. L. and X. C. wrote the first draft of the manuscript; and all of the authors analyzed the data and contributed to revising the manuscript.

Conflicts of interest

There are no conflicts to declare.

Acknowledgements

We are grateful to Prof. Dr Jonathan P. Reid at the University of Bristol for helping us set up the aerosol optical tweezer and for sharing the spectra inversion algorithm. This study was supported by the National Natural Science Foundation of China (No. 42205113, 42127806, 91544223, 91844301, and 21806169), and Beijing Institute of Technology Research Fund Program for Young Scholars.

References

- W. Wang, M. Liu, Y. Song and M. Ge, Sulfate formation is dominated by manganese-catalyzed oxidation of SO₂ on aerosol surfaces during haze events, *Nat. Commun.*, 2021, **12**, 1–10.
- J. P. D. Abbatt and O. Mohler, Solid ammonium sulfate aerosols as ice nuclei: A pathway for cirrus cloud formation, *Science*, 2006, **313**, 1770–1773.
- C. A. Pope and G. D. Thurston, Lung cancer, cardiopulmonary mortality, and long-term exposure to fine particulate air pollution, *JAMA*, 2002, **287**, 1132–1141.
- H. He and J. Hao, Mineral dust and NO_x promote the conversion of SO₂ to sulfate in heavy pollution days, *Sci. Rep.*, 2014, **4**, 4172.
- F. Yan, W. Chen and X. Wang, Stabilization for the secondary species contribution to PM_{2.5} in the Pearl River Delta (PRD) over the past decade, China: A meta-analysis, *Atmos. Environ.*, 2020, **242**, 117817.
- D. Shang, J. Peng, S. Guo, Z. Wu and M. Hu, Secondary aerosol formation in winter haze over the Beijing-Tianjin-Hebei Region, China, *Front. Environ. Sci. Eng.*, 2021, **15**, 1–13.
- X. Li, B. Zhao, W. Zhou and J. Jiang, Responses of gaseous sulfuric acid and particulate sulfate to reduced SO₂ concentration: a perspective from long-term measurements in Beijing, *Sci. Total Environ.*, 2020, **721**, 137700.



