

Cite this: *Chem. Sci.*, 2023, 14, 13367

## Recent advances in combining photo- and N-heterocyclic carbene catalysis

Xiaochen Wang, Senhui Wu, Rongxin Yang, Hongjian Song, Yuxiu Liu and Qingmin Wang \*

N-Heterocyclic carbenes (NHCs) are unique Lewis basic catalysts that mediate various organic transformations by means of polarity reversal. Although the scope of research on two-electron reactions mediated by NHC catalysts has been expanding, the types of these reactions are limited by the inability of NHCs to engage  $sp^3$ -electrophiles. However, the revival of photocatalysis has accelerated the development of free-radical chemistry, and combining photoredox catalysis and NHC catalysis to achieve NHC-mediated radical reactions under mild conditions could overcome the above-mentioned limitation. This review summarizes recent advances in combining photoredox and NHC catalysis, focusing on elucidation and exploration of mechanisms, with the aim of identifying challenges and opportunities to develop more types of catalytic models.

Received 28th June 2023  
Accepted 27th October 2023

DOI: 10.1039/d3sc03274d

rsc.li/chemical-science

### 1. Introduction

Organic small-molecule catalysts can simulate the activity of enzymes and are inexpensive, readily available, mild, and less toxic than metal catalysts. In particular, N-heterocyclic carbenes (NHCs) have been successfully used as organocatalysts over the past few decades because of their versatility and unique structure and been a research hotspot in the field of catalysis.<sup>1</sup> On the basis of their general properties and chemical applications, NHC-bound intermediates can be broadly divided into two types: electron-rich and electron-deficient (Scheme 1a and b,

respectively).<sup>1e</sup> The traditional mode of NHC catalysis typically involves direct participation of such intermediates in reactions to form chemical bonds through a two-electron process; however, the scope with respect to NHC-derived operators is restricted by their inability to engage  $sp^3$  electrophiles.<sup>2</sup> Therefore, the discovery of new reaction modes for NHC catalysts, such as single-electron transfer (SET) radical reactions, will bring about new reaction modes to NHC catalysis and also rejuvenate its vitality and vigour.

Recent advances in single-electron reactions involving photoredox catalysis<sup>3</sup> have enabled bond connections that were previously not possible by means of traditional methods, and these advances have been used for organic synthesis.<sup>4</sup> The revival of photocatalysis research has accelerated the development of free-radical chemistry, and visible-light catalysis has been combined with NHC catalysis to achieve NHC-mediated

State Key Laboratory of Elemento-Organic Chemistry, Research Institute of Elemento-Organic Chemistry, Frontiers Science Center for New Organic Matter, College of Chemistry, Nankai University, Tianjin 300071, People's Republic of China. E-mail: wangqm@nankai.edu.cn



Xiaochen Wang

Xiaochen Wang obtained her B.Sc. degree (2017) under the supervision of Prof. Zhiyu Dou at the Changchun University of Science and Technology. She then pursued an M.Sc. degree in 2018 under the supervision of Prof. Qingmin Wang at Nankai University and started her PhD studies in 2020. Her research focuses on photoredox catalyzed radical chemistry.



Senhui Wu

Senhui Wu obtained his B.Sc. degree (2022) from Jiangxi Agricultural University under the supervision of Prof. Ni Guorong. He is currently studying for a master's degree at Nankai University under the guidance of Professor Qingmin Wang. His main research focuses on synthesis and derivation of isoxazolin.



radical reactions under mild conditions. In recent years, the cocatalysis model has been in a state of explosion, and there are some reviews in related fields.<sup>5</sup> Although there have some big shoes to fill, our review focuses on the summary of the newly developed catalytic mechanism, especially in the last four years. At present, there are two main modes of NHC/photoredox dual catalysis: (1) single-electron oxidation of classical electron-rich NHC intermediates to generate ketyl radicals under photocatalytic conditions and (2) single-electron reduction of novel NHC-derived electron-deficient radical intermediates under photocatalytic conditions. Among them, the electron-rich or electron-deficient intermediates can be used as oxidizing or reducing agents after direct photoexcitation (Scheme 1c, top and bottom, respectively). And the generated ketyl radicals will engage in radical/radical cross-couplings; although not employing light activation directly, the pioneering work of Ohmiya and co-workers based on a NHC-derived ketyl radical using NHPI esters as oxidative radical precursors under thermal conditions will also be discussed.

This review is divided into two main sections. The first is dedicated to production of ketyl radicals by single-electron oxidation (*via* Breslow intermediates), and the second focuses

on generation of ketyl radicals by single-electron reduction (*via* acyl azolium intermediates).

## 2. Ketyl radicals generated *via* Breslow intermediates

Although not employing light activation directly, as pioneering work, in 2019, Ohmiya reported the first example of radical cross-coupling reactions based on a NHC-catalyzed decarboxylative coupling reaction between aryl aldehydes and tertiary or secondary alkyl carboxylic acid-derived redox-active esters to deliver aryl alkyl ketone (Scheme 2).<sup>6</sup> The Breslow intermediate formed from the aldehyde and NHC reduces the NHPI ester to afford a ketyl radical and an alkyl radical, and the radical-radical coupling between them followed by the elimination of NHC will afford the desired ketone product.

### 2.1 Reactions requiring an external photocatalyst

**2.1.1 Aldehydes as substrates.** For the oxidation pathway, NHC addition into an aldehyde generates a Breslow intermediate, which can then reduce a radical acceptor to generate an NHC-derived ketyl radical species. In 2012, the Rovis group realized the first combination of NHC catalysis and



Rongxin Yang

Rongxin Yang obtained her B.Sc. degree (2020) under the supervision of Prof. Jihui Li at Hebei Normal University. She is currently studying for a master's degree at Nankai University under the guidance of Professor Qingmin Wang. Her main research interests are the synthesis and derivation of natural products and photoredox catalytic free radical chemistry.



Yuxiu Liu

Dr. Yuxiu Liu obtained her B.Sc. degree in 1994 from Nankai University. She then obtained her PhD (2003) degree from Nankai University under the supervision of Prof. Runqiu Huang. Currently, she is a professor at Nankai University. Her research focuses on molecular design, synthesis, and application of new and efficient green chemical pesticides.



Hongjian Song

Dr. Hongjian Song obtained his B.Sc. degree in 2008 from Qufu Normal University. He then obtained his M.Sc. (2011) and PhD (2014) degrees under the supervision of Associate Prof. Yuxiu Liu and Prof. Qingmin Wang, respectively, at Nankai University, majoring in organic chemistry. Currently, he is an associate professor at Nankai University. His research focuses on natural pesticide discovery and new pesticide synthesis.



Qingmin Wang

Dr. Qingmin Wang is currently a professor at the State Key Laboratory of Elemento-Organic Chemistry, Nankai University. He obtained his B.Sc. degree (1994) from Lanzhou University and PhD degree (2000) from Nankai University under the supervision of Prof. Runqiu Huang. His research interests mainly focus on the isolation, total synthesis, structural optimization, and bioactivity research of natural products and the environmentally friendly green synthesis reactions through photocatalysis and electrocatalysis.







Scheme 1 (a) Typical electron-rich NHC-bound intermediates. (b) Typical electron-deficient NHC-bound intermediates. (c) Main modes of NHC/photoredox dual catalysis.

photocatalysis, and this dual-catalysis mode enables the catalytic asymmetric  $\alpha$ -acylation of cyclic tertiary amines to generate  $\alpha$ -amino ketones (Scheme 3).<sup>7</sup> Two chemically distinct activation modes eventually result in the formation of the desired C–C bond, with H<sub>2</sub> as the only by-product. Mechanistic studies indicated that irradiation of the photocatalyst with blue light populates the excited state, which is oxidized by *meta*-dinitrobenzene; the oxidized photocatalyst is reduced by the tertiary amine, and subsequent hydrogen atom abstraction results in the formation of an iminium ion. The reaction of the NHC with the aldehyde generates a nucleophilic Breslow intermediate, which is intercepted by the iminium ion to forge a new C–C bond. The elimination of the NHC provides the  $\alpha$ -amino ketone product and completes the catalytic cycle.

The Du group described modification of  $\alpha$ -amino acids and peptides with aldehydes by photoredox/NHC dual catalysis to access structurally diverse  $\alpha$ -amino ketones (Scheme 4).<sup>8</sup> Amino




Scheme 3 Asymmetric  $\alpha$ -acylation of tertiary amines.




Scheme 4 Decarboxylative carbonylation of  $\alpha$ -amino acids.

acids prepared in advance or generated *in situ* act as radical precursors. These reactions proceed by a mechanism similar to that described by Rovis *et al.*<sup>7</sup>

In 2020, Shu and a co-workers reported the direct synthesis of amides from aldehydes and imines under redox-neutral conditions at room temperature. The key to the success of this method is NHC/visible light dual catalysis, which enables photocatalytic reduction of imino esters to nitrogen-centered radicals, which react with radical cationic intermediates to form C–N bonds (Scheme 5).<sup>9</sup>



Scheme 2 Decarboxylative alkylation of aldehydes.



Scheme 5 Amide synthesis from aldehydes and imines.





Scheme 6 Direct coupling of aldehydes with alkenes.

Scheme 7 Synthesis of  $\alpha$ -arylated- $\beta$ -thiolated ketones.

Shortly thereafter, Shu's group reported intermolecular vinylation of aldehydes, which was achieved by two sequential C-H functionalization reactions involving alkenes as vinyllating reagents (Scheme 6).<sup>10</sup> The reaction conditions are compatible with aldehydes and olefins bearing a wide range of functional groups. It is worth mentioning that TTBDPB (3,3',5,5'-tetra-*tert*-butyldiphenylene), which is used in these reactions, acts both as an oxidant and as a reservoir for radical intermediates.

Along with this two-component coupling process, Shu and co-workers also developed a related three-component protocol for straightforward access to  $\beta$ -thiolated ketones from aldehydes, styrenes, and disulfides (Scheme 7).<sup>11</sup> First, a Breslow intermediate undergoes single-electron oxidation by the excited-state photocatalyst, and then reduction of a diaryl disulfide generates a sulfur radical. The addition of the sulfur radical to a styrene derivative forms a new carbon-centered radical intermediate, which undergoes radical cross-coupling with a ketyl radical to afford the  $\alpha$ -arylated- $\beta$ -thiolated ketone product. A more stable benzyl radical was obtained by the addition with styrene analogues.

Meanwhile, the Wang group also developed a method for three-component bisfunctionalization of unactivated olefins by



Scheme 8 (a) Bisfunctionalization of unactivated olefins by aldehydes and a diazoester. (b) Bisfunctionalization of unactivated olefins by aldehydes and 2-((difluoromethyl)-sulfonyl)benzo[d]thiazole.



Scheme 9 Synthesis of 1,3-disubstituted bicyclo[1.1.1]pentane ketones.

means of NHC/photoredox dual catalysis. Proton-coupled electron transfer (PCET) generates free radicals from a diazo ester, and the radicals add to olefins to generate new radicals that couple with ketyl radicals generated *via* photoredox-catalyzed oxidation of Breslow intermediates (Scheme 8a).<sup>12</sup> In addition, acyldifluoromethylation of inert alkenes can be achieved *via* a similar mechanism when 2-((difluoromethyl)-sulfonyl)benzo[d]thiazole is used as a radical precursor (Scheme 8b).<sup>13</sup>


Recently, the Jiang and Yu group also developed a three-component reaction to access 1,3-disubstituted bicyclo[1.1.1]pentane (BCP) ketones (Scheme 9).<sup>14</sup> The BCP scaffold is a bioisostere for the phenyl ring and could enhance the solubility and permeability of bioactive molecules. In their work, radicals derived from diazo esters perform an addition reaction onto [1.1.1]propellane to afford BCP radicals, which are then coupled with the ketyl radicals that are generated *via* oxidation of the Breslow intermediates by the photoredox catalyst.

**2.1.2 Enals as substrates.** In addition to aldehydes, enals have also been used as substrates to generate Breslow intermediates by a combination of NHC catalysis and photocatalysis.






Scheme 10 Chemoselective oxidative esterification of cinnamaldehydes.




Scheme 11  $\gamma$ - and  $\epsilon$ -alkylation of enals.

The distribution of radical features provided by conjugation in unsaturated Breslow intermediates can be further extended for remote functionalization. NHC addition into an enal generates a Breslow intermediate, which converted into its dienolate form by the elimination of the pre-added leaving group. For example, in 2018, Miyabe's group reported chemoselective oxidative esterification of cinnamaldehydes by oxidation of Breslow intermediates with  $O_2$  (Scheme 10).<sup>15</sup> This catalytic cycle is completed by continuous single-electron oxidation of the photocatalyst in the presence of methanol acting as a nucleophile.


In 2019, the Ye group reported the use of  $\gamma$ -oxidized enals as substrates for the synthesis of  $\gamma$ -multisubstituted- $\alpha,\beta$ -unsaturated esters by intramolecular alkylation reactions with alkyl halides bearing an electron-withdrawing group (Scheme 11).<sup>16</sup> An alkyl radical generated from the alkyl halide by photocatalysis reacts with a dienolate intermediate generated from the enal by NHC catalysis to give a homoenolate radical. The subsequent SET oxidation of the homoenolate radical mediated by the photocatalyst affords an  $\alpha,\beta$ -unsaturated acyl azolium intermediate and completes the photocatalytic cycle. Trapping of the acyl azolium intermediate by methanol gives the final product and regenerates the NHC catalyst.



Scheme 12 Synthesis of  $\gamma$ -difluoroalkyl- $\alpha,\beta$ -unsaturated esters.



Scheme 13 Synthesis of  $\gamma$ -alkyl- $\alpha,\beta$ -unsaturated esters.




Scheme 14 Synthesis of (*Z*)-allylic fluorides.

The Ye group also developed a method for  $\gamma$ -difluoroalkylation of  $\gamma$ -preoxidized enals to afford  $\gamma$ -difluoroalkyl- $\alpha,\beta$ -unsaturated esters (Scheme 12).<sup>17</sup> This method allows for efficient construction of  $C(sp^3)$ - $CF_2R$  bonds at the  $\gamma$  position of carbonyl compounds bearing an all-carbon quaternary center.

Subsequently, Ye and a colleague expanded the substrate scope of the reaction to include cyclopropane enals. Initial ring-opening by C-C bond cleavage and subsequent  $\gamma$ -alkylation with a halogenated compound *via* a radical process afford  $\gamma$ -alkyl- $\alpha,\beta$ -unsaturated esters (Scheme 13).<sup>18</sup> A variety of alkyl halides work well in the reaction, providing the desired  $\gamma$ -alkyl- $\alpha,\beta$ -unsaturated ester products in moderate to good yields.

In 2021, Huang and Chen *et al.* reported a method for photoinduced energy-transfer reactions of NHC-bound intermediates to yield (*Z*)-allylic fluorides with good stereochemical control (Scheme 14).<sup>19</sup> A homoenolate intermediate is generated by nucleophilic addition of an NHC to an enal substrate followed by a hydride shift. A dienolate formed by release of  $CO_2$



Scheme 15 Synthesis of  $\epsilon$ -benzyl- $\alpha,\beta$ - $\gamma,\delta$ -bisunsaturated esters.Scheme 16 Synthesis of pyrrolo[1,2-*d*][1,4]oxazepin-3(2*H*)-ones.

and methanol reacts with Selectfluor to give an acyl azolium intermediate. By energy transfer from the triplet excited state of the photocatalyst, the acyl azolium intermediate undergoes double bond isomerization to afford predominantly (*Z*)-acyl azolium, which reacts with MeOH to yield the product and regenerate the NHC catalyst.

Then Ye *et al.* extended the conjugate system further and realized regioselective  $\epsilon$ -benzylation of  $\gamma$ -alkenyl- $\gamma$ -oxidized enals to afford the corresponding  $\epsilon$ -benzyl- $\alpha,\beta$ - $\gamma,\delta$ -bisunsaturated esters in moderate to good yields (Scheme 15).<sup>20</sup>

Recently, the Chauhan group developed a stereoselective strategy to access pyrrolo[1,2-*d*][1,4]oxazepin-3(2*H*)-ones (Scheme 16).<sup>21</sup> Imine is formed upon one-electron oxidation, hydrogen atom transfer, and proton transfer of amine. Then the imine intermediate enters into the NHC catalytic cycle, where the Breslow intermediate acts as the homoenolate equivalent, which after tautomerization undergoes nucleophilic addition to imine to forge acyl azolium species. Eventually the azolium intermediate delivers the product.

## 2.2 Reactions not requiring an external photocatalyst

The process without an external photocatalyst is usually to reduce the radical precursor by generating an excited state of the Breslow intermediate under light excitation, directly generate free radicals through special radical precursors under light or generate EDA complexes through the Breslow intermediates and other reactants in the system. Various reactions



Scheme 17 Synthesis of protected amines.


Scheme 18 Synthesis of  $\alpha$ -iodo ketones.

can be achieved by combining light activation and NHC organocatalysis without an external photocatalyst. For example, in 2020, Chen's group demonstrated that transition-metal-free decarboxylative C(sp<sup>3</sup>)-X bond formation can be accomplished with photochemically active *N*-(acyloxy)phthalimide ester-NaI-NHC complexes; these reactions offer a convenient way to construct C(sp<sup>3</sup>)-C(sp<sup>2</sup>), C(sp<sup>3</sup>)-S, C(sp<sup>3</sup>)-O, and C(sp<sup>3</sup>)-Cl bonds (Scheme 17).<sup>22</sup> The key to these reactions is the electrostatic NHC-Na<sup>+</sup> interaction, which facilitates the formation of electron donor-acceptor complexes, irradiation of which generates radicals that go on to form the products. Later, the same research group described catalytic reactions of *N*-alkenoxypyridinium salts and NaI to give various  $\alpha$ -iodo ketones (Scheme 18).<sup>23</sup>

In 2022, the Larionov group developed a method for visible-light-induced NHC-catalyzed regioselective 1,2-diacylation reactions of alkenes that afford 1,4-diketones *via* three-component C-C-bond-forming radical coupling (Scheme 19).<sup>24</sup> Notably, in this system, the NHC catalyst plays two roles: an NHC-catalyst-derived intermediate forms an electron donor-acceptor complex, and the NHC catalyst acts as an acyl transfer reagent. Under irradiation by blue LEDs, intramolecular SET between a Breslow intermediate and an oxime generates nitrogen and ketyl radicals. The nitrogen radical fragments to release acetonitrile and an acyl radical, which adds to an olefin to generate an adduct radical. The adduct radical participates in a cross-coupling reaction with the ketyl radical to afford a 1,4-diketone in high yield.

The Hong group developed a catalytic method for enantioselective, C4-selective functionalization of pyridine derivatives (Scheme 20),<sup>25</sup> which proceeds by means of a mechanism





Scheme 19 Synthesis of 1,4-diketones from oximes, aldehydes, and alkenes.




Scheme 20 Synthesis of  $\beta$ -pyridylated esters.

similar to that reported by Larionov *et al.*<sup>24</sup> The key to these asymmetric  $\beta$ -pyridylation reactions of enals is an enhanced interactions between a chiral-NHC-bound homoenolate and a pyridinium salt in the presence of a pivalate salt and hexafluorobenzene, which effectively distinguishes between the two faces of a homoenolate radical. Because light-facilitated reactivity and rate acceleration were observed, Hong *et al.*<sup>25</sup> proposed an alternative mechanism involving photon absorption by a pyridine–pivalate electron donor–acceptor complex that triggers the formation of amidine radicals when the EDA complex is irradiated with visible light.

Other systems can, upon photoirradiation, directly generate radicals that can then participate in a catalytic cycle involving NHC. In 2020, the Hui group achieved stereoselective [4 + 2] cycloaddition reactions of 3-alkylenoxindoles and  $\alpha$ -diazoketones that proceed *via* this mechanism (Scheme 21).<sup>26</sup> Initially, a ketene is formed from the  $\alpha$ -diazoketone through a Wolff rearrangement reaction under blue light. Subsequently, the addition of NHC to the ketene generates an enolate, which



Scheme 21 Synthesis of tetrahydropyrano[2,3-*b*]indoles.



Scheme 22 Synthesis of dihydropyridinones.

participates in a [4 + 2] annulation reaction with the 3-alkylenoxindole to give an intermediate that is transformed into a tetrahydropyrano[2,3-*b*]indole upon release of the NHC catalyst. Subsequently, the Yao group developed a method for asymmetric [4 + 2] annulation reactions of saccharine-derived azadienes and  $\alpha$ -diazoketones, affording the corresponding sultam-fused dihydropyridinones in moderate to good yields with satisfactory to excellent enantio- and diastereo-selectivities (Scheme 22).<sup>27</sup>


In 2020, the Xuan group described a multicomponent reaction that relies on the different reactivities of two carbenes (Scheme 23).<sup>28</sup> One of the N-heterocyclic carbenes, acting as an organocatalyst, mediates the formation of a hydroxamic acid *in situ*, whereas the other carbene, which is formed by photolysis of diazoalkane, acts as a reactant. Then the hydroxamic acid and carbene participate in a solvent-dependent four-component reaction that provides hydroxamic acid esters, which are biologically important compounds.

## 3. Ketyl radicals generated *via* acyl azolium intermediates

### 3.1 Acyl imidazoles as acyl sources

**3.1.1 Direct formation of benzyl radicals.** Like processes involving single-electron oxidation of electron-rich Breslow intermediates, processes involving generation of electron-deficient acyl azolium intermediates and their single-electron reduction are currently a booming area of research. In 2020, Scheidt *et al.* reported a method for reductive single-electron alkylation of acyl azoliums to form ketones from aryl acyl






Scheme 23 Formation of hydroxamic acid esters by a four-component reaction in a cyclic ether as a reaction medium.



Scheme 24 Synthesis of ketones from HEs and aryl acyl imidazoles.



Scheme 25 Synthesis of ketones from HEs and alkyl acyl imidazoles.



Scheme 26 Synthesis of  $\alpha,\beta$ -disubstituted cyclohexanones from HEs and alkyl acyl imidazoles.



Scheme 27 Synthesis of sulfoxides from HEs and sulfinyl imidazoles.

imidazoles derived from carboxylic acids (Scheme 24).<sup>29</sup> Hantzsch esters (HEs), which can be easily synthesized from simple starting materials with a variety of structures, are used as alkylation reagents. The excited-state photocatalyst initially oxidizes the HE to an alkyl radical (most of the examples involve a benzyl radical), and single-electron reduction of the acyl azolium ion provides a ketyl radical while regenerating the ground-state photocatalyst. Radical–radical coupling and loss of the NHC afford the desired ketone. Note, however, that this reaction is applicable only to aryl carboxylic acids.

Scheidt *et al.* later expanded the reaction to alkyl acyl imidazoles by altering the structure of the NHC catalyst. Their findings revealed that both electronic and steric modifications of the NHC catalyst would affect the stability and accessibility of the radical intermediates, thereby controlling the reactivity of NHC catalysis (Scheme 25).<sup>30</sup>

In 2022, the Scheidt group developed a strategy for the construction of two contiguous C–C bonds *via* a formal [5 + 1] cycloaddition for the synthesis of  $\alpha,\beta$ -disubstituted cyclohexanones from HEs and alkyl acyl imidazoles (Scheme 26).<sup>31</sup> The

key to this transformation lies in the two photocatalytic cycles to achieve intramolecular cyclization. In the first cycle, the photoexcited photocatalyst oxidizes the HE to afford a benzyl radical, and single-electron reduction of an acyl azolium provides a ketyl radical. Intermolecular coupling of the two radicals and loss of the NHC give a linear ketone intermediate. In the second cycle, the corresponding enol or cesium enolate undergoes single-electron oxidation to produce a benzyl radical, which participates in an intramolecular cyclization reaction with the remote double bond. The resulting radical undergoes hydrogen atom transfer (HAT) or is reduced to the corresponding anion by the photocatalyst, and subsequent deprotonation by the solvent or bicarbonate affords the desired cyclohexanone product.

Subsequently, the Wu group reported the generation of sulfoxides from sulfinic acids and 4-substituted HEs in the presence of carbonyldiimidazole (Scheme 27).<sup>32</sup> Acting as free-radical precursors, HEs can generate not only benzyl radicals but also alkyl radicals, and a series of sulfinyl products were obtained by this method. This method further expands the





Scheme 28 Synthesis of ketones from alkylborates and alkyl acyl imidazoles.



Scheme 29 Synthesis of  $\gamma$ -aryloxyketones.

previous carboxylic acid category to sulfonic acid, and using sulfinyl imidazoles as pseudo-acyl sources.


**3.1.2 Indirect formation of benzyl radicals.** In addition to being directly accessible by single-electron oxidation of radical precursors, benzyl radicals can also be obtained indirectly by the addition of radicals to styrene analogues. In 2021, Ohmiya and co-workers designed a protocol that enables cross-coupling between alkylborates and alkyl acyl imidazoles in addition to radical relay-type alkylacylations of alkenes with alkylborates and alkyl acyl imidazoles, affording a diverse array of ketones (Scheme 28).<sup>33</sup> In a light-driven NHC catalytic cycle, an alkyl radical and a ketyl radical, generated from the borate or the acyl azolium intermediate by a SET process, participate in radical-radical coupling to yield the acylation product. Alternatively, the addition of the alkene to the reaction system results in a radical relay process, and subsequent radical-radical coupling between another alkyl radical and the ketyl radical affords the alkylacylation product.



Scheme 30 Synthesis of ketones from carboxylic acids and acyl imidazoles.



Scheme 31 Synthesis of substituted 3,4-dihydro-2H-pyrroles from oxime ethers and acyl imidazoles.



Scheme 32 Construction of  $\alpha$ -amino ketones from acyl imidazoles and amines.

In 2022, the Scheidt group reported a multicomponent reaction for the synthesis of  $\gamma$ -aryloxyketones *via* aryloxymethyl potassium trifluoroborate salts. An aryloxymethyl radical adds to a styrene derivative to provide a stabilized benzyl radical, and a subsequent radical-radical coupling reaction with an azolium radical affords the  $\gamma$ -aryloxyketone product (Scheme 29).<sup>34</sup>

In the same year, the Chi group published a method for coupling carboxylic acids and acyl imidazoles by means of a combination of NHC catalysis and photocatalysis (Scheme 30).<sup>35</sup> The carboxylic acids are directly used as radical precursors, and late-stage modification of commercial drugs and direct coupling of fragments of two medicinally active molecules were performed to demonstrate the utility of this method.

Recently, the Ye group reported the iminoacylation of alkenes *via* decarboxylation of  $\alpha$ -imino-oxy acids to generate iminyl radical intermediates. The addition of the iminyl radicals to a tethered alkene in a 5-*exo-trig* manner gives dihydropyrrole-derived carbon radicals (most of the examples were benzyl radicals), which couple with ketyl radicals generated from acyl azolium intermediates to form substituted 3,4-dihydro-2H-pyrroles (Scheme 31).<sup>36</sup>





Scheme 33 Construction of  $\beta,\gamma$ -unsaturated ketones from acyl imidazoles and alkenes.

**3.1.3 Formation of heteroatomic ortho radicals.** In 2022, the Wang group reported the direct acylation of  $\alpha$ -C(sp<sup>3</sup>)-H bonds of amines by acyl imidazoles to access  $\alpha$ -amino ketones (Scheme 32).<sup>37</sup> Single-electron oxidation of the amine and subsequent deprotonation afford an  $\alpha$ -amino radical, and then radical–radical cross-coupling with a ketyl radical provides the corresponding product.

**3.1.4 Formation of allyl radicals.** Recently, the Wang group achieved the direct allylic acylation *via* cross-coupling involving cooperative NHC, hydrogen atom transfer (HAT), and photoredox catalysis to synthesize  $\beta,\gamma$ -unsaturated ketones (Scheme 33).<sup>38</sup> The thiyl radical generated from single-electron oxidation of thiol serves as a powerful HAT catalyst, abstracting an allylic hydrogen from alkenes to generate allyl radicals. The coupling of allyl radicals with ketyl radicals affords desired  $\beta,\gamma$ -unsaturated ketones.

## 3.2 Acyl fluorides as acyl sources

**3.2.1 Direct formation of benzyl radicals.** In addition to imidazole, fluorine can also be used as a leaving group in single-electron reduction of acyl azolium intermediates to generate ketyl radicals. For example, in 2021, the Studer group developed a method for direct acylation of benzylic C–H bonds *via* NHC/



Scheme 34 Construction of benzyl aryl ketones by benzylic C(sp<sup>3</sup>)-arylation.



Scheme 35 Construction of 3-aryl-2-fluoro-2,3-dihydrobenzofurans from acyl fluorides and benzofurans.

photoredox dual catalysis (Scheme 34).<sup>39</sup> This mild method allows the preparation of a range of benzylic ketones and shows a good functional group. Under irradiation of blue LEDs, single-electron oxidation of electron-rich alkylarenes and subsequent deprotonation at the benzylic position give the corresponding benzyl radicals, which couple with ketyl radicals derived from acyl azolium intermediates to afford benzyl aryl ketones.


In addition, this research group also generated benzyl radicals by means of SET oxidation of the double bonds of benzofurans (Scheme 35).<sup>40</sup> Specifically, they reported fluoroarylation of benzofurans by acyl fluorides, which act as bifunctional reagents to incorporate both an aroyl moiety and fluorine into the product. Upon visible-light irradiation, the benzofuran is oxidized to a radical cation by a photoexcited photocatalyst. A ketyl radical is generated from an acyl azolium intermediate, and cross-coupling of the radical cation and the ketyl radical leads to an oxocarbenium ion. Diastereoselective trapping of this ion by a F<sup>−</sup> anion affords the 3-aryl-2-fluoro-2,3-dihydrobenzofuran product.

Like benzofurans, substituted styrenes can act as benzyl radical precursors by undergoing metal-mediated HAT. In 2022, the Wang group achieved Markovnikov-selective hydroacylation of alkenes by using a synergistic combination of cobalt, photoredox, and NHC catalysis (Scheme 36).<sup>41</sup> The cobalt catalytic cycle starts with SET oxidation of Co<sup>II</sup> to Co<sup>III</sup>, and then Co<sup>III</sup> is captured by phenylsilane to furnish a Co<sup>III</sup>–H intermediate. This intermediate engages in a HAT reaction with the substituted styrene to produce a benzyl radical. Meanwhile, SET reduction of an acyl azolium ion gives a ketyl radical, which undergoes radical–radical cross-coupling with the benzyl radical to generate the hydroacylation product.



Scheme 36 Markovnikov-selective hydroacylation of alkenes.






Scheme 37 Construction of ketones from acyl fluorides and alkyl trifluoroborates.



Scheme 39 Construction of  $\beta$ -trifluoromethyl- $\alpha$ -substituted ketones from acyl fluorides and styrenes.



Scheme 38 Synthesis of isochroman-1-one derivatives.



Scheme 40 Construction of  $\gamma$ -aryloxy ketones from acyl fluorides and aryl cyclopropanes.

Shortly thereafter, the Li group reported cross-coupling reactions of alkyl trifluoroborates with acid fluorides to generate various ketones; this method provides an alternative to the classical acylative Suzuki coupling chemistry (Scheme 37). Li *et al.* proposed the activation of an acyl azolium intermediate with a triplet-state photocatalyst, through an energy-transfer process, to form an excited-state acyl azolium intermediate.<sup>42</sup> This intermediate oxidizes the alkyl trifluoroborates to give a ketyl radical and a benzyl radical. The coupling of these two radicals affords the ketone product.


**3.2.2 Indirect formation of benzyl radicals.** In 2020, the Hopkinson group reported the use of acid fluorides and trifluoroacetophenones as substrates in a UVA-light-mediated photochemical transformation that leads to diverse isochroman-1-one derivatives (Scheme 38).<sup>43</sup> The azolium intermediate generated from the acid fluoride by NHC catalysis is excited by UVA irradiation to afford, after intersystem crossing, a triplet excited state. Then 1,5-HAT from the *o*-benzylic position to the radical-like carbonyl oxygen atom gives rise to a triplet-state dienol biradical. Rotation of this species before relaxation leads to a ground-state intermediate, which can react with the dienophile in a cycloaddition process. Finally, elimination of the NHC from the cycloadduct completes the catalytic cycle.

In 2020, the Studer group reported three-component coupling reactions of acyl fluorides, styrenes, and the Langlois reagent ( $\text{CF}_3\text{SO}_2\text{Na}$ ) to give various  $\beta$ -trifluoromethylated- $\alpha$ -substituted ketones (Scheme 39).<sup>44</sup> Reductive quenching of the excited-state photocatalyst by the trifluoromethanesulfinate anion gives a trifluoromethylsulfonyl radical that fragments to release  $\text{SO}_2$  and a trifluoromethyl radical, which then adds to the double bond of styrene to generate a transient benzylic radical. Meanwhile, SET reduction of an acyl azolium ion gives a ketyl radical, which undergoes radical-radical cross-coupling with the benzylic radical to give the  $\beta$ -trifluoromethyl- $\alpha$ -substituted ketones.

Shortly thereafter, Studer *et al.* reported a ring-opening/arylcarboxylation/acylation cascade reaction for the 1,3-difunctionalization of aryl cyclopropanes (Scheme 40).<sup>45</sup> The key to this transformation is that the aryl cyclopropane radical cation generated by SET oxidation of the aryl cyclopropane undergoes ring opening by a nucleophilic benzoate ion to give a benzylic radical. Radical-radical cross-coupling of the benzylic radical and a ketyl radical affords  $\gamma$ -aryloxy ketones.

The Feng and Fan group developed a method for intermolecular 1,2-diacylation of styrenes *via* cooperative NHC and photoredox catalysis with mediation by  $\text{PPh}_3$  and  $\text{Cs}_2\text{CO}_3$  (Scheme 41).<sup>46</sup> The mechanism is similar to that described by Studer *et al.*<sup>45</sup> An NHC-mediated reaction of a bisacyl carbonate





Scheme 41 Construction of symmetrical 1,4-dicarbonyl compounds from acyl fluorides and styrenes.




Scheme 42 Construction of unsymmetrical 1,4-dicarbonyl compounds from alkenes, acyl fluorides, and keto acids.


intermediate generated from an acyl fluoride produces a benzoate anion and an acyl azolium ion. The benzoate anion combines with a triphenylphosphine cation radical to form a phosphorus-centered radical, which undergoes  $\beta$ -scission to generate an acyl radical. The acyl radical attacks the styrene substrate to generate the corresponding benzyl radical. Next, intermolecular cross-coupling between a ketyl radical (derived from the acyl azolium ion) and the benzyl radical produces the desired product. In a similar manner, keto acids can also be used as acyl radical precursors *via* single-electron oxidation, and subsequent dicarbonylation of alkenes provides direct access to 1,4-dicarbonyl compounds (Scheme 42).<sup>47</sup>

Unlike trifluoromethylsulfonyl radicals, aryl sulfonyl radicals do not undergo  $\text{SO}_2$  extrusion to produce the corresponding aryl radicals. Therefore, in 2022, Studer and a colleague developed an arylsulfonate-catalyzed alkene acylation reaction. In this reaction, an arylsulfonate radical first adds to the alkene substrate to generate a carbon-centered radical, which then couples with a ketyl radical (generated from an acyl azolium), providing a three-component coupling intermediate. The subsequent base-mediated elimination of arylsulfinate forms the  $\alpha$ -acylated olefin product. Three catalytic cycles involving a carbene are interwoven (Scheme 43).<sup>48</sup>


The Zhang group described a similar reaction: 1,4-sulfonylacylation of 1,3-enynes to provide valuable structurally diverse tetrasubstituted allenyl ketones (Scheme 44).<sup>49</sup> The addition of a sulfonyl radical to a 1,3-enyne delivers a propargyl radical that can undergo reversible isomerization to generate a trisubstituted allenyl radical, which can participate in a radical-radical



Scheme 43 Construction of  $\alpha$ -substituted vinyl ketones by  $\alpha$ -acylation of alkenes.



Scheme 44 1,4-Sulfonylacylation of 1,3-enynes.



Scheme 45 1,2-Sulfonylacylation of allenes.


cross-coupling reaction with a ketyl radical generated by reduction of an acyl azolium ion. By replacing the 1,3-enyne acceptor of the sulfonyl radical with an allene, these investigators also achieved 1,2-sulfonylacylation of allenes to provide valuable sulfonyl-containing multisubstituted allyl ketones (Scheme 45).<sup>50</sup>

**3.2.3 Formation of phenyl radicals.** Recently, the Studer group reported a C–H Acylation of arenes and heteroarenes through cooperative photoredox/NHC radical catalysis (Scheme 46).<sup>51</sup> The cross coupling of arene radical cations, formed through the single electron oxidation of arenes, with an NHC-bound ketyl radical leads to ketone products. Importantly, if acylation occurs under classical Friedel–Crafts conditions using





Scheme 46 C–H Acylation of arenes.

Scheme 47 Construction of  $\alpha$ -amino ketones from acyl fluoride and oxamic acid.

$\text{AlCl}_3$  as a Lewis acid, different regioselectivity will be obtained. Therefore, by simply switching the reaction conditions, two different regional isomers can be obtained.

**3.2.4 Formation of carbamoyl radicals.** Recently, the Luo and Yang group reported a decarboxylative acylation of oxamic acid with acyl fluoride to produce  $\alpha$ -keto amides (Scheme 47).<sup>52</sup> A carbamoyl radical is generated through single-electron oxidation and then couples with a ketyl radical to produce the desired product.

### 3.3 Esters as acyl sources

In addition to imidazoles and fluorides, esters can also act as leaving groups. For example, in 2021, the Chi group developed a method for alkylation of aryl carboxylic esters with HES (Scheme 48).<sup>53</sup> The reaction starts with addition of an NHC catalyst to the ester to generate an acyl azolium intermediate. The photoexcitation of this intermediate converts it to an excited state that act as a single-electron oxidant. Subsequently, SET between the HE and the excited-state acyl azolium leads to a benzyl radical and a ketyl radical, and coupling of these two radicals affords the desired ketone product. Structurally sophisticated ketones, including ketones bearing medicinal fragments, could readily be prepared.



Scheme 48 Alkylation of carboxylic esters to generate ketones.

## 4. Conclusion and outlook

As complements to two-electron reaction modes, single-electron reaction modes give new vitality to NHC-catalyzed reactions. This review has focused on combinations of NHC catalysis with photocatalysis. The combinations have been classified on the basis of whether the ketyl radical is generated by single-electron oxidation of an electron-rich Breslow intermediate or by single-electron reduction of an electron-deficient acyl azolium intermediate. Reactions in the first category have been further categorized on the basis of their substrates and whether an external photocatalyst is required, whereas reactions in the second category have been categorized mainly on the basis of the type of leaving group and the type of radical that is generated. Through the combination of NHC catalysis and photocatalysis, many transformations that cannot be achieved by means of two-electron reactions can be realized.

Although NHC-mediated single-electron reactions have made great progress and breakthroughs, there are still some problems that remain to be solved, including: (1) acyl substrate scope: since the previously activated acyl precursors are mostly carboxylic acids, it would be important to explore whether the activation process of carboxylic acid derivatives, aldehyde derivatives and imines could be realized through the co-catalysis mode of light and NHC; (2) types of free radicals: since most of the literature reported was on the direct formation of benzyl radicals or the addition of alkyl radicals to aryl olefins to achieve the indirect formation of benzyl radicals, to explore other unstable radical such as selective and efficient acylation of unactivated C–H bonds through this catalytic mode will be meaningful; (3) enantioselectivity control: how to build a quaternary or tertiary stereocenter at the  $\alpha$ -position of a carbonyl group with high enantioselectivity should be considered; (4) catalytic mode: the development of new catalytic modes through merging NHC catalysis with otherwise established chemistry protocols such as electrochemistry or metal catalysis should be explored as well.



## Author contributions

Xiaochen Wang: conceptualization, visualization, and writing – original draft. Senhui Wu and Rongxin Yang: Investigation. Hongjian Song and Yuxiu Liu: writing – review & editing. Qingmin Wang: supervision, conceptualization, writing – review & editing, and funding acquisition. All authors have given approval to the final version of the manuscript.

## Conflicts of interest

There are no conflicts to declare.

## Acknowledgements

We are grateful to the National Natural Science Foundation of China (22271166 and 22077071) and the Frontiers Science Center for New Organic Matter, Nankai University (63181206) for generous financial support for our programs.

## References

- (a) D. Enders, O. Niemeier and A. Henseler, Organocatalysis by N-Heterocyclic Carbenes, *Chem. Rev.*, 2007, **107**, 5606–5655; (b) X. Y. Chen and S. Ye, Enantioselective Cycloaddition Reactions of Ketenes Catalyzed by N-Heterocyclic Carbenes, *Synlett*, 2013, **24**, 1614–1622; (c) X. Y. Chen and S. Ye, N-heterocyclic carbene-catalyzed reactions of C–C unsaturated bonds, *Org. Biomol. Chem.*, 2013, **11**, 7991–7998; (d) M. N. Hopkinson, C. Richter, M. Schedler and F. Glorius, An overview of N-heterocyclic carbenes, *Nature*, 2014, **510**, 485–496; (e) D. M. Flanagan, F. Romanov-Michailidis, N. A. White and T. Rovis, Organocatalytic Reactions Enabled by N-Heterocyclic Carbenes, *Chem. Rev.*, 2015, **115**, 9307–9387; (f) C. Zhang, J. F. Hooper and D. W. Lupton, N-Heterocyclic Carbene Catalysis via the  $\alpha,\beta$ -Unsaturated Acyl Azolium, *ACS Catal.*, 2017, **7**, 2583–2596; (g) M. Zhao, Y. T. Zhang, J. Chen and L. Zhou, Enantioselective Reactions Catalyzed by N-Heterocyclic Carbenes, *Asian J. Org. Chem.*, 2018, **7**, 54–69; (h) S. Mondal, S. R. Yetra, S. Mukherjee and A. T. Biju, NHC-Catalyzed Generation of  $\alpha,\beta$ -Unsaturated Acylazoliums for the Enantioselective Synthesis of Heterocycles and Carbocycles, *Acc. Chem. Res.*, 2019, **52**, 425–436; (i) X. Y. Chen, Z. H. Gao and S. Ye, Bifunctional N-Heterocyclic Carbenes Derived from L-Pyroglutamic Acid and Their Applications in Enantioselective Organocatalysis, *Acc. Chem. Res.*, 2020, **53**, 690–702; (j) H. Ohmiya, N-Heterocyclic Carbene-Based Catalysis Enabling Cross-Coupling Reactions, *ACS Catal.*, 2020, **10**, 6862–6869.
- (a) R. Breslow, On the Mechanism of Thiamine Action. IV.1 Evidence from Studies on Model Systems. On the Mechanism of Thiamine Action. IV.1 Evidence from Studies on Model Systems, *J. Am. Chem. Soc.*, 1958, **80**, 3719–3726; (b) B. E. Maki, A. Chan, E. M. Phillips and K. A. Scheidt, Tandem Oxidation of Allylic and Benzylic Alcohols to Esters Catalyzed by N-Heterocyclic Carbenes, *Org. Lett.*, 2007, **9**, 371–374; (c) T. Ukai, R. Tanaka and T. Dokawa, *J. Pharm. Soc. Jpn.*, 1943, **63**, 296–300; (d) For selected reviews, see: T. Ishii, K. Nagao and H. Ohmiya, Recent advances in N-heterocyclic carbene-based radical catalysis, *Chem. Sci.*, 2020, **11**, 5630–5636.
- (a) N. R. Patel, C. B. Kelly, A. P. Siegenfeld and G. A. Molander, Mild, Redox-Neutral Alkylation of Imines Enabled by an Organic Photocatalyst, *ACS Catal.*, 2017, **7**, 1766–1770; (b) J. Jia, Q. Lefebvre and M. Rueping, Reductive Coupling of Imines with Redox-active Esters by Visible Light Photoredox Organocatalysis, *Org. Chem. Front.*, 2020, **7**, 602–608; (c) P. Ji, Y. Zhang, Y. Wei, H. Huang, W. Hu, P. A. Mariano and W. Wang, Visible-Light-Mediated, Chemo- and Stereoselective Radical Process for the Synthesis of C-Glycoamino Acids, *Org. Lett.*, 2019, **21**, 3086–3092; (d) H.-H. Zhang and S. Yu, Radical Alkylation of Imines with 4-Alkyl-1,4-dihydropyridines Enabled by Photoredox/Brønsted Acid Cocatalysis, *J. Org. Chem.*, 2017, **82**, 9995–10006; (e) D. P. Plasko, C. J. Jordan, B. E. Ciesa, M. A. Merrill and J. M. Hanna, Visible light-promoted alkylation of imines using potassium organotrifluoroborates, *Photochem. Photobiol. Sci.*, 2018, **17**, 534–538; (f) K. Cao, S. M. Tan, R. Lee, S. Yang, H. Jia, X. Zhao, B. Qiao and Z. Jiang, Catalytic Enantioselective Addition of Prochiral Radicals to Vinylpyridines, *J. Am. Chem. Soc.*, 2019, **141**, 5437–5443; (g) Y. Li, K. Zhou, Z. Wen, S. Cao, X. Shen, M. Lei and L. Gong, Copper(II)-Catalyzed Asymmetric Photoredox Reactions: Enantioselective Alkylation of Imines Driven by Visible Light, *J. Am. Chem. Soc.*, 2018, **140**, 15850–15858.
- For reviews and selected work on photochemistry(a) T. P. Yoon, M. A. Ischay and J. Du, Visible light photocatalysis as a greener approach to photochemical synthesis, *Nat. Chem.*, 2010, **2**, 527–532; (b) J. Xuan and W.-J. Xiao, Visible-Light Photoredox Catalysis, *Angew. Chem., Int. Ed.*, 2012, **51**, 6828–6838; (c) C. K. Prier, D. A. Rankic and D. W. C. MacMillan, Visible Light Photoredox Catalysis with Transition Metal Complexes: Applications in Organic Synthesis, *Chem. Rev.*, 2013, **113**, 5322–5363; (d) R. Francke and R. D. Little, Redox catalysis in organic electrosynthesis: basic principles and recent developments, *Chem. Soc. Rev.*, 2014, **43**, 2492–2521; (e) N. A. Romero and D. A. Nicewicz, Organic Photoredox Catalysis, *Chem. Rev.*, 2016, **116**, 10075–10166; (f) K. L. Skubi, T. R. Blum and T. P. Yoon, Dual Catalysis Strategies in Photochemical Synthesis, *Chem. Rev.*, 2016, **116**, 10035–10074; (g) M. H. Shaw, J. Twilton and D. W. C. MacMillan, Photoredox Catalysis in Organic Chemistry, *J. Org. Chem.*, 2016, **81**, 6898–6926.
- For reviews and selected work on radical NHC catalysis(a) K. Liu, M. Schwenzer and A. Studer, Radical NHC Catalysis, *ACS Catal.*, 2022, **12**, 11984–11999; (b) R. Song, Z. Jin and R. Y. Chi, NHC-catalyzed covalent activation of heteroatoms for enantioselective reactions, *Chem. Sci.*, 2021, **12**, 5037–5043; (c) K.-Q. Chen, H. Sheng, Q. Liu, P.-L. Shao and X.-Y. Chen, N-heterocyclic carbene-catalyzed radical reactions, *Sci. China: Chem.*, 2021, **64**, 7–





- Electrostatic Interaction of N-Heterocyclic Carbenes, *Org. Lett.*, 2020, **22**, 8059–8064.
- 24 S. Jin, X. Sui, G. C. Haug, V. D. Nguyen, H. T. Dang, H. D. Arman and O. V. Larionov, N-Heterocyclic Carbene-Photocatalyzed Tricomponent Regioselective 1,2-Diacylation of Alkenes Illuminates the Mechanistic Details of the Electron Donor–Acceptor Complex Mediated Radical Relay Process, *ACS Catal.*, 2022, **12**, 285–294.
- 25 H. Choi, G. R. Mathi, S. Hong and S. Hong, Enantioselective functionalization at the C4 position of pyridinium salts through NHC catalysis, *Nat. Commun.*, 2022, **13**, 1776–1783.
- 26 C. Wang, Z. Wang, J. Yang, S. Shi and X. Hui, Sequential Visible-Light and N-Heterocyclic Carbene Catalysis: Stereoselective Synthesis of Tetrahydropyrano[2,3-b]indoles, *Org. Lett.*, 2020, **22**, 4440–4443.
- 27 Y. Chen, B. Shi, H. Yin, Y. Liu, C. Yu, K. Zhang, T. Li and C. Yao, Stereoselective synthesis of chiral sultam-fused dihydropyridinones *via* photopromoted NHC catalyzed [4 + 2] annulation, *Org. Chem. Front.*, 2022, **9**, 5191–5196.
- 28 B. Cai, Q. Li, C. Empel, L. Li, R. M. Koenigs and J. Xuan, Dark and Light Reactions of Carbenes–Merging Carbene Transfer Reactions with N-Heterocyclic Carbene Catalysis for the Synthesis of Hydroxamic Acid Esters, *ACS Catal.*, 2022, **12**, 11129–11136.
- 29 A. V. Bay, K. P. Fitzpatrick, R. C. Betori and K. A. Scheidt, Combined Photoredox and Carbene Catalysis for the Synthesis of Ketones from Carboxylic Acids, *Angew. Chem., Int. Ed.*, 2020, **59**, 9143–9148.
- 30 A. V. Bay, K. P. Fitzpatrick, G. A. González-Montiel, A. O. Farah, P. H. Cheong and K. A. Scheidt, Light-Driven Carbene Catalysis for the Synthesis of Aliphatic and  $\alpha$ -Amino Ketones, *Angew. Chem., Int. Ed.*, 2021, **60**, 17925–17931.
- 31 A. V. Bay, E. J. Farnam and K. A. Scheidt, Synthesis of Cyclohexanones by a Tandem Photocatalyzed Annulation, *J. Am. Chem. Soc.*, 2022, **144**, 7030–7037.
- 32 X. Wang, Y. Tang, S. Ye, J. Zhang, Y. Kuang and J. Wu, Access to Sulfoxides under NHC/Photocatalysis *via* a Radical Pathway, *Org. Lett.*, 2022, **24**, 2059–2063.
- 33 Y. Sato, Y. Goto, K. Nakamura, Y. Miyamoto, Y. Sumida and H. Ohmiya, Light-Driven N-Heterocyclic Carbene Catalysis Using Alkylborates, *ACS Catal.*, 2021, **11**, 12886–12892.
- 34 P. Wang, K. P. Fitzpatrick and K. A. Scheidt, Combined Photoredox and Carbene Catalysis for the Synthesis of  $\gamma$ -Aryloxy Ketones, *Adv. Synth. Catal.*, 2022, **364**, 518–524.
- 35 S. Ren, X. Yang, B. Mondal, C. Mou, W. Tian, Z. Jin and Y. R. Chi, Carbene and photocatalyst-catalyzed decarboxylative radical coupling of carboxylic acids and acyl imidazoles to form ketones, *Nat. Commun.*, 2022, **13**, 2846–2855.
- 36 Y. Dong, C. Zhang, Z. Gao and S. Ye, Iminoacylation of Alkenes *via* Photoredox N-Heterocyclic Carbene Catalysis, *Org. Lett.*, 2023, **25**, 855–860.
- 37 X. Wang, B. Zhu, Y. Liu and Q. Wang, Combined Photoredox and Carbene Catalysis for the Synthesis of  $\alpha$ -Amino Ketones from Carboxylic Acids, *ACS Catal.*, 2022, **12**, 2522–2531.
- 38 X. Wang, R. Yang, B. Zhu, Y. Liu, H. Song, J. Dong and Q. Wang, Direct Allylic Acylation *via* Cross-Coupling Involving Cooperative N-Heterocyclic Carbene, Hydrogen Atom Transfer, and Photoredox Catalysis, *Nat. Commun.*, 2023, **14**, 2951–2960.
- 39 Q. Meng, L. Lezius and A. Studer, Benzylic C–H acylation by cooperative NHC and photoredox catalysis, *Nat. Commun.*, 2021, **12**, 2068–2075.
- 40 X. Yu, Q. Meng, C. G. Daniliuc and A. Studer, Aroyl Fluorides as Bifunctional Reagents for Dearomatizing Fluoroarylation of Benzofurans, *J. Am. Chem. Soc.*, 2022, **144**, 7072–7079.
- 41 X. Tao, Q. Wang, L. Kong, S. Ni, Y. Pan and Y. Wang, Branched-Selective Hydroacylation of Alkenes *via* Photoredox Cobalt and N-Heterocyclic Carbene Cooperative Triple Catalysis, *ACS Catal.*, 2022, **12**, 15241–15248.
- 42 H. Huang, Q. Dai, H. Leng, Q. Li, S. Yang, Y. Tao, X. Zhang, T. Qia and J. Li, Suzuki-type cross-coupling of alkyl trifluoroborates with acid fluoride enabled by NHC/photoredox dual catalysis, *Chem. Sci.*, 2022, **13**, 2584–2590.
- 43 (a) A. Mavroskoufis, K. Rajes, P. Golz, A. Agrawal, V. Ruß, P. J. Götze and M. N. Hopkinson, N-Heterocyclic Carbene Catalyzed Photoenolization/Diels–Alder Reaction of Acid Fluorides, *Angew. Chem., Int. Ed.*, 2020, **59**, 3190–3194; (b) A. Mavroskoufis, M. Lohani, M. Weber, M. N. Hopkinson and J. P. Götze, A. (TD)-DFT study on photo-NHC catalysis: photoenolization/Diels–Alder reaction of acid fluorides catalyzed by N-heterocyclic carbenes, *Chem. Sci.*, 2023, **14**, 4027–4037.
- 44 Q. Meng, N. Döben and A. Studer, Cooperative NHC and Photoredox Catalysis for the Synthesis of *b*-Trifluoromethylated Alkyl Aryl Ketones, *Angew. Chem., Int. Ed.*, 2020, **59**, 19956–19960.
- 45 Z. Zuo, C. G. Daniliuc and A. Studer, Cooperative NHC/Photoredox Catalyzed Ring-Opening of Aryl Cyclopropanes to 1-Aroyloxy-3-Acylated Alkanes, *Angew. Chem., Int. Ed.*, 2021, **60**, 25252–25257.
- 46 S. Li, H. Shu, S. Wang, W. Yang, F. Tang, X. Li, S. Fan and Y. Feng, Cooperative NHC and Photoredox Catalysis for the Synthesis of 1,4-Dicarbonyl Compounds *via* Diacylation of Alkenes, *Org. Lett.*, 2022, **24**, 5710–5714.
- 47 L. Wang, J. Sun, J. Xia, M. Li, L. Zhang, R. Ma, G. Zheng and Q. Zhang, Visible light-mediated NHCs and photoredox co-catalyzed radical 1,2-dicarbonylation of alkenes for 1,4-diketones, *Sci. China: Chem.*, 2022, **65**, 1938–1944.
- 48 K. Liu and A. Studer, Direct  $\alpha$ -Acylation of Alkenes *via* N-Heterocyclic Carbene, Sulfinate, and Photoredox Cooperative Triple Catalysis, *J. Am. Chem. Soc.*, 2021, **143**, 4903–4909.
- 49 L. Wang, R. Ma, J. Sun, G. Zheng and Q. Zhang, NHC and visible light-mediated photoredox cocatalyzed 1,4-sulfonylacylation of 1,3-enynes for tetrasubstituted allenyl ketones, *Chem. Sci.*, 2022, **13**, 3169–3175.
- 50 L. Wang, J. Sun, J. Xia, R. Ma, G. Zheng and Q. Zhang, Visible light-mediated NHC and photoredox co-catalyzed 1,2-



- sulfonylacylation of allenes *via* acyl and allyl radical cross-coupling, *Org. Chem. Front.*, 2023, **10**, 1047–1055.
- 51 J. Reimler, X. Yu, N. Spreckelmeyer, C. G. Daniliuc and A. Studer, Regiodivergent C–H Acylation of Arenes by Switching from Ionic- to Radical-Type Chemistry Using NHC Catalysis, *Angew. Chem., Int. Ed.*, 2023, **62**, e202303222.
- 52 H. Yang, X. Jin, H. Jiang and W. Luo, Construction of C(CO)–C(CO) Bond *via* NHC-Catalyzed Radical Cross-Coupling Reaction, *Org. Lett.*, 2023, **25**, 1829–1833.
- 53 S. Ren, W. Lv, X. Yang, J. Yan, J. Xu, F. Wang, L. Hao, H. Chai, Z. Jin and Y. R. Chi, Carbene-Catalyzed Alkylation of Carboxylic Esters *via* Direct Photoexcitation of Acyl Azolium Intermediates, *ACS Catal.*, 2021, **11**, 2925–2934.

