



Cite this: *Environ. Sci.: Adv.*, 2023, 2, 1600

The emission of low pH water from Gulf of Mexico seeps as revealed by $\delta^{13}\text{C}$ - CO_2 and methane oxidation data†

Sydney I. Loudon * and John D. Kessler

Seawater was collected around the MC118 hydrocarbon seep in the Gulf of Mexico and was used previously for a study of aerobic methane (CH_4) oxidation. During that experiment, changes in the dissolved concentrations and $\delta^{13}\text{C}$ isotopes of CH_4 and CO_2 were recorded. Originally, the CO_2 concentrations and isotopes were recognized to qualitatively follow trends supporting the microbial conversion of CH_4 to CO_2 via aerobic oxidation, however, no attempt was made to quantitatively explain this CO_2 data. The present study models the $\delta^{13}\text{C}$ - CO_2 changes that occur as a result of CH_4 oxidation, accounting for the carbon already present as dissolved inorganic carbon (DIC), DIC added via CH_4 oxidation, and the pH of seawater. This study discovers that to accurately model the measured concentration and isotopic data for CO_2 , the seawater emitted from this seep site must have a pH which is at most between 6.49 and 7.24, and possibly up to 0.43 ± 0.08 pH units lower. These results are corroborated by direct measurements of pH from seeps in the Mediterranean Sea. A first-order extrapolation indicates that while cold seeps in the Gulf of Mexico may be a source of low pH water influencing the carbon dynamics of the deep ocean environment, this influence is likely less than that of current surface ocean acidification caused by the infiltration of atmospheric CO_2 .

Received 3rd May 2023
Accepted 9th October 2023

DOI: 10.1039/d3va00117b

rsc.li/esadvances

Environmental significance

Methane seeps have been widely studied to understand the amount and fate of released methane, a potent greenhouse gas. However, this study reveals that seeps also emit low pH water, broadening the environmental significance of seep emissions to include ocean acidification. Using samples from a Gulf of Mexico seep, measurements and models of aerobic methane oxidation were conducted. These analyses revealed the pH of water emitted from this seep must be between 6.49 and 7.24, much lower than the average ocean pH of 8.1. This low pH water likely has important, localized influences. But a first-order extrapolation suggests the potential rate of deep ocean acidification in this environment is likely lower than the rate of seawater acidification from atmospheric CO_2 .


Introduction

Seeps emit methane (CH_4) and other hydrocarbons from the seafloor and are concentrated along continental margins. One area with many seeps that has been well documented and studied is the Gulf of Mexico, which contains an estimated 914 natural hydrocarbon seeps.^{1,2} Both the abundance of hydrocarbons underneath the seafloor and the geology of the basin contribute to the number of seeps found in the Gulf of Mexico.³



Methane is emitted from seeps in the form of dissolved CH_4 and bubble streams⁴ which can dissolve into the water column during ascent.⁵ The rate at which these CH_4 bubbles dissolve into the surrounding water depends on a variety of factors including, but not limited to, bubble size and depth of

emission⁶ with substantial amounts of dissolution occurring near the seafloor.⁷

Once dissolved in the overlying waters, methanotrophic bacteria use CH_4 as a carbon and energy source.⁸ These methanotrophs are present in the water surrounding CH_4 seeps and aerobically oxidize CH_4 via the overall reaction shown in eqn (1).⁹



The CO_2 produced from aerobic CH_4 oxidation then mixes and equilibrates with the dissolved inorganic carbon (DIC) pool already present in the seawater (eqn (2) and (3)),



Department of Earth and Environmental Science, University of Rochester, Rochester, NY, USA. E-mail: slouden2@ur.rochester.edu

† Electronic supplementary information (ESI) available. See DOI: <https://doi.org/10.1039/d3va00117b>



where $[\text{CO}_2]$ represents the sum of dissolved $[\text{CO}_2]$ and $[\text{H}_2\text{CO}_3]$, and $[\text{DIC}]$ is the sum of the inorganic carbon components.

The specific rate of microbial oxidation of CH_4 is faster for the light isotopologue ($^{12}\text{CH}_4$) than the heavy isotopologue ($^{13}\text{CH}_4$), resulting in a kinetic isotope effect and producing measurable changes in the isotopic composition of the residual CH_4 . Thus, the CO_2 produced *via* CH_4 oxidation is enriched in the ^{12}C isotope compared to the residual CH_4 .¹⁰ These isotopic differences change as a function of the extent of oxidation and can be used to confirm that CH_4 oxidation is occurring. Additionally, these changes can quantitatively determine the extent of the starting CH_4 pool that has been oxidized and have been further used to determine the oxidation rate.¹¹

The investigation presented here began with data measured during an experiment exploring aerobic CH_4 oxidation in seawater.⁹ The experiment measured changes in the dissolved concentrations and $\delta^{13}\text{C}$ isotopes of CH_4 and CO_2 in a closed seawater incubation system experiencing significant amounts of aerobic CH_4 oxidation.¹² The goal of the present study was to develop an isotopic model to quantitatively explain changes in the measured $\delta^{13}\text{C}$ - CO_2 data as a function of the extent of CH_4 oxidation, the results of which discovered that seeps emit relatively low pH water alongside CH_4 . Variables included in this model were the isotopic fractionation associated with aerobic CH_4 oxidation, the background concentrations of CO_2 and DIC present in the system prior to this CH_4 oxidation event, and the added CO_2 from CH_4 oxidation. This investigation reveals that the model and measurements only agree at pH values significantly below those of the background deep ocean, indicating that seeps are a source of low pH waters to the deep ocean.

Experimental

Sample collection

The full details describing sample collection and analysis can be found in Chan *et al.*⁹ In brief, samples were collected in the Gulf of Mexico at site MC118 (28°51'N, 88°29.5'W) during a research expedition from 12–17 April 2015 onboard the E/V *Nautilus*. Samples were collected at depths of 794 and 888 m using the Suspended-Particle Rosette sampler¹³ mounted to the remotely operated vehicle (ROV) *Hercules*. Water samples were taken just above the seafloor in water visibly impacted by CH_4 bubbles. Other hydrocarbons, including oil, were present as well.⁹

The collected water samples were incubated at near *in situ* temperatures using a mesocosm incubation system developed by Chan *et al.*¹⁴ The mesocosm incubation system was connected to a dissolved gas analysis system which measured the concentrations and stable isotopes of CH_4 and CO_2 throughout the incubation period. Samples for DNA analysis and cell counts were isolated periodically during incubation to characterize the microbial communities and ensure methanotrophic bacteria were present.⁹

Isotope modeling procedure

Since this experiment incubated seawater samples in a closed vessel, the isotopic fractionation caused by CH_4 oxidation was modeled following closed system or Rayleigh fractionation

equations.¹² We note that other studies of CH_4 oxidation in open natural seawater environments still displayed isotopic fractionation following closed system kinetics.^{11,15} The closed nature of the incubation vessel meant that any CO_2 produced from CH_4 oxidation accumulated in the vessel. Thus, to model the $\delta^{13}\text{C}$ - CO_2 data, an accumulated product model was incorporated (eqn (4)).¹⁶

$$\delta_x = \frac{\delta_{\text{CH}_4,0} + 1000}{f} \left[1 - (1-f)^{1/\alpha} \right] - 1000 \quad (4)$$

Here, δ_x represents the isotopic values of the accumulated CO_2 produced from CH_4 oxidation, $\delta_{\text{CH}_4,0}$ represents an average of the isotopic values of CH_4 at the start of the incubation, and α represents the isotopic fractionation factor, defined as the ratio of the first-order rate constants for the oxidation of $^{12}\text{CH}_4$ and $^{13}\text{CH}_4$ ($\alpha = k_{12}/k_{13}$). Previously, Chan *et al.* confirmed that CH_4 oxidation followed first-order oxidation kinetics and that α ranged from 1.016 to 1.025.^{9,12} The variable f in eqn (4) represents the fraction of the starting CH_4 pool that has been oxidized. Here, we determine f using the CH_4 concentration data from Chan *et al.*^{9,12} as shown in eqn (5),

$$f = 1 - \frac{[\text{CH}_4]_i}{[\text{CH}_4]_0} \quad (5)$$

where $[\text{CH}_4]_i$ is the concentration of CH_4 measured at different times throughout the incubation period, and $[\text{CH}_4]_0$ is the averaged initial CH_4 concentration. To account for variability in the data prior to the onset of more rapid CH_4 oxidation, initial values of CH_4 and CO_2 concentrations and isotopes were averaged across several measurements.

While δ_x represents the isotopic values of the accumulated CO_2 produced during this CH_4 oxidation experiment, this CO_2 is added to a large pool of CO_2 and DIC initially present in the seawater sample. Thus, to accurately model the measured $\delta^{13}\text{C}$ - CO_2 data, a weighted isotopic average was used (eqn (6)).

$$\delta_{\text{DIC}} = \frac{[\text{DIC}]_B \delta_{\text{DIC},B} + \Delta[\text{CH}_4] \delta_x}{[\text{DIC}]_B + \Delta[\text{CH}_4]} \quad (6)$$

Here, δ_x represents the accumulated CO_2 from eqn (4). $\Delta[\text{CH}_4]$ represents the amount of CO_2 added to the incubation from CH_4 oxidation and was determined from $[\text{CH}_4]_0 - [\text{CH}_4]_i$. This definition assumes that all CH_4 oxidized is converted to CO_2 , which is likely an overestimate as some will be used to generate biomass. $\delta_{\text{DIC},B}$ and $[\text{DIC}]_B$ represent the isotopic composition and concentration of DIC, respectively, in the seawater before this CH_4 oxidation event. This mixing calculation is conducted with background DIC rather than background CO_2 to account for any equilibration between the newly produced CO_2 and the DIC system (eqn (2)). The value of $\delta_{\text{DIC},B}$ was determined from the initial $\delta^{13}\text{C}$ - CO_2 value measured using a mass balance relationship¹⁷ and experimentally determined isotopic fractionation factors.^{18,19} At a given temperature, the isotopic fractionation between the different inorganic carbon species is a function of pH.¹⁷ An initial pH was chosen to determine a possible isotopic offset between CO_2 and DIC. (A full description of the calculation of the isotopic offset is provided in the ESI.†) The value of $\delta_{\text{DIC},B}$ was determined by adding the



calculated isotopic offset value to the initial, measured $\delta^{13}\text{C}-\text{CO}_2$ value for each incubation experiment. On average, the values of δ_{CO_2} were 9.38‰ lighter than the values of δ_{DIC} .

$[\text{DIC}]_{\text{B}}$ was determined based on the dissolved CO_2 concentration measurements and the pH of the seawater sample. Combining eqn (3) with the equilibrium relationships of K_1 and K_2 (eqn (2)) produces an equation for $[\text{DIC}]_{\text{B}}$ as a function of CO_2 concentration and pH (eqn (7)),

$$[\text{DIC}]_{\text{B}} = [\text{CO}_2]_{\text{B}} \left(1 + \frac{K_1}{[\text{H}^+]} + \frac{K_1 K_2}{[\text{H}^+]^2} \right) \quad (7)$$

where $K_1 = 9.558 \times 10^{-7} \text{ mol kg}^{-1}$ and $K_2 = 5.521 \times 10^{-10} \text{ mol kg}^{-1}$ at the temperature and salinity of the incubation (7 °C and 35 ppt).^{20,21} The value of $[\text{CO}_2]_{\text{B}}$ is set equal to $[\text{CO}_2]$ at the start of the incubation experiment. The pH was chosen to be equal to the value used in the isotopic offset between δ_{CO_2} and δ_{DIC} .

This model calculates the isotopic value of DIC in the system as a mixture of the background DIC and the DIC added from CH_4 oxidation (eqn (6)). However, since values of δ_{CO_2} were measured rather than δ_{DIC} , the modeled values of δ_{DIC} (eqn (6)) must be reverted to values of δ_{CO_2} for comparison with the data. This is accomplished by subtracting the corresponding isotopic offset value calculated previously.

For each incubation experiment, the pH values incorporated into these calculations were varied until the residuals between the modeled values of δ_{CO_2} and the measured data were minimized.

Results and discussion

Data and results

The modeled and measured values of δ_{CO_2} for the four incubation experiments are shown in Fig. 1. Incubation experiments

S2 and S3 show the best agreement between the modeled and the measured values, especially during more rapid CH_4 oxidation, characterized by a large drop in $\delta^{13}\text{C}-\text{CO}_2$. In S2 and S3, a period of positive isotopic values is followed by a rapid change to negative isotopic values. This follows the expected values where $\delta^{13}\text{C}-\text{CO}_2$ is positive before more rapid CH_4 oxidation begins and quickly becomes negative during more rapid CH_4 oxidation. Incubation experiments S1 and S4 also show agreement between the data and the model, however, these samples follow a slightly different trend from S2 and S3. S1 and S4 both begin with negative isotopic values and display a steadier decline.

The pH values required to produce the isotopic models shown in Fig. 1 represent the pH at the beginning of the incubation experiment. The pH at the end of the incubation experiment was calculated using eqn (8).

$$[\text{DIC}]_{\text{F}} = [\text{CO}_2]_{\text{F}} \left(1 + \frac{K_1}{[\text{H}^+]} + \frac{K_1 K_2}{[\text{H}^+]^2} \right) \quad (8)$$

K_1 and K_2 are the same as in eqn (7). $[\text{CO}_2]_{\text{F}}$ is $[\text{CO}_2]$ at the end of the incubation experiment, and $[\text{DIC}]_{\text{F}}$ is $[\text{DIC}]_{\text{B}}$ plus the change in CH_4 , assuming all CH_4 removed was added to DIC *via* CH_4 oxidation. The pH values calculated at the beginning and end of each incubation experiment are shown in Table 1.

The average difference between final and initial pH values for all four incubation experiments is 0.1 ± 0.07 pH units. Overall, S1 is the most basic, and S3 is the most acidic. S2 shows no change in pH over the duration of the incubation experiment, while S4 shows the largest change. Data from all four incubation experiments suggests a pH considerably lower than the average ocean pH of 8.1.²²




Fig. 1 The modeled $\delta^{13}\text{C}-\text{CO}_2$ data (blue) compared to the measured values of $\delta^{13}\text{C}-\text{CO}_2$ (black) over the incubation time for all four samples collected at MC118.



Table 1 The estimated pH range for each of the four samples

Sample	Estimated pH range
MC118-S1	7.13–7.24
MC118-S2	6.67
MC118-S3	6.49–6.62
MC118-S4	6.69–6.84

Disagreement between the modeled and measured data can be attributed to the natural variation in the measured data, differences in initial CH₄ concentration, oxidation of other carbon compounds in the system, and/or analytical uncertainties. The variability in the measured data is not as present in the modeled values due to the averaging of initial data before it was used in the model. Each of the four incubations had different starting concentrations of CH₄. In S2 and S3, initial CH₄ concentrations were approximately 150 μM; in S1 and S4, initial CH₄ concentrations were approximately 50 μM. Methane oxidation rates were higher in S2 and S3 compared to S1 and S4. Higher initial CH₄ concentrations and faster oxidation rates indicate CH₄ oxidation was contributing more CO₂ to the DIC pool than in experiments S1 and S4. Oxidation of seep-derived dissolved organic carbon (DOC) would also add CO₂ to the DIC pool in these experiments.²³ Other geologic hydrocarbons were present in the incubations, which was visible at the sample collection site. Sequencing of the 16S rRNA gene at the end of each of the four incubations showed the presence of many species of hydrocarbon oxidizing bacteria, with CH₄ oxidizing bacteria making up a relatively low percentage of the microbial community.⁹

Discussion

A pH measurement was not taken at the sampling site of this dataset, however, there is strong evidence that the water directly above the CH₄ seep is relatively acidic. Isotope and concentration data suggest that the pH of the sample site is between 6.49 and 7.24, if not lower. One assumption made in modeling the isotope data was that all CH₄ removed during the incubation experiment was converted to CO₂. Based on eqn (1), two moles of O₂ should be removed for every mole of CH₄. Dissolved O₂ (DO) data from Chan *et al.* indicate a DO to CH₄ ratio less than 2 to 1.⁹ Chan *et al.* hypothesize that this discrepancy is due to the formation of biomass that has not been fully oxidized to CO₂ by the end of the incubation experiment.⁹ Thus, if we assume that only half of the CH₄ is fully oxidized to CO₂ while the rest remains as biomass at the conclusion of this experiment, an assumption supported by the DO to CH₄ ratio,⁹ the estimated pH of the water emitted from this seep is approximately 0.43 ± 0.08 pH units lower.

The pH of fluids emitted from hydrothermal vents has been well documented^{24,25} with the development of *in situ* measurement techniques,^{26,27} however, few measurements of pH at CH₄ seep sites have been published. The limited published data that exist indicate that water surrounding and coming from cold CH₄ seeps has an approximately neutral pH.^{1,28} Sisma-Ventura *et al.* report measurements of pH as low as 6.83 in the water

column above a hydrocarbon seep in the Southeast Mediterranean Sea with low pH values of 6.8 to 7.4 recorded throughout the water column up to 50 m above the seafloor.²⁸

The data and conclusions of Sisma-Ventura *et al.* support the findings of this study.²⁸ The measurement of a pH of 6.83 in the water above a hydrocarbon seep falls within the calculated pH range for the MC118 seep. The lower pH values calculated for MC118 could be attributed to the fact that water samples were collected directly above the seep emission *via* ROV whereas Sisma-Ventura *et al.* collected water up to 50 m above the seep using a carousel of Niskin bottles.²⁸

Sisma-Ventura *et al.* conclude that the low pH values measured above hydrocarbon seeps are a result of the oxidation of CH₄ and other hydrocarbons to CO₂ which is added to the DIC pool of the background bottom water, lowering the pH.^{28,29} Sisma-Ventura *et al.* hypothesize that hydrocarbon seeps have a substantial impact on bottom water chemistry.²⁸ The results of the present study support this conclusion and suggest that CH₄ seeps are a source of low pH water to the deep Gulf of Mexico.

To estimate the potential impact of this low pH water, the modeled results from this study were extrapolated to the entire Gulf of Mexico to estimate the amount of low pH water emitted and its influence on the total pH of the Gulf of Mexico. For these calculations, the Gulf of Mexico was modeled as a cylinder with a total volume of $2.434 \times 10^{15} \text{ m}^3$ and a height equivalent to the average depth, 1615 m.³⁰ It was assumed that all estimated 914 seeps in the Gulf of Mexico² emit the same low pH water at the same, constant rate. Estimates of water flux from Gulf of Mexico seeps range from 9.4 to 30 mm per year, and seep diameters from which water is emitted are estimated to be between 0.2 and 1.2 km.^{31,32} Maximum values were chosen for this calculation to prevent against underestimating the potential impact on bottom water pH. Therefore, the lowest pH value determined of 5.91 was used, which assumes half of the oxidized CH₄ remained as biomass. It was assumed that the background pH of the deep Gulf of Mexico was 8.1.²² This also maximizes the estimated pH impact as the effect of low pH water emitted from seeps would be reduced in less basic surrounding water. The amount of low pH water emitted by all seeps over the span of one year was then calculated, and the resulting pH changes were determined assuming this low pH seep water impacts (i) the total volume of the Gulf of Mexico, (ii) the bottom 50 m of the Gulf of Mexico, and (iii) the bottom 10 m of the Gulf of Mexico.

Any pH change was negligible assuming the total volume or the bottom 50 m were influenced by low pH water emitted from cold seeps. When we assumed that low pH seep water only impacts the bottom 10 m of the Gulf of Mexico, the pH decreased by 1.38×10^{-4} in one year. For comparison, the average yearly decrease in pH from surface ocean acidification from the infiltration of atmospheric CO₂ is 2.00×10^{-3} .²² Only if the water flow from all seeps in the Gulf of Mexico was higher by a factor of 10 would a similar decrease in pH be observed as in the surface waters, assuming only the bottom 10 m are impacted by seep water. A hypothetical increase in water flow by a factor of 100 would be necessary for the bottom 50 m to display a similar annual pH decrease to the surface waters. The



- 38 J. Blouet, S. Arndt, P. Imbert and P. Regnier, Are seep carbonates quantitative proxies of CH₄ leakage? Modeling the influence of sulfate reduction and anaerobic oxidation of methane on pH and carbonate precipitation, *Chem. Geol.*, 2021, 557, 120254.
- 39 B. B. Bernard, Methane in marine sediments, *Deep-Sea Res., Part A*, 1979, 26, 429–443.
- 40 R. Luff, K. Wallmann and G. Aloisi, Numerical modeling of carbonate crust formation at cold vent sites: significance for fluid and methane budgets and chemosynthetic biological communities, *Earth Planet. Sci. Lett.*, 2004, 221, 337–353.

