



# Modified minimal-size fragments of heparan sulfate as inhibitors of endosulfatase-2 (Sulf-2)<sup>†</sup>

 Cite this: *Chem. Commun.*, 2024, 60, 436

 Received 26th June 2023,  
 Accepted 3rd November 2023

DOI: 10.1039/d3cc02565a

rsc.li/chemcomm

 Alice Kennett,<sup>a</sup> Sven Epple,<sup>a</sup> Gabriella van der Valk,<sup>a</sup> Irene Georgiou,<sup>a</sup>  
 Evelyne Gout,<sup>b</sup> Romain R. Vivès<sup>ib</sup> and Angela J. Russell<sup>ib</sup>\*<sup>a,c</sup>

**Sulf-2 has been identified as a putative target for anticancer therapies. Here we report the design and synthesis of sulfated disaccharide inhibitors based on IdoA(2S)-GlcNS(6S). Trisulfated disaccharide inhibitor IdoA(2S)-GlcNS(6Sulfamate) demonstrated potent Sulf-2 inhibition. The IC<sub>50</sub> value was determined to be 39.8 μM ± 18.3, which is comparable to a tetrasaccharide inhibitor of HSulf-1 reported in the literature. We propose that the disaccharide IdoA(2S)-GlcNS(6S) is the shortest fragment size required for effective inhibition of the Sulfs.**

Endosulfatases (Sulf-1 and Sulf-2) are located in the extracellular matrix and are responsible for the selective desulfation of the sulfate group on the glucosamine 6-*O*-sulfate residues within heparan sulfate (HS) proteoglycans and have a strong substrate specificity for the [Glc/IdoA(2S)-GlcNS(6S)] trisulfated disaccharide (Fig. 1).<sup>1a,b</sup> The trisulfated disaccharide [Glc/IdoA(2S)-GlcNS(6S)] has a low abundance within HS, and therefore seemingly subtle modifications by Sulf activity result in major functional consequences.<sup>2</sup> This highlights the importance of Sulf activity and indicates how targeting the Sulfs could have significant downstream effects on HS-mediated processes. Sulf-2 inhibitors are putative anticancer therapeutics because the sulfs have been linked to the regulation of signalling pathways such as Wnt and FGF *via* the modulation of the 6-*O*-sulfation status of HS.<sup>3</sup> Sulf-2 expression is induced or upregulated in various cancers and its role has been identified as being pro-tumourigenic, with Sulf-2 gene silencing or knock-out leading to decreased tumour formation. Therefore, Sulf-2 inhibition has been identified as a potential therapeutic target for many cancers.<sup>4a,b</sup> For this reason, the development of endosulfatase inhibitors has gained attention over the past decade.

Scheilwies *et al.* reported glucosamine-based small molecule inhibitors substituting the 6-*O*-sulfate (–OSO<sub>3</sub><sup>–</sup>) with the sulfamate motif (–OSO<sub>2</sub>NH<sub>2</sub>). This preliminary work utilised the smallest, most relevant unit of HS, α-GlcNS(6S) to template inhibitor design.<sup>5</sup> The biochemical characterisation of this compound in a competition assay with fluorogenic substrate 4-methylumbelliferyl sulfate (4-MUS), revealed that the sulfamate inhibitor had an IC<sub>50</sub> values of 95 μM against HSulf-1 and 130 μM against HSulf-2, and importantly was more selective for the Sulfs than other sulfatases investigated. In 2015, Miller *et al.* aimed to replicate the inhibitory activity of the glucosamine-6-sulfamate inhibitors and develop a structure activity relationship. All compounds synthesised were found to have minimal inhibition of Sulf-2 at 1 mM.<sup>6</sup> However, there were some discrepancies in assay protocol between the two papers that may explain the different inhibition potencies reported, so the question remains of whether 1 is a true inhibitor of Sulfs. Recently, Chiu *et al.* reported the design and synthesis of di-, tri- and tetra-saccharide fragments of HS with the sulfamate modification as inhibitors of Sulf-1.<sup>7</sup> The disaccharide, GlcNS(6Sulfamate)-IdoA(2S) only caused 20% Sulf-1 inhibition at 0.7 mM (IC<sub>50</sub> value not determined), and the trisaccharide and tetrasaccharide analogues were more potent with IC<sub>50</sub> values of 0.53 and 29.6 μM, respectively.



**Fig. 1** Structure of HS highlighting the disaccharide residue, IdoA(2S)-GlcNS(6S), that Sulfs have a preference for.

<sup>a</sup> Department of Chemistry, University of Oxford, Oxford OX1 3TA, UK.

E-mail: angela.russell@chem.ox.ac.uk

<sup>b</sup> Univ. Grenoble Alpes, CNRS, CEA, IBS, Grenoble, France


<sup>c</sup> Department of Pharmacology, University of Oxford, Oxford OX1 3QT, UK

<sup>†</sup> Electronic supplementary information (ESI) available. See DOI: <https://doi.org/10.1039/d3cc02565a>






Scheme 2 Synthesis of inhibitor 4.



Scheme 3 Synthesis of inhibitor 5.

conditions reported by Miller *et al.*, to give sulfamate **21** in 66% yield. Multiple conditions were trialed for the global debenzoylation and deprotection of the amino-Cbz group, and the optimal conditions were found to be catalytic transfer hydrogenation using cyclohexene as the hydrogen donor with 20% Pd(OH)<sub>2</sub> in refluxing methanol.<sup>11</sup> Under these conditions, methyl ester **22** was isolated in 58% yield. Finally, the primary amine was sulfated using sulfur trioxide–pyridine complex in basic aqueous medium. Purification by ion-paired reverse-phase HPLC using 2 M triethylammonium bicarbonate and acetonitrile gradient, followed by elution through a Dowex<sup>®</sup> 50WX8 Na<sup>+</sup>-form column, gave **4** in 22% isolated yield.

It was originally envisioned that the synthesis of putative inhibitor **5** could diverge from the synthesis of **4**, *via* benzylation of intermediate alcohol **19**. However, all attempts at benzylation of the ido 4-OH of **19** were unsuccessful and therefore idose glycosyl donor **23** was synthesised according to Hu *et al.* (Scheme 3).<sup>12</sup> With the alternative ido-glycosyl donor in hand, the glycosylation reaction between **23** and glycosyl acceptor **16** was activated using TMSOTf and proceeded

effectively to afford the desired disaccharide **24** in 85% yield. Desilylation of the 6-O-TBDPS group of **24** using TBAF buffered in acetic acid proceeded to afford **25** in 64% yield. Subsequent sulfamoylation of the primary alcohol to **26** was achieved in 74% yield by altering the previous conditions to use 2 equivalents of sulfamoyl chloride at 0 °C. Subsequently, the labile protecting groups of compound **26** were removed by catalytic NaOCH<sub>3</sub> in CH<sub>3</sub>OH to produce diol **27** in 94% yield. The resulting diol **27** was then subjected to oxidation with the TEMPO/PIDA reagent system to afford lactone **28** in 58% yield. **28** was immediately hydrolysed in basic aqueous medium, and the resulting 2-OH moiety was treated with sulfur trioxide–pyridine complex under microwave irradiation. After elution through a Dowex<sup>®</sup> 50WX8 Na<sup>+</sup>-form column, **29** was isolated in 66% yield over two steps. Finally, the hydrogenolysis-labile protecting groups of **29** were cleaved by Pd(OH)<sub>2</sub>/C catalysed hydrogenation in methanol and aqueous phosphate buffered saline (20 mM, pH 7.4), to give a primary amine intermediate, which was successively subjected to sulfur trioxide pyridine



