



Cite this: *Chem. Commun.*, 2024, 60, 632

Unlocking novel therapies: cyclic peptide design for amyloidogenic targets through synergies of experiments, simulations, and machine learning

Daria de Raffele^{ab} and Ioana M. Ilie *^{ab}

Existing therapies for neurodegenerative diseases like Parkinson's and Alzheimer's address only their symptoms and do not prevent disease onset. Common therapeutic agents, such as small molecules and antibodies struggle with insufficient selectivity, stability and bioavailability, leading to poor performance in clinical trials. Peptide-based therapeutics are emerging as promising candidates, with successful applications for cardiovascular diseases and cancers due to their high bioavailability, good efficacy and specificity. In particular, cyclic peptides have a long *in vivo* stability, while maintaining a robust antibody-like binding affinity. However, the *de novo* design of cyclic peptides is challenging due to the lack of long-lived druggable pockets of the target polypeptide, absence of exhaustive conformational distributions of the target and/or the binder, unknown binding site, methodological limitations, associated constraints (failed trials, time, money) and the vast combinatorial sequence space. Hence, efficient alignment and cooperation between disciplines, and synergies between experiments and simulations complemented by popular techniques like machine-learning can significantly speed up the therapeutic cyclic-peptide development for neurodegenerative diseases. We review the latest advancements in cyclic peptide design against amyloidogenic targets from a computational perspective in light of recent advancements and potential of machine learning to optimize the design process. We discuss the difficulties encountered when designing novel peptide-based inhibitors and we propose new strategies incorporating experiments, simulations and machine learning to design cyclic peptides to inhibit the toxic propagation of amyloidogenic polypeptides. Importantly, these strategies extend beyond the mere design of cyclic peptides and serve as template for the *de novo* generation of (bio)materials with programmable properties.

Received 18th September 2023,
 Accepted 6th December 2023

DOI: 10.1039/d3cc04630c

rsc.li/chemcomm

^a University of Amsterdam, van 't Hoff Institute for Molecular Sciences, Science Park 904, P.O. Box 94157, 1090 GD Amsterdam, The Netherlands. E-mail: i.m.ilie@uva.nl

^b Amsterdam Center for Multiscale Modeling (ACMM), University of Amsterdam, P.O. Box 94157, 1090 GD Amsterdam, The Netherlands

1 Introduction

Neurodegenerative disorders, such as Alzheimer's, Parkinson's and Creutzfeldt-Jakob disease, affect over 50 million people world-wide, with over 10 million new cases a year. The world



Daria de Raffele

Daria De Raffele is a postdoctoral researcher in the group of Dr Ioana M. Ilie. She obtained her PhD from Universitat Jaume I where she focused on the QM/MM theoretical study of enzyme promistuity. Her research focuses on the design of cyclic peptides to inhibit the toxic propagation of the prion protein.



Ioana M. Ilie

Dr Ioana M. ILIE is an Assistant Professor in the Computational Chemistry group at the University of Amsterdam. The research in her group focusses on multiscale simulations of biomolecular systems. In particular, her team relies on rational design and multiscale simulations to (1) understand and control the aggregation mechanisms of polypeptides and their response to the biological environment, (2) to design of peptide-based therapeutics for neurodegenerative diseases and (3) to develop smart (bio)materials with responsive properties.





Fig. 1 (a) Ferulic aldehyde (MW 194 Da) inhibits the $A\beta_{42}$ multimerization. (b) Cartoon representation of the Aducanumab antibody (PDB ID: 6CO3⁴¹) (MW 146 kDa). Highlighted are the heavy chain (cyan), light chain (light blue). The area enclosed by the red circle represents the binding interface between the antibody and the N-terminus of the $A\beta_{42}$ peptide. (c) Naturally-occurring cyclotide kalata B1 (MW 2.92 kDa), derived from residues 306–311 of tau.⁴²

preventing amyloid formation and promoting the formation of “off-pathway” soluble oligomers and prefibrillar aggregates;^{65,66} it can also disrupt $A\beta_{40}$ fibrils and break down tau tangles.^{67,68} Additionally, the degradation products of curcumin – ferulic acid and vanillin – have a better solubility than curcumin. Ferulic acid can destabilize $A\beta_{40}$ and $A\beta_{42}$ fibrils⁶⁹ and vanillin can prevent amyloid aggregation in living organisms.⁷⁰ Likewise cholic acid, a primary bile acid, presents an anti-amyloidogenic behavior, inhibiting amyloid formation and preventing secondary nucleation.⁵⁹ Vitamin k3, known for its anti-inflammatory and anti-cancer properties, inhibits the $A\beta_{42}$ aggregation process and has a positive impact on reducing cytotoxicity in human neuronal cell line.⁶² We refer the interested reader to a comprehensive review of small molecule inhibitors for amyloid- β ⁷¹ for amyloidogenic polypeptides in general.⁷²

Antibodies are Y-shaped proteins of larger molecular weight than small molecules (>150 kDa), which can recognize and bind to protein targets with high specificity and modulate their toxic behavior.⁷³ In particular, monoclonal antibodies have been designed both for therapeutic and diagnostic applications. They bind to amyloidogenic polypeptides and/or their aggregates to stabilize a desired conformations and make the substrate unavailable for conversion^{10,43} and/or sterically interfere with the aggregation process.^{11–13} For instance, the DesAbs single-domain antibodies targeting $A\beta_{42}$ epitopes⁷⁴ interact with the monomeric peptide, bind with high affinity to the oligomeric species, but not the fibrillar structures, can inhibit secondary nucleation¹² and suppress $A\beta_{42}$ -mediated toxicity in *C. elegans*.⁷⁴ Aducanumab and lecanemab, approved anti- $A\beta_{42}$ agents, are monoclonal antibodies effective for patients in the early stages of AD due to their ability to reduce amyloid deposits in the brain.⁴³ Aducanumab (Fig. 1(b)) binds at the N-terminal residues 3 to 7 and can discriminate between monomers and aggregated species,⁴⁴ while lecanemab binds to soluble $A\beta_{42}$ protofibrils.⁷⁵ Cinpanemab and prasinezumab, two monoclonal antibodies directed against α -synuclein aggregates failed in clinical trials due to the lack of positive effects

in disease progression.^{16,17} The POM-family of antibodies (POMologues) has been developed to recognize a variety of epitopes along the sequence of the cellular prion protein¹⁰ and modulate its toxic effects.⁷⁶ Notwithstanding, no drug against prion diseases is currently in clinical trials. Despite the recent success in the Alzheimer's field with aducanumab⁴³ and the potential of gantenerumab,⁷⁷ antibodies have limitations as therapeutics, including stability and immunogenicity,^{14,15} which can impact clinical efficiency.

2.2 Cyclic peptides

Cyclic peptides are naturally occurring or chemically synthesized macrocycles consisting of circular sequences of amino acids,^{78–81} Fig. 1(c). Compared to small molecules, they can bind larger, more polar and solvent exposed protein surfaces.²⁰ To mimic naturally occurring cyclic peptides, the macrocycles can be experimentally synthesized from linear precursors by connecting their N- and C-terminal residues *via* covalent bonds (head-to-tail cyclization).⁸² Chemically, cyclization can be achieved through lactamization or *via* disulfide bond formation ensuring the link between the two termini of the linear precursor.⁸³ Head-to-tail cyclization restricts the dynamics of a peptide and can stabilize the formation of β -hairpins. Alternative approaches to favor other conformations such as α -helices can be achieved *via* stapling, *i.e.* *via* cross-linking of two or more side-chains. We refer the interested reader to detailed reviews on the synthesis of cyclic peptides⁸⁴ and stapled peptides,^{85,86} and focus in the following paragraphs on head-to-tail cyclized peptides.

Because of their physicochemical properties, cyclic peptides present a series of advantages as compared to their linear precursors, small molecules and biological therapeutic agents such as antibodies. First, the rigidity obtained through cyclization provides increased stability, higher resistance to proteolysis²⁷ and enhanced cell permeability as compared to linear peptides.^{20,87,88} Second, their size and functional properties ensure that the contact area is large enough to provide high selectivity, and their ability to form salt-bridges and hydrogen bonds can lead to strong binding affinities.²⁶ Hence they can maintain a robust antibody-like binding to (undruggable) interfaces with high affinity,^{20,79,89} due to their larger surface and implicitly the higher number of hydrogen bonding partners. Third, cyclic peptides have good *in vivo* stability, which contributes to enhanced retention and circulation, particularly if they are rich in non-canonical amino acids.²⁰

Cyclic peptide-based therapeutics also face a series of challenges. Orally administered cyclic peptide-rich drugs struggle with poor oral bioavailability,⁷⁸ because of the susceptibility of cyclic peptides to resist proteolytic degradation in the gastrointestinal tract.²⁷ Nevertheless, different routes of administration, such as subcutaneous or *via* intravenous injections, overcome these difficulties and aid in the efficient delivery of the peptide-drug to the target.⁸⁴ Another obstacle involves preventing off-target interactions, a challenge often tackled by selectively modifying natural amino acids in the sequence to non-natural ones.⁸⁴



Highlight

In the amyloid world, the cyclic peptide development has been growing over the past decade.^{30,78,80} Typical methods involve designing peptides rich in aromatic moieties, hydrophobic amino acids, or D-amino acids (due to the stereoselectivity for L-amino acids of proteases) that disrupt the aggregation process, *i.e.* β -breakers or agents that bind to monomeric and oligomeric species, competing with the responsible polypeptide to hinder its aggregation and/or toxic transformation.⁹⁰ For instance, the RD2D3 D-peptide (H-ptlhthnrrrrrrprtrllhthnr-NH₂[†]), designed to modulate the binding of PrP^C to A β ₄₂ oligomers, interferes with A β ₄₂-PrP^C heteroassembly in a concentration-dependent manner.⁹¹ Its cyclic successor presents better *in vitro* potency and pharmacokinetic properties⁹² and could potentially alter A β ₄₂ aggregation. The bicyclic DesBP peptide (RAACKLGIKACTSVYHACGGKRR) was rationally designed to bind monomeric A β ₄₂ at residues 31–36 and 38–42^{24,93} and was shown to alter the morphology of A β ₄₂ aggregates in a dose dependent manner. In particular, higher peptide concentrations lead to increased aggregate disorder and reduced cytotoxicity.⁹³ Similarly, the BD1 cyclic peptide (O-ySGLIKWTTALLR[‡]TYC-NH₂) was shown to inhibit α -synuclein fibril formation *in vitro*.⁹⁴ The D,L- α -peptide CP-2 cyclic peptide (IjwHsK[‡]) prevents α -syn aggregation into toxic oligomers by an “off-pathway” mechanism.⁹⁵ Particularly, it interacts with the N-terminus and the non-amyloidogenic region, altering the protein’s membrane interaction properties and fibril morphologies, thereby preventing the toxic membrane disruption. The macrocyclic inhibitory peptides (MCIPs), were designed to bind to amyloids by mimicking human IAPP (hIAPP) interaction surfaces while maintaining only minimal hIAPP-derived self-/cross-recognition elements.⁹⁶ Inhibitor selectivity was tuned by chirality, which lead to nanomolar binding affinities to hIAPP, to both amyloid- β ₄₀ and amyloid- β ₄₂ peptides, high proteolytic stability in human plasma and human brain-blood-barrier crossing ability.⁹⁶ Also, disulfide-rich macrocyclic peptides are versatile scaffolds for stable biochemical tool development. Two examples are SFTI-1 (GRCTKSIPPICFPD, disulfide connectivity: Cys3–Cys11), a cyclic peptide that inhibits trypsin, and the kB1 cyclotide (GLPVCGETCVGGTCNTPGCTCSWPVCTR[‡]N, disulfide connectivity: Cys5–Cys19, Cys9–Cys21 and Cys14–Cys26), which have an inherent ability to inhibit the fibril growth of the tau-derived hexapeptide Ac-VQIVYK-NH₂ (AcPHF6).⁴² Particularly, kB1 is a stronger inhibitor of tau fibrillization than SFTI-1, enabling better binding and/or disruption of AcPHF6 fibrils. Recently, tau mimetic peptides (β -bracelets) have been designed starting from the high-resolution structure of the tau fibril fold by extracting β -strand sequences linked by β -arcs.⁹⁷ The newly generated peptides self-assemble into parallel β -sheet fibrils and can serve as templates for the design of soluble inhibitors of tau seeding.

In terms of the cellular prion protein, no progress has been made on the therapeutic cyclic peptide market, despite its well defined secondary structure in the soluble form. Potential causes are the lack of druggable pockets or a stable unique binding region in the globular domain, and the intrinsically disordered nature of the tail. Though, the existence of

monoclonal antibodies that bind in the nanomolar regime to PrP^C indicate that putative interaction sites are available.¹⁰ We hypothesise that the rational design of cyclic peptides starting from available high resolution structures of PrP^C in complex with monoclonal antibodies may serve as starting points for the design of cyclic peptides that can potentially stabilize the soluble isoform of the protein and therefore prevent its toxic transformation. Alternatively, by tweaking the environmental conditions through mild solvent alteration, *e.g.* by replacing water with D₂O⁹⁸ or by adding organic compounds,^{99,100} one can delicately alter the conformational landscape of the protein to reveal new (allosteric) druggable pockets without disturbing the protein’s secondary structure. We refer the interested reader to a series of reviews on peptide-based strategies to interfere with protein misfolding and aggregation,^{101,102} a review on the therapeutic potential of cyclic⁹⁰ and bicyclic peptides.¹⁰³ Studies older than 10 years focusing on anti-amylin cyclic peptides and peptide-based inhibitors have been reviewed elsewhere.¹⁰⁴

3 Design methods and pitfalls

3.1 Conventional peptide design approaches

To design a soluble peptide-based binder with simultaneously high target specificity, binding affinity, cell permeability and proteolytic stability requires prior knowledge of the molecular target and its environmental conditions. Experimentally, genetically encoded methods such as phage display¹⁰⁵ or mRNA display^{106,107} allow the generation of libraries of cyclic peptides that bind with high affinity to the target.^{81,108} While these libraries offer the generation of a vast array of molecules, the chemical synthesis step as well as the numerous experimental trials are time and resource consuming. Furthermore, translating cyclic-peptide hits obtained through display technologies into clinical applications has proven challenging due to potential shortcomings in their pharmacological properties, including limited oral bioavailability, cell permeability, and solubility.¹⁰⁸ Other approaches, such as directed evolution mimic the natural evolution process of a peptide by creating a diverse library to screen for mutants with improved characteristics.¹⁰⁹ Directed evolution does not require information on the structure–function relationship of the substrate, and relies on an iterative procedure of random mutations and artificial selection to discover new and useful proteins, but is limited by the exhaustive pool of possibilities to be tested.

Over the past years, rational design approaches for *de novo* peptide design have gained momentum. Rational design relies on a detailed understanding of the amino acid sequence, protein high resolution structure, function and interaction mechanisms.³³ It involves the identification and mutation of key residues associated with protein stability to improve targeted physical and catalytic properties.^{33,110–112} Rational design relies on human intervention, which often offers an informed and efficient means to narrow down the search space for amino acids, resulting in a smaller and more manageable pool of

[†] Small letters indicate D-enantiomeric amino acids.

[‡] J is the norleucine amino acid.



effective peptides. *De novo* rational cyclic peptide design requires (a) high resolution three dimensional structures and biochemical/biophysical information of the target protein, and/or (b) detailed information of the ligand properties (*e.g.* hydrogen bonding abilities to the target, hydrophobicity, cyclization chemistry, existence of natural and non-natural residues) and conformations (*i.e.* the designed peptide may assume different conformations in the bound and unbound states).¹¹³ Recent advancements in cryo-electron microscopy (cryo-EM) have enabled the determination of the three-dimensional (3D) high resolution structures of new amyloidogenic aggregates and their monomeric precursors.¹¹⁴ These 3D structures corroborated with a comprehensive understanding of molecular interactions and structure–function relationships could enable the rational design of (cyclic) peptides tailored for amyloidogenic targets. As a matter of fact, rational design has been successfully used to generate the DesAbs antibodies targeting amyloid- β ¹² or specific α -synuclein and hIAPP epitopes.¹¹⁵

Starting from the high resolution structures of amyloid fibrils of tau, α -synuclein, and amyloid- β , miniproteins, ranging from 35 to 48 residues, were successfully designed to bind to the fibrillar tips of the targets and inhibit aggregation *in vitro* and *in vivo*.¹¹⁶ First a library of peptide-based inhibitors was created using Rosetta. Subsequently, Rosetta's MotifGraft protocol¹¹⁷ was used to dock the inhibitors onto the fibrils and energy minimized. The top-ranking inhibitors, *i.e.* the best binders, were subjected to molecular dynamics simulations to assess the stability of the complexes. Lastly, Rosetta's *ab initio* structure prediction algorithm¹¹⁸ was employed for the final screening of inhibitors. Inhibitors with the most favorable energy predictions and the smallest root mean squared deviations from the original design were selected for experimental validation.

From a computational perspective, virtual screening allows fast screening of millions of compounds prior to experimental testing, thereby reducing cost and saving time. Virtual screening using cyclic peptides is limited by the availability of three-dimensional structures of the targets, by the absence of drugable pockets and by the lack of information on the structure of the designed cyclic peptide. To overcome some of the limitations, different computational techniques have been combined with machine learning to predict protein structures and complexes thereof. Notable examples include HADDOCK (High Ambiguity Driven protein–protein Docking),^{119–121} RosettaFold¹²² and AlphaFold2.^{123,124} HADDOCK uses biochemical and biophysical interaction data, such as nuclear magnetic resonance titration experiments or mutagenesis data, to facilitate the protein–protein docking process.¹¹⁹ Recent developments include the generation of cyclic peptide conformations and docking to the protein target using knowledge of the binding site on the protein side to drive the modeling.¹²⁵ AlphaFold2 is a deep-learning algorithm that incorporates neural network architectures inspired by the physical and geometric aspects of protein structures.¹²⁶ It employs insights from evolutionary conservation through the analysis of multiple sequence alignments. These alignments are generated by considering information from evolutionary related proteins, along with

the 3D coordinates of a few homologous structures known as templates. Similarly, RoseTTAFold also utilizes multiple sequence alignments and a set of initial templates to accurately predict folded structures¹²² and protein–protein complexes.^{40,122} These technological advancements contribute significantly to the prediction protein structures through computational means. The intrinsic disorder associated with amyloidogenic polypeptides implies that the target protein lacks a stable structure and that its native state is better described by a diverse conformational ensemble rich in disordered structures.^{2,127,128} In this context, AlphaFold2 fails to predict such conformations, which often gives rise to unrealistic structures that do not accurately capture the states in the ensemble^{127–129} (Fig. 2). The lack of realistic and physically accurate ensembles of structures hampers the design of any type of inhibitor, which represents a limitation of these novel deep learning techniques.

3.2 Molecular dynamics simulations in peptide design

Computer simulations are powerful tools that enable the generation of quantitative conformational ensembles for the target (intrinsically disordered) protein with properties comparable to experimental results.^{127,128,130} Moreover, molecular dynamics simulations can go beyond experimental resolution to provide valuable insights into the stability and dynamics of cyclic peptides,¹³¹ structural detail on their membrane permeability,¹³² as well as quantitative distributions of their target-bound and target-unbound states.¹²⁸

Recently extensive molecular dynamics simulations at full atomistic resolution (Table 1) have been used to successfully identify transient monomeric A β ₄₂ conformations that have characteristics of fibrillar structures.¹³³ States of the monomeric, dimeric, oligomeric and fibrillar amyloidogenic polypeptides have been thoroughly characterized and have been reviewed elsewhere.^{2,128,130} The identified pool of structures could be potentially used for small molecule or cyclic peptide docking and design. Ideally, access to a well organized, reliable, and consistently maintained database of molecular dynamics trajectories of amyloidogenic polypeptides would avoid the repeated generation of similar trajectories and enable more



Fig. 2 AlphaFold2 predictions for (a) amyloid- β ₄₂, (b) α -synuclein, (c) hIAPP, (d) PrP^C and (e) tau. The structural elements are color-coded according to the confidence level of their AlphaFold2 prediction, red to blue for low to high confidence intervals, respectively.



Table 1 Computational studies of amyloidogenic polypeptides in complex with different agents

Ref.	Peptide	Agent	Model	Solvent	Method	Sampling ^a
Barz <i>et al.</i> ¹³³	A β ₄₂ monomer	—	Charmm36m	TIP3P	H-REMD	40.8 μ s
Jakubowski <i>et al.</i> ⁵²	A β ₄₂ fibril	94 small molecules	Charmm36m	TIP3P	MD	10.4 μ s
Dehabadi <i>et al.</i> ⁵⁰	A β ₄₂ dimer	Ferulic aldehyde	Charmm36m	TIP3P	MD	2.6 μ s
		Vanillin				2.6 μ s
Sciacca <i>et al.</i> ⁵¹	A β ₄₀ monomer	SilA, SilB	Charmm36	TIP3P	MD	3 μ s
Garcia-Vinuales <i>et al.</i> ⁵³	hIAPP monomer	—	Charmm36m	TIP3P	MD	6 μ s
		SilA				6.5 μ s ^b
		SilB				6.5 μ s ^b
Dubey <i>et al.</i> ¹³⁶	hIAPP fibril	Myricetin	Amber99sb	TIP3P	MD	1.05 μ s
Chen & Krishnan ¹³⁷	p53-NTD	EGCG	OPLS-AA 2005	TIP4D	MD	500 ns
Frost & Zacharias ⁴¹	A β ₂₋₇	AduFab	Charmm22*	TIP3P	MD	500 ns
	A β ₄₂	AduFab				1 μ s
	A β ₄₂ dimer					781 ns
	A β ₄₂ hexamer					1 μ s
	A β ₄₂ fubril					254 ns
Ilie & Caflisch ²⁸	PrP ^C	—	Charmm36m	TIP3P	MD	5 μ s
		POM1			MD	5 μ s
	PrP ^C	POM6			MD	5 μ s
Ilie <i>et al.</i> ²⁹	PrP ^C	—	Charmm36	ABSINTH	MC/MD	4.8 μ s MD + 240M MC
	PrP ^C	POM1			MC/MD	4.8 μ s MD + 240M MC
	PrP ^C	POM6			MC/MD	4.8 μ s MD + 240M MC
Kalmankar <i>et al.</i> ¹⁴⁰	A β ₄₂ monomer	Cter-M cyclotide	OPLS3e	TIP4P	RESPA	900 ns
	A β ₁₇₋₄₂ pentamer					900 ns
	A β ₁₁₋₄₂ fibril					900 ns
	A β ₄₂ double fibril					900 ns

^a Cumulative sampling over all replicas. Abbreviations. MD, molecular dynamics; H-REMD, Hamiltonian replica exchange molecular dynamics; MC/MD, hybrid Monte Carlo/molecular dynamics. RESPA, reversible multiple time scale molecular dynamics. ^b Two sets of simulations at different concentrations.

rapid and consistent advancement in amyloid-related drug discovery. Example of such a publicly available database is the *Molecular Dynamics Data Bank*. The *European Repository for Biosimulation Data*.

For small molecule docking, snapshots from molecular dynamics simulations of the A β ₄₂ monomer,¹³⁴ dimer⁵⁰ or multimers^{52,135} have been clustered to generate representative ensembles to be prepared for docking, which can be experimentally validated.⁶² Briefly, curcumin and a set of curcumin derivatives were docked onto A β ₄₂ multimeric conformations generated with molecular dynamics simulations.^{50,52} Results revealed that the small molecules interact with high probability with the amyloidogenic driving domains ¹⁶KLVFF²⁰ and ²⁹GAIG³³ of A β ₄₂ and disrupt their secondary structure in the hexameric⁵² and dimeric arrangements.⁵⁰ Interestingly, Silybin A (Sil A) and Silybin B (Sil B), two diastereoisomers of silybinin were shown to have different interaction preferences to A β ₄₀ and distinct biological response.⁵¹ Sil A binding the aromatic residues F19 and F20 slowed down aggregation, while Sil B interacting primarily with the C-terminus of the polypeptide fully abolished amyloid aggregation. Compelling evidence suggests that Silybin B is a powerful inhibitor also against the toxic self-assembly of hIAPP.⁵³ Simulation and experimental work, revealed that the frequent interactions of Sil B with the S20–S29 sequence induces disorder in the amyloidogenic core and attenuates hIAPP toxicity and aggregation propensity.⁵³ Myricetin, another polyphenolic flavonoid was shown to bind hIAPP at the amyloidogenic core and its C-terminus preventing

aggregation and distorting the fibrils.¹³⁶ The differential binding score (DIBS) was introduced to determine the binding preferences of ligands to an ensemble of IDP conformations by comparison against random coil ensembles of the same protein extracted from MD simulations.¹³⁷ The validation was performed on epigallocatechin-3-gallate (EGCG) binding to the unstructured N-terminus of the tumor suppressor p53 protein, which compared favorably to experimental results. The predictive ability of simulations has been demonstrated in a translational study, in which atomistic simulations were used to design new polythiophene derivatives against prion aggregation, prior to *in vivo* testing.⁵⁴ The compounds subsequently showed substantial prophylactic and therapeutic potency in prion-infected mice. Hence, simulations are powerful tools to generate conformational ensembles of the target polypeptide, which can act as scaffolds for the docking and design of molecules to target specific amyloid-forming regions.

The effects of antibodies on the structural and dynamic properties of amyloidogenic polypeptides have also revealed valuable insight into their modulating properties. Specifically, molecular dynamics simulations of Aducanumab in complex with A β ₄₂ revealed that the antibody sterically binds to monomeric, oligomeric and fibrillar species, with the binding site at the N-terminus (residues 2–7) preserved across all systems.⁴¹ Additionally, the results showed that the monomer unfolds and hydrophobically collapses on the antibody's surface, while in the complexes with aggregated species, the β -sheet structure of the peptide remains conserved.⁴¹ All-atom simulations of PrP^C



in complex with the neurotoxic POM1 and the innocuous POM6 antibodies revealed that the two antibodies, despite targeting similar epitopes, modulate differently the intrinsic flexibility of the protein²⁸ and its orientation with respect to the cellular membrane.²⁹ The information extracted from the simulations of amyloidogenic polypeptides in complex with antibodies could serve as starting points for the optimization and design of agents (*e.g.* antibodies, peptides) to bind with higher affinity towards selected species or for the rational design of cyclic peptides to modulate the target's conformational landscape enabling access to new binding sites.

Aside from the structure and the conformational landscape of the target, the conformations of the designed cyclic peptide in the target-bound and target-unbound states play an important role. Essentially, the designed cyclic peptides often adopt different conformations in solution as compared to the target-bound state. To design an efficient peptide-based inhibitor one needs to understand the conformational transitions of the cyclic peptides between the different states. While some peptide-protein complex structures are available, obtaining high resolution structures of cyclic peptides in solution is hampered by their low core-to-surface ratio, absence of specific couplings (*e.g.* NH-H α) and diverse conformations in solution.¹³¹ Hence, molecular dynamics simulations have been successfully used to predict the energetically relevant conformational ensembles of cyclic peptides in solution, which compare favorably to available experimental data (*e.g.* NMR chemical shifts).¹³⁸ We refer the interested reader to a comprehensive review of computational methods to characterize the behavior of cyclic peptides in solution¹³¹ and underline the synergistic effects of experimental and computational works.

Regarding the implications to the cyclic peptide design aspect, molecular dynamics simulations exceed experimental resolutions and can provide insight into the structural interactions between the peptide and the target at atomistic level of detail. For instance, macrocyclic peptides found in plants (cyclotides) have been experimentally shown to inhibit the aggregation of tau and amyloid- β_{42} fibrils.¹³⁹ The peptide was subsequently docked onto 3D structures of A β_{42} fibrils and subjected to molecular dynamics simulations.¹⁴⁰ The results explained experimental observations to reveal that the Cter-M cyclotide from *C. ternatea* (GLPTCGETCTLGTCYVPDCSCSW-PICMKN) binds the A β_{42} fibril *via* hydrogen bonding, hydrophobic, electrostatic and π - π interactions, thereby inhibiting aggregation.¹⁴⁰ Particularly, the peptide disrupts intermolecular hydrogen bonds and salt bridges in the A β_{42} fibril, which are crucial for its structural integrity. The effects occur within the first 50 ns of the simulations with disruptions in the fibril secondary structure at residues 2–7 and 38–41, resulting in the loss of extended β -sheet conformations. Importantly, the A β_{42} fibril in absence of the peptide maintains stable extended β -sheet conformations throughout the simulation trajectory.

Other approaches rely on available high resolution structures of protein complexes to identify linear interface motifs with an appropriate distances between residues to facilitate subsequent cyclization.¹⁴¹ In particular, backbone motifs of

epitopes within protein-protein interfaces were identified and compared against available cyclic peptide databases to pinpoint promising candidates with desired structural features.¹⁴¹ Subsequently, the generated cyclic peptide-protein complexes underwent refinement through molecular dynamics simulations in explicit solvent to determine the binder with the highest target affinity. To validate the efficacy of this method, initial tests were conducted on a complex formed by the bovine trypsin inhibitor (BPTI) protein and the trypsin protease. The method identified a cyclic peptide that resembled the BPTI protein backbone at the interface, which is in agreement with experimentally known structures.

Despite extensive simulations, challenges remain when exploring the conformational space of IDPs both in the presence and in absence of modulators.¹⁴² Convergence is an issue due to the rugged free energy landscapes of the polypeptides, their size (which at times imposes the use of large simulation boxes) and/or kinetic traps. Some of these difficulties are overcome by using enhanced sampling techniques, implicit solvents and/or coarse-grained models, which together with advances in computing power enable the access to longer time- and length-scales. Alternative approaches, include reducing the size of the system by simulating fragments of the polypeptide of interest and using statistical mechanics tools to derive the conformational free energies of the full IDP.¹⁴³ Current force fields struggle with over- or underestimating the properties of an IDP as compared to experimental quantities. Here, the IDP-tailored choices are the all-atom additive Charmm36m¹⁴⁴ and Amber ff14IDPSFF,¹⁴⁵ which have been fine tuned to reach experimental agreement and improve the conformational sampling of intrinsically disordered proteins.¹⁴⁶ More recently, machine learning has been integrated into the development and improvement of force fields^{147,148} and novel techniques are emerging for IDP-specific force fields. An example is Charmm-NN, which uses atom-type based neural networks to calculate energies and forces¹⁴⁹ and is subject to further improvements. A detailed overview of the challenges associated with IDP simulations and their reconciliation with experimental data have been reviewed in ref. 2 and 142. On the methodology side, the determination of the binding free energies of the cyclic peptide to the target also require special attention. For instance, using perturbation free energy calculations, a popular method with small molecules, one can determine the relative binding free energies and mechanistic detail, while preserving the flexibility of the complex.¹⁵⁰ Nevertheless, the convergence still remains an issue. Alternatively, umbrella sampling, a technique that provided valuable insight into the thermodynamics of monomer attachment to amyloid fibrils,^{143,151,152} would be a suitable choice for the determination of the binding free energies of a peptide to the target.

3.3 Computational methods for cyclic peptide design

Various computational methods have proven essential in the design of cyclic peptides for amyloidogenic targets. TANGO,¹⁵³ an algorithm developed to identify amyloidogenic sequences in proteins, was used to guide the search for cyclic peptides with improved binding affinity to A β_{40} oligomers.¹⁵⁴ Residues 102–



Highlight

117 (PRRYTIAALLSPYSWS) from the G strand of protein trans-thyretin (TTR) were used as starting point to generate a peptide that binds A β ₄₀ and redirects it towards protease-sensitive, nonfibrillar aggregates. The peptide was head-to-tail cyclized and TANGO was used to select specific mutations that would retain or stabilize the antiparallel two-stranded β -sheet, resulting in a series of cyclic G (CG) peptides. Out of the five newly synthesized peptides, CG8 (TKVVTpPRYTIAKLSSPYSYSQ)§ had the most pronounced affinity towards A β ₄₀, results confirmed by ThT fluorescence analyses.¹⁵⁴ Cyclization of CG8 *via* a disulfide bond using the simple cyclic peptide application (SCPA)¹⁵⁵ within ROSETTA and the addition of a second D-proline (TKVVTpPRYTIAKLSSpPSYSQ) lead to increased peptide stability, enhanced conformational uniformity, and a higher β -sheet content.¹⁵⁶ These findings highlight the added value of cyclization and conformational homogeneity as design strategies.

Des3PI (design of peptides targeting protein–protein interactions) is a novel computational fragment-based approach for designing cyclic peptides with high target specificity.¹⁵⁷ This algorithm performs docking calculations of an amino acid library onto the targeted protein surface and subsequently connects residues with favorable target binding affinities to generate novel cyclic peptide sequences and structures. We envision that the potential of this method can be exploited to the maximum when combined with quantitative representations from molecular dynamics simulations to generate novel amyloid-binding cyclic peptides.

Among the computational methods employed in designing peptides, FoldX emerges as a powerful tool due to its ability to determine the free energy contributions of each atom at protein interfaces based on its own position relative to neighbours in the complex.¹⁵⁸ It can thereby predict the impact of mutations on protein stability and optimize protein sequences for improved stability and desired functional properties. Relying on FoldX to perform an exhaustive thermodynamic profiling, the tandem peptide CAP1 was designed to inhibit tau aggregation.¹⁵⁹ Both *in vitro* and *in vivo* experiments confirmed computational predictions by showing that CAP1 binds with high specificity and affinity (EC₅₀ = 145 ± 49 nM) to tau aggregates, impeding their spread within cells. Additionally, CAP1 proved effective in hindering the ability of tau polymorphs obtained from the brains of Alzheimer's disease patients to initiate aggregation.

4 Machine learning for cyclic peptide design, property and activity prediction

Machine learning enables the rapid *in silico* screening and development of small molecules with therapeutic applications.³⁶ On the peptide engineering side, machine learning has found recent applications in antibody optimization^{160,161} and enzyme evolution.³⁴ The potential for cyclic peptide design is still in its initial stages and requires accurate and reliable training data. For

instance, using conformational ensembles from molecular dynamics, data can be generated and incorporated in training sets to create machine learning models able to accurately predict structural ensembles of peptides and their complexes or to generate peptide sequences with improved physico-chemical properties. For improved performance and increased accuracy, experimental data (*e.g.* binding, toxicity) can be incorporated.

In fact, by using data from molecular dynamics simulations of cyclic pentapeptides with diverse sequences and structural attributes as training datasets, machine learning models have been trained to predict structural ensembles for novel cyclic-peptide sequences, a method known as structural ensembles achieved by molecular dynamics and machine learning (StrEAMM).¹⁶² Alternative methods rely on generating comprehensive training datasets comprising sequences of blood–brain barrier penetrating linear peptides (BBPs) sourced from established databases and scientific literature, alongside non-BBPs peptides from UniProt to predict and explore novel BBPs with improved properties.¹⁶³ AbDiffuser introduced a diffusion model tailored for the generation of three-dimensional antibody structures and corresponding sequences for biotechnological applications.¹⁶⁴ Large protein families can be reliably mapped to a sequence ordinate using sequence alignment. AbDiffuser is an equivariant diffusion model designed to take advantage of these properties. The model adheres to physics-based constraints (*e.g.* bonds, torsional angles) and can accommodate different sequence lengths, thereby reducing the memory complexity. AbDiffuser relies on the Aligned Protein Mixer (APMixer), a neural network operating within the SE(3) equivariance framework to ensure consistent behavior, when subjected to rotations and translations in the three-dimensional space. Validation of the predictions through *in silico* and *in vitro* work underlines the importance of computational and experimental synergies when designing molecules with tailored properties.

Within the landscape of neurodegenerative diseases, MobiDB emerges as a resource that provides a comprehensive view of polypeptide disorder.¹⁶⁵ This repository compiles an array of comprehensive data related to intrinsically disordered proteins (IDPs) and regions (IDRs) encompassing both experimental and computational information on protein disorder, (*e.g.* sequences, structures, and functional annotations). Its utility is extended to experimental scientists seeking detailed information about individual protein systems, as well as bioinformaticians who require substantial, unified protein datasets for building statistical classifiers. More recently, MobiDB integrates AlphaFold predictions sourced from AlphaFoldDB.¹⁶⁶

A recent study highlighting the synergy between modern computational techniques and experiments, focused on developing a versatile method for designing proteins capable of targeting specific peptide sequences derived from armadillo proteins.¹⁶⁷ Using no known structure, Monte Carlo simulations were employed to construct a hash table for bidentate side-chain-backbone interactions, to ensure the stability of the desired protein–peptide interface. Identified key residues were optimized using Rosetta to construct both the protein and

§ Small letters indicate D-enantiomeric amino acids.



Highlight



Fig. 3 Schematic overview of the proposed *de novo* peptide design strategy against amyloidogenic targets. The first steps involve the preparation of the structures, the identification of quantitative conformational distributions of the target (top left panel) and the binder (bottom left panel). After preparing the two structures, the peptide can be docked onto the monomeric or multimeric target to determine the structure of the complex. Alternatively, the complex can be rationally designed starting from high resolution structures (middle panel). Next, the binding of the peptides is computationally and experimentally tested. Importantly, the extracted data (e.g. binding free energies, kinetics) can be incorporated in feedback loops powered by machine learning engines (e.g. active learning cycle) to improve the peptide sequences and/or properties. After several cycles, the best binders are advanced into pre-clinical validation. Created with BioRender.com.

enable to extract the statistically relevant states of the target, to be used in subsequent steps.

Second, quantitative characterization of the conformations populated by the cyclic peptides in the target-bound *versus* target-unbound states are factors to be accounted for. The size of the peptides (below 20 residues) and their cyclic nature often limits their structure generation or even the complex prediction *via* deep learning approaches such as AlphaFold-Multimer¹⁸² or *via* experimental techniques.¹³¹ Assuming that the initial peptide–protein complex is known, e.g. from crystal structures¹⁸³ or from *de novo* design,¹⁸⁴ one can isolate the peptide from the complex and explore its conformational space in the unbound state *via* (enhanced sampling) molecular dynamics simulations at full atomistic resolution.¹³¹ For a cyclic peptide with unknown bound and unbound conformations, a convenient approach to obtain statistically meaningful conformations in solution is to generate its sequence by building its residues in an excluded volume-obeying manner,¹⁸⁵ and sampling its conformational space *via* Monte Carlo simulations and/or relaxing it using (enhanced sampling) molecular dynamics simulations. Nevertheless, the latter contains no information on the conformations sampled by the peptide when bound to the target, which may represent a bottleneck when trying to dock to the target.

Third, a key aspect is the structure of the complex, which aids in understanding what type of interactions drive the assembly and which residues contribute the most to peptide binding and complex stability. Experimentally, a series of crystal structures of cyclic peptide–protein complexes have been resolved¹⁸³ but none in complex with amyloidogenic targets. The thermodynamics and kinetics of peptide binding can be tested using methods such as surface plasmon resonance or isothermal titration calorimetry but none provides specific information on the binding epitope. Computationally, if the representative 3D structures are known, the peptide can be rationally designed and/or docked onto the target and enhanced sampling or deep learning techniques are employed

to extract its binding free energy.¹⁸⁶ Alternatively, in absence of known structures and/or unstably bound complexes, long molecular dynamics simulations could potentially reveal new binding sites. This approach may be efficient if the amyloidogenic target has a well defined secondary structure as is the case for PrP^C, or has druggable pockets. However, for polypeptides with a high degree of plasticity this is a resource intensive and potentially ineffective strategy, which would only slow down peptide design. Machine learning can facilitate the design of peptides, and corroborated with simulations and/or experiments, can aid in the estimation of binding affinities,¹⁸⁷ and improve the peptide sequence for optimal binding to the target.^{171,188} Hence, if combined in an effective manner, computer simulations and machine learning can considerably increase peptide design and optimization efficiency, and can therefore speed up drug development.

The fourth ingredient prior to clinical advancement is the experimental *in vitro* and *in vivo* validation. Given the complementarity of computational and experimental work, an attractive approach would be to integrate the trio, *i.e.* simulations, machine learning and experiments, into a dynamic and iterative engine. For instance, molecular dynamics simulations and deep learning, could be first used to predict and optimize protein and peptide conformations, stability, binding affinities, aiding in the selection of lead candidates prior to experimental validation. Then results from the trio can be incorporated into feedback loops^{37,189,190} that would allow the design of novel and improved peptide sequences, prediction of cyclic peptide bioactivity, better target selectivity, and off-target effects, thus aiding in the faster identification of potent and safe candidates. Hence, the unique integration of such methodologies can aid the design and optimization of novel experiments and computational work. Furthermore, an approach as such can significantly reduce the number of experiments that are required for validation and can increase the homogeneity across the experimental data sets (e.g. environmental conditions).³⁹



6 Perspectives and outlook

In the last 20 years more than 16 cyclic peptide based therapeutics have been FDA and EMA-approved, mainly as antibiotics, anti-cancer therapeutics, antifungals and immunomodulators.^{80,191} Despite extensive research, no cyclic peptide-based drug for neurodegenerative diseases has passed clinical trials. The challenges arise due to the intrinsic disorder of the targets lacking traditional long-lived druggable pockets,⁹⁰ the limited understanding of the associated mechanisms and the often ineffective integration and active feedback between disciplines. Here, we reviewed the design strategies for cyclic peptide design against amyloid-forming targets from a computational perspective and emphasized on the potential of the interconnection between computer simulations, experiments and machine learning in anti-amyloid cyclic peptide design for therapeutic, imaging and diagnostic applications. As such, we proposed a recipe, which can function like a digital twin *i.e.* creating scenarios relying on available information to improve performance and prevent design flaws, allowing for rapid analysis and accurate predictions.¹⁹² Essentially in cyclic peptide design, the digital twin would rely on information from computational and experimental findings to simulate the effect of a cyclic peptide-based drug on an amyloidogenic target, while enhancing the design and optimization of future peptide-based drugs, with better targeting abilities, reduced risk and lower cost. Clearly, the integration is not effortless and requires the efficient incorporation of extensive data from both experiments and simulations (*e.g.* binding constants, toxicity assays, morphological effects *etc.*) to be constantly exchanged between the physical and virtual machine. Importantly, while simulations essentially act as digital twins by themselves, the incorporation of homogeneous experimental data *via* machine learning powered engines can improve predictions, making the next substantial step in (peptide-based) drug design.

The concepts and proposed strategies extend beyond drug design for therapeutic applications and hold the potential to aid in adjacent fields such as (bio)material design or controlled drug delivery.¹⁹³ In essence, it all boils down to the gathering and the smart processing information from diverse sources to create a digital correspondent of a material capturing its composition, structure, responsiveness to external stimuli *etc.*¹⁹⁴ to generate design rules for programmable and adaptable materials.¹⁹⁵

Author contributions

Daria de Raffelle – data curation, formal analysis, investigation, visualization, writing – original draft, writing – review & editing. Ioana M. Ilie – conceptualization, funding acquisition, investigation, methodology, project administration, resources, software, supervision, validation, writing – original draft, writing – review & editing.

Conflicts of interest

There are no conflicts to declare.

Acknowledgements

I. M. I. acknowledges support from the Sectorplan Bèta & Techniek of the Dutch Government and the Dementia Research – Synapsis Foundation Switzerland.

Notes and references

- 1 K. Gammon, *Nature*, 2014, **515**, 299–300.
- 2 I. M. Ilie and A. Cafilisch, *Chem. Rev.*, 2019, **119**, 6956–6993.
- 3 C. Hoffmann, R. Sansevino, G. Morabito, C. Logan, R. M. Vabulas, A. Ulusoy, M. Ganzella and D. Milovanovic, *J. Mol. Biol.*, 2021, **433**, 166961.
- 4 T. P. J. Knowles, M. Vendruscolo and C. M. Dobson, *Nat. Rev. Mol. Cell Biol.*, 2014, **15**, 384–396.
- 5 C. M. Dobson, *Trends Biochem. Sci.*, 1999, **24**, 329–332.
- 6 F. Chiti and C. M. Dobson, *Annu. Rev. Biochem.*, 2017, **86**, 27–68.
- 7 F. Chiti and C. M. Dobson, *Annu. Rev. Biochem.*, 2006, **75**, 333–366.
- 8 A. Aguzzi, *Nature*, 2009, **459**, 924–925.
- 9 O. V. Galzitskaya, S. Y. Grishin, A. V. Glyakina, N. V. Dovidchenko, A. V. Konstantinova, S. V. Kravchenko and A. K. Surin, *Int. J. Mol. Sci.*, 2023, **24**, 3781.
- 10 M. Polymenidou, R. Moos, M. Scott, C. Sigurdson, Y.-Z. Shi, B. Yajima, I. Hafner-Bratkovic, R. Jerala, S. Hornemann, K. Wuthrich, A. Bellon, M. Vey, G. Garen, M. N. G. James, N. Kav and A. Aguzzi, *PLoS One*, 2008, **3**, 1–17.
- 11 R. H. Scannevin, *Curr. Opin. Chem. Biol.*, 2018, **44**, 66–74.
- 12 F. A. Aprile, P. Sormanni, M. Podpolny, S. Chhangur, L.-M. Needham, F. S. Ruggeri, M. Perni, R. Limbocker, G. T. Heller, T. Sneideris, T. Scheidt, B. Mannini, J. Habchi, S. F. Lee, P. C. Salinas, T. P. J. Knowles, C. M. Dobson and M. Vendruscolo, *Proc. Natl. Acad. Sci. U. S. A.*, 2020, **117**, 13509–13518.
- 13 E. Gibbs, J. Silverman, B. Zhao, X. Peng, J. Wang, C. Wellington, I. Mackenzie, S. Plotkin, J. Kaplan and N. Cashman, *Sci. Rep.*, 2019, **9**, 9870.
- 14 P. Chames, M. Van Regenmortel, E. Weiss and D. Baty, *Br. J. Pharmacol.*, 2009, **157**, 220–233.
- 15 H. Samaranyake, T. Wirth, D. Schenkwein, J. K. Rätty and S. Ylä-Herttua, *Ann. Med.*, 2009, **41**, 322–331.
- 16 A. E. Lang, A. D. Siderowf, E. A. Macklin, W. Poewe, D. J. Brooks, H. H. Fernandez, O. Rascol, N. Giladi, F. Stocchi, C. M. Tanner, R. B. Postuma, D. K. Simon, E. Tolosa, B. Mollenhauer, J. M. Cedarbaum, K. Fraser, J. Xiao, K. C. Evans, D. L. Graham, I. Sapir, J. Inra, R. M. Hutchison, M. Yang, T. Fox, S. B. Haeblerlein and T. Dam, *N. Engl. J. Med.*, 2022, **387**, 408–420.
- 17 G. Pagano, K. I. Taylor, J. Anzures-Cabrera, M. Marchesi, T. Simuni, K. Marek, R. B. Postuma, N. Pavesi, F. Stocchi, J.-P. Azulay, B. Mollenhauer, L. López-Manzanares, D. S. Russell, J. T. Boyd, A. P. Nicholas, M. R. Luquin, R. A. Hauser, T. Gasser, W. Poewe, B. Ricci, A. Boulay, A. Vogt, F. G. Boess, J. Dukart, G. D'Urso, R. Finch, S. Zanigni, A. Monnet, N. Pross, A. Hahn, H. Svoboda, M. Britschgi, F. Lipsmeier, E. Volkova-Volkmar, M. Lindemann, S. Dziadek, S. Holiga, D. Rukina, T. Kustermann, G. A. Kerchner, P. Fontoura, D. Umbrecht, R. Doody, T. Nikolcheva and A. Bonni, *N. Engl. J. Med.*, 2022, **387**, 421–432.
- 18 L. Wang, N. Wang, W. Zhang, X. Cheng, Z. Yan, G. Shao, X. Wang, R. Wang and F. Caiyun, *Signal Transduction Targeted Ther.*, 2022, **7**, 1–48.
- 19 K. Fosgerau and T. Hoffmann, *Drug Discovery Today*, 2015, **20**, 122–128.
- 20 P. Dougherty, A. Sahni and D. Pei, *Chem. Rev.*, 2019, **119**, 10241–10287.
- 21 K. Kurrikoff, D. Aphkhasava and U. Langel, *Curr. Opin. Pharmacology*, 2019, **47**, 27–32.
- 22 C. Flores, G. Fouquet, I. C. Moura, T. T. Maciel and O. Hermine, *Front. Immunol.*, 2019, **10**, 1–7.
- 23 V. Armiento, K. Hille, D. Naltsas, J. Lin, A. Barron and A. Kapurniotu, *Angew. Chem., Int. Ed.*, 2020, **59**, 3372–3384.
- 24 T. Ikenoue, F. Aprile, P. Sormanni and M. Vendruscolo, *Front. Neurosci.*, 2021, **15**, 1–9.
- 25 D. S. Nielsen, N. E. Shepherd, W. Xu, A. J. Lucke, M. J. Stoermer and D. P. Fairlie, *Chem. Rev.*, 2017, **117**, 8094–8128.



- 26 K. Patel, L. J. Walport, J. L. Walshe, P. D. Solomon, J. K. K. Low, D. H. Tran, K. S. Mouradian, A. P. G. Silva, L. Wilkinson-White, A. Norman, C. Franck, J. M. Matthews, J. M. Guss, R. J. Payne, T. Passioura, H. Suga and J. P. Mackay, *Proc. Natl. Acad. Sci. U. S. A.*, 2020, **117**, 26728–26738.
- 27 H. Etayash, D. Pletzer, P. Kumar, S. K. Straus and R. E. W. Hancock, *J. Med. Chem.*, 2020, **63**, 9228–9236.
- 28 I. M. Ilie and A. Cafflich, *Biochim. Biophys. Acta, Proteins Proteomics*, 2022, **1870**, 11–12.
- 29 I. M. Ilie, M. Bacci, A. Vitalis and A. Cafflich, *Biophys. J.*, 2022, **121**, 2813–2825.
- 30 X. Jing and K. Jin, *Med. Res. Rev.*, 2020, **40**, 753–810.
- 31 J. Santos, P. Gracia, S. Navarro, S. Peña-Díaz, J. Pujols, N. Cremades, I. Pallarès and S. Ventura, *Nat. Commun.*, 2021, **12**, 1–10.
- 32 S. Yamaguchi, S. Ito, T. Masuda, P.-O. Couraud and S. Ohtsuki, *J. Controlled Release*, 2020, **321**, 744–755.
- 33 Q. Liu, G. Xun and Y. Feng, *Biotechnol. Adv.*, 2019, **37**, 530–537.
- 34 Z. Wu, S. B. J. Kan, R. D. Lewis, B. J. Wittmann and F. H. Arnold, *Proc. Natl. Acad. Sci. U. S. A.*, 2019, **116**, 8852–8858.
- 35 B. J. Wittmann, Y. Yue and F. H. Arnold, *Cell Syst.*, 2021, **12**, P1026–1045.
- 36 K. K. Yang, Z. Wu and F. H. Arnold, *Nat. Met.*, 2019, **16**, 687–694.
- 37 B. Mohr, K. Shmilovich, I. S. Kleinwächter, D. Schneider, A. L. Ferguson and T. Bereau, *Chem. Sci.*, 2022, **13**, 4498–4511.
- 38 S. Ovchinnikov and P.-S. Huang, *Curr. Opin. Struct. Biol.*, 2021, **65**, 136–144.
- 39 H. Narayanan, F. Dingfelder, A. Butte, N. Lorenzen, M. Sokolov and P. Arosio, *Trends Pharmacol. Sci.*, 2021, **42**, 151–165.
- 40 M. Baek and D. Baker, *Nat. Med.*, 2022, **19**, 13–14.
- 41 C. V. Frost and M. Zacharias, *Proteins: Struct., Funct., Bioinf.*, 2020, **88**, 1592–1606.
- 42 C. K. Wang, S. E. Northfield, Y.-H. Huang, M. C. Ramos and D. J. Craik, *Eur. J. Med. Chem.*, 2016, **109**, 342–349.
- 43 J. Sevigny, P. Chiao, T. Bussière, P. H. Weinreb, L. Williams, M. Maier, R. Dunstan, S. Salloway, T. Chen, Y. Ling, J. O’Gorman, F. Qian, M. Arastu, M. Li, S. Chollate, M. S. Brennan, O. Quintero-Monzon, R. H. Scannevin, H. M. Arnold, T. Engber, K. Rhodes, J. Ferrero, Y. Hang, A. Mikulskis, J. Grimm, C. Hock, R. M. Nitsch and A. Sandrock, *Nature*, 2016, **537**, 50–56.
- 44 J. Arndt, F. Qian, B. Smith, C. Quan, K. Kilambi, M. Bush, T. Walz, R. Pepinsky, T. Bussière, S. Hamann, T. Cameron and P. Weinreb, *Sci. Rep.*, 2018, **8**, 6412.
- 45 N. J. Yang and M. J. Hinner, *Getting Across the Cell Membrane: An Overview for Small Molecules, Peptides, and Proteins*, Springer, New York, 2014, pp. 29–53.
- 46 S. Stegemann, C. Moreton, S. Svanbäck, K. Box, G. Motte and A. Paudel, *Drug Discovery Today*, 2022, **28**, 103344.
- 47 P. Kulkarni, S. Bhattacharya, S. Achuthan, A. Behal, M. K. Jolly, S. Kotnala, A. Mohanty, G. Rangarajan, R. Salgia and V. Uversky, *Chem. Rev.*, 2022, **122**, 6614–6633.
- 48 C. J. Oldfield and A. K. Dunker, *Annu. Rev. Biochem.*, 2014, **83**, 553–584.
- 49 L. L. N. Ngoc, S. G. Itoh, P. Sompornpisut and H. Okumura, *Chem. Phys. Lett.*, 2020, **758**, 137913.
- 50 M. H. Dehabadi, A. Cafflich, I. M. Ilie and R. Firouzi, *J. Phys. Chem. B*, 2022, **126**, 7627–7637.
- 51 M. F. M. Sciacca, V. Romanucci, A. Zarrelli, I. Monaco, F. Lolicato, N. Spinella, C. Galati, G. Grasso, L. D’Urso, M. Romeo, L. Diomedede, M. Salmona, C. Bongiorno, G. D. Fabio, C. L. Rosa and D. Milardi, *ACS Chem. Neurosci.*, 2017, **8**, 1767–1778.
- 52 J. M. Jakubowski, A. A. Orr, D. A. Le and P. Tamamis, *J. Chem. Inf. Model.*, 2019, **60**, 289–305.
- 53 S. García-Viñuales, I. M. Ilie, A. M. Santoro, V. Romanucci, A. Zarrelli, G. D. Fabio, A. Cafflich and D. Milardi, *Biochim. Biophys. Acta, Proteins Proteomics*, 2022, **1870**, 140772.
- 54 U. S. Herrmann, A. K. Schütz, H. Shirani, D. Huang, D. Saban, M. Nuvolone, B. Li, B. Ballmer, A. K. O. Åslund, J. J. Mason, E. Rushing, H. Budka, S. Nyström, P. Hammarström, A. Böckmann, A. Cafflich, B. H. Meier, K. P. R. Nilsson, S. Hornemann and A. Aguzzi, *Sci. Transl. Med.*, 2015, **7**, 299ra123.
- 55 J. Zhao, A. Blayney, X. Liu, L. Gandy, W. Jin, L. Yan, J.-H. Ha, A. J. Canning, M. Connelly and C. Yang, *et al.*, *Nat. Commun.*, 2021, **12**, 986.
- 56 P. Velander, L. Wu, F. Henderson, S. Zhang, D. R. Bevan and B. Xu, *Biochem. Pharmacol.*, 2017, **139**, 40–55.
- 57 H. Ashrafiyan, E. H. Zadeh and R. H. Khan, *Int. J. Biol. Macromol.*, 2021, **167**, 382–394.
- 58 M. Perni, C. Galvagnion, A. Maltsev, G. Meisl, M. B. Müller, P. K. Challa, J. B. Kirkegaard, P. Flagmeier, S. I. Cohen and R. Cascella, *et al.*, *Proc. Natl. Acad. Sci. U. S. A.*, 2017, **114**, E1009–E1017.
- 59 N. Majid, M. K. Siddiqi, A. N. Khan, S. Shabnam, S. Malik, A. Alam, V. N. Uversky and R. H. Khan, *ACS Chem. Neurosci.*, 2019, **10**, 4704–4715.
- 60 S. K. Chaturvedi, N. Zaidi, P. Alam, J. M. Khan, A. Qadeer, I. A. Siddique, S. Asmat, Y. Zaidi and R. H. Khan, *PLoS One*, 2015, **10**, e0136528.
- 61 M. K. Siddiqi, P. Alam, S. K. Chaturvedi, M. V. Khan, S. Nusrat, S. Malik and R. H. Khan, *Biochim. Biophys. Acta, Proteins Proteomics*, 2018, **1866**, 549–557.
- 62 P. Alam, S. K. Chaturvedi, M. K. Siddiqi, R. K. Rajpoot, M. R. Ajmal, M. Zaman and R. H. Khan, *Sci. Rep.*, 2016, **6**, 26759.
- 63 P. Alam, M. K. Siddiqi, S. K. Chaturvedi, M. Zaman and R. H. Khan, *Int. J. Biol. Macromol.*, 2017, **99**, 477–482.
- 64 Z. Gazova, K. Siposova, E. Kurin, P. Mucaji and M. Nagy, *Proteins: Struct., Funct., Bioinf.*, 2013, **81**, 994–1004.
- 65 I. Morales, C. Cerda-Troncoso, V. Andrade and R. B. Maccioni, *J. Alzheimers Dis.*, 2017, **60**, 451–460.
- 66 R. Banerjee, *Spectrochim. Acta, Part A*, 2014, **117**, 798–800.
- 67 J. S. Rane, P. Bhaumik and D. Panda, *J. Alzheimers Dis.*, 2017, **60**, 999–1014.
- 68 A. Thapa, S. D. Jett and E. Y. Chi, *ACS Chem. Neurosci.*, 2016, **7**, 56–68.
- 69 K. Ono, M. Hirohata and M. Yamada, *Biochem. Biophys. Res. Commun.*, 2005, **336**, 444–449.
- 70 A. Kundu and A. Mitra, *Plant Foods Hum. Nutr.*, 2013, **68**, 247–253.
- 71 S. Joker, S. Khazaei, H. Behnammanesh, A. Shamloo, M. Erfani, D. Beiki and O. Bavi, *Biophys. Rev.*, 2019, **11**, 901–925.
- 72 R. Roy and S. Paul, *J. Phys. Chem. B*, 2023, **127**, 600–615.
- 73 G. J. Weiner, *Nat. Rev. Cancer*, 2015, **15**, 361–370.
- 74 F. A. Aprile, P. Sormanni, M. Perni, P. Arosio, S. Linse, T. P. J. Knowles, C. M. Dobson and M. Vendruscolo, *Sci. Adv.*, 2017, **3**, e1700488.
- 75 L. Söderberg, M. Johannesson, P. Nygren, H. Laudon, F. Eriksson, G. Osswald, C. Möller and L. Lannfelt, *Neurotherapeutics*, 2022, **20**, 195–206.
- 76 T. Sonati, R. R. Reimann, J. Falsig, P. K. Baral, T. O’Connor, S. Hornemann, S. Yaganoglu, B. Li, U. S. Herrmann, B. Wieland, M. Swayampakula, M. H. Rahman, D. Das, N. Kav, R. Riek, P. P. Liberski, M. N. G. James and A. Aguzzi, *Nature*, 2013, **501**, 102–106.
- 77 R. J. Bateman, J. Cummings, S. Schobel, S. Salloway, B. Vellas, M. Boada, S. E. Black, K. Blennow, P. Fontoura, G. Klein, S. S. Assunção, J. Smith and R. S. Doody, *Alzheimer’s Res. Therapy*, 2022, **14**, 178.
- 78 A. Zorzi, K. Deyle and C. Heinis, *Curr. Opin. Chem. Biol.*, 2017, **38**, 24–29.
- 79 E. Mihara, S. Watanabe, N. K. Bashiruddin, N. Nakamura, K. Matoba, Y. Sano, R. Maini, Y. Yin, K. Sakai, T. Arimori, K. Matsumoto, H. Suga and J. Takagi, *Nat. Commun.*, 2021, **12**, 1543.
- 80 H. Zhang and S. Chen, *RSC Chem. Biol.*, 2022, **3**, 18–31.
- 81 T. Passioura, T. Katoh, Y. Goto and H. Suga, *Annu. Rev. Biochem.*, 2014, **83**, 727–752.
- 82 H. Y. Chow, Y. Zhang, E. Matheson and X. Li, *Chem. Rev.*, 2019, **119**, 9971–10001.
- 83 C. White and A. Yudin, *Nat. Chem.*, 2011, **3**, 509–524.
- 84 M. Erak, K. Bellmann-Sickert, S. Els-Heindl and A. G. Beck-Sickinger, *Bioorg. Med. Chem.*, 2018, **26**, 2759–2765.
- 85 M. T. J. Bluntzer, J. O’Connell, T. S. Baker, J. Michel and A. N. Hulme, *Pept. Sci.*, 2020, **113**, e24191.
- 86 P. M. Cromm, J. Spiegel and T. N. Grossmann, *ACS Chem. Biol.*, 2015, **10**, 1362–1375.
- 87 V. Digiesi, V. de la Oliva Roque, M. Vallaro, G. Caron and G. Ermondi, *Eur. J. Pharm. Biopharm.*, 2021, **165**, 259–270.
- 88 P. Matsson, B. C. Doak, B. Over and J. Kihlberg, *Adv. Drug Delivery Rev.*, 2016, **101**, 42–61.
- 89 A. A. Vinogradov, Y. Yin and H. Suga, *J. Am. Chem. Soc.*, 2019, **141**, 4167–4181.
- 90 B. D. Khairnar, A. Jha and J. M. Rajwade, *J. Mater. Sci.*, 2023, **58**, 9834–9860.



- 91 N. S. Rösener, L. Gremer, E. Reinartz, A. König, O. Brener, H. Heise, W. Hoyer, P. Neudecker and D. Willbold, *J. Biol. Chem.*, 2018, **293**, 15748–15764.
- 92 S. Schemmert, L. C. Camargo, D. Honold, I. Gering, J. Kutzsche, A. Willuweit and D. Willbold, *Int. J. Mol. Sci.*, 2021, **22**, 6553.
- 93 T. Ikenoue, F. A. Aprile, P. Sormanni, F. S. Ruggeri, M. Perni, G. T. Heller, C. P. Haas, C. Middel, R. Limbocker, B. Mannini, T. C. T. Michaels, T. P. J. Knowles, C. M. Dobson and M. Vendruscolo, *Sci. Rep.*, 2020, **10**, 1–15.
- 94 T. Ikenoue, M. Oono, M. So, H. Yamakado, T. Arata, R. Takahashi, Y. Kawata and H. Suga, *ChemBioChem*, 2023, **24**, 1–7.
- 95 M. Chemerovski-Glikman, E. Rozentur-Shkop, M. Richman, A. Grupi, A. Getler, H. Y. Cohen, H. Shaked, C. Wallin, S. K. T. S. Wärmländer, E. Haas, A. Gräslund, J. H. Chill and S. Rahimipour, *Chem. – Eur. J.*, 2016, **22**, 14236–14246.
- 96 A. Spanopoulou, L. Heidrich, H.-R. Chen, C. Frost, D. Hrle, E. Malideli, K. Hille, A. Grammatikopoulos, J. Bernhagen, M. Zacharias, G. Rammes and A. Kapurniotu, *Angew. Chem., Int. Ed.*, 2018, **57**, 14503–14508.
- 97 B. H. Rajewski, K. M. Makwana, I. J. Angera, D. K. Geremia, A. R. Zepeda, G. I. Hallinan, R. Vidal, B. Ghetti, A. L. Serrano and J. R. Del Valle, *J. Am. Chem. Soc.*, 2023, **145**, 23131–23142.
- 98 G. Giubertoni, L. Feng, K. Klein, G. Giannetti, Y. Choi, A. van der Net, G. Castro-Linares, F. Caporaletti, D. Micha, J. Hunger, A. Deblais, D. Bonn, A. Šarić, I. M. Ilie, G. H. Koenderink and S. Woutersen, *bioRxiv*, 2023.
- 99 I. M. Ilie, C. Ehrhardt, A. Cafilisch and G. Weitz-Schmidt, *J. Chem. Inf. Model.*, 2023, **63**, 3878–3891.
- 100 A. Borsatto, O. Akkad, I. Galdadas, S. Ma, S. Damfo, S. Haider, F. Kozielski, C. Estarellas and F. L. Gervasio, *eLife*, 2022, **11**, 1–25.
- 101 V. Armiento, A. Spanopoulou and A. Kapurniotu, *Angew. Chem., Int. Ed.*, 2019, **59**, 3372–3384.
- 102 J. L. Arenas, J. Kaffy and S. Ongeri, *Curr. Opin. Chem. Biol.*, 2019, **52**, 157–167.
- 103 S. Ullrich and C. Nitsche, *Protein Sci.*, 2023, DOI: [10.1002/pep2.24326](https://doi.org/10.1002/pep2.24326).
- 104 R. K. Saini, D. Goyal and B. Goyal, *Chem. Res. Tox.*, 2020, **33**, 2719–2738.
- 105 G. P. Smith, *Science*, 1985, **228**, 1315–1317.
- 106 N. Nemoto, E. Miyamoto-Sato, Y. Husimi and H. Yanagawa, *FEBS Lett.*, 1997, **414**, 405–408.
- 107 R. W. Roberts and J. W. Szostak, *Proc. Natl. Acad. Sci. U. S. A.*, 1997, **94**, 12297–12302.
- 108 C. Sohrabi, A. Foster and A. Tavassoli, *Nat. Rev. Chem.*, 2020, **4**, 90–101.
- 109 J. D. Bloom and F. H. Arnold, *Proc. Natl. Acad. Sci. U. S. A.*, 2009, **106**, 9995–10000.
- 110 Y.-W. Lin, *Coord. Chem. Rev.*, 2017, **336**, 1–27.
- 111 J. Franceus, S. Dhaene, H. Decadt, J. Vandepitte, J. Caroen, J. Van der Eycken, K. Beerens and T. Desmet, *Chem. Commun.*, 2019, **55**, 4531–4533.
- 112 T. M. Caradonna and A. G. Schmidt, *npj Vaccines*, 2021, **6**, 154.
- 113 V. Ramakrishnan, K. Patel and R. Goyal, *De Novo Peptide Design: Principles and Applications*, 2022.
- 114 P. Ragonis-Bachar and M. Landau, *Curr. Opin. Struc. Biol.*, 2021, **68**, 184–193.
- 115 P. Sormanni, F. A. Aprile and M. Vendruscolo, *Proc. Natl. Acad. Sci. U. S. A.*, 2015, **112**, 9902–9907.
- 116 K. A. Murray, C. J. Hu, S. L. Griner, H. Pan, J. T. Bowler, R. Abskharon, G. M. Rosenberg, X. Cheng, P. M. Seidler and D. S. Eisenberg, *Proc. Natl. Acad. Sci. U. S. A.*, 2022, **119**, e2206240119.
- 117 D.-A. Silva, B. E. Correia and E. Procko, in *Motif-Driven Design of Protein-Protein Interfaces*, ed. B. L. Stoddard, 2016, vol. 1414, pp. 285–304.
- 118 P. Bradley, K. M. Misura and D. Baker, *Science*, 2005, **309**, 1868–1871.
- 119 G. Van Zundert, J. Rodrigues, M. Trellet, C. Schmitz, P. Kastritis, E. Karaca, A. Melquiond, M. van Dijk, S. De Vries and A. Bonvin, *J. Mol. Biol.*, 2016, **428**, 720–725.
- 120 G. C. van Zundert and A. M. Bonvin, *Protein Struct. Predict.*, 2014, **1137**, 163–179.
- 121 A. Vangone, J. Rodrigues, L. Xue, G. van Zundert, C. Geng, Z. Kurkuoglu, M. Nellen, S. Narasimhan, E. Karaca and M. van Dijk, *et al.*, *Proteins: Struct., Funct., Bioinf.*, 2017, **85**, 417–423.
- 122 M. Baek, F. DiMaio, I. Anishchenko, J. Dauparas, S. Ovchinnikov, G. R. Lee, J. Wang, Q. Cong, L. N. Kinch and R. D. Schaeffer, *et al.*, *Science*, 2021, **373**, 871–876.
- 123 J. Jumper, R. Evans, A. Pritzel, T. Green, M. Figurnov, O. Ronneberger, K. Tunyasuvunakool, R. Bates, A. Ždek and A. Potapenko, *et al.*, *Nature*, 2021, **596**, 583–589.
- 124 G. Casadevall, C. Duran and S. Osuna, *JACS Au*, 2023, **3**, 1554–1562.
- 125 V. Charitou, S. C. Van Keulen and A. M. Bonvin, *J. Chem. Theory Comput.*, 2022, **18**, 4027–4040.
- 126 J. Jumper, R. Evans, A. Pritzel, T. Green, M. Figurnov, O. Ronneberger, K. Tunyasuvunakool, R. Bates, A. Ždek and A. Potapenko, *et al.*, *Nature*, 2021, **596**, 583–589.
- 127 K. M. Ruff and R. V. Pappu, *J. Mol. Biol.*, 2021, **433**, 167208.
- 128 L. M. Pietrek, L. S. Stelzl and G. Hummer, *Curr. Opin. Struc. Biol.*, 2023, **78**, 102501.
- 129 F. Pinheiro, J. Santos and S. Ventura, *J. Mol. Biol.*, 2021, **433**, 167059.
- 130 W. Wang, *Phys. Chem. Chem. Phys.*, 2021, **23**, 777–784.
- 131 J. Damjanovic, J. Miao, H. Huang and Y.-S. Lin, *Chem. Rev.*, 2021, **121**, 2292–2324.
- 132 M. Sugita, S. Sugiyama, T. Fujie, Y. Yoshikawa, K. Yanagisawa, M. Ohue and Y. Akiyama, *J. Chem. Inf. Model.*, 2021, **61**, 3681–3695.
- 133 B. Barz, A. K. Buell and S. Nath, *Chem. Commun.*, 2021, **57**, 947–950.
- 134 M. Zhu, A. D. Simone, D. Schenk, C. M. Dobson and M. Vendruscolo, *J. Chem. Phys.*, 2013, **139**, 0351011–03510110.
- 135 Y. Chebaro, P. Jiang, T. Zang, Y. Mu, P. H. Nguyen, N. Mousseau and P. Derreumaux, *J. Phys. Chem. B*, 2012, **116**, 8412–8422.
- 136 R. Dubej, S. H. Kulkarni, S. C. Dantu, R. Panigrahi, D. M. Sardesai, N. Malik, J. D. Acharya, J. Chugh, S. Sharma and A. Kumar, *Biol. Chem.*, 2021, **402**, 179–194.
- 137 Q.-H. Chen and V. Krishnan, *Sci. Rep.*, 2021, **11**, 22583.
- 138 Q. N. N. Nguyen, J. Schwochert, D. J. Tantillo and R. S. Lokey, *Phys. Chem. Chem. Phys.*, 2018, **20**, 14003–14012.
- 139 N. V. Kalmankar, H. Hari, R. Sowdhamini and R. Venkatesan, *J. Med. Chem.*, 2021, **64**, 7422–7433.
- 140 N. V. Kalmankar, B. R. Gehi and R. Sowdhamini, *Front. Mol. Biosci.*, 2022, **9**, 986704.
- 141 B. L. Santini and M. Zacharias, *Front. Chem.*, 2020, **8**, 573259.
- 142 K. Kasahara, H. Terazawa, T. Takahashi and J. Higo, *Comput. Struct. Biotechnol. J.*, 2019, **17**, 712–720.
- 143 I. M. Ilie, D. Nayar, W. K. den Otter, N. F. A. van der Vegt and W. J. Briels, *J. Chem. Theory Comput.*, 2018, **14**, 3298–3310.
- 144 J. Huang, S. Rauscher, G. Nawrocki, T. Ran, M. Feig, B. L. de Groot, H. Grubmueller and A. D. MacKerell Jr, *Nat. Met.*, 2016, **14**, 71–73.
- 145 D. Song, R. Luo and H.-F. Chen, *J. Chem. Inf. Model.*, 2017, **57**, 1166–1178.
- 146 J. Mu, H. Liu, J. Zhang, R. Luo and H.-F. Chen, *J. Chem. Inf. Model.*, 2021, **61**, 1037–1047.
- 147 I. Poltavsky and A. Tkatchenko, *J. Phys. Chem. Lett.*, 2021, **12**, 6551–6564.
- 148 T. Mueller, A. Hernandez and C. Wang, *J. Chem. Phys.*, 2020, **152**(5), 050902.
- 149 P. Zhang and W. Yang, *J. Chem. Phys.*, 2023, **159**, 024118.
- 150 K. Wallraven, F. L. Holmelin, A. Glas, S. Hennig, A. I. Frolov and T. N. Grossmann, *Chem. Sci.*, 2020, **11**, 2269–2276.
- 151 T. Gurry and C. M. Stultz, *Biochem. J.*, 2014, **53**, 6981–6991.
- 152 N. Schwierz, C. V. Frost, P. L. Geissler and M. Zacharias, *J. Phys. Chem. B*, 2017, **121**, 671–682.
- 153 A.-M. Fernandez-Escamilla, F. Rousseau, J. Schymkowitz and L. Serrano, *Nat. Biotechnol.*, 2004, **22**, 1302–1306.
- 154 X. Lu, C. R. Brickson and R. M. Murphy, *ACS Chem. Neurosci.*, 2016, **7**, 1264–1274.
- 155 G. Bhardwaj, V. K. Mulligan, C. D. Bahl, J. M. Gilmore, P. J. Harvey, O. Cheneval, G. W. Buchko, S. V. Pulavarti, Q. Kaas and A. Eletsky, *et al.*, *Nature*, 2016, **538**, 329–335.
- 156 C. B. Est, P. Mangrolia and R. M. Murphy, *Prot. Eng., Design and Sel.*, 2019, **32**, 47–57.
- 157 M. Delaunay and T. Ha-Duong, *J. Comp.-Aided Mol. Des.*, 2022, **36**, 605–621.
- 158 J. Schymkowitz, J. Borg, F. Stricher, R. Nys, F. Rousseau and L. Serrano, *Nucleic Acids Res.*, 2005, **33**, W382–W388.
- 159 N. Louros, M. Ramakers, E. Michiels, K. Konstantoulou, C. Morelli, T. Garcia, N. Moonen, S. DHaeyer, V. Goossens and D. R. Thal, *et al.*, *Nat. Commun.*, 2022, **13**, 1351.



- 160 D. M. Mason, S. Friedensohn, C. R. Weber, C. Jordi, B. Wagner, S. Meng, P. Gainza, B. E. Correia and S. T. Reddy, *bioRxiv*, 2019.
- 161 D. Mason, S. Friedensohn, C. Weber, C. Jordi, B. Wagner, S. Meng, R. Ehling, L. Bonati, J. Dahinden, P. Gainza, B. Correia and S. Reddy, *Nat. Biomed. Eng.*, 2021, **5**, 1–13.
- 162 J. Miao, M. L. Descoteaux and Y.-S. Lin, *Chem. Sci.*, 2021, **12**, 14927–14936.
- 163 X. Chen, Q. Zhang, B. Li, C. Lu, S. Yang, J. Long, B. He, H. Chen and J. Huang, *Front. Genet.*, 2022, **13**, 845747.
- 164 K. Martinkus, J. Ludwiczak, K. Cho, W.-C. Lian, J. Lafrance-Vanasse, I. Hotzel, A. Rajpal, Y. Wu, R. Bonneau and V. Gligorijevic *et al.*, *arXiv*, 2023, preprint, arXiv:2308.05027.
- 165 D. Piovesan, A. Del Conte, D. Clementel, A. M. Monzon, M. Bevilacqua, M. C. Aspromonte, J. A. Iserte, F. E. Orti, C. Marino-Buslje and S. C. Tosatto, *Nucleic Acids Res.*, 2023, **51**, D438–D444.
- 166 M. Varadi and S. Velankar, *Proteomics*, 2022, 1–10.
- 167 K. Wu, H. Bai, Y.-T. Chang, R. Redler, K. E. McNally, W. Sheffler, T. Brunette, D. R. Hicks, T. E. Morgan and T. J. Stevens, *et al.*, *Nature*, 2023, **616**, 581–589.
- 168 I. Anishchenko, S. J. Pellock, T. M. Chidyausiku, T. A. Ramelot, S. Ovchinnikov, J. Hao, K. Bafna, C. Norn, A. Kang and A. K. Bera, *et al.*, *Nature*, 2021, **600**, 547–552.
- 169 Z. Du, H. Su, W. Wang, L. Ye, H. Wei, Z. Peng, I. Anishchenko, D. Baker and J. Yang, *Nat. Protoc.*, 2021, **16**, 5634–5651.
- 170 J. Ingraham, M. Baranov, Z. Costello, V. Frappier, A. Ismail, S. Tie, W. Wang, V. Xue, F. Obermeyer and A. Beam, *et al.*, *BioRxiv*, 2022, 1–12.
- 171 N. R. Bennett, B. Coventry, I. Goreschnik, B. Huang, A. Allen, D. Vafeados, Y. P. Peng, J. Dauparas, M. Baek and L. Stewart, *et al.*, *Nat. Commun.*, 2023, **14**, 2625.
- 172 M. Baek, F. DiMaio, I. Anishchenko, J. Dauparas, S. Ovchinnikov, J. Wang and D. Baker, *RoseTTAFold: The first release of RoseTTAFold*, <https://zenodo.org/record/5068265>, 2021.
- 173 J. Ho, A. Jain and P. Abbeel, *Adv. Neural Inf. Process*, 2020, **33**, 6840–6851.
- 174 J. L. Watson, D. Juergens, N. R. Bennett, B. L. Trippe, J. Yim, H. E. Eisenach, W. Ahern, A. J. Borst, R. J. Ragotte and L. F. Milles, *et al.*, *Nature*, 2023, 1–3.
- 175 J. Dauparas, I. Anishchenko, N. Bennett, H. Bai, R. J. Ragotte, L. F. Milles, B. I. Wicky, A. Courbet, R. J. de Haas and N. Bethel, *et al.*, *Science*, 2022, **378**, 49–56.
- 176 C. A. Grambow, H. Weir, N. L. Diamant, A. M. Tseng, T. Biancalani, G. Scalia and K. V. Chuang, *RINGER: Rapid Conformer Generation for Macrocycles with Sequence-Conditioned Internal Coordinate Diffusion*, 2023.
- 177 S. A. Rettie, K. V. Campbell, A. K. Bera, A. Kang, S. Kozlov, J. De La Cruz, V. Adebomi, G. Zhou, F. DiMaio, S. Ovchinnikov and G. Bhardwaj, *bioRxiv*, 2023.
- 178 A. Cabezón, M. Calvelo, J. R. Granja, N. Piñeiro and R. Garcia-Fandino, *J. Colloid Interface Sci.*, 2023, **642**, 84–99.
- 179 J. Ingraham, M. Baranov, Z. Costello, V. Frappier, A. Ismail, S. Tie, W. Wang, V. Xue, F. Obermeyer, A. Beam and G. Grigoryan, 2022.
- 180 J. Roel-Touris and A. M. Bonvin, *Comput. Struct. Biotechnol. J.*, 2020, **18**, 1182–1190.
- 181 W. G. Noid, *J. Phys. Chem. B*, 2023, **127**, 4174–4207.
- 182 R. Evans, M. O'Neill, A. Pritzel, N. Antropova, A. Senior, T. Green, A. Židek, R. Bates, S. Blackwell, J. Yim, O. Ronneberger, S. Bodenstern, M. Zielinski, A. Bridgland, A. Potapenko, A. Cowie, K. Tunyasuvunakool, R. Jain, E. Clancy, P. Kohli, J. Jumper and D. Hassabis, *bioRxiv*, 2021, preprint, DOI: [10.1101/2021.10.04.463034](https://doi.org/10.1101/2021.10.04.463034).
- 183 A. K. Malde, T. A. Hill, A. Iyer and D. P. Fairlie, *Chem. Rev.*, 2019, **119**, 9861–9914.
- 184 M. Wendt, R. Bellavita, A. Gerber, N.-L. Efrém, T. Ramshorst, N. M. Pearce, P. R. J. Davey, I. Everard, M. Vazquez-Chantada, E. Chiarparin, P. Grieco, S. Hennig and T. N. Grossmann, *Angew. Chem., Int. Ed.*, 2021, **60**, 13937–13944.
- 185 I. M. Ilie and A. Cafilisch, *J. Phys. Chem. B*, 2018, **122**, 11072–11082.
- 186 M. Ciemny, M. Kurcinski, K. Kamel, A. Kolinski, N. Alam, O. Schueler-Furman and S. Kmiecik, *Drug Discovery Today*, 2018, **23**, 1530–1537.
- 187 S. Romero-Molina, Y. B. Ruiz-Blanco, J. Mieres-Perez, M. Harms, J. Münch, M. Ehrmann and E. Sanchez-Garcia, *J. Prot. Res.*, 2022, **21**, 1829–1841.
- 188 F. Wan, D. Kontogiorgos-Heintz and C. de la Fuente-Nunez, *Drug Discovery Today*, 2022, **1**, 195–208.
- 189 K. Shmilovich, S. S. Panda, A. Stouffer, J. D. Tovar and A. L. Ferguson, *Digital Discovery*, 2022, **1**, 448–462.
- 190 A. F. Rama Ranganathan, *Method and apparatus using machine learning for evolutionary data-driven design of proteins and other sequence defined biomolecules*, US20220348903A1, 2022.
- 191 J. Li, K. Yanagisawa, M. Sugita, T. Fujie, M. Ohue and Y. Akiyama, *J. Chem. Inf. Model.*, 2023, **63**, 2240–2250.
- 192 A. Fuller, Z. Fan, C. Day and C. Barlow, *IEEE Access*, 2020, **8**, 108952–108971.
- 193 R. Merindol and A. Walther, *Chem. Soc. Rev.*, 2017, **46**, 5588–5619.
- 194 D. Topolsky, A. Belyakov, V. Pochinskaia, I. Topolskaya, N. Yumagulov and D. Fedorov, 2022 IEEE International Multi-Conference on Engineering, Computer and Information Sciences (SIBIRCON), 2022.
- 195 C. Kaspar, B. J. Ravoo, W. G. van der Wiel, S. V. Wegner and W. H. P. Pernice, *Nature*, 2021, **594**, 345–355.

