

peptidoglycan.^{8,9} An important implication of this glycan-targeting concept is that a new class of antitubercular agents can be developed with specific recognition motifs designed to interact directly with key components of the mycobacterial cell envelope to kill *Mtb*. This is a major advance in the quest for novel TB drugs since this strategy bypasses the requirement for a therapeutic agent to cross the 'waxy' *Mtb* cell envelope, and is distinct from conventional approaches. Indeed, glycan-targeting antibodies have reached the clinic, highlighting the potential of this approach.¹³

In this work, we sought to demonstrate the interaction of dimeric boronic acids with the mycobacterial cell envelope through direct visualisation *via* a click-based fluorescence strategy (Fig. 1B). We designed and synthesised two new dimeric boronic acid derivatives bearing either an azido- or alkynyl-handle for post-mycobacterial engagement labelling. The compounds retain antimycobacterial activity following incorporation of the azido and alkyne modifications into the scaffold and provide direct evidence that the dimeric boronic acids are localised to the mycobacterial cell surface. This confirms that extracellular therapeutics or, potentially, diagnostics can be developed using this platform.


Based on our previous findings,⁸ we rationalised that incorporation of 'clickable' capture units into the middle of the linker unit whilst retaining the essential spacing between the two 3-carboxy-phenyl boronic acids to mimic the structure of our most potent dimeric boronic acids⁸ would not interfere with antimycobacterial potency. Our approach to add the fluorogenic probe in the second-step, rather than incorporating this functionality directly into the dimeric boronic acid scaffold, is crucial since fluorophore reporter dyes, which tend to be hydrophobic, could promote non-specific interactions or engagement with the mycobacterial cell envelope.¹⁴

The synthetic route to the new probes containing an azido or alkyne group is shown in Scheme 1 and Scheme S1 (ESI†). **B2-alkyne** and **B2-N₃** were prepared from commercial bis-MPA (2,2-bis(hydroxymethyl)propionic acid) cores containing either an azide or alkyne unit, which were used as the scaffolds to

install both 3-carboxy-phenyl boronic acid units in two steps and in ~30% yield. First the *N*-Boc bis-MPA precursors were deprotected with trifluoroacetic acid followed by an acyl-chloride coupling promoted by triethylamine to install the 3-carboxy-phenyl boronic acid groups in one-step, giving access to the **B2-alkyne** and **B2-N₃** dimeric boronic acid probes (see ESI† for full experimental and characterisation data).

With the two boronic acid probes (**B2-alkyne** and **B2-N₃**) in hand, our first objective was to determine whether the addition of the azido or alkyne modifications impacted on antimycobacterial potency. Evaluation of the minimum inhibitor concentrations (MICs) using the resazurin reduction assay¹⁵ found that both compounds killed *Mycobacterium smegmatis* with similar MIC values to those reported using related dimeric boronic acids with a PEG-based linker (Fig. 1 and Scheme 1),^{8,9} indicating that the clickable handles do not interfere with function. Interestingly, the azido derivative was more potent than the alkyne derivative with a MIC of 0.39 mM compared to 3.1 mM for the alkyne derivative (Scheme 1).

The first approach to directly visualise the interaction between the dimeric boronic acids with mycobacteria was with **B2-alkyne** and the fluorogenic azide probe CalFluor 488 azide (Az488), Fig. 2A. Az488 is only fluorescent following the reaction with an alkyne *via* Cu-catalysed alkyne-azide [3+2] cycloaddition (CuAAC).¹⁶ A potential advantage of this approach is that no wash steps are required after the fluorescence generating CuAAC reaction to remove unreacted probe, which may be



Scheme 1 Synthesis of antimycobacterial 'clickable' dimeric boronic acids. (i) TFA, 3 h rt. (ii) 3 eq. 3-(chlorocarbonyl)phenyl boronic acid, 6 eq. Et₃N, DCM, 0 °C 30 min, rt 16 h. Minimum inhibitory concentration (MIC) against *Mycobacterium smegmatis*.




Fig. 2 (A) Strategy for 'click'-capture of **B2-alkyne** with fluorogenic CalFluor 488 azide. (B) Fluorescence microscopy of **B2-alkyne** labelled *Mycobacterium smegmatis* with Az488 *via* CuAAC. Scale bars are 5 μm.



preferred for either low affinity, transient or non-covalent interactions. *M. smegmatis*, as a fast-growing non-pathogenic model system of *Mtb*, were cultured to logarithmic phase and exposed to sub-MIC of **B2-alkyne** (1 mM) for 15 min followed by incubation with phenylboronic acid (2 mM) lacking the alkyne targeting moiety to compete with the **B2-alkyne**. Under these assay conditions, no killing of mycobacteria was observed. The cells were fixed before a second labelling step with Az488 and imaged directly, Fig. 2B. Additionally, control cells were incubated in the absence of **B2-alkyne** and exposed to unlabelled phenylboronic acid and the same Cu-'click' labelling procedure. Fluorescence microscopy revealed cell surface labelling of the mycobacterial cells treated with the **B2-alkyne** compared to control cells (Fig. 2B), indicating that the dimeric boronic acid **B2-alkyne** is readily incorporated into the mycobacterial cell envelope. We were not able to quantify Az488 labelled mycobacterial cells with flow cytometry analysis although a labelled sub-population was present (Fig. S3, ESI[†]), perhaps suggesting that non-covalent boronic acid interactions coupled with flow cytometry dilution effects leads to the release of the boronic acid from the cell envelope.

In order to investigate the interaction of dimeric boronic acids on live mycobacterial cells, next we evaluated the more potent **B2-N₃** reporter with the brighter DBCO-Cy3 dye (Fig. 3A). *M. smegmatis* cells were cultured to mid-logarithmic phase and in this instance exposed to sub-MIC levels of **B2-N₃** (100 μM) for 30 min. After this time the cells were washed to remove any free **B2-N₃** and then exposed to the DBCO-Cy3 secondary label to enable strain promoted Cu-free azide-alkyne cycloaddition (SPAAC) reaction of DBCO-Cy3 with **B2-N₃**.¹⁷ The cells were washed to remove unreacted DBCO-Cy3 and analysed by microscopy (Fig. 3B) and flow cytometry. As described before, untreated cells exposed to DBCO-Cy3 in the absence of **B2-N₃** and untreated mycobacterial cells were also evaluated. Interestingly, the majority of *M. smegmatis* cells incubated with **B2-N₃** showed fluorescent labelling (Fig. 3B), with a higher efficiency than **B2-alkyne** and, as expected, no background fluorescence was observed for the control cells. Super-resolution live cell imaging with an Airyscan detector¹⁸ found the fluorescent signal was predominantly localised at the cell surface (Fig. 3B), confirming the interaction of **B2-N₃** with components of the mycobacterial cell envelope. The higher resolution imaging of **B2-N₃** can be ascribed to a combination of increased antimycobacterial potency of **B2-N₃**, suggesting stronger interactions with the cell envelope and the higher fluorescence quantum yield of DBCO-Cy3 (0.15)¹⁹ compared to CalFluor Az488-triazole conjugate (0.07).¹⁶ Similar to **B2-alkyne**, we were unable to quantify the extent of **B2-N₃** mycobacterial labelling with flow cytometry but also observed a sub-population of fluorescently labelled cells (Fig. S4, ESI[†]), supporting that boronic acids have a pivotal role in modifying the mycobacterial cell surface.

In conclusion, we have developed a strategy to provide direct evidence that antitubercular dimeric boronic acids target extracellular mycobacterial cell surface components. This was achieved through the rational design and synthesis of new




Fig. 3 (A) Strategy for 'click'-capture of **B2-N₃** with DBCO-Cy3 (B) fluorescence microscopy of **B2-N₃** labelled *Mycobacterium smegmatis* with DBCO-Cy3 via SPAAC. Scale bars are 5 μm.

modular dimeric boronic acids with precision spacing between the glycan-chelating boronic acid moieties and a 'clickable'-recruitment handle in the centre of the linker unit. A pro-fluorescence 'click' strategy revealed binding but did not provide sufficient resolution. In contrast, copper-free click recruitment of DBCO-Cy3 on live mycobacterial cells revealed substitutional cell labelling as shown by microscopy. Importantly, this two-step 'click' capture approach confirms that dimeric boronic acids are retained on the exterior of the mycobacterial cells, thus demonstrating modification the cell envelope. Taken together, these results validate that new drugs, and diagnostics, for mycobacteria can be discovered by targeting the unique extracellular glycans rather than the traditional approaches of targeting specific intracellular pathways, which often fail due to the low permeability of mycobacteria. It is our hope that this approach will lead to the development of novel TB therapeutics alongside new imaging and diagnostic platforms.

This work was supported by a Sir Henry Dale Fellowship to EF jointly funded by the Wellcome Trust and Royal Society (104193/Z/14/Z and 104193/Z/14/B), research grants from the Royal Society (RG120405) and the Leverhulme Trust (RPG-2019-087) and the



EPSRC MAS doctoral training centre for a studentship to RMFT (EP/L015307). We thank Ian Hands-Portman for technical assistance with the microscopy studies. For the purpose of open access, the author has applied a Creative Commons Attribution (CC BY) licence to any Author Accepted Manuscript version arising from this submission.

Conflicts of interest

There are no conflicts to declare.

Notes and references

- 1 World Health Organisation 2021 Global Tuberculosis Report https://www.who.int/tb/publications/global_report/en/.
- 2 M. Pai, T. Kasaeva and S. Swaminathan, *N. Engl. J. Med.*, 2022, **386**, 1490–1493.
- 3 G. B. Migliori, S. Tiberi, A. Zumla, E. Petersen, J. M. Chakaya, C. Wejse, M. Munoz Torrico, R. Duarte, J. W. Alffenaar, H. S. Schaaf, B. J. Marais, D. M. Cirillo, R. Alagna, A. Rendon, E. Pontali, A. Piubello, J. Figueroa, G. Ferlazzo, A. Garcia-Basteiro, R. Centis, D. Visca, L. D'Ambrosio, G. Sotgiu and Members of the Global Tuberculosis Network, *Int. J. Infect. Dis.*, 2020, **92S**, S15–S25.
- 4 K. Dheda, T. Gumbo, N. R. Gandhi, M. Murray, G. Theron, Z. Udwadia, G. B. Migliori and R. Warren, *Lancet Respir. Med.*, 2014, **2**, 321–338.
- 5 P. J. Brennan and H. Nikaido, *Annu. Rev. Biochem.*, 1995, **64**, 29–63.
- 6 S. M. Batt, D. E. Minnikin and G. S. Besra, *Biochem. J.*, 2020, **477**, 1983–2006.
- 7 E. Fullam and R. J. Young, *RSC Med. Chem.*, 2020, **12**, 43–56.
- 8 C. S. Guy, M. I. Gibson and E. Fullam, *Chem. Sci.*, 2019, **10**, 5935–5942.
- 9 C. S. Guy, K. Murray, M. I. Gibson and E. Fullam, *Org. Biomol. Chem.*, 2019, **17**, 9524–9528.
- 10 J. Brem, R. Cain, S. Cahill, M. A. McDonough, I. J. Clifton, J. C. Jimenez-Castellanos, M. B. Avison, J. Spencer, C. W. Fishwick and C. J. Schofield, *Nat. Commun.*, 2016, **7**, 12406.
- 11 S. G. Kurz, S. Hazra, C. R. Bethel, C. Romagnoli, E. Caselli, F. Prati, J. S. Blanchard and R. A. Bonomo, *ACS Infect. Dis.*, 2015, **1**, 234–242.
- 12 F. L. Rock, W. Mao, A. Yaremchuk, M. Tukalo, T. Crepin, H. Zhou, Y. K. Zhang, V. Hernandez, T. Akama, S. J. Baker, J. J. Plattner, L. Shapiro, S. A. Martinis, S. J. Benkovic, S. Cusack and M. R. Alley, *Science*, 2007, **316**, 1759–1761.
- 13 E. Sterner, N. Flanagan and J. C. Gildersleeve, *ACS Chem. Biol.*, 2016, **11**, 1773–1783.
- 14 Z. Zhang, D. Yomo and C. Gradinaru, *Biochim. Biophys. Acta, Biomembr.*, 2017, **1859**, 1242–1253.
- 15 J. C. Palomino, A. Martin, M. Camacho, H. Guerra, J. Swings and F. Portaels, *Antimicrob. Agents Chemother.*, 2002, **46**, 2720–2722.
- 16 P. Shieh, V. T. Dien, B. J. Beahm, J. M. Castellano, T. Wyss-Coray and C. R. Bertozzi, *J. Am. Chem. Soc.*, 2015, **137**, 7145–7151.
- 17 J. M. Baskin, J. A. Prescher, S. T. Laughlin, N. J. Agard, P. V. Chang, I. A. Miller, A. Lo, J. A. Codelli and C. R. Bertozzi, *Proc. Natl. Acad. Sci. U. S. A.*, 2007, **104**, 16793–16797.
- 18 X. Wu and J. A. Hammer, *Methods Mol. Biol.*, 2021, **2304**, 111–130.
- 19 <https://www.aatbio.com/products/dbco-cy3>.

