Dalton Transactions



PAPER View Article Online
View Journal | View Issue



Cite this: *Dalton Trans.*, 2017, **46**, 5680

Received 30th November 2016, Accepted 4th February 2017 DOI: 10.1039/c6dt04527h

rsc.li/dalton

Modulation of the CO₂ fixation in dinickel azacryptands†

F. Möller, ^a L. Castañeda-Losada, ^a J. R. C. Junqueira, ^a R. G. Miller, ^a M. L. Reback, ^a B. Mallick, ^a M. van Gastel ^b and U.-P. Apfel ^a

While bimetallic azacryptands are known to selectively coordinate CO_2 , there is little knowledge on how different substitution patterns of the azacryptand cage structure influence CO_2 coordination. Stopped-flow UV-vis spectroscopy, electrochemical analysis and DFT calculations were performed on a series of dinickel azacryptands and showed different rates of CO_2 coordination to the complexes. We herein present data showing that the different flexibility of the azacryptands is directly responsible for the difference in the CO_2 uptake capability of dinickel azacryptand complexes.

Introduction

The fixation and utilization of CO2 as a C1-building block is an important research field towards the recycling of the potent greenhouse gas CO2. While enzymes like CO-dehydrogenases9 and ureases¹⁰ allow selective CO₂ fixation under mild and aqueous conditions, 7,8 they are not cost effective for industrial use. Therefore, it is vital to develop materials that are as selective and efficient as enzymes but at a much lower cost. Indeed a number of synthetic approaches for the fixation and transportation of CO2 have been reported, such as metal organic frameworks (MOFs),² covalent organic frameworks (COFs)^{3,4} or even inorganic carbonates.^{5,6} Although they show promising properties, most of them have low selectivity towards CO2 in the presence of other atmospheric gases and also low stability in the presence of moisture.7,8 Cryptands, such as bis-Tren azacryptands (Tren = tris(2-aminoethyl)amine), have been shown to allow selective fixation and transportation of small molecules (e.g. bicarbonates, azides or thiocyanates). 9-13 By using cryptands, nowadays frequently used for anion recognition as well as in metal chelation, an attempt for a comparable strategy for CO₂ fixation was made. ¹⁴ The stability and selectivity of the resulting compounds for the fixation of small molecules depend on the cage size as well as on the effective size of the small molecules. 9,10,13,15-18 As a result, the

Scheme 1 Azacryptands $L_A{}^R$ and imines $L_I{}^R$ with different linker molecules L^R . The linkers are arranged according to their (i) different steric bulks on the central benzyl unit, (ii) capability to directly alter the electron density within the azacryptand cavity and (iii) different cage sizes. 5,14,15,17,20

uptake and binding properties of small molecules (*e.g.* halogenides and pseudohalogenides) can be selectively tuned by increasing the size of the cryptand and/or altering the binding motifs. The size of the binding cavity of cryptands for a potential small molecule to enter can be rationally designed by using different linker molecules connecting both Tren-moieties (Scheme 1). ¹⁹ Along this line, Nelson *et al.* recently showed crystallographic evidence for different CO_2 coordination in dicobalt-azacryptand complexes. ²⁰ While $[CO_2L_A^{Fur}](CIO_4)_2$ reveals a short Co–Co distance of 4.2924(3) Å and a M– O_{CO_2} bond length of 2.145(9) Å, the Co–Co distance

^aInorganic Chemistry I/Bioinorganic Chemistry, Ruhr University Bochum, Universitätsstraße 150, 44801 Bochum, Germany. E-mail: ulf.apfel@rub.de ^bMax-Planck-Institut für Chemische Energiekonversion, Stiftstraße 34-36, 45470 Mülheim. Germany

[†] Electronic supplementary information (ESI) available: Synthesis and characterization of compounds, X-ray crystallographic analysis, UV-Vis spectra, SQUID and kinetic data. CCDC 1517780–1517787 and 1517950. For ESI and crystallographic data in CIF or other electronic format see DOI: 10.1039/c6dt04527h

Dalton Transactions


in [Co₂L_A^H](ClO₄)₂ is significantly increased to 5.939(2) Å and the M-O_{CO} bond length is decreased to 1.920(6) Å. Mechanistic insight into the CO₂ fixation in [Cu₂L_A^H](ClO₄)₃ and [Cu₂L_A^{H,para}](ClO₄)₃ was provided by Chen as well as Mooney et al., where they highlighted the necessity of an additional hydroxyl group on one metal site for successful CO₂ uptake. 19,21 Likewise, DFT-calculations were reported for $[Cu_2L_A^{Thio}]^{4+}$, $[Cu_2L_A^{\dot{H}}]^{4+}$ and $[Cu_2L_A^{Fur}]^{4+}$, with the lowest energy barrier for CO₂ binding reported for [Cu₂L_A^{Fur}]⁴⁺. 19 Notably, alteration of the linker not only had an effect on the substrate binding but also had a strong influence on the metal binding strength as was shown for $L_A^{\ Thio},\,L_A^{\ Fur}$ and $L_A^{\ Py}.^{22}$ We recently showed that [Ni₂L_A^H](Cl)(ClO₄)₃ is capable of performing rapid CO₂ uptake from air $(k = 0.067 \pm 0.005 \text{ M}^{-1} \text{ s}^{-1})$ and was able to reversibly bind CO₂ by substitution with azides.²³ Furthermore, we could show that the azide ligand could be replaced by atmospheric CO2 in a quasi-reversible process upon irradiation with UV light. Surprisingly, [Ni₂L_A^{tBu}](Cl) (ClO₄)₃ did not reveal any notable CO₂ fixation. Inspired by our initial results on the different binding capabilities of $[Ni_2L_A^H]^{4+}$ and $[Ni_2L_A^{tBu}]^{4+}$ as well as the opposing theoretical reports on altered CO2 binding by varied linker moieties, we set out to further experimentally and theoretically elucidate the effects of linker variations in azacryptand cages on the CO₂ uptake capability and kinetics.

Results and discussion

Synthesis and characterization


The azacryptands L_A^R were synthesized *via* a two-step synthesis according to literature procedures (Scheme 1). 9,19,20,23-26 In a first step, [2 + 3]-Schiff-base condensation of the respective dialdehydes and Tren afforded the imines L_{I}^{R} in good yields (57-77%). The reaction of L_I^R with KBH₄ yielded the azacryptands $L_A^{\ R}$ in good to excellent yields (66-99%). The molecular structures of the hexa-imines $L_I^{\ F},\ L_I^{\ OMe}$ and $L_I^{\ Me},$ and the hexa-amines $L_A^{\ Py}$ and $L_A^{\ OH}$ are presented in Fig. S1 and S2.† A simple way to investigate the influence of the linker molecule on the metal cryptand properties during CO2 uptake is the application of the imine species L_I^R since they are structurally more rigid than their amine counterparts. We therefore attempted complex formation of the hexa-imines with Ni(ClO₄)₂·6H₂O. The reaction solely afforded [(Tren)Ni(CH₃CN)₂] (ClO₄)₂ in 33% yield by a Ni-catalyzed imine hydrolysis in the presence of water of crystallization (Scheme 2).27 The molecular structure was unequivocally confirmed by X-ray crystallography (Fig. S3†). Similar decomposition results were also observed for other hexa-imine azacryptands by ESI-MS. The only exception was L_I^{OH,Me}, which afforded a mononuclear complex when reacted with MnCl2; similar results are reported in the literature.²⁸

Due to the instability of the imine complexes in the presence of moisture, we did not further investigate the imines towards the possibility of CO2 coordination and instead focused on the hexa-amines $L_A^{\ R}$ as ligands. The coordination



Scheme 2 Formation of [(Tren)Ni(CH₃CN)₂](ClO₄)₂ through Ni-catalyzed imine-cleavage.

reactions of L_A^R with Ni(ClO₄)₂·6H₂O were performed under a nitrogen atmosphere to avoid any bicarbonate formation (Scheme 3). While the UV-vis spectra of [Ni₂L_A^R](Cl)(ClO₄)₃ $(L_A^R = L_A^H, L_A^F, L_A^{OMe})$ and L_A^{Me} showed absorption bands at \sim 390, 480, 560, and 620 nm, the spectra of $[Ni_2L_A^R](Cl)(ClO_4)_3$ $(L_A^R = L_A^{fBu}, L_A^{OH}, L_A^{H,para})$ and L_A^{Fur} solely revealed a broad band at ~580 nm containing a shoulder at lower wavelengths (Fig. 1, S4 and S5†). While a detailed band assignment has not been possible, recent crystallographic studies on [Ni₂L_A^H](Cl) $(ClO_4)_3$ and $[Ni_2L_A^{tBu}](Cl)(ClO_4)_3$ suggest an overall unchanged



Scheme 3 Reaction of the azacryptands LAR with Ni(ClO₄)₂·6H₂O.




Fig. 1 Molecular structures of [Ni₂L_A^{OH,Me}](ClO₄)₂ (left) and [Ni₂L_A^{Thio}] (ClO₄)₄ (right). Hydrogen atoms and counter ions were omitted for

Paper Dalton Transactions

shared structural motif of the above-mentioned complexes.²³ Likewise, ESI-MS experiments support a similar composition of the compounds by showing comparable mass patterns (Fig. S6†). Structural analysis revealed that one of the two Ni(II)-centers is coordinated by a water molecule and an acetonitrile, while the other is coordinated to a chloride.²³ Notably, different molecular assemblies were observed upon the reaction of Ni(ClO₄)₂·6H₂O with either L_A^{Thio} or L_A^{OH,Me}. The molecular structure of $[Ni_2L_A^{OH,Me}](ClO_4)_2$ (Fig. 1) reveals two octahedrally coordinated Ni-centers. Each Ni-center is surrounded by four nitrogen-donor atoms of the Tren-ligand with an average Ni-N distance of 2.14 Å. Additionally, one phenolate linker is coordinated in a μ^2 -fashion, bridging both Ni-atoms with bonding distances of 2.164(2) [Ni(1)-O(1)] and 2.177(2)[Ni(2)-O(1)] Å. The coordination sphere of each Ni-center is completed by a non-bridging phenolate linker with Ni(1)-O(3) and Ni(2)-O(2) distances of 2.033(3) and 2.021(3) Å, respectively. While this compound appears to be a mixed valent $Ni(\pi)$ / Ni(III)-species, its low synthetic yield (13%) stopped us from elucidating the exact electronic nature. It is possible that protonation of the ligand framework takes place to afford an all Ni(II)-species. Surprisingly, LAThio revealed a different coordination behavior in the solid state where both Ni-atoms are coordinated on the outside of the cryptand cavity, as was reported for the structurally related complexes [Cu₂L_A^{Thio}] (O₃SOCF₃)₂ and [Ag₂L_A^{Thio}](ClO₄)₂.²⁹ Each Ni(II)-center is octahedrally coordinated to only three N-donor atoms of the Trenmoiety as well as two additional acetonitriles and one water ligand. We assume that the different structure of [Ni₂L_A^{Thio}] (ClO₄)₄ compared to e.g. [Ni₂L_A^H](Cl)(ClO₄)₃ can be explained by a different metal binding affinity and therefore an altered complex stabilization, as was reported by Nelson and coworkers.²² We next investigated the CO₂ fixation behavior of all [Ni₂L_A^R](Cl)(ClO₄)₃ complexes by UV-vis spectroscopy and ESI-MS. A clear change in the UV-vis spectra upon purging the $[Ni_2L_A^R](Cl)(ClO_4)_3$ (R = H, F, OMe, Me and Fur) solutions with CO₂ is observed, showing a decrease in absorption intensity of the bands between 430-580 nm with the formation of an intense absorption band at about 610 nm (Fig. 2 and S4†). The amplitude of the absorption band as well as the disappearance of the original bands between 430-580 nm depends on the substitution pattern at the linker unit.

In analogy to our recent finding for [Ni₂L_A^H](Cl)(ClO₄)₃, such changes can be attributed to the coordination of CO2 within the cavity of the cryptand to afford a bicarbonate dinickel complex.23 ESI-MS analysis further supports the fixation of either 12CO2 or 13CO2 for the reported complexes by the appearance of the [Ni₂L_A^R(HCO₃)] mass-peak (Fig. S7 and S8†).²³ Notably, while the color changes of the complexes upon reaction with CO2 are usually unclear from dark to light blue, [Ni₂L_A^F](Cl)(ClO₄)₃ reveals a distinct color change upon CO₂ fixation from blue to red (Fig. S9 and S10†). In contrast, no apparent changes could be observed for the complexes comprising the L_A^{Thio} , L_A^{OH} , L_A^{Py} , or $L_A^{OH,Me}$ moiety. Likewise, ESI-MS analysis solely revealed the mass peaks of the starting complexes. It can thus be assumed that these complexes do not possess the capability to fixate CO2 under the described reaction conditions, although a small shift of the main band from 564 nm to 571 nm was observed in the UV-vis spectrum upon CO₂ addition to [Ni₂L_A^{H,para}](Cl)(ClO₄)₃. ESI-MS showed a new mass-peak at m/z = 756, which clearly indicates a reaction of [Ni₂L_A^{H,para}](Cl)(ClO₄)₃ to afford a new complex. This behavior can most likely be attributed to the formation of [Ni₂L_A^{H,para}(CN)](ClO₄)₃ comprising a bridging CN⁻-ligand but no coordinated bicarbonate. Further evidence for the presence of a CN⁻ ligand was provided by IR spectroscopy showing a signal at 2022 cm⁻¹ that can be assigned to a bridging CN⁻ moiety. A similar observation was recently reported for [Cu₂L_A^{H,para}(CN)](ClO₄)₃, which was obtained via C-C bond cleavage of a coordinated acetonitrile.30

Recent DFT calculations by Mooney et al. suggested an even better CO_2 -fixation in $\left[Cu_2L_A^{Thio}\right]$ than in the respective [Cu₂L_A^H] complex. 19 In contrast, [Ni₂L_A^{Thio}](ClO₄)₄ does not perform any CO2 uptake. Our finding is supported by a recent experimental investigation of Fabbrizzi and co-workers who report on the inability of [Cu₂L_A^{Thio}](ClO₄)₃ to bind the HCO₃⁻ anion.15

Kinetic analysis

To further evaluate the differences of the azacryptand platform we performed UV-vis stopped-flow investigations. We expected a significant alteration of the CO2 uptake kinetics with different substitution patterns. The time-dependent absorption changes were measured at different temperatures (15-45 °C) and CO₂ concentrations. The obtained absorption changes were then fitted using a pseudo-first order equation of the type $A = A_0 \exp(-k_{\text{obs}} \cdot t)$ (A = absorbance, A_0 = initial absorbance, t = time in s (Tables 1, S1-S3 and Fig. 3, S11†). The obtained data clearly demonstrate that the CO₂ uptake rate is dependent upon the substitution pattern of the dinickel aza-




Fig. 2 UV-vis spectra (MeCN/MeOH 4:1, RT) of the reaction of [Ni₂L_A^R] (Cl)(ClO₄)₃ with CO₂: (a) [Ni₂L_A^F], (b) [Ni₂L_A^{OMe}], (c) [Ni₂L_A^{H,para}] and (d) $[Ni_2L_A^{Py}]$

Dalton Transactions

Table 1 k_2 values obtained from the slope of the plot of k_{obs} vs. the CO₂ concentration at 298.15 K

$\left[Ni_{2}L_{A}^{\ R}\right]$	$k_2 \left[M^{-1} \text{ s}^{-1} \right]$	$\left[Ni_{2}L_{A}^{R}\right]$	$k_2 \left[\mathbf{M}^{-1} \ \mathbf{s}^{-1} \right]$
$egin{array}{c} L_A^{\ H} \ L_A^{\ F} \ L_A^{\ OMe} \end{array}$	$6.7 \times 10^{-2} \pm 5.0 \times 10^{-3}$ $2.0 \times 10^{-2} \pm 3.1 \times 10^{-3}$ $1.9 \times 10^{-2} \pm 2.7 \times 10^{-3}$	$L_A^{Me} \ L_A^{Fur}$	$1.6 \times 10^{-2} \pm 1.1 \times 10^{-3}$ $9.8 \times 10^{-4} \pm 1.1 \times 10^{-4}$




Fig. 3 Plot of $k_{\rm obs}$ vs. the CO₂ concentration of $L_{\rm A}^{\rm R}$ in MeCN at 298.15 K for the reaction of [Ni₂L_A^R] with CO₂.

cryptand complex. Notably, the CO2 fixation is slower in $[Ni_2L_A^R](Cl)(ClO_4)_3$ complexes with sterically more bulky groups according to $L^H > L^F > L^{OMe} > L^{Me} \gg L^{tBu}$. In addition, the application of the furan linker molecules results in a significant decrease of the rate of CO₂ uptake. This kinetic trend is valid for all temperatures measured (Tables S2 and S3†). The small ΔH^{\ddagger} and ΔS^{\ddagger} values obtained from the Eyringplots indicate that the coordination of CO₂ in all substrates proceeds in more than one step.²³ Therefore these small values reflect a more complex association of CO2 and thus preequilibrium and activation enthalpies compose the apparent ΔG^{\ddagger} value. Comparable changes of the reaction rates upon alteration of the reactive site environment were previously reported by Holm and co-workers on [Ni^{II}(pyN₂^{R2})(OH)]^{-.31} In contrast, we present an example that exhibits no obvious alteration of the steric bulk on the metal center.

Azide fixation

A likely explanation for the alteration of the CO₂ uptake kinetics is a decisive change of the Ni-Ni distance and cavity size due to the influence of the linker. A similar hypothesis was reported by Nelson and co-workers.³² Likewise, the particular shape of the formed HCO₃⁻ anion can play a significant part in the destabilization within the dinickel complex as one C-O bond is directed towards the opening of the cavity and can interfere with the ligand periphery. As such, the uptake of linear molecules, e.g. azides, should not be dramatically influ-




Fig. 4 Molecular structure of [Ni₂L_A^F(N₃)](ClO₄)₃. Hydrogen atoms, solvent molecules and counter anions were omitted for clarity.

enced by the alteration of the substitution pattern. Along this line we and others have shown structural evidence for the successful azide coordination into the cavity of LAR. 9-11,23 Even when no CO2 binding was observed, such complexes, e.g. [Ni₂L_A^{tBu}](Cl)(ClO₄)₃, allowed for rapid coordination of N₃ between both nickel atoms. Correspondingly, we tested the capability of [Ni₂L_A^R](Cl)(ClO₄)₃ to allow azide coordination. All investigated azacryptands, except [Ni₂L_AOH,Me](ClO₄)₂, show fixation of N₃-, which is obvious from the changes in their UVvis spectra by the formation of a new common absorption band at about 350 nm (Fig. S12†).23 Additionally, ESI-MS analysis further confirms the formation of an azide complex and reveals the respective $[Ni_2L_A^R(N_3)]$ mass peaks. Crystals suitable for X-ray crystallography were obtained for $[Ni_2L_A^F(N_3)]$ (ClO₄)₃ (Fig. 4) and the results confirm the incorporation of N_3 between the two Ni-centers. It is notable that the Ni-Ni distance (6.275 Å) is significantly larger than in [Ni₂L_A^H(N₃)] $(ClO_4)_3$ (6.129 Å) and $[Ni_2L_A^{tBu}(N_3)](ClO_4)_3$ (6.119 Å). (23,33

The general coordination of azides within the cavity and the alteration of the Ni-Ni distances within structurally comparable metal complexes underline the influence of the substitution pattern. It also shows that CO₂ is a key component in the different uptake kinetics. Furthermore, the successful incorporation of negatively charged azides additionally shows that the lone pairs of the furan, pyridine or thiophene linker cannot be a major reason for the weak or no CO₂ binding in [Ni₂L_A^{Fur}](Cl) (ClO₄)₃ or [Ni₂L_A^{Py}](Cl)(ClO₄)₃, respectively. In light of the acidic properties of CO₂ in an aqueous environment, changes of the substitution pattern might also alter the basicity of the coordinating N-donors and thus the nucleophilicity of the metal atoms. This hypothesis, however, has to be ruled out since the redox potentials of the complexes did not show a trend when electron withdrawing groups (e.g. F) or electron donating groups (e.g. ^tBu, Me) were installed. For all complexes, multiple irregular electron transfer steps can be observed at ~1.5 V vs. Fc/Fc⁺, which we were not able to assign (Fig. S13†).

Theoretical analysis

In order to rationalize the experimentally observed differences in CO2 binding of [Ni2LAR](Cl)(ClO4)3 complexes, DFT calcu**Paper**

lations were performed for complexes with R = H, F, Me and tBu (Fig. 5a). Both Ni^{2+} centers were found to be in the highspin state (S = 1) consistent with the observed octahedral and trigonal-bipyramidal coordinations of the Ni^{2+} centers and this finding is in line with previous SQUID measurements. 23 The two triplet states were found to be exchange-uncoupled by broken-symmetry calculations. Though the substituents differ in their electron-donating and withdrawing capacities, no electronic effect was observed at the nickel ions as well as at all amines, as became apparent from unchanged Mulliken charges, bond distances and orbital compositions. The latter is in-line with the experimental observation that no clear correlation between the CO_2 uptake kinetics and electron donating capacity of the substituent could be found.

However, a noticeable steric effect was observed in the calculations in that rotation of the phenyl groups leads to a steric clash with the bulky t Bu-substituent (Fig. 5b) while the smaller CH₃ groups allow larger rotational flexibility of the ligand. For R = H and F, the phenyl rotation is essentially unhindered (Fig. 5c).

The bulkiness and flexibility of the substituent correlates with the observed CO₂ uptake kinetics, suggesting that these two factors help in tuning the kinetics and that it is most likely the rate-determining step in the reaction mechanism. Moreover, since the electronic structure at the nickel centers and the amines are the same for all of the complexes, every complex should in principle be able to take up and convert CO₂. This was tested by performing an additional calculation in which Cl⁻ and MeCN solvent molecules were removed, hydroxide was inserted at the position where the crystal structure contains a




Fig. 5 (a) Geometry optimized structure of $[Ni_2L_A^F]$ with Cl^- , H_2O and MeCN; (b) side view of $[Ni_2L_A^{fBu}]$, showing the steric hindrance of tBu with the neighboring phenyl group; (c) side view of $[Ni_2L_A^H]$ where steric effects are absent; (d) intermediate structure with bent CO_2 where CO_2 has been introduced near Ni^{2+} at 2.2 Å, leading to barrierless C-O bond formation towards HCO_3^- .

H₂O molecule and CO₂ was introduced near the Ni-ions. Geometry optimization indeed leads to a barrier-less formation of HCO₃⁻ with the driving force being C-O bond formation. An intermediate structure of this mechanism is shown in Fig. 5d, which is in agreement with the mechanism proposed for copper cryptates. 19 Of note here is that contrary to the experimental findings, the Ni-Ni distance is largely independent of the substituent and ranges from 6.09 to 6.11 Å and the starting position of CO₂ has to be chosen such that the Ni-O bond is short and amounts to 2.2 Å. A starting structure with a longer Ni-O distance does not lead to the formation of HCO₃⁻. It may thus be conceivable that the different flexibility and dynamics related to the size of the ligand $(L_A^H < L_A^F <$ $L_A^{Me} \ll \ll L_A^{\ tBu}$) modulates the initial binding of CO_2 and thereby the Ni-O bond distance as well as the activation and the kinetics towards C-O bond formation. Contrary to the fixation of CO₂, DFT calculations for the fixation of N₃ reproduce the experimentally observed difference in the Ni-Ni distance (6.129 Å for L_A^H and 6.261 Å for L_A^F), which clearly is a consequence of a different geometric arrangement of the phenyl groups caused by different substitution patterns.

Conclusion

The coordination of CO2 in dinickel azacryptands can be manipulated through the presence of different linker molecules comprising Tren cages. UV-vis spectroscopic analyses, as well as ESI-MS analyses clearly show an influence of different functional groups on the CO2 uptake. Functional groups pointing into the cryptand cavity, as in $L_A^{\ Fur},\ L_A^{\ Py}$ or $L_A^{\ Thio},$ and LAOH,Me significantly slow down or even prohibit a coordination of CO2. In contrast to this, functional groups pointing out of the cavity show an increasing CO2-fixation rate with decreasing steric demand ($L_A^{\ tBu}$ «« $L_A^{\ OMe}$ < $L_A^{\ Me}$ < $L_A^F < L_A^H$). Both DFT calculations and cyclic voltammetry demonstrate that there are no electronic effects at the nickel centers as a result of the different substituents. Therefore, we attribute the observed changes in reactivity to structural changes. Furthermore, the DFT calculations performed herein show that with increasing steric demand of the linker, the flexibility of the azacryptand core is decreased, providing a kinetic barrier to the initial coordination of CO₂. In contrast to the binding of CO2, all dinickel complexes show fixation of azides. The results clearly show that controlling the flexibility of the cryptand can regulate binding of different substrates. With this in hand, new applications might be accessible for azacryptands, e.g. within catalysis or gas separation utilizing cryptands as the ligand platform.

Experimental

General techniques

All reactions were performed under either a dry N_2 atmosphere using standard Schlenk techniques or in a glovebox. All

Dalton Transactions Paper

solvents were dried according to standard methods. ¹H, ¹³C NMR spectra were recorded on a Bruker DPX-200 NMR, Bruker DPX-250 NMR or a DPX-400 NMR spectrometer at room temperature. Peaks were referenced to residual ¹H signals from the deuterated solvent and are reported in parts per million (ppm). IR spectra were measured with a Bruker Tensor 27 FT-IR spectrometer as a KBr pellet and are reported in cm⁻¹. Mass spectra were measured with a Shimadzu OP-2010 instrument. The dialdehydes 17,34-37 as well as the azacryptands L_A^{tBu} , L_A^{Tur} , L_A^{Tur} , L_A^{Thio} , L_A^{Thio} , $L_A^{H,para}$, L_A^{Py} , L_A^{Py} , and $L_A^{OH,Mo^{31}}$ were synthesized according to literature procedures. All other chemicals were used as received from commercial vendors.

Caution! Perchlorate salts of metal complexes with organic ligands are potentially explosive. They should be handled with care, and prepared only in small quantities.

X-ray data collection and structure solution refinement

Single crystals suitable for X-ray analysis were coated with Paratone-N oil, mounted on a fiber loop, and placed in a cold, gaseous N_2 stream on the diffractometer. L_I^F and L_A^{Py} were measured on an Oxford XCalibur diffractometer performing φ and ω scans at 170(2) K. Diffraction intensities were measured using graphite-monochromatic Mo K α radiation (λ = 0.71073 Å). $[Ni_2L_A^F(N_3)](ClO_4)_3$, $[Ni_2L_A^{Thio}](ClO_4)_4$ and L_A^{OH} were measured on a SuperNova diffractometer performing φ and ω scans at 100(2) K. Diffraction intensities were measured using graphite monochromatic Cu K α radiation (λ = 1.54184 Å). $[Ni_2L_A^{OH,Me}](ClO_4)_2$, L_I^{Me} and L_I^{OMe} were measured on a STOE IPDS I diffractometer performing ω scans at 170(2) K. Diffraction intensities were measured using graphite-monochromatic Mo K α radiation ($\lambda = 0.71073$ Å). Data collection, indexing, initial cell refinements, frame integration, final cell refinements, and absorption corrections were accomplished with the program CrysAlis Pro (Agilent Technologies, Version 1.171.37.34, 2014) and X-Area, respectively. Space groups were assigned by analysis of the metric symmetry and systematic absences (determined by XPREP) and were further checked by PLATON^{38,39} for additional symmetry. Structures were solved by direct methods and refined against all data in the reported 2Θ ranges by full-matrix least squares on F^2 with the SHELXL program suite^{40,41} using the OLEX2 interface.⁴² The program PLATON SQUEEZE was used for the structures L_A^{Py} and [Ni₂L_A^{Thio}](ClO₄)₄ to eliminate non-refinable solvent molecules. 43 Crystallographic data as well as refinement parameters are presented in Tables S4-S7 in the ESI.†

Stopped-flow measurements

Time-dependent spectrophotometry was measured with a UV-Vis spectrophotometer S600 from Analytik Jena and a SFA-20 Rapid Kinetics Accessory from Hi-Tech Scientific. Temperature control was obtained with an attached cryostat and a cuvette-holder with a temperature-unit. The used MeCNsolutions were prepared from a stock-solution of MeCN saturated with CO_2 ($[CO_2]_{298K} = 0.28 \text{ mol L}^{-1}$)⁴⁴ and degassed MeCN. The complex was synthesized in situ in degassed MeCN under an N₂ atmosphere.

Electrochemical analysis

The electrochemical studies were performed on a Gamry Reference 600 in 100 mM tetrabutylammonium hexafluorophosphate (TBAPF₆) as a supporting electrolyte, 20 mM Ni(ClO₄)₆·6H₂O and 10 mM L_A^R in degassed MeCN. Glassy carbon, Pt wire and Ag|AgNO₃ (10 mM) in MeCN were used as working, counter and reference electrodes respectively. Cyclic voltammograms (CV) were recorded between -2.0 and +1.5 V at 100 mV s⁻¹ in degassed MeCN and after CO₂ purging. The working electrode was polished with alumina paste 0.3 µm (Buehler) before each measurement. The solutions were purged for 10, 20, 60 and 120 seconds with CO2. The results are reported versus Fc/Fc⁺.

DFT calculations

All calculations have been performed with the ORCA program.45 The BP86 functional46 was used along with the Def2-svp basis set. 47 The resolution of the identity (RI) approximation has been employed to speed up the calculation time.48,49 Scalar relativistic effects are included in zero order regular approach (ZORA).50,51 Solvent effects were taken into account by using the COSMO solvation model.⁵²

General synthetic procedure for L_I^R

In a typical experiment, the respective dialdehyde (3.6 mmol) was dissolved in MeCN (100 mL). A solution of Tren (2.5 mmol) in MeCN (20 mL) was added dropwise to the solution within 3 h. The solution was stirred at RT for 12 h and the formed solid was filtered off, washed with MeCN and dried in a vacuum to give the hexa-imine L_I^R .

L_I^{OMe}: white solid, 85% yield. ¹H NMR (400 MHz, CDCl₃): δ [ppm] = 7.73 (d, 6H), 7.60 (s, 6H), 5.17 (s, 3H), 3.90 (s, 9H), 3.77 (s, 6H), 3.32 (s, 6H), 2.80 (s, 12H). ¹³C NMR (100 MHz, $CDCl_3$: δ [ppm] = 160.8, 160.6, 138.4, 125.9, 112.8, 60.1, 56.2, 56.0. ESI-MS calc. for $[C_{39}H_{48}N_8O_3]^+$: m/z = 677.38. Found: m/z = 677.08. IR (KBr, cm⁻¹): 2948, 2875, 2823, 1643, 1590, 1454, 1372, 1197, 1156, 1063, 1030, 863, 697, 657. Anal. calc. for $[C_{39}H_{60}N_8K + MeOH + H_2O]$: N, 15.38; C, 65.80; H, 9.11. Found: N, 15.75; C, 65.73; H, 8.9.

L_I^{Me}: white solid, 85% yield. ¹H NMR (400 MHz, CDCl₃): δ [ppm] = 8.00 (s, 6H), 7.57 (s, 6H), 5.18 (s, 3H), 3.76 (s, 6H), 3.31 (s, 6H), 3.05-2.62 (m, 12H), 2.54 (s, 9H). ¹³C NMR (100 MHz, CDCl₃): δ [ppm] = 161.2, 139.1, 136.9, 130.2, 128.0, 60.1, 56.1, 21.4. ESI-MS calc. for $[C_{39}H_{49}N_8]^+$: m/z = 629.41. Found: m/z = 629.24. IR (KBr, cm⁻¹): 2947, 2871, 2802, 1641, 1439, 1372, 1333, 1289, 1158, 1068, 1032, 921, 866, 739, 691. Anal. calc. for $[C_{39}H_{48}N_8 + \frac{1}{2}H_2O]$: N, 17.57; C, 73.44; H, 7.74. Found: N, 17.77; C, 73.38; H, 7.75.

 $L_{\rm I}^{\rm F}$: white solid, 62% yield. ¹H NMR (400 MHz, CDCl₃): δ [ppm] = 7.88 (dd, 6H), 7.58 (dd, 6H), 5.30 (s, 3H), 3.73 (s, 6H), 3.32 (s, 6H), 2.80 (s, 12H). ¹³C NMR (100 MHz, CDCl₃): δ [ppm] = 165.0, 162.5, 159.3 (d, J = 2.7 Hz), 139.5 (d, J = 7.4 Hz), 127.8, 114.2, 113.9, 60.0, 55.8. ESI-MS calc. for $[C_{36}H_{40}F_3N_8]^+$: m/z = 641.33. Found: m/z = 641.00. IR (KBr, cm⁻¹): 3070, 2945, 2905, 2885, 2838, 2807, 2734, 1644, 1611, 1432, 1383,

Dalton Transactions Paper

1364, 1287, 1123, 1154, 1067, 1032, 656. Anal. calc. for [C₃₆H₃₉F₃N₈]: N, 17.49; C, 67.48; H, 6.14. Found: N, 17.42; C, 67.19; H, 6.26.

L_I^{OH}: yellow solid, 57% yield. ¹H NMR (200 MHz, MeOD) δ [ppm]: 7.60 (d, 12H), 5.19 (s, 3H). 3.69 (br s, 3H, OH), 2.73 (m, 12H), 2.56 (m, 12H). ¹³C NMR (50 MHz, MeOD) δ (ppm): 163.1, 160.8, 139.1, 125.8, 116.1, 60.7, 57.6, 57.1, 40.0. ESI-MS calc. for $[C_{36}H_{44}N_8ONa]^+$: m/z = 659.34. Found: m/z = 659.23. IR (KBr, cm⁻¹): 3352, 2948, 2885, 2832, 1640, 1569, 1435, 1368, 1335, 1240, 1173, 1065, 1024, 924, 876, 733, 692.

General synthetic procedures for L_A^R

The hexa-imine L_I^R (1.0 mmol) was dissolved in dry MeOH (50 mL) and heated to reflux. KBH4 (11.7 mmol) was then added in small portions. The reaction mixture was heated under reflux overnight, cooled to RT and the solvent was removed under reduced pressure. The residue was suspended in 2 M NH₄Cl solution (40 mL), extracted with DCM (3 × 40 mL) and dried over MgSO₄. The organic solvent was removed under reduced pressure and the remaining solid was dried in a vacuum to give L_A^R .

 L_A^{OMe} : white solid, 92%. ¹H NMR (400 MHz, CDCl₃): δ [ppm] = 6.73 (d, 9H), 4.33 (s, 6H), 3.77 (s, 9H), 3.65 (s, 12H), 2.65 (d, 24H). ¹³C NMR (100 MHz, CDCl₃): δ [ppm] = 159.7, 140.7, 119.9, 113.1, 55.5, 54.5, 53.0, 47.5. ESI-MS calc. for $[C_{30}H_{61}N_8O_3]^+$: m/z = 689.49. Found: m/z = 689.29. IR (KBr, cm⁻¹): 3415, 2944, 2831, 1600, 1462, 1338, 1292, 1160, 1056, 846, 707. Anal. calc. for C₃₉H₄₈N₈O₃: N, 16.56; C, 69.21; H, 7.15. Found: N, 16.58; C, 68.88; H, 7.05.

L_A^{Me}: white solid, 99% yield. ¹H NMR (400 MHz, CDCl₃): δ [ppm] = 7.01 (s, 6H), 6.92 (s, 3H), 3.93 (s, 6H), 3.59 (s, 12H), 2.68 (dd, 24H), 2.27 (s, 9H). ¹³C NMR (100 MHz, CDCl₃): δ [ppm] = 139.5, 138.3, 128.4, 124.7, 54.7, 53.1, 47.6, 21.4. ESI-MS calc. for $[C_{39}H_{619}N_8]^+$: m/z = 641.50. Found: m/z = 641.50641.30. IR (KBr, cm⁻¹): 3443 (br s), 3006, 2919, 2832, 1644, 1607, 1460, 1291, 1162, 1110, 1076, 846, 714. Anal. calc. for $[C_{39}H_{60}KN_8 + \frac{1}{2}H_2O + MeOH]$: N, 15.35; C, 65.80; H, 9.11. Found: N, 15.57; C, 65.73; H, 8.90.

L_A^F: white solid, 99% yield. ¹H NMR (400 MHz, CDCl₃): δ [ppm] = 6.95 (d, 6H), 6.70 (s, 3H), 3.58 (s, 12H), 2.69–2.59 (m, 24H), 2.03 (s, 6H). ¹³C NMR (100 MHz, CDCl₃): δ [ppm] = 164.3, 161.9, 143.2 (d, J = 7.2 Hz), 122.2, 113.1, 112.9, 55.5, 53.3, 48.0. ESI-MS calc. for $[C_{36}H_{52}F_3N_8]^+$: m/z = 653.43. Found: m/z = 653.20. IR (KBr, cm⁻¹): 3420, 3299, 3075, 2946, 2887, 2818, 1643, 1623, 1453, 1361, 1283, 1149, 1129, 1111, 1000, 788, 662. Anal. calc. for [C₃₆H₅₁F₃N₈]: N, 17.16; C, 65.23; H, 7.87. Found: N, 16.95; C, 65.34; H, 7.50.

L_A^{OH,Me}: brown solid, 27% yield. ¹H NMR (400 MHz, CDCl₃) [ppm]: 6.68 (s, 6H), 3.70 (s, 12H), 2.65 (d, J = 6.4 Hz), 2.156 (s, 9H). ¹³C NMR (100 MHz, CDCl₃): δ [ppm] = 155.6, 129.0, 127.0, 123.8, 54.8, 50.9, 46.8, 20.5. ESI-MS calc. for $\left[C_{39}H_{61}N_8O_3\right]^+$: m/z = 689.49. Found: m/z = 689.31, 711.15 [M + Na]⁺. IR (KBr, cm⁻¹): 3442, 3309, 2964, 2912, 2842, 1735, 1612, 1477, 1262, 1097, 1024, 866, 803, 695. Anal. calc. for $[C_{39}H_{60}N_8K_2O_3 + 4CH_3OH]$: N, 12.52; C, 57.69; H, 8.56. Found: N, 11.93; C, 57.4; H, 8.23.

L_A^{OH}: no extraction with DCM was required. The residue was washed with MeOH and precipitates were filtered off. The solvent was removed under reduced pressure to afford a brownish solid, 84% yield. ¹H NMR (200 MHz, MeOD) δ [ppm]: 6.96 (s, 3H), 6.84 (s, 6H), 3.81 (s, 12H), 3.08 (m, 12H), 2.78 (m, 12H). ¹³C NMR (50 MHz, MeOD) δ [ppm]: 159.3, 138.7, 121.5, 117.0, 53.8, 52.9, 47.7. ESI-MS calc. for $[C_{36}H_{55}N_8O_3]^+$: m/z = 647.43. Found: m/z = 647.45. IR (KBr, cm⁻¹): 3244, 2956, 2836, 1655, 1597, 1456, 1306, 1165, 1101, 1000, 846, 713.

General synthetic procedures for $[Ni_2L_A^R](Cl)_v(ClO_4)_x$ (x = 2-4; $y = 0, 1)^{23}$

The respective compound LAR (0.038 mmol) was dissolved in 2 mL degassed MeCN/MeOH or MeCN/EtOH 4:1. A solution of Ni(ClO₄)₂·6H₂O (0.078 mmol) in MeCN/EtOH 4:1 was added and the mixture was stirred for 12 h at room temperature. The solvent was removed under reduced pressure and the residue was crystallized from MeCN/Et2O, or through slow evaporation of the solvent.

[Ni₂L_A^{Me}](Cl)(ClO₄)₃: blue solid, 69% yield. ESI-MS calc. for $[C_{39}H_{61}N_8Ni_2 + 3ClO_4]^+$: m/z = 1054.22. Found: m/z = 1054.7. IR (KBr, cm⁻¹): 3421, 2957, 1637, 1613, 1443, 1385, 1146, 1115, 1086, 874, 842, 753, 629.

 $[Ni_2L^F](Cl)(ClO_4)_3$: blue solid, 60% yield. ESI-MS calc. for $[C_{36}H_{51}F_3N_8Ni_2 + CH_3CN]^+$: m/z = 809.32. Found: m/z = 809.3. IR (KBr, cm⁻¹): 3422, 2933, 2875, 1665, 1626, 1598, 1459, 1342, 1300, 1144, 1115, 1085, 982, 878, 841, 664, 628.

[Ni₂L_AOMe](Cl)(ClO₄)₃: green solid, 90% yield. ESI-MS calc. for $[C_{39}H_{55}N_8Ni_2O_3 + 2ClO_4^- + CH_3CN]^+$: m/z = 1038.23. Found: m/z = 1038.4. IR (KBr, cm⁻¹): 3413, 2955, 2843, 1651, 1605, 1467, 1440, 1340, 1302, 1143, 1112, 1084, 842, 710,

 $[Ni_2L_A^{Fur}](Cl)(ClO_4)_3$: green solid, 42% yield. ESI-MS calc. for $[C_{30}H_{49}N_8Ni_2O_3 + 3ClO_4]^+$: m/z = 982.11. Found: m/z =982.63. IR (KBr, cm⁻¹): 3458, 3250, 2858, 1653, 1450, 1344, 1092, 928, 816, 627.

[Ni₂L_A^{Py}](Cl)(ClO₄)₃: dark blue solid, 98% yield. ESI-MS calc. for $[C_{33}H_{51}N_{11}Ni_2 + CH_3CN]^+$: m/z = 758.33, 1015.16 $[+3ClO_4]$. Found: m/z = 758.06, 1015.08 $[+3ClO_4]$. IR (KBr, cm⁻¹): 3418, 1940, 1883, 1641, 1608, 1458, 1143, 1085, 794,

[Ni₂L_A^{Thio}](ClO₄)₄: blue crystals, 26% yield. ESI-MS calc. for $[C_{30}H_{48}N_8Ni_2S_3 + 3ClO_4]^+$: m/z = 1029.0. Found: m/z = 1028.5. IR (KBr, cm⁻¹): 3416 (br), 3256, 3061, 2949, 2835, 1629, 1448, 1294, 1147, 1112, 1088, 1007, 885, 742, 683, 629.

[Ni₂L_AOH](ClO₄)₃: green solid, 87% yield. ESI-MS calc. for $[C_{36}H_{55}Cl_2N_8Ni_2O_7]^+$: m/z = 897.22. Found: m/z = 896.8. IR (KBr, cm⁻¹): 3387, 3230, 2965, 2781, 1642, 1603, 1460, 1310, 1140, 1086, 995, 628.

[Ni₂L_AOH,Me](ClO₄)₂ was obtained by crystallization from MeCN. Blue crystals, 13% vield. ESI-MS calc. for $[C_{39}H_{59}N_8Ni_2O_3]^+$: m/z = 803.35. Found: m/z = 803.25. IR (KBr, cm⁻¹): 3422, 3272, 2919, 2808, 1637, 1473, 1309, 1261, 1085, 954, 821, 626.

Acknowledgements

This work was supported by the Fonds der Chemischen Industrie (Liebig grant to U.-P. A.) and through the Deutsche Forschungsgemeinschaft (Emmy Noether grant to U.-P. A., AP242/2-1) as well as the Max Planck Gesellschaft (M. v. G.). F. M. gratefully acknowledges financial support by the Deutsche Bundesstiftung Umwelt. We thank Prof. Christian Herrmann, Physical Chemistry, Ruhr-University Bochum, for providing the stopped flow equipment and for valuable discussions. Open Access funding provided by the Max Planck Society.

Notes and references

- 1 V. Begoña and S. B. Salvador, Eur. J. Inorg. Chem., 2008, 1, 84-97.
- C. E. Wilmer, O. K. Farha, Y.-S. Bae, J. T. Hupp and R. Q. Snurr, *Energy Environ. Sci.*, 2012, 5, 9849–9856.
- 3 J. Fu, S. Das, G. Xing, T. Ben, V. Valtchev and S. Qiu, *J. Am. Chem. Soc.*, 2016, **138**, 7673–7680.
- 4 Y. Zeng, R. Zou and Y. Zhao, Adv. Mater., 2016, 28, 2855-2873.
- 5 J. Jaschik, M. Jaschik and K. Warmuziński, *Chem. Process Eng.*, 2016, 37, 29–39.
- 6 M. Tong, Q. Yang and C. Zhong, *Microporous Mesoporous Mater.*, 2015, **210**, 142–148.
- 7 E. S. Sanz-Pérez, C. R. Murdock, S. A. Didas and C. W. Jones, *Chem. Rev.*, 2016, 116, 11840–11876.
- 8 Y. Jia, H. Su, Z. Wang, Y.-L. E. Wong, X. Chen, M. Wang and T. W. D. C. Chan, *Anal. Chem.*, 2016, **88**, 9364–9367.
- 9 L. Fabbrizzi, P. Pallavicini, L. Parodi, A. Perotti, N. Sardone and A. Taglietti, *Inorg. Chim. Acta*, 1996, 7–9.
- 10 M. G. Drew, J. Hunter, D. J. Marrs, J. Nelson and C. Harding, J. Chem. Soc., Dalton Trans., 1992, 3235–3242.
- 11 C. J. Harding, F. E. Mabbs, E. J. L. MacInnes, V. McKee and J. Nelson, *J. Chem. Soc., Dalton Trans.*, 1996, 3227–3230.
- 12 M. G. Basallote, J. Durán, M. J. Fernández-Trujillo and M. A. Máñez, *J. Chem. Soc., Dalton Trans.*, 2002, 2074–2079.
- 13 J.-M. Lehn, Pure Appl. Chem., 1980, 11, 2442-2459.
- 14 J. M. Lehn, Pure Appl. Chem., 1978, 50, 871-892.
- 15 V. Amendola, L. Fabbrizzi, C. Mangano, P. Pallavicini and M. Zema, *Inorg. Chim. Acta*, 2002, 337, 70–74.
- 16 V. Amendola, M. Bonizzoni, D. Esteban-Gómez, L. Fabbrizzi, M. Licchelli, F. Sancenón and A. Taglietti, Coord. Chem. Rev., 2006, 250, 1451–1470.
- 17 A. A. C. Wild, K. Fennell, G. G. Morgan, C. M. Hewage and J. P. G. Malthouse, *Dalton Trans.*, 2014, 43, 13557.
- 18 P. Mateus, R. Delgado, V. André and M. T. Duarte, *Inorg. Chem.*, 2015, **54**, 229–240.
- 19 M. M. El-Hendawy, S. Bandaru, N. J. English and D. A. Mooney, *Catal. Sci. Technol.*, 2013, 3, 2234–2243.
- 20 Y. Dussart, C. Harding, P. Dalgaard, C. McKenzie, R. Kadirvelraj, V. McKee and J. Nelson, *J. Chem. Soc., Dalton Trans.*, 2002, 1704–1713.

- 21 J.-M. Chen, W. Wei, X.-L. Feng and T.-B. Lu, *Chem. Asian J.*, 2007, 2, 710–719.
- 22 F. Arnaud-Neu, S. Fuangswasdi, B. Maubert, J. Nelson and V. McKee, *Inorg. Chem.*, 2000, **39**, 573–579.
- 23 F. Möller, K. Merz, C. Herrmann and U.-P. Apfel, *Dalton Trans.*, 2016, **45**, 904–907.
- 24 P. J. Cragg, A Practical Guide to Supramolecular Chemistry, Wiley-VCH, England, 2005.
- 25 R. M. Town, V. McKee, M. Arthurs and J. Nelson, *J. Chem. Educ.*, 2001, 78, 1269.
- 26 D. Chen and A. E. Martell, *Tetrahedron*, 1991, 47, 6895–6902.
- 27 M. F. Mahon, J. McGinley, A. Denise Rooney and J. M. D. Walsh, *Inorg. Chim. Acta*, 2009, 362, 2353– 2360
- 28 J. Nelson, V. McKee and G. G. Morgan, *Prog. Inorg. Chem.*, 1998, 47, 167–316.
- 29 M. G. Drew, C. J. Harding, O. W. Howarth, Q. Lu, D. J. Marrs, G. G. Morgan, V. McKee and J. Nelson, *J. Chem. Soc., Dalton Trans.*, 1996, 3021–3030.
- 30 T. Lu, X. Zhuang, Y. Li and S. Chen, *J. Am. Chem. Soc.*, 2004, 15, 4760–4761.
- 31 D. Huang, O. V. Makhlynets, L. L. Tan, S. C. Lee, E. V. Rybak-Akimova and R. H. Holm, *Inorg. Chem.*, 2011, 50, 10070–10081.
- 32 V. McKee, J. Nelson and R. M. Town, *Chem. Soc. Rev.*, 2003, 32, 309–325.
- 33 A. Escuer, C. J. Harding, Y. Dussart, J. Nelson, V. McKee and R. Vicente, *J. Chem. Soc., Dalton Trans.*, 1999, 223–228.
- 34 R. P. Jimenez, M. Parvez, T. C. Sutherland and J. Viccars, *Eur. J. Org. Chem.*, 2009, 5635–5646.
- 35 F. Banishoeib, A. Henckens, S. Fourier, G. Vanhooyland, M. Breselge, J. Manca, T. J. Cleij, L. Lutsen, D. Vanderzande, L. H. Nguyen, H. Neugebauer and N. S. Sariciftci, *Thin Solid Films*, 2008, 516, 3978–3988.
- 36 J. Bao, Z. Zhang, R. Tang, H. Han and Z. Yang, *J. Lumin.*, 2013, **136**, 68–74.
- 37 X.-Y. Yang, W. S. Tay, Y. Li, S. A. Pullarkat and P.-H. Leung, *Organometallics*, 2015, **34**, 1582–1588.
- 38 A. L. Spek, J. Appl. Crystallogr., 2003, 36, 7-13.
- A. L. Spek, *Acta Crystallogr.*, *Sect. D: Biol. Crystallogr.*, 2009,
 5, 148–155.
- 40 G. M. Sheldrick, Acta Crystallogr., Sect. A: Fundam. Crystallogr., 2008, 64, 112–122.
- 41 C. B. Hübschle, G. M. Sheldrick and B. Dittrich, *J. Appl. Crystallogr.*, 2011, 44, 1281–1284.
- 42 O. V. Dolomanov, L. J. Bourhis, R. J. Gildea, J. A. K. Howard and H. Puschmann, *J. Appl. Crystallogr.*, 2009, 42, 339–341.
- 43 A. L. Spek, Acta Crystallogr., Sect. C: Cryst. Struct. Commun., 2015, 71, 9–18.
- 44 A. Gennaro, A. A. Isse and E. Vianello, *J. Electroanal. Chem.*, 1990, **289**, 203–215.
- 45 F. Neese, Wiley Interdiscip. Rev.: Comput. Mol. Sci., 2012, 2, 73–78.
- 46 A. D. Becke, Phys. Rev. A, 1988, 38, 3098-3100.

Paper

- 47 D. A. Pantazis, X.-Y. Chen, C. R. Landis and F. Neese, *J. Chem. Theor. Comput.*, 2008, 4, 908–919.
- 48 K. Eichkorn, O. Treutler, H. Öhm, M. Häser and R. Ahlrichs, *Chem. Phys. Lett.*, 1995, **240**, 283–290.
- 49 K. Eichkorn, F. Weigend, O. Treutler and R. Ahlrichs, *Theor. Chem. Acc.*, 1997, **97**, 119–124.
- 50 E. van Lenthe, A. van der Avoird and P. E. S. Wormer, J. Chem. Phys., 1998, 108, 4783–4796.
- 51 E. van Lenthe, J. G. Snijders and E. J. Baerends, *J. Chem. Phys.*, 1996, **105**, 6505–6516.
- 52 A. Klamt and G. Schüürmann, *J. Chem. Soc., Perkin Trans.* 2, 1993, 799–805.