



Cite this: *J. Mater. Chem. C*, 2018, **6**, 668

Correction: Bright and efficient light-emitting diodes based on MA/Cs double cation perovskite nanocrystals

Bing Xu,^{ab} Weigao Wang,^a Xiaoli Zhang,^{ab} Wanyu Cao,^c Dan Wu,^d Sheng Liu,^b Haitao Dai,^c Shuming Chen,^{*a} Kai Wang^{*a} and Xiaowei Sun^{*a}

DOI: 10.1039/c7tc90204b

rsc.li/materials-c

Correction for 'Bright and efficient light-emitting diodes based on MA/Cs double cation perovskite nanocrystals' by Bing Xu *et al.*, *J. Mater. Chem. C*, 2017, **5**, 6123–6128.

The authors regret a mistake in reporting the solvent used in the fabrication of MAPbBr₃ in the Experimental section.

The sentence "For the MAPbBr₃ synthesis, a 200 µL solution containing 0.1 mmol of PbBr₂, 0.16 mmol of MABr, 20 µL of oleylamine and 500 µL of oleic acid in dimethylformamide were added to 10 mL **acetone** at room temperature to produce a colloidal solution." should read "For the MAPbBr₃ synthesis, a 200 µL solution containing 0.1 mmol of PbBr₂, 0.16 mmol of MABr, 20 µL of oleylamine and 500 µL of oleic acid in dimethylformamide were added to 10 mL **toluene** at room temperature to produce a colloidal solution."

The Royal Society of Chemistry apologises for these errors and any consequent inconvenience to authors and readers.

^a Department of Electrical & Electronic Engineering, Southern University of Science and Technology, Shenzhen, 518055, China. E-mail: chensm@sustc.edu.cn

^b School of Power and Mechanical Engineering, Wuhan University, Wuhan, 430072, China

^c Tianjin Key Laboratory of Low Dimensional Materials Physics and Preparing Technology, School of Science, Tianjin University, Tianjin 300072, China

^d School of Electrical & Electronic Engineering, Nanyang Technological University, 50 Nanyang Avenue, 639798, Singapore

