

Allylic sp^3 C–H borylation of alkenes via allyl-Pd intermediates: an efficient route to allylboronates†

 Hong-Ping Deng,^a Lars Eriksson^b and Kálmán J. Szabó*^a

 Cite this: *Chem. Commun.*, 2014, 50, 9207

 Received 30th May 2014,
 Accepted 26th June 2014

DOI: 10.1039/c4cc04151h

www.rsc.org/chemcomm

Palladium catalyzed allylic C–H functionalization was performed using exocyclic alkene substrates. Multi-component synthesis of stereodefined homoallylic alcohols could be performed using a reaction sequence involving allylic C–H borylation and allylation of aldehydes.

Catalytic C–H borylation has become a practically useful synthetic method for preparation of organoboronates.¹ The main reason is that these transition metal catalyzed C–H functionalization reactions can be performed under relatively mild conditions with remarkably high selectivity^{1b,c} usually using B_2Pin_2 as a boronate source. The largest efforts have been focused on sp^2 C–H borylation of aromatic and alkene substrates to obtain aryl/heteroaryl² and vinyl³ boronates. However, in the last couple of years increased attention has been focused on development of sp^3 C–H functionalization methods.⁴ These studies involved functionalization of aliphatic C–H bonds^{4d–h,l,m,o–q} usually directed by heteroatoms, benzylic C–H bonds^{4i–k} and there are a few examples of allylic C–H borylation^{4a–c} as well. A selective allylic C–H borylation^{4a–c} is particularly challenging to achieve due to two main reasons: (i) under catalytic conditions allylboronates very easily rearrange to the more stable vinylboronates.^{3b,c,f,4c} Thus, even if the kinetic product is an allylboronate, the thermodynamic (final) product of the C–H borylation of alkenes is vinylboronate; *i.e.* an overall sp^2 instead of sp^3 C–H bond functionalization; and (ii) for non-symmetrical organometallic intermediates (*e.g.* allyl or alkyl–metal species) the regioselectivity of the borylation is difficult to control. Therefore, only a very few transition metal catalyzed methods are available for allylic C–H borylation of alkenes^{4a–c} and because of (i) and (ii) the substrate scope is also very narrow.

The previously developed procedures providing predominantly allylboronate products are based on C–H functionalization of simple cycloalkenes (Fig. 1). Sabo-Etienne and Caballero^{4a} have shown that





Fig. 1 Overview of catalytic allylic C–H borylations.

cycloheptene undergoes hydroboration and allylic C–H borylation in the presence of catalytic amounts of bis(dihydrogen)Ru complex. We have shown^{4b,c} that simple cycloalkenes can be reacted with B_2Pin_2 in the presence of Ir-catalysts to give allyl-Bpin compounds.

Interestingly, other catalytic conditions with the above *endo*-cyclic alkene substrates using rhodium^{3g} or palladium^{3c,f} catalysts also give allyl-Bpin products in varying amounts. However, for substrates with an *exo*-cyclic double bond allylic C–H borylation has never been reported. This can probably be explained by the mechanistic features of the currently available Ru, Rh, Ir and Pd catalyzed methods. In all cases initial formation of an M–Bpin complex can be postulated (eqn (1)), which undergoes a *syn* insertion into the double bond followed by a *syn* selective β -hydride elimination. However, acyclic compounds can undergo unhindered rotation of the σ -bonds, and therefore the β -hydride elimination may easily result in the thermodynamically more stable vinyl-Bpin form.^{4c}



Therefore, we decided to develop a new sp^3 allylic C–H borylation reaction based on an alternative mechanistic concept. We hypothesized that a Pd-catalyzed process based on initial formation of an allyl-Pd complex followed by transmetalation⁵ with B_2Pin_2 may

^a Department of Organic Chemistry, Stockholm University, Sweden

^b Department of Inorganic and Structural Chemistry, Stockholm University, Sweden.

 E-mail: kalmans@organ.su.se; Web: <http://www.organ.su.se/ks/>;

Fax: +46-8-15 49 08

† CCDC 1000349. For crystallographic data in CIF or other electronic format see DOI: 10.1039/c4cc04151h

avoid the termination of the reaction with β -hydride elimination. The realization of this idea is very challenging, as closing the catalytic cycle (see below) requires use of oxidants, while B_2Pin_2 is a reductant and allylboronates are sensitive to oxidation.⁶

We directed the initial studies to C–H functionalization of β -pinene **1a**, as these compounds readily form⁷ (η^3 -allyl)palladium complexes with stoichiometric amounts of Pd-salts. Indeed, when **1a**, **3a**, an appropriate oxidant and catalytic amounts of Pd(TFA)₂ were mixed in (CD₃)₂CO, formation of borylated β -pinene was observed (Fig. 1c). Using deuterated acetone enabled us to follow the reaction by ¹H NMR. The ¹H NMR of the reaction mixture showed that the reaction was not completed, probably because of product inhibition. When the allylboronate product was quenched with nitro-benzaldehyde (**2**), the corresponding homoallylic alcohol **4a** was formed selectively as a single, regio-stereoisomer.

Gratifyingly, the entire procedure with **1a**, **2**, **3a**, the oxidant (BQ), TFA and the Pd-catalyst could be performed as a multi-component⁸ (or genuine one-pot) reaction (Table 1, entry 1). As we used optically active β -pinene, the multicomponent C–H borylation–allylation sequence gave an enantiomerically pure product (**4a**). The structure of **4a** was assigned on the basis of single crystal X-ray diffraction. Subsequently, we studied the synthetic scope of the reaction. We have found that cyclic substrates with an exocyclic double bond give synthetically useful yields in the C–H borylation based allylation of aldehydes (Table 1). In most cases (except **1a**) we obtained complex mixtures and low yields, when we used BQ as an oxidant. However, 2,6-dimethyl BQ (DMBQ) successfully replaced BQ. Deuterated acetone proved to be an ideal solvent in most cases as it allowed us to study the crude mixtures by ¹H NMR. In some cases, the process was slow in acetone (e.g. entries 4–6 and 10) and therefore the solvent was changed to trifluoro toluene, which gave a higher reaction rate.

Nitro-benzaldehyde **2** could be replaced by benzaldehyde or aliphatic aldehyde (entries 2 and 3). The multicomponent reaction is still very selective but the yield was dropped (*cf.* entries 1–3). The reaction with six-membered ring based substrates **1b–d** gave exclusively the branched allylic products **4d–e** (entries 4 and 5). There are three stereogenic carbons in product **4f**, thus statistically four diastereomers could be formed. However, the reaction proceeds with a remarkably high stereoselectivity, as only two diastereomers were obtained in a ratio of 9:1 (entry 6). The reactions for the five membered ring based substrate **1e** proceeded faster than for the six membered ring analogs, and therefore the reactions could be conducted at rt. The best yield and selectivity were obtained, when DMBQ was replaced by tetramethyl BQ as an oxidant (entry 7). The seven membered ring based substrate **1f** reacted with high yield (entry 8) but the regioselectivity was also lower than for the six-membered ring based substrates. In the case of **1g** containing an eight-membered ring the regioselectivity drops to 2.5:1 (entry 9). We had a limited success with borylation of heterocyclic substrates, such as **1h** (entry 10). This compound can also be transformed into **4j** with high selectivity but the yield is poor and we could not improve it by extensive optimization. For acyclic analogs the yield and the selectivity drop dramatically (entry 11). For example **1i** reacts very slowly and with low conversion probably because of inhibition of the Pd-catalyst. Interestingly,

Table 1 Allylic C–H borylation of alkenes^a


Entry	Alkene	Temp. (°C)/solvent	Yield ^b (%)	B:L	
				linear (L)	branched (B)
1		rt/(CD ₃) ₂ CO		4a 78	>50:1
2	1a	rt/(CD ₃) ₂ CO		4b ^c 58	>26:1
3	1a	rt/(CD ₃) ₂ CO		4c ^c 56	>50:1
4		40/PhCF ₃		4d 53	>50:1
5		40/PhCF ₃		4e 51	>50:1
6		40/PhCF ₃		4f 43	>50 ^d :1
7 ^e		rt/(CD ₃) ₂ CO		4g 51	>50:1
8		40/(CD ₃) ₂ CO		4h 69	8:1
9		40/(CD ₃) ₂ CO		4i 80	2.5:1
10 ^f		40/PhCF ₃		4j 38	>50:1
11		40/(CD ₃) ₂ CO		4k 14	15 ^g :1

^a Unless otherwise stated the reactions were carried out with **1** (0.1 mmol), **2**, nitro-benzaldehyde (0.2 mmol), **3a** (0.2 mmol), Pd(TFA)₂ (0.01 mmol), DMBQ (0.2 mmol) and TFA (0.05 mmol) in solvent (0.2–0.5 mL) for 24 h. ^b Isolated yields for the linear (L) and branched (B) products together. Unless otherwise stated the branched product was isolated as a single diastereomer. ^c PhCHO (entry 2) and *n*-C₆H₁₃CHO (entry 3) were used instead of nitro-benzaldehyde. ^d d.r. = 9:1. ^e Tetramethyl-benzoquinone instead of DMBQ. ^f Reaction was carried out with **1h** (0.2 mmol) and **2** (0.1 mmol). ^g d.r. = 3:2. Ar = 4-NO₂C₆H₄, DMBQ = 2,6-dimethylbenzoquinone.

the reaction can be performed with diboronic acid (**3b**) instead of B_2Pin_2 with a slight alteration of the reaction conditions (eqn (2)). This is a remarkable result, as it shows that highly oxidation



sensitive allylboronic acids⁶ can also be reaction intermediates under oxidative allylic C–H borylation conditions.



We suggest that the first step of the process is formation of allyl–palladium complex **5** by deprotonation and palladation of the allylic position of the substrate, such as **1b** (Fig. 2).⁷ The subsequent step is transmetalation by B_2pin_2 . It was shown⁵ that these reactions proceed easily, when weakly coordinating ligands are on Pd. This could explain that $\text{Pd}(\text{TFA})_2$ is an excellent catalyst for the process, while $\text{Pd}(\text{OAc})_2$ with the chelating acetate group is inefficient. Iwasawa and co-workers⁹ have recently shown that Pd–Bpin complexes are stable species. The reductive elimination of the Bpin group in **6** is supposed to be fast⁵ due to the strong *trans* influence of the Bpin ligand.¹⁰ It proceeds with a very high regioselectivity leading to the linear allylboronate product. This high regioselectivity is a prerequisite of the high selectivity of the allylation of aldehyde **2** affording the branched homoallylic product **4d**. The reductive elimination involves formation of $\text{Pd}(0)$, which has to be reoxidized at the closing of the catalytic cycle. The main role of the used quinone is reoxidation of $\text{Pd}(0)$ to $\text{Pd}(\text{II})$. Added trifluoroacetic acid increases the oxidation potential of the quinones and also catalyzes the allylboronation of the aldehydes.¹¹

In summary, we have shown for the first time that allylic C–H borylation can be performed with exocyclic alkenes. Multicomponent reaction involving this new C–H borylation–allylboronation sequence can be performed to obtain stereodefined homoallylic alcohols. The reaction proceeds *via* regioselective borylation and a subsequent regio- and stereoselective allylation. The mechanistically novel element in this reaction is that it proceeds *via* initial formation of an allyl–palladium intermediate, which then undergoes transmetalation with B_2pin_2 and a subsequent regioselective reductive elimination of the allylboronate product.




Fig. 2 Suggested catalytic cycle exemplified by substrate **1b**.

Support from the Swedish Research Council and the Knut och Alice Wallenbergs Foundation, as well as a post-doctoral fellowship for H.-P. Deng from the Wenner-Gren foundation, is gratefully acknowledged. The generous gift of B_2pin_2 from Allychem is appreciated.

Notes and references

- (a) D. G. Hall, *Boronic Acids*, Wiley, Weinheim, 2011; (b) I. A. I. Mkhaldid, J. M. Murphy, J. H. Barnard, T. B. Marder and J. F. Hartwig, *Chem. Rev.*, 2010, **110**, 890; (c) J. F. Hartwig, *Chem. Soc. Rev.*, 2011, **40**, 1992; (d) T. Ishiyama and N. Miyaura, *Pure Appl. Chem.*, 2006, **78**, 1369.
- (a) J.-Y. Cho, M. K. Tse, D. Holmes, R. E. Maleczka and M. R. Smith, *Science*, 2002, **295**, 305; (b) I. I. B. A. Vanchura, S. M. Preshlock, P. C. Roosen, V. A. Kallepalli, R. J. Staples, J. R. E. Maleczka, D. A. Singleton and I. I. I. M. R. Smith, *Chem. Commun.*, 2010, **46**, 7724; (c) T. Ishiyama, J. Takagi, K. Ishida, N. Miyaura, N. R. Anastasi and J. F. Hartwig, *J. Am. Chem. Soc.*, 2002, **124**, 390; (d) T. Ishiyama, J. Takagi, J. F. Hartwig and N. Miyaura, *Angew. Chem., Int. Ed.*, 2002, **41**, 3056; (e) T. A. Boebel and J. F. Hartwig, *J. Am. Chem. Soc.*, 2008, **130**, 7534; (f) D. W. Robbins, T. A. Boebel and J. F. Hartwig, *J. Am. Chem. Soc.*, 2010, **132**, 4068; (g) C. C. Tzschucke, J. M. Murphy and J. F. Hartwig, *Org. Lett.*, 2007, **9**, 761; (h) I. A. I. Mkhaldid, D. N. Coventry, D. Albesa-Jove, A. S. Batsanov, J. A. K. Howard, R. N. Perutz and T. B. Marder, *Angew. Chem., Int. Ed.*, 2006, **45**, 489; (i) T. Ishiyama, H. Isou, T. Kikuchi and N. Miyaura, *Chem. Commun.*, 2010, **46**, 159; (j) S. Kawamorita, H. Ohmiya and M. Sawamura, *J. Org. Chem.*, 2010, **75**, 3855; (k) K. Yamazaki, S. Kawamorita, H. Ohmiya and M. Sawamura, *Org. Lett.*, 2010, **12**, 3978; (l) A. Ros, R. López-Rodríguez, B. Estepa, E. Álvarez, R. Fernández and J. M. Lassaletta, *J. Am. Chem. Soc.*, 2012, **134**, 4573; (m) H.-X. Dai and J.-Q. Yu, *J. Am. Chem. Soc.*, 2012, **134**, 134; (n) B. Xiao, Y.-M. Li, Z.-J. Liu, H.-Y. Yang and Y. Fu, *Chem. Commun.*, 2012, **48**, 4854; (o) Y. Kuninobu, T. Iwanaga, T. Omura and K. Takai, *Angew. Chem., Int. Ed.*, 2013, **52**, 4431.
- (a) I. A. I. Mkhaldid, R. B. Coapes, S. N. Edes, D. N. Coventry, F. E. S. Souza, R. L. Thomas, J. J. Hall, S. W. Bi, Z. Y. Lin and T. B. Marder, *Dalton Trans.*, 2008, 1055; (b) V. J. Olsson and K. J. Szabó, *Org. Lett.*, 2008, **10**, 3129; (c) N. Selander, B. Willy and K. J. Szabó, *Angew. Chem., Int. Ed.*, 2010, **49**, 4051; (d) T. Ohmura, Y. Takasaki, H. Furukawa and M. Suginome, *Angew. Chem., Int. Ed.*, 2009, **48**, 2372; (e) J. Takaya, N. Kirai and N. Iwasawa, *J. Am. Chem. Soc.*, 2011, **133**, 12980; (f) N. Kirai, S. Iguchi, T. Ito, J. Takaya and N. Iwasawa, *Bull. Chem. Soc. Jpn.*, 2013, **86**, 784; (g) A. Kondoh and T. F. Jamison, *Chem. Commun.*, 2010, **46**, 907; (h) T. Kikuchi, J. Takagi, H. Isou, T. Ishiyama and N. Miyaura, *Chem. – Asian J.*, 2008, **3**, 2082; (i) T. Kikuchi, J. Takagi, T. Ishiyama and N. Miyaura, *Chem. Lett.*, 2008, **37**, 664; (j) I. Sasaki, H. Doi, T. Hashimoto, T. Kikuchi, H. Ito and T. Ishiyama, *Chem. Commun.*, 2013, **49**, 7546.
- (a) A. Caballero and S. Sabo-Etienne, *Organometallics*, 2007, **26**, 1191; (b) V. J. Olsson and K. J. Szabó, *Angew. Chem., Int. Ed.*, 2007, **46**, 6891; (c) V. J. Olsson and K. J. Szabó, *J. Org. Chem.*, 2009, **74**, 7715; (d) H. Y. Chen, S. Schlecht, T. C. Semple and J. F. Hartwig, *Science*, 2000, **287**, 1995; (e) J. M. Murphy, J. D. Lawrence, K. Kawamura, C. Incarvito and J. F. Hartwig, *J. Am. Chem. Soc.*, 2006, **128**, 13684; (f) C. W. Liskey and J. F. Hartwig, *J. Am. Chem. Soc.*, 2012, **134**, 12422; (g) C. W. Liskey and J. F. Hartwig, *J. Am. Chem. Soc.*, 2013, **135**, 3375; (h) S. H. Cho and J. F. Hartwig, *J. Am. Chem. Soc.*, 2013, **135**, 8157; (i) T. A. Boebel and J. F. Hartwig, *Organometallics*, 2008, **27**, 6013; (j) S. Shimada, A. S. Batsanov, J. A. K. Howard and T. B. Marder, *Angew. Chem., Int. Ed.*, 2001, **40**, 2168; (k) T. Ishiyama, K. Ishida, J. Takagi and N. Miyaura, *Chem. Lett.*, 2001, 1082; (l) T. Ohmura, T. Torigoe and M. Suginome, *J. Am. Chem. Soc.*, 2012, **134**, 17416; (m) T. Ohmura, T. Torigoe and M. Suginome, *Organometallics*, 2013, **32**, 6170; (n) T. Ohmura, T. Torigoe and M. Suginome, *Chem. Commun.*, 2014, **50**, 6333; (o) S. Kawamorita, T. Miyazaki, T. Iwai, H. Ohmiya and M. Sawamura, *J. Am. Chem. Soc.*, 2012, **134**, 12924; (p) S. Kawamorita, R. Murakami, T. Iwai and M. Sawamura, *J. Am. Chem. Soc.*, 2013, **135**, 2947; (q) T. Mita, Y. Ikeda, K. Michigami and Y. Sato, *Chem. Commun.*, 2013, **49**, 5601.



5 J. M. Larsson and K. J. Szabó, *J. Am. Chem. Soc.*, 2013, **135**, 443.
6 M. Raducan, R. Alam and K. J. Szabó, *Angew. Chem., Int. Ed.*, 2012, **51**, 13050.
7 B. M. Trost, P. E. Streege, L. Weber, T. J. Fullerton and T. J. Dietsche, *J. Am. Chem. Soc.*, 1978, **100**, 3407.
8 K. J. Szabó, in *Science of Synthesis Reference Library: Multicomponent Reactions*, ed. T. J. J. Müller, Thieme, Stuttgart, 2013, p. 345.
9 N. Kirai, J. Takaya and N. Iwasawa, *J. Am. Chem. Soc.*, 2013, **135**, 2493.
10 J. Zhu, Z. Lin and T. B. Marder, *Inorg. Chem.*, 2005, **44**, 9384.
11 (a) V. Rauniyar and D. G. Hall, *Angew. Chem., Int. Ed.*, 2006, **45**, 2426;
(b) S. H. Yu, M. J. Ferguson, R. McDonald and D. G. Hall, *J. Am. Chem. Soc.*, 2005, **127**, 12808; (c) N. Selander, S. Sebelius, C. Estay and K. J. Szabó, *Eur. J. Org. Chem.*, 2006, 4085; (d) N. Selander, A. Kipke, S. Sebelius and K. J. Szabó, *J. Am. Chem. Soc.*, 2007, **129**, 13723.

