ChemComm



COMMUNICATION

View Article Online



A synthetic approach to chrysophaentin F†

Cite this: Chem. Commun., 2019, 55 4837

Received 27th February 2019, Accepted 1st April 2019

DOI: 10.1039/c9cc01666i

rsc.li/chemcomm

Jean-Baptiste Vendeville, ab Rebecca F. Matters, Angi Chen, Mark E. Light, a Graham J. Tizzard, Christina L. L. Chai ** and David C. Harrowven ** a

The chrysophaentins are a newly discovered natural product family displaying promising anti-infective activity. Herein we describe an approach to chrysophaentin F that uses an array of metal catalysed coupling reactions (Cu, Ni, Pd, W, Mo) to form key bonds.

The chrysophaentins were discovered by Bewley et al. in the methanoic extract of Chrysophaeum taylori alga during a screening study to identify promising new anti-infectives. Eight closely related bioactive macrocycles were identified in the extract, named chrysophaentins A-H (1-5, 8-10), which were subsequently joined by the linear chrysophaentins E2 (6) and E3 (7) (Fig. 1).2 From a structural perspective they define a new class of marine natural products, the bis-diarylbutenes. Their core resembles that of the bis-bibenzyl family of natural products,^{3,4} but with two additional carbon centres and unsaturation in each of the chains linking the diaryl ether units.

In assays against Staphylococcus aureus (SA), methicillinresistant SA (MRSA) and multidrug-resistant SA (MDR-SA), chrysophaentins A (1), F (8) and H (10) were found to have useful potency, with minimum inhibitory concentrations (MIC₅₀) in the range of 0.8-9.5 μ g mL⁻¹. By contrast, the acyclic chrysophaentin E 5, and those with a bromide on arene A or arene C (2-4 and 9), showed greatly reduced activity. Chrysophaentin A (1) was also found to inhibit the bacterial cell division protein FtsZ, a popular target in antimicrobial drug discovery programmes.5

From a synthetic perspective the chrysophaentins have proven to be elusive targets. To date, all of the approaches




Fig. 1 The chrysophaentins and our approach to their total synthesis.

described have sought to make one of the symmetrical chrysophaentins using a cyclodimerization strategy, yet none has succeeded in gaining access to the macrocyclic core. 2,4-6 Herein we describe our work on the development of a general synthetic approach to the natural product family, exemplified by syntheses of the dehalogentated core 11 and, tentatively, of chrysophaentin F (8) in impure form. Our approach, summarized in Fig. 1, has both divergent and convergent phases and uses an array of metal catalyzed reactions (Cu, Ni, Pd, W and Mo) to construct key bonds.

To test the validity of our approach, we first targeted the chrysophaentin F-H core 11 lacking all halogen substituents. Our synthesis began with the preparation of benzo-1,3-dioxane 15 and phenol 16 using standard protocols (Scheme 1).⁷⁻⁹ Their union to diaryl ether 18 was then achieved using a Chan-Lam-Evans coupling procedure. 10 After examining a range

^a Chemistry, University of Southampton, Highfield, Southampton, SO17 1BJ, UK. E-mail: dch2@soton ac uk

^b Institute of Chemical and Engineering Sciences, Agency for Science, Technology and Research (A*STAR), 8 Biomedical Grove, Neuros, #07-01, 138665, Singapore ^c Department of Pharmacy, National University of Singapore, 18 Science Drive 4, 117543, Singapore. E-mail: phacllc@nus.edu.sg

[†] Electronic supplementary information (ESI) available: Experimental accounts with spectral details and copies of NMR spectra. CCDC 1899398 and 1898679. For ESI and crystallographic data in CIF or other electronic format see DOI: 10.1039/ c9cc01666j

Communication ChemComm

Scheme 1 Synthesis of the pivotal diaryl ether 20

of catalysts, solvents and reaction conditions, we found that it was best to employ Cu(OTf)₂ with 4 Å molecular sieves in ethanol under an oxygen atmosphere. In that way diaryl ether **18** could be formed reliably in 76% yield. Sequential hydrolysis of the acetal to phenol **17**; protection as benzyl ether **19** and conversion of the benzylic alcohol to the corresponding chloride, ¹¹ gave the pivotal diaryl ether **20** in high overall yield.

At this juncture our synthesis entered a divergent phase where diaryl ether 20 was advanced to both the B-O-C and A-O-D subunits, 26 and 28 respectively (Scheme 2). Thus, using a Pd-catalysed procedure developed by Buchwald et al. 12 diaryl ether 20 was coupled with TMS-acetylene and 1-hexyne respectively, to afford alkynes 21 and 22 in excellent yield. The esters in each of these products were then reduced with LiAlH4 to facilitate their conversion to halides 25 and 26.11 Alas, attempts to effect the coupling of benzyl bromide 25 and 1-hexyne using the aforementioned Buchwald procedure also induced alkyne to allene isomerisation.¹² However, by switching to a nickel catalyzed coupling reaction developed by Gau et al., 13 it was successfully coupled with hexynylAlEt2 to give diyne 28 in good yield after deprotection of the silyl acetylene 27 with catalytic silver triflate.14 Pleasingly, the Buchwald procedure proved effective for the coupling of diaryl ether fragments 26 and 28 providing the macrocyclic precursor **29** in 50% yield. ¹²

The stage was now set to enact our endgame strategy, which sought to use alkyne metathesis for the critical macrocyclization reaction. ^{15,16} Pleasingly, this proved remarkably efficient as, after some optimization, triyne **29** was transformed into macrocyclic diyne **30** in near quantitative yield using Schrock's alkylidyne catalyst in toluene at 80 °C for 12 h under high dilution. ¹⁶ Success was confirmed by X-ray crystallographic analysis (Fig. 2). Selective hydrogenation of **30** using Lindlar's catalyst in the presence of quinoline next provided bis-*cis*-alkene **31**, leaving us the task of unmasking the six phenol

Scheme 2 Synthesis of dehalogenated chrysophaentin 11.




Fig. 2 X-ray crystal structure of macrocyclic diyne 30.

residues. Although the double bonds in **31** proved sensitive to an array of standard deprotection protocols, a combination of BCl₃ and tetrabutylammonium iodide in DCM gave our target **11** in 92% yield (Scheme 2).¹⁷

The total synthesis of chrysophaentin F 8 became our next target with the preparation of the keystone diaryl ether 38 becoming the immediate goal (Scheme 3). To that end, benzoic acid 32 was reduced with LiAlH₄ to diol 33, which in turn was

ChemComm Communication

Scheme 3 A tentative synthesis of chrysophaentin F.

protected as its benzyl ether 34. Arene bromination to 35 then facilitated a Miyaura borylation to 37 enabling its coupling to phenol 16 using a Chan-Lam-Evans procedure. Finally, conversion of the resulting benzyl alcohol 36 to the corresponding chloride 38 delivered the required diaryl ether.

Chrysophaentin F. 8

The divergent strategy used to prepare dehalochrysophaentin 11 was now applied to the synthesis of chrysophaentin F 8. While most of the steps were easy to replicate, it proved advantageous to use a Stille coupling to advance chloride 38 to alkyne 39 as it proceeded reliably in near quantitative yield. 18 All other steps in the sequence mirrored those demonstrated previously, readily providing the B-O-C and A-O-D diaryl ether subunits 44 and 46 (Scheme 3). These were coupled using the Buchwald procedure to give triyne 47, setting the stage for our end-game strategy.

Pleasingly, after some optimization, trivne 47 was transformed into macrocyclic diyne 48 in modest yield using Fürstner's molybdenum-phenanthroline pre-catalyst in toluene at 40 °C for 12 h. 15 Hydrozirconation of 48 with Schwartz reagent (generated in situ by reduction of Cp2ZrCl2) followed by chlorination with NCS gave a complex mixture of products, 19,20 that was partially separated by column chromatography (see ESI† p. S124 and S125). The main fractions were then combined and treated with BCl3 and tetrabutylammonium iodide to unmask the phenolic residues. Purification of the product mixture by column chromatography gave a major fraction exhibiting spectral characteristics consistent with the formation of chrysophaentin F 8 in an impure state (see ESI† for LRMS, ¹H and ¹³C NMR data). Alas, attempts to purify the natural product further by preparative TLC and HPLC proved unrewarding with material losses putting paid to further endeavours.

In summary, we have developed a synthetic approach to chrysophaentin F 8 with the flexibility to allow all members of this family to be targeted. The approach demonstrates the value of (i) Chan-Lam-Evans coupling reactions for the preparation of diaryl ethers with high steric demand; (ii) Buchwald's, Gau's and Stille's procedures for effecting sp-sp3 coupling reactions between alkynes and benzyl halides;12,13 and (iii) Fürtsner's alkylidyne pre-catalyst for macrocyclisation through ring-closing alkyne metathesis. 15,16

We thank Prof. A. Fürstner for his kind donation of the molybdenumphenanthroline pre-catalyst. The University of Southampton VC Scholarship Scheme, the A*STAR ARAP Scheme, the ERDF (LabFact: InterReg V project 121) and the EPSRC [EP/K039466/1, EP/P013341/1 and EP/L003325/1] are thanked for their financial support.

Conflicts of interest

There are no conflicts to declare.

Notes and references

- 1 A. Plaza, J. Keffer, G. Bifulco, J. Lloyd and C. Bewley, J. Am. Chem. Soc., 2010, 132, 9069; J. R. Davison and C. Bewley, J. Nat. Prod., 2019, 82, 148
- 2 J. L. Keffer, J. T. Hammill, J. R. Lloyd, A. Plaza, P. Wipf and C. A. Bewley, Mar. Drugs, 2012, 10, 1103.
- D. C. Harrowven and S. L. Kostiuk, Nat. Prod. Rep., 2012, 29, 223. For our work on macrocyclic bis-bibenzyl total synthesis see: F. A. Almalki and D. C. Harrowven, Eur. J. Org. Chem., 2016, 5738; S. L. Kostiuk, T. Woodcock, L. F. Dudin, P. D. Howes and D. C. Harrowven, Chem. -Eur. J., 2011, 17, 10906; D. C. Harrowven, T. Woodcock and P. D. Howes, Angew. Chem., Int. Ed., 2005, 44, 3899.
- 4 H. P. Erikson, Cell, 1995, 80, 367; M. H. Foss, Y.-J. Eun and D. B. Weibel, Biochemistry, 2011, 50, 7719.
- 5 A. Brockway, C. Grove, M. Mahoney and J. Shaw, Tetrahedron Lett., 2015, 56, 3396.
- 6 In common with others, our first approach to chrysophaentin F was based on a cyclodimerization strategy, e.g. $49 \rightarrow 50$. Indeed, after

much optimization this gave a cyclodimer in 6% yield that could not be assigned unambiguously to either dimer **50** or its regioisomer **51** using the spectral data attained.

- 7 N. Gisch, J. Ballzarini and C. Meier, J. Med. Chem., 2007, 50, 1658.
- 8 T. Ishiyama, M. Murata and N. Miyaura, J. Org. Chem., 1995, 60, 7508.
- 9 K. Bao, A. Fan, Y. Dai, L. Zhang, W. Zhang, M. Cheng and X. Yao, Org. Biomol. Chem., 2009, 7, 5084.
- D. Chan, K. Monaco, R. Wang and M. Winters, Tetrahedron Lett., 1998, 39, 2933; D. Evans, J. Katz and T. West, Tetrahedron Lett., 1998, 39, 2937; P. Lam, C. Clark, S. Saubern, J. Adams, M. Winters, D. Chan and A. Combs, Tetrahedron Lett., 1998, 39, 2941. See also S. V. Ley and A. W. Thomas, Angew. Chem., Int. Ed., 2003, 42, 5400.
- 11 R. Appel, Angew. Chem., Int. Ed. Engl., 1975, 14, 801.

- 12 C. H. Larsen, K. W. Anderson, R. E. Tundel and S. L. Buchwald, *Synlett*, 2006, 2941. The optimised procedure reported used the alkyne in slight excess (1.3 equiv.) and was adopted for advancing benzyl chloride 20 to alkynes 21 and 22. However, it was not deemed appropriate to use alkynes 28 and 46 in excess in their respective couplings with benzyl chlorides 26 and 44, as each was prepared by multi-step sequences. When using equimolar quantities of each component, some benzyl chloride was usually recovered (~25%) but no alkyne. The known isomerisation of the alkyne product to an allene was also observed as a side reaction in many cases.
- 13 D. Biradar and H. Gau, Chem. Commun., 2011, 47, 10467.
- 14 A. Orsini, A. Vitérisi, A. Bodlenner, J.-M. Weibel and P. Pale, *Tetrahedron Lett.*, 2005, 46, 2259.
- J. Heppekausen, R. Stade, R. Goddard and A. Fürstner, *J. Am. Chem. Soc.*, 2010, 132, 11045; A. Fürstner, *Chem. Commun.*, 2011, 47, 6505;
 A. Fürstner and P. W. Davies, *Chem. Commun.*, 2005, 2307.
- 16 M. L. Listemann and R. R. Schrock, Organometallics, 1985, 4, 74.
- 17 P. R. Brooks, M. C. Wirtz, M. G. Vetelino, D. M. Rescek, G. F. Woodworth, B. P. Morgan and J. W. Coe, *J. Org. Chem.*, 1999, 64, 9719.
- 18 R. R. Singidi and T. V. RajanBabu, Org. Lett., 2008, 10, 3351; D. C. Harrowven, D. P. Curran, S. L. Kostiuk, I. L. Wallis-Guy, S. Whiting, K. J. Stenning, B. Tang, E. Packard and L. Nanson, Chem. Commun., 2010, 46, 6335; D. C. Harrowven and I. L. Guy, Chem. Commun., 2004, 1968.
- 19 Z. Huang and E. Negishi, Org. Lett., 2006, 8, 3675.
- D. W. Hart and J. Schwartz, J. Am. Chem. Soc., 1974, 96, 8115;
 D. W. Hart, T. F. Blackburn and J. Schwartz, J. Am. Chem. Soc., 1975, 97, 679;
 J. Schwartz and J. A. Labinger, Angew. Chem., Int. Ed. Engl., 1976, 15, 333.