



Cite this: *Chem. Soc. Rev.*, 2025, **54**, 3819

## Bis(trimethylsilyl)phosphide chemistry: a half-century of advances across the periodic table

Jack Baldwin and David P. Mills \*

Whilst bis(trimethylsilyl)amide has been used extensively as a ligand across the periodic table, the chemistry of its heavier group 15 congeners is relatively underdeveloped. However, bis(trimethylsilyl)phosphide coordination chemistry has provided unique structural motifs and has also shown potential applications in catalysis, materials science, and bioinorganic chemistry. This review, which marks 55 years since the first report of a bis(trimethylsilyl)phosphide complex, provides a comprehensive overview of the synthesis, characterisation and reactivity of structurally authenticated s-, p-, d- and f-block metal complexes of this ligand, focusing on salient single crystal XRD and NMR spectroscopic data. We discuss the factors influencing the diverse coordination modes and reactivity profiles of bis(trimethylsilyl)phosphide complexes, together with an overview of their potential as precursors for novel solid-state materials, aiming to inspire future research endeavours using this ligand. We also review the small number of bis(triisopropylsilyl)phosphide complexes, in order to provide motivation for the future study of other bis(silyl)phosphide ligands.

Received 13th November 2024

DOI: 10.1039/d4cs01141d

[rsc.li/chem-soc-rev](https://rsc.li/chem-soc-rev)

### 1. Introduction

Bis(trimethylsilyl)amide,  $\{N(\text{SiMe}_3)_2\}$  ( $N''$ ), and derivatives thereof, have been used extensively across the periodic table to provide landmark complexes<sup>1–7</sup> since the first s-block examples were reported by Wannagat in 1961.<sup>8</sup> The popularity of  $N''$  in coordination chemistry can be attributed to a combination of: (i) kinetic protection of the metal coordination

sphere and inter-ligand dispersion force stabilisation provided by sterically demanding  $\text{SiMe}_3$  groups; (ii) the absence of potential  $\beta$ -hydride elimination decomposition pathways; (iii) the high lipophilicity of  $\text{SiMe}_3$  groups increasing complex solubility in non-polar solvents and facilitating crystal growth for structure determination; (iv) negative hyperconjugation by the silyl groups making the ligand charge more diffuse, resulting in softer N-donor atoms and promoting additional stabilising  $M \cdots \text{Si-C}$  and  $M \cdots \text{C-H}$  interactions; (v) NMR spectra that are typically easy to interpret; and, (vi) the ease of preparation and commercial availability of s-block ligand transfer agents.<sup>1–7</sup>

Department of Chemistry, The University of Manchester, Oxford Road, Manchester, M13 9PL, UK. E-mail: david.mills@manchester.ac.uk



Jack Baldwin

Jack Baldwin is from Beverley, East Yorkshire, in the UK. He completed his MChem (2019) at Newcastle University and his PhD (2024) at The University of Manchester, where he also carried out postdoctoral work. His research is centred around the synthesis and characterisation of low coordinate f-block metal complexes with silylphosphide ligands.



David P. Mills

David P. Mills is from Caerphilly, South Wales, in the UK. He did his MChem (2004) and PhD (2008) degrees at Cardiff University, and postdoctoral work at the University of Nottingham. He moved to the University of Manchester in 2012 as a lecturer, and was promoted to Professor of Inorganic Chemistry in 2021. His research group focuses on the synthesis of metal complexes with unusual bonding motifs and electronic structures, predominantly those containing f-block metals.





Fig. 1 Graph depicting the number of structurally authenticated N'' vs. P'' metal complexes containing elements from each block of the periodic table.<sup>9</sup>

The coordination chemistry of bis(trimethylsilyl)phosphide,  $\{P(\text{SiMe}_3)_2\}$  (P''), is immature in comparison to that of the lighter congener N'', despite it having many of the same advantages. This is evident from a survey of the Cambridge Structural Database (CSD), which shows 210 entries for P'' vs. 4062 entries for N'' structurally authenticated metal complexes (non-metals and the metalloids B, Si, As, Sb and Te were excluded from this search and are outside the scope of this review); the 210 P'' metal complexes consist of 94 p-, 64 d-, 30 s- and 22 f-block examples (Fig. 1).<sup>9</sup> This can be ascribed to difficulties in handling s-block P'' salts, which are pyrophoric, malodorous and toxic, and the synthesis of these ligand transfer agents typically using  $P(\text{SiMe}_3)_3$  as a starting material (Scheme 1).<sup>10–13</sup> As well as having similar handling concerns to P'',  $P(\text{SiMe}_3)_3$  is relatively expensive, and thus is typically synthesised from red phosphorus, sodium and chlorotrimethylsilane on large scales in dimethoxyethane (DME) (Scheme 1); this synthetic procedure can discourage investigations as it is inherently hazardous.<sup>14</sup> However, an advantage of P'' over N'' is that the 100% abundant  $I = \frac{1}{2}$  metal-bound  $^{31}\text{P}$  nuclei can provide a useful spectroscopic handle, e.g. by using NMR or EPR spectroscopy. P'' complexes could also potentially deliver unique applications as P'' is softer than N'' and has a higher propensity to bridge metal ions as M–P bonds are longer than M–N bonds.<sup>15–17</sup> Since the first reported syntheses of s-block P'' complexes<sup>18</sup> they have been used in a plethora of reactions, ranging from nucleophilic additions to radical-mediated processes;<sup>19</sup> they have also been shown to be versatile reducing agents,<sup>20</sup> and a precursor to phosphorus-centred radicals.<sup>19</sup>


Here we provide a review of P'' coordination chemistry, focusing on structurally authenticated examples of complexes

containing direct M–P bonds. When appropriate we make comparisons to complexes of the lighter congener, N''; previous reviews of metal phosphido chemistry have tended to cover a wider range of ligand substituents and a narrower range of elements.<sup>21–23</sup> The review is divided into sections of s-, p-, d-, and f-block metal P'' complexes; these sections are subdivided by group number, with heterometallic 'ate' complexes containing s-block elements included in the p-, d- or f-block sections as appropriate. Where M–P bond lengths from single crystal XRD studies and chemical shifts and coupling constants from  $^{31}\text{P}$  and  $^{29}\text{Si}$  (4.7% abundant,  $I = \frac{1}{2}$ ) NMR spectroscopic data have been reported they are compiled in Table 1 (calculation of the standard error of the mean (SEM) =  $\text{SD}/\sqrt{N}$ , where SD = standard deviation and  $N$  = sample size). We focus on discussions of these data in this review, which we intend to provide a useful resource to the community and to inspire future research in P'' coordination chemistry. We also provide coverage of nascent bis(triisopropylsilyl)phosphide chemistry, in order to show that other bis(silyl)phosphide ligands should provide new avenues to explore in future.

## 2.1 s-Block P'' complexes

Alkali metal and alkaline earth metal P'' complexes are used extensively as ligand transfer agents, and therefore are important starting materials to p-, d- and f-block P'' complexes. Surprisingly, to date there are no structurally characterised examples of Na P'' complexes on the CSD,<sup>9</sup> though various synthetic routes to group 1 P'' salts including Na derivatives have been reported extensively.<sup>11,12</sup> A Be P'' complex has yet to be structurally authenticated to the best of our knowledge,<sup>9</sup> thus there are obvious gaps in s-block P'' chemistry that can be relatively easily addressed.

**2.1.1 Group 1 complexes.** To the best of our knowledge, the first report of a Li P'' complex was by Bürger in 1970, but this was not structurally authenticated.<sup>18</sup> The first isolated Li P'' complex,  $[\text{Li}(\text{P}'')(\text{DME})]$ , was reported in 1974 by Höldrich,<sup>11,118</sup> but the first structurally characterised Li P'' complexes were not published until 1987 by Rai.<sup>24</sup> In Rai's report the reaction of *in situ*-formed  $P(\text{SiMe}_3)_3$  with  $^n\text{BuLi}$  in THF was shown to give the dimeric complex  $[\text{Li}(\mu\text{-P}'')(\text{THF})_2]_2$  (1, Fig. 2); this undergoes oligomerisation when placed under vacuum at 20 °C for 20 hours to yield the ladder-like tetranuclear complex  $[\text{Li}_4(\mu_2\text{-P}'')_2(\mu_3\text{-P}'')_2(\text{THF})_2]$  (3, Fig. 2).<sup>24</sup> Similarly, the DME-solvated dimeric Li P'' complex  $[\text{Li}(\mu\text{-P}'')(\text{DME})]_2$  (2, Fig. 2) was later synthesised in 1989 by Becker from the reaction of  $P(\text{SiMe}_3)_3$  with MeLi in DME.<sup>25</sup> The solvent-free hexanuclear ladder-like complex,  $[\text{Li}_6(\mu_2\text{-P}'')_2(\mu_3\text{-P}'')_4]$  (4, Fig. 2), was prepared in 1992 by Hey-Hawkins following treatment of  $\text{HP}''$  with  $^n\text{BuLi}$  in cyclopentane,



Scheme 1 Synthesis of  $\text{KP}''$  from red phosphorus, sodium and trimethylsilylchloride.<sup>10–13</sup>



**Table 1** Mean M–P bond lengths (Å), and chemical shifts ( $\delta$ ) and coupling constants (Hz) for  $^{31}\text{P}$  and  $^{29}\text{Si}$  NMR spectroscopic data, of structurally authenticated  $\text{P}''$  complexes<sup>9</sup>

Complex	Mean M–P (Å)	Solvent	$^{31}\text{P}$ Chemical shift ( $\delta$ )	$^{31}\text{P}$ Coupling constant (Hz)	$^{29}\text{Si}$ Chemical shift ( $\delta$ )	$^{29}\text{Si}$ Coupling constant (Hz)	Ref.
s-Block	2.62(2)	$d_8$ -Toluene	–297.6, s	—	—	—	24
1 [Li( $\mu$ - $\text{P}''$ )(THF) <sub>2</sub> ] <sub>2</sub>	2.559(4)	—	—	—	—	—	25
2 [Li( $\mu$ - $\text{P}''$ )(DME) <sub>2</sub> ] <sub>2</sub>	2.50(4) <sup>a</sup>	$d_8$ -Toluene	–297.7, s	—	—	—	24
3 [Li <sub>4</sub> ( $\mu_2$ - $\text{P}''$ ) <sub>2</sub> ( $\mu_3$ - $\text{P}''$ ) <sub>2</sub> (THF) <sub>2</sub> ]	2.56(4) <sup>b</sup>	—	—	—	—	—	26
4 [Li <sub>6</sub> ( $\mu_2$ - $\text{P}''$ ) <sub>2</sub> ( $\mu_3$ - $\text{P}''$ ) <sub>4</sub> ]	2.51(1) <sup>a</sup>	—	—	—	—	—	26
	2.517(16) <sup>b</sup>	—	—	—	—	—	26
5-K [K <sub>4</sub> ( $\mu_2$ - $\text{P}''$ ) <sub>2</sub> ( $\mu_3$ - $\text{P}''$ ) <sub>2</sub> ] <sub>∞</sub>	3.4168(11) <sup>a</sup>	THF	–293.4, s	—	0.70, s	$^1J_{\text{P}''\text{Si}} = 54$	27
	3.171(6) <sup>b</sup>	—	—	—	—	—	27
5-Rb [Rb <sub>4</sub> ( $\mu_2$ - $\text{P}''$ ) <sub>2</sub> ( $\mu_3$ - $\text{P}''$ ) <sub>2</sub> ] <sub>∞</sub>	3.485(2) <sup>a</sup>	THF	–287.1, s	—	0.41, s	$^1J_{\text{P}''\text{Si}} = 55$	27
	3.4157(12) <sup>b</sup>	—	—	—	—	—	27
5-Cs [Cs <sub>4</sub> ( $\mu_2$ - $\text{P}''$ ) <sub>2</sub> ( $\mu_3$ - $\text{P}''$ ) <sub>2</sub> ] <sub>∞</sub>	3.656(2) <sup>a</sup>	THF	–270.0, s	—	0.50, s	$^1J_{\text{P}''\text{Si}} = 52$	27
	3.5816(12) <sup>b</sup>	—	—	—	—	—	27
6 [Cs( $\mu$ - $\text{P}''$ )( $\mu$ -1,4-dioxane) <sub>3</sub> (1,4-dioxane)] <sub>∞</sub>	3.6137(17)	$d_8$ -THF	–276.1, s	—	—	—	28
7 [Mg( $\mu$ - $\text{P}''$ )(Br)(THF) <sub>2</sub> ]	2.5624(16)	$d_6$ -Benzene	–296.9, br	—	2.9, br	—	20
8 [Mg <sub>3</sub> ( $\mu_2$ - $\text{P}''$ ) <sub>4</sub> ( $\text{P}''$ ) <sub>2</sub> ]	2.594(7) <sup>c</sup>	$d_8$ -Toluene	–242.55, t <sup>c</sup>	$^2J_{\text{PP}} = 17.9$	3.74, s	—	29
	2.455(4) <sup>d</sup>	—	–275.44, t <sup>d</sup>	—	—	—	29
9 [Mg( $\text{P}''\text{C}(\text{Ph})\text{C}(\text{C}_2\text{Ph})$ )( $\mu$ - $\text{P}''$ ) <sub>2</sub> ]	2.5576(19) <sup>c</sup>	$d_8$ -Toluene	–262.29, t <sup>c</sup>	$^2J_{\text{PP}} = 5$	4.56, s <sup>c</sup>	$^1J_{\text{P}''\text{Si}} = <2^c$	30
	2.6998(17)	—	–98.4, t	—	3.63, d	$^1J_{\text{P}''\text{Si}} = 6.3$	30
10 [Mg( $\text{P}''$ ) <sub>2</sub> (DME)]	2.487(2)	—	—	—	—	—	31
11 [Mg( $\text{P}''$ ) <sub>2</sub> (THF) <sub>2</sub> ]	2.5023(19)	$d_8$ -Toluene	–294.7, s	—	1.81, s	$^1J_{\text{P}''\text{Si}} = 33.0$	32
12-Ca [Ca( $\text{N}''$ )( $\mu$ - $\text{P}''$ )(sol)] <sub>2</sub>	2.9269(18)	$d_6$ -Benzene	–229.96, s <sup>e</sup>	—	5.42, d <sup>e</sup>	$^1J_{\text{P}''\text{Si}} = 18.7^e$	33
	—	—	–254.82, s <sup>f</sup>	—	–1.77, d <sup>f</sup>	$^1J_{\text{P}''\text{Si}} = 22.0^f$	33
12-Ba [Ba( $\text{N}''$ )( $\mu$ - $\text{P}''$ )(sol)] <sub>2</sub>	3.289(2)	$d_8$ -THF	–222.05, s	—	1.03, s	$^1J_{\text{P}''\text{Si}} = 34.8$	34
13-Ca <i>trans</i> -[Ca( $\text{P}''$ ) <sub>2</sub> (THF) <sub>4</sub> ]	2.917(4)	—	—	—	—	—	33
13-Sr <i>trans</i> -[Sr( $\text{P}''$ ) <sub>2</sub> (THF) <sub>4</sub> ]	3.021(10)	$d_8$ -Toluene	–256.2, s	—	1.11, s	$^1J_{\text{P}''\text{Si}} = 34.1$	35
13-Ba <i>trans</i> -[Ba( $\text{P}''$ ) <sub>2</sub> (THF) <sub>4</sub> ]	3.175(10)	$d_8$ -THF	–251.0, s	$^2J_{\text{PP}} = 23.5$	1.08, s	$^1J_{\text{P}''\text{Si}} = 44.5$	36
	—	—	—	—	—	$^3J_{\text{P}''\text{Si}} = 1.0$	36
14 [Sr( $\text{P}''$ )( $\mu$ - $\text{P}''$ ) <sub>3</sub> Sr(THF) <sub>3</sub> ]	3.118(8) <sup>c</sup>	$d_8$ -Toluene	–276.78, s <sup>d</sup>	$^2J_{\text{PP}} = 22.5^d$	1.60, s	$^1J_{\text{P}''\text{Si}} = 38.4^d$	35
	3.042(4) <sup>d</sup>	—	—	—	—	—	35
15 [Ca( $\text{P}''$ ) <sub>2</sub> (TMTA) <sub>2</sub> ]	2.995(2)	$d_8$ -THF	–276.8, s <sup>d</sup>	$^2J_{\text{PP}} = 14.5$	0.87, s	$^1J_{\text{P}''\text{Si}} = 34.4$	36
	—	—	—	—	—	$^3J_{\text{P}''\text{Si}} = -2.7$	36
p-Block							
16 [Al( $\text{P}''$ )(tmp) <sub>2</sub> ]	2.361(2)	$d_6$ -Benzene	–238.00, s	—	—	—	37
17-ALMe [Al(Me) <sub>2</sub> ( $\text{P}''$ )(dmap)]	2.3768(9)	$d_6$ -Benzene	—	—	—	—	38
17-GaMe [Ga(Me) <sub>2</sub> ( $\text{P}''$ )(dmap)]	2.372(1)	$d_8$ -Toluene	–273.80, s	—	—	—	39
17-Ga <sup>t</sup> Bu [Ga( <sup>t</sup> Bu) <sub>2</sub> ( $\text{P}''$ )(dmap)]	2.3948(6)	$d_6$ -Benzene	–281.00, s	—	—	—	40
18 [Ga( $\text{P}''$ )( <sup>t</sup> Bu-DAB)]	2.2991(11)	—	—	—	—	—	41
19 [Ga( $\text{P}''$ ) <sub>2</sub> ( <sup>t</sup> Bu-DAB)]	2.3426(18)	—	—	—	—	—	41
20-Ga [Ga( <sup>Dipp</sup> Pnacnac)( $\text{P}''$ )(Cl)]	2.3310(9)	$d_6$ -Benzene	–255.00, s	—	—	—	42
20-In [In( <sup>Dipp</sup> Pnacnac)( $\text{P}''$ )(Cl)]	2.4806(8)	$d_6$ -Benzene	–252.90, s	—	—	—	42
21-ALMe [Al(Me) <sub>2</sub> ( $\mu$ - $\text{P}''$ ) <sub>2</sub> ]	2.457(2)	$d_8$ -Toluene	–226.7, s	—	6.24, Vir. t	$^1J_{\text{P}''\text{Si}} = 9.9$	43 and 44
21-AIEt [Al(Et) <sub>2</sub> ( $\mu$ - $\text{P}''$ ) <sub>2</sub> ]	2.458(1)	$d_6$ -Benzene	–246.90, s	—	—	—	45
21-ALCH <sub>2</sub> <sup>1</sup> Pr [Al(CH <sub>2</sub> <sup>1</sup> Pr) <sub>2</sub> ( $\mu$ - $\text{P}''$ ) <sub>2</sub> ]	2.476(2)	$d_6$ -Benzene	–245.30, s	—	—	—	43
21-ALCH <sub>2</sub> SiMe <sub>3</sub> [Al(CH <sub>2</sub> SiMe <sub>3</sub> ) <sub>2</sub> ( $\mu$ - $\text{P}''$ ) <sub>2</sub> ]	2.483(1)	$d_8$ -Toluene	–231.49, s	—	6.38, s	$^1J_{\text{P}''\text{Si}} = 6.2$	46
21-GaMe [Ga(Me) <sub>2</sub> ( $\mu$ - $\text{P}''$ ) <sub>2</sub> ]	2.450(1)	$d_6$ -Benzene	–219.20, s	—	6.41, s	—	40 and 47
21-GaEt [Ga(Et) <sub>2</sub> ( $\mu$ - $\text{P}''$ ) <sub>2</sub> ]	2.4558(7)	$d_8$ -Toluene	–227.80, s	—	—	—	48
21-Ga <sup>t</sup> Bu [Ga( <sup>t</sup> Bu) <sub>2</sub> ( $\mu$ - $\text{P}''$ ) <sub>2</sub> ]	2.4533(6)	$d_6$ -Benzene	–227.00, s	—	—	—	49
21-GaCH <sub>2</sub> <sup>1</sup> Bu [Ga(CH <sub>2</sub> <sup>1</sup> Bu) <sub>2</sub> ( $\mu$ - $\text{P}''$ ) <sub>2</sub> ]	2.517(3)	$d_6$ -Benzene	–215.24, s	—	—	—	50
21-GaCH <sub>2</sub> SiMe <sub>3</sub> [Ga(CH <sub>2</sub> SiMe <sub>3</sub> ) <sub>2</sub> ( $\mu$ - $\text{P}''$ ) <sub>2</sub> ]	2.4887(16)	$d_6$ -Benzene	–205.88, s	—	—	—	51
21-InMe [In(Me) <sub>2</sub> ( $\mu$ - $\text{P}''$ ) <sub>2</sub> ]	2.630(1)	$d_8$ -Toluene	–239.80, s	—	—	—	52
21-InEt [In(Et) <sub>2</sub> ( $\mu$ - $\text{P}''$ ) <sub>2</sub> ]	2.645(1)	$d_6$ -Benzene	–242.90, s	—	—	—	53
21-InPh [In(Ph) <sub>2</sub> ( $\mu$ - $\text{P}''$ ) <sub>2</sub> ]	2.612(1)	$d_6$ -Benzene	–221.59, s	—	—	—	54
21-InCH <sub>2</sub> Ph [In(CH <sub>2</sub> Ph) <sub>2</sub> ( $\mu$ - $\text{P}''$ ) <sub>2</sub> ]	2.6123(6)	$d_2$ -DCM	–220.30, s	—	—	—	55
21-InCH <sub>2</sub> SiMe <sub>3</sub> [In(CH <sub>2</sub> SiMe <sub>3</sub> ) <sub>2</sub> ( $\mu$ - $\text{P}''$ ) <sub>2</sub> ]	2.655(3)	$d_6$ -Benzene	–227.36, s	—	—	—	56
21-TlMe [Tl(Me) <sub>2</sub> ( $\mu$ - $\text{P}''$ ) <sub>2</sub> ]	2.692(3)	$d_6$ -Benzene	–234.00, s	$^1J_{\text{TlP}} = 2462$	—	—	57
22 [Al(Me) <sub>2</sub> (dmap)( $\mu$ - $\text{P}''$ )(Ga(Me) <sub>3</sub> )]	2.416(1)	$d_8$ -Toluene	–262.40, s	—	—	—	58
23-CH <sub>2</sub> SiMe <sub>3</sub> [In(Me)(CH <sub>2</sub> SiMe <sub>3</sub> )( $\mu$ - $\text{P}''$ ) <sub>2</sub> ]	2.635(2)	$d_6$ -Benzene	–234.32, s <sup>g</sup>	—	—	—	59
	—	—	–234.67, s <sup>h</sup>	—	—	—	59
23-CH <sub>2</sub> <sup>t</sup> Bu [In(Me)(CH <sub>2</sub> <sup>t</sup> Bu)( $\mu$ - $\text{P}''$ ) <sub>2</sub> ]	2.637(3)	$d_6$ -Benzene	–239.56, s <sup>g</sup>	—	—	—	60
	—	—	–239.42, s <sup>h</sup>	—	—	—	60
24 [{In(CH <sub>2</sub> <sup>t</sup> Bu) <sub>2</sub> ] <sub>2</sub> ( $\mu$ - $\text{P}''$ )}( $\mu$ -PH(SiMe <sub>3</sub> ))] ]	2.650(5)	$d_6$ -Benzene	–209.92, s	$^1J_{\text{PH}} = 473$	—	—	60
	—	—	–209.12, s	—	—	—	60



Table 1 (continued)

Complex	Mean M–P (Å)	Solvent	<sup>31</sup> P Chemical shift (δ)	<sup>31</sup> P Coupling constant (Hz)	<sup>29</sup> Si Chemical shift (δ)	<sup>29</sup> Si Coupling constant (Hz)	Ref.
25-AlBrCH <sub>2</sub> SiMe <sub>3</sub> [Al(CH <sub>2</sub> SiMe <sub>3</sub> )(Br)(μ-P'')] <sub>2</sub>	2.436(4)	d <sub>6</sub> -Benzene	-215.15, s	—	—	—	61
25-GaClMe [Ga(Me)(Cl)(μ-P'')] <sub>2</sub>	2.4106(10)	—	—	—	—	—	62
25-GaClCH <sub>2</sub> <sup>t</sup> Bu [Ga(Cl)(CH <sub>2</sub> <sup>t</sup> Bu)(μ-P'')] <sub>2</sub>	2.422(3)	d <sub>6</sub> -Benzene	-233.86, s	—	—	—	50
25-GaBrCH <sub>2</sub> SiMe <sub>3</sub> [Ga(CH <sub>2</sub> SiMe <sub>3</sub> )(Br)(μ-P'')] <sub>2</sub>	2.424(3)	d <sub>6</sub> -Benzene	-227.61, s	—	—	—	50
25-InClCH <sub>2</sub> SiMe <sub>3</sub> [In(CH <sub>2</sub> SiMe <sub>3</sub> )(Cl)(μ-P'')] <sub>2</sub>	2.593(2)	d <sub>6</sub> -Benzene	-241.43, s	—	—	—	59
25-InClCp* [In(Cp*)(Cl)(μ-P'')] <sub>2</sub>	2.621(2)	d <sub>6</sub> -Benzene	-148.60, s	—	—	—	63
26-GaPh [{Ga(Ph) <sub>2</sub> (μ-P'')}(μ-Cl)]	2.391(4)	d <sub>6</sub> -Benzene	-214.74, s	—	—	—	64
26-GaCH <sub>2</sub> <sup>t</sup> Bu [{Ga(CH <sub>2</sub> <sup>t</sup> Bu) <sub>2</sub> (μ-P'')}(μ-Cl)]	2.451(3)	d <sub>6</sub> -Benzene	-210.22, s	—	—	—	50
26-GaCH <sub>2</sub> SiMe <sub>3</sub> [{Ga(CH <sub>2</sub> SiMe <sub>3</sub> ) <sub>2</sub> (μ-P'')}(μ-Cl)]	2.416(6)	d <sub>6</sub> -Benzene	-213.27, s	—	—	—	50
26-InCH <sub>2</sub> <sup>t</sup> Bu [{In(CH <sub>2</sub> <sup>t</sup> Bu) <sub>2</sub> (μ-P'')}(μ-Cl)]	2.622(5)	d <sub>6</sub> -Benzene	-227.34, s	—	—	—	60
26-InCH <sub>2</sub> SiMe <sub>3</sub> [{In(CH <sub>2</sub> SiMe <sub>3</sub> ) <sub>2</sub> (μ-P'')}(μ-Cl)]	2.603(4)	d <sub>6</sub> -Benzene	-218.99, s	—	—	—	56
27-Al [{Al(Et) <sub>2</sub> (μ-P'')}(μ-As'')]	2.497(1)	d <sub>6</sub> -Benzene	-240.93, s -245.83, s	—	—	—	65
27-In [{In(CH <sub>2</sub> SiMe <sub>3</sub> ) <sub>2</sub> (μ-P'')}(μ-As'')]	2.691(3)	d <sub>8</sub> -Toluene	-229.84, s -230.72, s	—	—	—	66
29 [{Ga(Et) <sub>2</sub> (μ-P'')}(μ-Sb'')]	2.574(2)	d <sub>8</sub> -Toluene	—	—	—	—	48
30-ClP [{Ga(Cl) <sub>2</sub> (μ-P'')}(μ-P'')]	2.379(3)	d <sub>8</sub> -Toluene	-158.40, s	—	—	—	67
30-BrP [{Ga(Br) <sub>2</sub> (μ-P'')}(μ-P'')]	2.386(3)	d <sub>8</sub> -Toluene	-158.40, s	—	—	—	68
30-IP [{Ga(I) <sub>2</sub> (μ-P'')}(μ-P'')]	2.398(4)	d <sub>8</sub> -Toluene	-158.40, s	—	—	—	68 and 69
30-IAs [{Ga(I) <sub>2</sub> (μ-P'')}(μ-As'')]	2.443(3)	d <sub>6</sub> -Benzene	-261.25, s	—	—	—	69
31-Al [Al(H) <sub>2</sub> (μ-P'')] <sub>3</sub>	2.398(2)	d <sub>8</sub> -Toluene	-273.50, s	—	—	—	70
31-Ga [Ga(H) <sub>2</sub> (μ-P'')] <sub>3</sub>	2.392(3)	d <sub>8</sub> -Toluene	-265.80, s	—	—	—	71
32-Al [{Al(Me) <sub>2</sub> }(μ-P'')}{HP(SiMe <sub>3</sub> ) <sub>2</sub> }]	2.4287(13)	d <sub>8</sub> -Toluene	-253.00, d <sup>g</sup> -252.46, d <sup>h</sup>	[ <sup>1</sup> J <sub>PSi</sub> + <sup>2</sup> J <sub>PP</sub> = 85.0] <sup>g</sup> [ <sup>1</sup> J <sub>PSi</sub> + <sup>2</sup> J <sub>PP</sub> = 91.1] <sup>h</sup>	—	—	72
32-Ga [{Ga(Me) <sub>2</sub> }(μ-P'')}{HP(SiMe <sub>3</sub> ) <sub>2</sub> }]	2.415(2)	d <sub>8</sub> -Toluene	-241.74, t <sup>g</sup> -241.01, t <sup>h</sup>	<sup>2</sup> J <sub>PP</sub> = 89.2 <sup>g</sup> <sup>2</sup> J <sub>PP</sub> = 44.0 <sup>g</sup> <sup>2</sup> J <sub>PP</sub> = 84.7 <sup>h</sup> <sup>2</sup> J <sub>PP</sub> = 42.3 <sup>h</sup> <sup>2</sup> J <sub>PP</sub> = 30.5	5.53, dt <sup>h</sup> 6.0, dVir.t <sup>h</sup>	<sup>1</sup> J <sub>PSi</sub> = 8.2 <sup>h</sup> <sup>3</sup> J <sub>PSi</sub> = 6.9 <sup>h</sup> [ <sup>1</sup> J <sub>PSi</sub> + <sup>3</sup> J <sub>PSi</sub> = 7.2] <sup>h</sup>	72
33 [{Ga(Me) <sub>2</sub> (μ-P'')}(μ-P(Me))] <sub>2</sub>	2.443(9)	d <sub>3</sub> -Chloroform	-108.7, t -238.2, t	—	—	—	73
34-Al [Al(H) <sub>2</sub> (μ-P'') <sub>2</sub> Li(Et <sub>2</sub> O) <sub>2</sub> ]	2.4001(13)	d <sub>8</sub> -Toluene	-282.0, br, s	—	—	—	74
34-Ga [Ga(H) <sub>2</sub> (μ-P'') <sub>2</sub> Li(Et <sub>2</sub> O) <sub>2</sub> ]	2.4122(12)	d <sub>8</sub> -Toluene	-277.7, s	—	—	—	75
35 [Al(H) <sub>2</sub> (μ-P'') <sub>2</sub> Na(DME) <sub>2</sub> ]	2.405(1)	d <sub>6</sub> -Benzene	-283.0, s	—	3.7, t	<sup>1</sup> J <sub>PSi</sub> = 9.8	76
36 [Ge(P'')(Ar <sup>MCS</sup> ) <sub>2</sub> ]	2.329(1)	d <sub>6</sub> -Benzene	-18.2, s	—	—	—	77
37-Ge [Ge( <sup>Dipp</sup> Nacnac)(P'')]	2.3912(8)	d <sub>6</sub> -Benzene	-192.7, s	—	2.0, d	<sup>1</sup> J <sub>PSi</sub> = 17.1	78 and 79
37-Sn [Sn( <sup>Dipp</sup> Nacnac)(P'')]	2.5526(7)	d <sub>6</sub> -Benzene	-183.5, s	<sup>1</sup> J <sub>PSi</sub> = 17.1 <sup>1</sup> J <sub>SnP</sub> = 1453	4.0, d	<sup>1</sup> J <sub>PSi</sub> = 17	78
37-Pb [Pb( <sup>Dipp</sup> Nacnac)(P'')]	2.715(2)	d <sub>6</sub> -Benzene	-116.6, s	<sup>1</sup> J <sub>PbP</sub> = 2852	7.4, d	<sup>1</sup> J <sub>PSi</sub> = 36.0	78 and 80
38 [Ge(Se)( <sup>Dipp</sup> Nacnac)(P'')]	2.2976(7)	d <sub>6</sub> -Benzene	-172.6, s	<sup>1</sup> J <sub>PSi</sub> = 26.0 <sup>1</sup> J <sub>SeP</sub> = 52.0	—	—	81
39-Ge [Ge(Ph*)(P'')]	2.291(4)	d <sub>6</sub> -Benzene	-48.6, s	—	—	—	82
39-Sn [Sn(Ph*)(P'')]	2.527(1)	d <sub>6</sub> -Benzene	-123.1, s	<sup>1</sup> J <sub>117SnP</sub> = 1396 <sup>1</sup> J <sub>119SnP</sub> = 1453	4.08, d	<sup>1</sup> J <sub>PSi</sub> = 38.5	82
40 [Pb(P'')(μ-P'')] <sub>2</sub>	2.77(1) <sup>c</sup> 2.70(1) <sup>d</sup>	d <sub>6</sub> -Benzene	-218.0, s <sup>dg</sup> -217.3, s <sup>dh</sup> -281.4, s <sup>cg</sup> -302.4, s <sup>ch</sup>	<sup>1</sup> J <sub>PbP</sub> = 1264 <sup>dg</sup> <sup>1</sup> J <sub>PbP</sub> = 1183 <sup>dh</sup> <sup>1</sup> J <sub>PbP</sub> = 1658 <sup>cg</sup> <sup>1</sup> J <sub>PbP</sub> = 1598 <sup>ch</sup>	—	—	83
41 [Sn(P'')(μ-P'') <sub>2</sub> Ca(μ-P'') <sub>2</sub> Ca(N'')]	Sn–P: 2.695(4) <sup>c</sup> 2.795(9) <sup>d</sup>	d <sub>6</sub> -Benzene	-220.0, br, s (SnP) -232.97, s (CaP)	—	1.75, s (SnPSi <sub>2</sub> ) 5.70, s (CaPSi <sub>2</sub> )	—	84
42 [{Sn(P'')(μ <sub>2</sub> -P'') <sub>2</sub> Ba}]	Ca–P: 2.892(7) <sup>c</sup> Sn–P: 2.684(7) <sup>c</sup> 2.600(6) <sup>d</sup> Ba–P: 3.239(7) <sup>c</sup>	d <sub>8</sub> -Toluene	-235.35, s	<sup>1</sup> J <sub>SnP</sub> = 1068.13	4.5, s	—	34



Table 1 (continued)

Complex	Mean M–P (Å)	Solvent	<sup>31</sup> P Chemical shift (δ)	<sup>31</sup> P Coupling constant (Hz)	<sup>29</sup> Si Chemical shift (δ)	<sup>29</sup> Si Coupling constant (Hz)	Ref.
<b>43-Ca</b> [Ca(THF) <sub>2</sub> {Sn(μ <sub>2</sub> -P'')(μ <sub>3</sub> -P') <sub>2</sub> }]	Ca–P: 2.903(2) Sn–P: 2.642(1)	<i>d</i> <sub>6</sub> -Benzene	–223.58, s (SnP) –295.15, s (Sn <sub>2</sub> P)	<sup>1</sup> J <sub>SnP</sub> = 710.7 <sup>3</sup> J <sub>SnP</sub> = 74.1 (SnP) <sup>1</sup> J <sub>SnP</sub> = 578.6 <sup>2</sup> J <sub>PP</sub> = 77.3 (SnP)	—	—	33
<b>43-Ba</b> [Ba(THF) <sub>x</sub> {Sn(μ <sub>2</sub> -P'')(μ <sub>3</sub> -P') <sub>2</sub> }]	Ba–P: 3.302(12) Sn–P: 2.685(9)	<i>d</i> <sub>8</sub> -Toluene	–220.30, s	<sup>1</sup> J <sub>SnP</sub> = 730.0	—	—	34
d-Block							
<b>37-Cr</b> [Cr(DIPPnacnac)(P'')]	2.3641(3)	—	—	—	—	—	85
<b>37-Mn</b> [Mn(DIPPnacnac)(P'')(THF)]	2.461(1)	—	—	—	—	—	85
<b>37-Zn</b> [Zn(DIPPnacnac)(P'')]	2.2728(3)	<i>d</i> <sub>6</sub> -Benzene	–288.85, s	<sup>1</sup> J <sub>Psi</sub> = 28.5	3.49, d	<sup>1</sup> J <sub>Psi</sub> = 28.5	85
<b>44</b> [{Y(P'') <sub>2</sub> }(μ-P'') <sub>2</sub> ]	2.849(4) <sup>c</sup>	<i>d</i> <sub>6</sub> -Benzene	–107.80, dp <sup>c</sup>	<sup>1</sup> J <sub>YP</sub> = 56.7, <sup>2</sup> J <sub>PP</sub> = 5.0 <sup>c</sup>	—	—	86 and 87
	2.678(4) <sup>d</sup>	—	–104.80, dt <sup>d</sup>	<sup>1</sup> J <sub>YP</sub> = 122.4, <sup>2</sup> J <sub>PP</sub> = 5.0 <sup>d</sup>	—	—	
<b>45</b> [Sc{C(PPh <sub>2</sub> S) <sub>2</sub> }(P'')(py) <sub>2</sub> ]	2.618(14)	<i>d</i> <sub>8</sub> -Toluene	–176.40, br, s	—	—	—	88
<b>46-La</b> [La{P(SiMe <sub>3</sub> ) <sub>2</sub> }(THF) <sub>2</sub> ]	2.886(2)	<i>d</i> <sub>6</sub> -Benzene	–113.0, br	—	2.66, d	<sup>1</sup> J <sub>SiP</sub> = 22.4	89
<b>47</b> [Ti(Cp) <sub>2</sub> (P'')]	2.468(1)	—	251.0, br, s	—	—	—	90
<b>48-Hf</b> [Hf(Cp) <sub>2</sub> (P'') <sub>2</sub> ]	2.553(1)	<i>d</i> <sub>6</sub> -Benzene	–98.83, s	—	—	—	91
<b>48-Zr</b> [Zr(Cp) <sub>2</sub> (P'') <sub>2</sub> ]	—	<i>d</i> <sub>6</sub> -Benzene	–72.18, s	—	—	—	91
<b>49-Hf</b> [Hf(Cp) <sub>2</sub> (Cl)(P'')]	—	<i>d</i> <sub>6</sub> -Benzene	–153.97, s	—	—	—	91
<b>49-Zr</b> [Zr(Cp) <sub>2</sub> (Cl)(P'')]	2.547(6)	<i>d</i> <sub>8</sub> -Toluene	–108.90, s	—	—	—	92
<b>50-Hf</b> [Hf(Cp) <sub>2</sub> (Me)(P'')]	—	<i>d</i> <sub>6</sub> -Benzene	–141.92, s	—	—	—	91
<b>50-Zr</b> [Zr(Cp) <sub>2</sub> (Me)(P'')]	2.629(3)	—	—	—	—	—	91
<b>51</b> [Zr(C <sub>5</sub> H <sub>4</sub> Me) <sub>2</sub> (P'') <sub>2</sub> ]	2.617(3)	<i>d</i> <sub>6</sub> -Benzene	–71.20, s	—	—	—	93
<b>52</b> [Zr(Cp) <sub>2</sub> (Cl){P(SiMe <sub>3</sub> ) <sub>2</sub> C(H)C(Ph)}]	2.855(4)	<i>d</i> <sub>6</sub> -Benzene	–179.50, s	—	—	—	94
<b>53-Me</b> [Zr(Cp) <sub>2</sub> (Me){P(SiMe <sub>3</sub> ) <sub>2</sub> -C(H)C(Ph)}]	2.915(3)	<i>d</i> <sub>6</sub> -Benzene	–179.30, s	—	—	—	94
<b>53-Bu</b> [Zr(Cp) <sub>2</sub> ( <sup>n</sup> Bu){P(SiMe <sub>3</sub> ) <sub>2</sub> -C(H)C(Ph)}]	—	<i>d</i> <sub>6</sub> -Benzene	–176.60, s	—	—	—	94
<b>53-C</b> ≡ <b>CPh</b> [Zr(Cp) <sub>2</sub> (C ≡ CPh){P(SiMe <sub>3</sub> ) <sub>2</sub> C(H)C(Ph)}]	2.774(3)	<i>d</i> <sub>6</sub> -Benzene	–189.90, s	—	—	—	95
<b>54-Cr</b> [Cr(Cp)(μ-P'') <sub>2</sub> ]	2.3839(13)	<i>d</i> <sub>6</sub> -Benzene	No resonance observed	—	—	—	96
<b>54-Mn</b> [Mn(Cp)(μ-P'') <sub>2</sub> ]	2.5099(7)	—	—	—	—	—	85
<b>55</b> [Li(12-crown-4) <sub>2</sub> ][Mo(Cp)(CO) <sub>2</sub> (μ-P'')]	2.4304(6)	—	—	—	—	—	97
<b>56</b> [1,2-W <sub>2</sub> (P'')(NMe <sub>2</sub> ) <sub>4</sub> ]	2.423(3)	<i>d</i> <sub>6</sub> -Benzene	–106.70, s <sup>i</sup> –88.30, s <sup>j</sup>	<sup>1</sup> J <sub>WP</sub> = 243 <sup>i</sup> <sup>1</sup> J <sub>WP</sub> = 250 <sup>j</sup>	—	—	98
<b>57</b> [1,2-W <sub>2</sub> (PCy <sub>2</sub> )(P'')(NMe <sub>2</sub> ) <sub>4</sub> ]	2.425(4)	<i>d</i> <sub>6</sub> -Benzene	–106.90, s <sup>i</sup> –96.30, s <sup>j</sup>	<sup>1</sup> J <sub>WP</sub> = 239 <sup>i</sup> <sup>1</sup> J <sub>WP</sub> = 234 <sup>j</sup>	—	—	98
<b>58</b> [{Mn(P'')(μ-P'') <sub>2</sub> }{Mn(P'')(THF)}]	2.526(5) <sup>c</sup> 2.439(4) <sup>d</sup>	—	—	—	—	—	82
<b>59-EtMe<sub>4</sub></b> [Fe(C <sub>5</sub> EtMe <sub>4</sub> )(CO) <sub>2</sub> (P'')]	2.359(3)	<i>d</i> <sub>6</sub> -Benzene	–219.10, s	—	6.84, d	<sup>1</sup> J <sub>Psi</sub> = 46.7	99
<b>59-BuMe<sub>4</sub></b> [Fe(C <sub>5</sub> <sup>n</sup> BuMe <sub>4</sub> )(CO) <sub>2</sub> (P'')]	—	<i>d</i> <sub>6</sub> -Benzene	–218.60, s	—	6.84, d	<sup>1</sup> J <sub>Psi</sub> = 46.4	99
<b>59-1,3-Bu<sub>2</sub>H<sub>3</sub></b> [Fe(C <sub>5</sub> H <sub>3</sub> <sup>n</sup> Bu <sub>2</sub> -1,3)(CO) <sub>2</sub> (P'')]	—	<i>d</i> <sub>6</sub> -Benzene	–266.40, s	—	7.63, d	<sup>1</sup> J <sub>Psi</sub> = 53.0	99
<b>60-Et</b> [Ni(κ <sup>2</sup> -depe)(P'')]	—	—	—	—	—	—	100
<b>60-Cy</b> [Ni(κ <sup>2</sup> -dcpe)(P'')]	2.225(2)	—	—	—	—	—	100
<b>60-Ph</b> [Ni(κ <sup>2</sup> -dppe)(P'')]	—	—	—	—	—	—	100
<b>61-Ni<sup>n</sup>Bu</b> [Ni(Br)( <sup>n</sup> Bu <sub>2</sub> -bimy) <sub>2</sub> (P'')]	—	<i>d</i> <sub>6</sub> -Benzene	–197.30, s	—	—	—	101
<b>61-Ni<sup>t</sup>Pr</b> [Ni(t)( <sup>t</sup> Pr <sub>2</sub> -bimy) <sub>2</sub> (P'')]	—	<i>d</i> <sub>6</sub> -Benzene	–178.60, s	—	—	—	101
<b>61-Pd<sup>n</sup>Bu</b> [Pd(t)( <sup>n</sup> Bu <sub>2</sub> -bimy) <sub>2</sub> (P'')]	2.3648(17)	<i>d</i> <sub>6</sub> -Benzene	–192.80, s	—	—	—	101
<b>61-Pd<sup>t</sup>Pr</b> [Pd(t)( <sup>t</sup> Pr <sub>2</sub> -bimy) <sub>2</sub> (P'')]	2.3442(12)	<i>d</i> <sub>6</sub> -Benzene	–181.70, s	—	—	—	101
<b>62</b> [Ni(PMe <sub>3</sub> )(μ-P'') <sub>2</sub> ]	2.381(1) <sup>c</sup> 2.129(2) <sup>d</sup>	—	—	—	—	—	102
<b>63-Pd</b> [{Pd(PPh <sub>3</sub> ) <sub>2</sub> (P'')}{Si[N( <sup>t</sup> Bu) <sub>2</sub> CPh]}]	—	<i>d</i> <sub>8</sub> -THF	–173.00, t	<sup>2</sup> J <sub>PP</sub> = 5.2	0.01, d	<sup>1</sup> J <sub>Psi</sub> = 6.0	103
<b>63-Pt</b> [{Pt(PPh <sub>3</sub> ) <sub>2</sub> (P'')}{Si[N( <sup>t</sup> Bu) <sub>2</sub> CPh]}]	2.364(3)	<i>d</i> <sub>6</sub> -Benzene	–94.60, t	<sup>2</sup> J <sub>PP</sub> = 33.3 <sup>1</sup> J <sub>Psi</sub> = 219.2 <sup>1</sup> J <sub>PtP</sub> = 1633	8.3, d	<sup>1</sup> J <sub>Psi</sub> = 7.4	103
<b>64</b> [Cu( <sup>t</sup> Pr <sub>2</sub> -bimy) <sub>2</sub> (P'')]	2.291(7)	<i>d</i> <sub>6</sub> -Benzene	–261.00, s	—	—	—	104
<b>65</b> [Au( <sup>t</sup> Pr <sub>2</sub> -bimy)(P'')]	—	<i>d</i> <sub>6</sub> -Benzene	–235.00, s	—	—	—	105
<b>66-Cu</b> [Cu( <sup>t</sup> Pr)(P'')]	2.1913(15)	<i>d</i> <sub>6</sub> -Benzene	–268.00, s	—	—	—	104
<b>66-Au</b> [Au( <sup>t</sup> Pr)(P'')]	2.3174(10)	<i>d</i> <sub>6</sub> -Benzene	–235.70, s	—	—	—	105
<b>67</b> [Au(CAAC <sup>MEt</sup> )(P'')]	2.3253(5)	<i>d</i> <sub>6</sub> -Benzene	–233.10, s	—	—	—	106
<b>68-Cu</b> [Cu(μ-P'') <sub>6</sub> ]	2.210(4)	<i>d</i> <sub>6</sub> -Benzene	–149.00, s	—	—	—	104



Table 1 (continued)

Complex	Mean M–P (Å)	Solvent	<sup>31</sup> P Chemical shift (δ)	<sup>31</sup> P Coupling constant (Hz)	<sup>29</sup> Si Chemical shift (δ)	<sup>29</sup> Si Coupling constant (Hz)	Ref.
<b>68-Ag</b> [Ag(μ-P'')] <sub>6</sub>	2.404(4)	<i>d</i> <sub>6</sub> -Benzene	–236.00, s	—	—	—	104
<b>69</b> [Hg(P'')] <sub>2</sub>	2.406(1)	<i>d</i> <sub>6</sub> -Benzene	–162.00, s	—	—	—	83
<b>70-Zn</b> [Zn(P'')(μ-P'')] <sub>2</sub>	2.420(1) <sup>c</sup> 2.295(1) <sup>d</sup>	<i>d</i> <sub>6</sub> -Benzene	–183.00, br, s <sup>c</sup> –237.30, br, s <sup>d</sup>	—	—	—	83 83
<b>70-Cd</b> [Cd(P'')(μ-P'')] <sub>2</sub>	2.594(1) <sup>c</sup> 2.459(1) <sup>d</sup>	<i>d</i> <sub>6</sub> -Benzene	–180.10, br, s <sup>c</sup> –229.50, br, s <sup>d</sup>	—	—	—	83 83
<b>71-<sup>t</sup>Bu</b> [Zn( <sup>t</sup> Bu)(μ-P'')] <sub>2</sub>	—	<i>d</i> <sub>6</sub> -Benzene	–216.70, s <sup>c</sup>	—	4.17, s	—	107
<b>71-<sup>i</sup>Pr</b> [Zn( <sup>i</sup> Pr)(μ-P'')] <sub>2</sub>	2.411(4)	<i>d</i> <sub>6</sub> -Benzene	–216.30, s	—	4.24, s	—	107
<b>71-CH<sub>2</sub>SiMe<sub>3</sub></b> [Zn(CH <sub>2</sub> SiMe <sub>3</sub> )(μ-P'')] <sub>2</sub>	2.415(1)	<i>d</i> <sub>6</sub> -Benzene	–205.80, s	—	4.38, s	—	107
<b>72-Zn</b> [Zn(P''Pr <sub>3</sub> )Cl(μ-P'')] <sub>2</sub>	2.437(1) <sup>c</sup> 2.419(1) <sup>d</sup>	—	—	—	—	—	108
<b>72-Cd</b> [Cd(P''Pr <sub>3</sub> )Br(μ-P'')] <sub>2</sub>	2.590(1) <sup>c</sup> 2.578(1) <sup>d</sup>	—	—	—	—	—	108
<b>73</b> [{Zn(Cl)(MeCN)(μ-P'')] <sub>2</sub> }{Zn(μ-Cl)} <sub>2</sub>	2.374(3)	—	—	—	—	—	108
<b>74</b> [N <sup>t</sup> Bu <sub>4</sub> ] <sub>2</sub> [Cd <sub>4</sub> (μ) <sub>8</sub> (P'')] <sub>2</sub>	2.507(7)	—	—	—	—	—	108
<b>75-Me</b> [Zn(Me)(P'')] <sub>3</sub>	2.390(3)	<i>d</i> <sub>6</sub> -Benzene	–246.80, s	—	4.62, s	—	107
<b>75-Et</b> [Zn(Et)(P'')] <sub>3</sub>	—	<i>d</i> <sub>6</sub> -Benzene	–246.90, s	—	4.61, s	—	107
<b>75-<sup>i</sup>Pr</b> [Zn( <sup>i</sup> Pr)(P'')] <sub>3</sub>	2.408(6)	<i>d</i> <sub>6</sub> -Benzene	–243.60, s	—	—	—	107
<b>75-<sup>n</sup>Bu</b> [Zn( <sup>n</sup> Bu)(P'')] <sub>3</sub>	2.388(4)	<i>d</i> <sub>6</sub> -Benzene	–246.00, s	—	4.67, s	—	107
<b>76-Cr</b> [Cr(CO) <sub>5</sub> (μ-P'')]{Al(dmap)(Me <sub>2</sub> )}	2.528(1)	<i>d</i> <sub>2</sub> -DCM	–278.10, s	—	—	—	39
<b>76-Fe</b> [Fe(CO) <sub>4</sub> (μ-P'')]{Al(dmap)(Me <sub>2</sub> )}	2.377(1)	<i>d</i> <sub>8</sub> -Toluene	–258.40, s	—	—	—	39
<b>76-Ni</b> [Ni(CO) <sub>3</sub> (μ-P'')]{Al(dmap)(Me <sub>2</sub> )}	2.315(2)	<i>d</i> <sub>6</sub> -Benzene	–275.20, s	—	—	—	39 and 109
<b>77-Cr</b> [Zr(C <sub>5</sub> H <sub>4</sub> Me) <sub>2</sub> (μ-P'')] <sub>2</sub> Cr(CO) <sub><i>n</i></sub>	2.656(6)	<i>d</i> <sub>6</sub> -Benzene	–38.40, s	—	—	—	110
<b>77-Mo</b> [Zr(Cp) <sub>2</sub> (μ-P'')] <sub>2</sub> Mo(CO) <sub>4</sub>	2.6648(11)	<i>d</i> <sub>6</sub> -Benzene	–57.90, s	—	—	—	110
<b>77-Ni</b> [Zr(Cp) <sub>2</sub> (μ-P'')] <sub>2</sub> Ni(CO) <sub>2</sub>	2.654(1)	<i>d</i> <sub>6</sub> -Benzene	–42.10, s	—	—	—	111
f-Block							
<b>46-Ce</b> [Ce(P'') <sub>3</sub> (THF) <sub>2</sub> ]	2.849(3)	<i>d</i> <sub>6</sub> -Benzene	616.7, br	—	5.30, s	—	89
<b>46-Pr</b> [Pr(P'') <sub>3</sub> (THF) <sub>2</sub> ]	2.837(3)	<i>d</i> <sub>6</sub> -Benzene	1894.2, br	—	15.65, s	—	89
<b>46-Nd</b> [Nd(P'') <sub>3</sub> (THF) <sub>2</sub> ]	2.818(2)	<i>d</i> <sub>6</sub> -Benzene	2570.3, br	—	42.94, s	—	89 and 112
<b>46-Sm</b> [Sm(P'') <sub>3</sub> (THF) <sub>2</sub> ]	2.789(3)	<i>d</i> <sub>6</sub> -Benzene	–259.2, br	—	0.52, s	—	89
<b>46-Tm</b> [Tm(P'') <sub>3</sub> (THF) <sub>2</sub> ]	2.707(16)	<i>d</i> <sub>6</sub> -Benzene	—	—	—	—	113
<b>78</b> [Sm(P'')(μ-P'')] <sub>3</sub> Sm(THF) <sub>3</sub>	3.039(5) <sup>c</sup> 3.027(3) <sup>d</sup>	<i>d</i> <sub>8</sub> -THF	—	—	—	—	114
<b>79-Sm</b> [{Sm(P'') <sub>3</sub> (THF)} <sub>2</sub> (μ-I)K <sub>3</sub> (THF)]	3.033(5)	<i>d</i> <sub>6</sub> -Benzene	—	—	—	—	115
<b>79-Eu</b> [{Eu(P'') <sub>3</sub> (THF)} <sub>2</sub> (μ-I)K <sub>3</sub> (THF)]	3.0316(4)	<i>d</i> <sub>6</sub> -Benzene	—	—	—	—	115
<b>80</b> [KYb(P'') <sub>3</sub> (μ-K)(P'')] <sub>2</sub> ∞	2.952(10)	<i>d</i> <sub>6</sub> -Benzene	—	—	—	—	115
<b>81-Sm</b> <i>trans</i> -[Sm(P'') <sub>2</sub> (py) <sub>4</sub> ]	3.0342(9)	<i>d</i> <sub>6</sub> -Benzene	—	—	—	—	115
<b>81-Eu</b> <i>trans</i> -[Eu(P'') <sub>2</sub> (py) <sub>4</sub> ]	3.0364(7)	<i>d</i> <sub>6</sub> -Benzene	—	—	—	—	115
<b>81-Yb</b> <i>trans</i> -[Yb(P'') <sub>2</sub> (py) <sub>4</sub> ]	2.928(13)	<i>d</i> <sub>6</sub> -Benzene	–253.93, s	<sup>1</sup> J <sub>YbP</sub> = 925	1.58, Vir. t	<sup>1</sup> J <sub>PSi</sub> = 16	115
<b>82-Sm</b> [Sm(P'') <sub>2</sub> (18-crown-6)]	3.089(3)	<i>d</i> <sub>6</sub> -Benzene	—	—	—	—	115
<b>82-Eu</b> [Eu(P'') <sub>2</sub> (18-crown-6)]	3.096(4)	<i>d</i> <sub>6</sub> -Benzene	—	—	—	—	115
<b>82-Yb</b> [Yb(P'') <sub>2</sub> (18-crown-6)]	2.9662(10)	<i>d</i> <sub>6</sub> -Benzene	–265.58, s	<sup>1</sup> J <sub>YbP</sub> = 977	1.94, Vir. t	<sup>1</sup> J <sub>PSi</sub> = 17	115
<b>83-Th</b> [Th(P'')(Cp*) <sub>2</sub> (Cl)]	—	<i>d</i> <sub>6</sub> -Benzene	109.0, s	—	—	—	116
<b>83-U</b> [U(P'')(Cp*) <sub>2</sub> (Cl)]	2.788(4)	<i>d</i> <sub>6</sub> -Benzene	—	—	—	—	116
<b>84-Th</b> [Th(P'')(Cp*) <sub>2</sub> (Me)]	2.888(4)	<i>d</i> <sub>6</sub> -Benzene	115.22, s	—	—	—	116
<b>84-U</b> [U(P'')(Cp*) <sub>2</sub> (Me)]	2.893(4)	<i>d</i> <sub>6</sub> -Benzene	—	—	—	—	116
<b>85-Th</b> [Th{P(SiMe <sub>3</sub> )(SiMe <sub>2</sub> CH <sub>2</sub> )}(Cp*) <sub>2</sub> ]	—	<i>d</i> <sub>6</sub> -Benzene	95.88, s	—	—	—	116
<b>85-U</b> [U{P(SiMe <sub>3</sub> )(SiMe <sub>2</sub> CH <sub>2</sub> )}(Cp*) <sub>2</sub> ]	2.655(6)	<i>d</i> <sub>6</sub> -Benzene	—	—	—	—	116
<b>86-Th</b> [Th(Tren <sup>DMBS</sup> )(P'')]	2.9406(11)	<i>d</i> <sub>6</sub> -Benzene	–100.09, s	—	—	—	117
<b>86-U</b> [U(Tren <sup>DMBS</sup> )(P'')]	2.8646(14)	<i>d</i> <sub>6</sub> -Benzene	2055.21, s	—	—	—	117
<b>87-Th</b> [Th(Tren <sup>TIPS</sup> )(P'')]	2.9020(13)	<i>d</i> <sub>6</sub> -Benzene	–66.45, s	—	8.49, s	—	117
<b>87-U</b> [U(Tren <sup>TIPS</sup> )(P'')]	2.8391(9)	<i>d</i> <sub>6</sub> -Benzene	—	—	–97.09, s	—	117

<sup>a</sup> Refers to stile. <sup>b</sup> Refers to rung bonds of ladder complexes. <sup>c</sup> Refers to bridging. <sup>d</sup> Refers to terminal P'' ligands. <sup>e</sup> Refers to *syn*-isomers. <sup>f</sup> Refers to *anti*-isomers. <sup>g</sup> Refers to *cis*-isomers. <sup>h</sup> Refers to *trans*-isomers. <sup>i</sup> Refers to the *anti*-conformation. <sup>j</sup> Refers to the *gauche*-conformation.

showing how decreasing the amount of donor solvent molecules gives rise to an increase in aggregation.<sup>26</sup> The mean Li–P bond lengths in **1** (2.62(2) Å) are longer than those found in **2** (2.559(4) Å) and the ladder complex **3** (2.50(4) Å for the Stiles and 2.56(4) Å for the rungs of the ladder).<sup>24,25</sup> The six-step ladder complex **4** exhibits equal Li–P bond lengths for the Stiles of the ladder of 2.51(1) Å,<sup>26</sup> but the Li–P distances of the rungs of the ladder

vary considerably (2.38(1)–2.63(1) Å),<sup>26</sup> the shortest Li–P bond lengths are seen for the μ<sub>2</sub>-P'' at the termini as these Li ions are only two-coordinate, and unusually exhibit bent geometries.<sup>119</sup> Complex **4** dissolves in THF and disaggregates to form complex **3**.<sup>26</sup> The related solvent-free Li N'' complexes exist as cyclic trimers, [Li(μ-N'')]<sub>3</sub>, in which the ring is planar,<sup>120,121</sup> or as dimers in [Li(μ-N'')]<sub>2</sub>.<sup>122</sup>





Fig. 2 Group 1  $M P''$  complexes  $[\text{Li}(\mu\text{-}P'')(\text{THF})_2]_2$  (**1**),  $[\text{Li}_4(\mu_2\text{-}P'')_2(\mu_3\text{-}P'')_2(\text{THF})_2]$  (**2**),  $[\text{Li}(\mu\text{-}P'')(DME)_2]_2$  (**3**),  $[\text{Li}_6(\mu_2\text{-}P'')_2(\mu_3\text{-}P'')_4]$  (**4**),  $[\text{M}_4(\mu_2\text{-}P'')_2(\mu_3\text{-}P'')_2]_\infty$  (**5-M**;  $M = \text{K}, \text{Rb}, \text{Cs}$ ),  $[\text{Cs}(\mu\text{-}P'')(\mu\text{-}1,4\text{-dioxane})_3(1,4\text{-dioxane})]_\infty$  (**6**).

Similar ladder-like structures were obtained for  $P''$  complexes of the larger group 1 metals K, Rb and Cs by Ruhlandt-Senge and Uhlig in 1998.<sup>27</sup> The polymeric heavy alkali metal  $P''$  complexes  $[\text{M}_4(\mu_2\text{-}P'')_2(\mu_3\text{-}P'')_2]_\infty$  (**5-M**;  $M = \text{K}, \text{Rb}, \text{Cs}$ , Fig. 2) were synthesised by the treatment of  $\text{P}(\text{SiMe}_3)_3$  with the parent metal alkoxide  $\text{MO}^t\text{Bu}$  ( $M = \text{K}, \text{Rb}, \text{Cs}$ ) in THF, and were found to contain four-coordinate metal centres and five-coordinate phosphorus atoms in the solid state.<sup>27</sup> Whilst complexes **5-K** and **5-Rb** were found to react rapidly with air and moisture, **5-Cs** spontaneously ignites upon contact with air, owing to the increase in reactivity with the size of the alkali metal ion.<sup>27</sup> A higher degree of oligomerisation was observed for the heavy group 1  $P''$  complexes in the solid state in comparison to their lighter Li analogues;<sup>26,27</sup> this is limited to growth of the Stiles of the ladder, signifying that the steric bulk of  $P''$  is sufficient to prevent oligomer growth in other directions. The  $M\text{-P}$  bond lengths of **5-M** increase as the size of the metal increases from K-Cs; however, the  $\text{K-P}$  bond lengths of **5-K** (3.3169(7), 3.4063(8), and 3.4273(8) Å) are slightly longer than expected from ionic radii trends. This was attributed to the steric demand of  $P''$  having a greater effect on the smaller K ion than

for Rb or Cs, with **5-Rb** and **5-Cs** having mean  $M\text{-P}$  bond lengths of 3.486(2) Å and 3.656(2) Å, respectively.<sup>27,123</sup>

A 1,4-dioxane-bridged polymeric Cs  $P''$  complex  $[\text{Cs}(\mu\text{-}P'')(\mu\text{-}1,4\text{-dioxane})_3(1,4\text{-dioxane})]_\infty$  (**6**, Fig. 2) was synthesised in 2008 by Ionkin by the reaction of  $\text{P}(\text{SiMe}_3)_3$  with  $\text{CsF}$  in 1,4-dioxane at 90 °C for 3 hours.<sup>28</sup> This study focussed on using  $\text{CsF}$ ,  $\text{P}(\text{SiMe}_3)_3$  and 2,4,6-tri-*tert*-butylphenylchloride ( $\text{Mes}^*\text{Cl}$ ) in order to form  $\text{Mes}^*P''$ , as this reaction does not occur at room temperature without reactive alkali metal precursors present.<sup>28</sup> The reaction progress was monitored by  $^{31}\text{P}\{^1\text{H}\}$  NMR spectroscopy as **6** shows a distinctive chemical shift of  $-276.10$  ppm.<sup>27,28</sup> As seen previously for the ladder-like oligomers **5**, larger alkali metals like Cs give higher coordination numbers and higher order aggregates.<sup>28</sup> The mean  $\text{Cs-P}$  bond length of **6** (3.6137(17) Å) is slightly longer than those seen for **5-Cs**.<sup>27,28</sup>

**2.1.2 Group 2 complexes.** Alkali metal ligand transfer agents often act as reducing agents and so are not always well-suited for straightforward salt metathesis reactions, therefore alkaline earth complexes such as Grignard reagents are often employed.<sup>124</sup> The dimeric Mg  $P''$  complex  $[\text{Mg}(\mu\text{-}P'')(\text{Br})(\text{THF})_2]$  (**7**, Fig. 3) was prepared by Coles in 2010 *via* the protonolysis reaction of  $\text{HP}''$



Fig. 3 Mg  $P''$  complexes  $[\text{Mg}(\mu\text{-}P'')(\text{Br})(\text{THF})_2]$  (**7**),  $[\text{Mg}_3(\mu\text{-}P'')_4(P'')_2]$  (**8**),  $[\text{Mg}(P''\text{C}(\text{Ph})\text{C}(\text{C}_2\text{Ph}))(\mu\text{-}P'')]_2$  (**9**),  $[\text{Mg}(P'')_2(\text{DME})]$  (**10**),  $[\text{Mg}(P'')_2(\text{THF})_2]$  (**11**).



with MeMgBr in Et<sub>2</sub>O.<sup>20</sup> The related alkyl complex [Mg(μ-P'')(Me)(THF)]<sub>2</sub> was also identified in the reaction mixture by <sup>1</sup>H NMR spectroscopy from the protonolysis reaction of HP'' with MgMe<sub>2</sub>, which is present due to the Schlenk equilibrium of MeMgBr.<sup>20</sup> A trinuclear Mg P'' complex, [Mg<sub>3</sub>(μ-P'')<sub>4</sub>(P'')<sub>2</sub>] (**8**, Fig. 3), was synthesised by Westerhausen in 1998 by a similar protocol; in solution an equilibrium was observed with a dimeric form of this complex, formulated as [Mg<sub>2</sub>(μ-P'')<sub>2</sub>(P'')<sub>2</sub>], as evidenced by two sets of triplets in the <sup>31</sup>P{<sup>1</sup>H} NMR spectrum (δ<sub>P</sub> = -242.55 and -275.44 ppm; <sup>2</sup>J<sub>PP</sub> = 17.9 Hz) for the bridging and terminal P'' ligands, respectively.<sup>29</sup> A variable temperature (VT) experiment was performed in order to deconvolute the spectrum and show that the dinuclear complex is favoured at higher temperatures.<sup>29</sup> The mean Mg-P<sub>terminal</sub> bond lengths of the two periphery three-coordinate Mg ions in **8** (2.455(4) Å) are shorter than their mean Mg-P<sub>bridging</sub> distances (2.546(5) Å), whilst the central four-coordinate Mg ion exhibits a range of Mg-P<sub>bridging</sub> bond lengths (2.605(3)–2.678(2) Å). This is due to the large steric effect of four P'' ligands about a single Mg ion in **8**; the bridging Mg-P bond lengths in **7** (2.5624(7) Å) are shorter than those seen in **8**.<sup>20,29</sup>

In 1998, Westerhausen reported the reaction of [Mg(P'')<sub>2</sub>]<sub>n</sub> with diphenylbutadiyne to give the dimeric [2 + 2]-cycloaddition product [Mg{P''C(Ph)C(C<sub>2</sub>Ph)}(μ-P'')]<sub>2</sub> (**9**, Fig. 3), which contains MgPC<sub>2</sub> four-membered rings with mean Mg-P<sub>bridging</sub> bond lengths of 2.5576(19) Å.<sup>30</sup> The formation of **9** was confirmed by <sup>31</sup>P{<sup>1</sup>H} NMR spectroscopy, with two triplets at -98.4 and -262.3 ppm for the phosphine and bridging phosphide groups, respectively (<sup>2</sup>J<sub>PP</sub> = 5 Hz).<sup>30</sup> Evidence of weak conjugation of the triple and double bonds within **9** was provided by their respective stretching bands in the IR spectrum at 2131 and 1596 cm<sup>-1</sup>.<sup>30</sup>

The monomeric solvated Mg P'' complexes [Mg(P'')<sub>2</sub>(DME)] (**10**, Fig. 3) and [Mg(P'')<sub>2</sub>(THF)]<sub>2</sub> (**11**, Fig. 3) were synthesised by Westerhausen in 1994 and 1995, respectively, by the separate protonolysis reactions of two equivalents of HP'' with MgR<sub>2</sub> (R = <sup>n</sup>Bu, <sup>sec</sup>Bu, N'') in the respective donor solvent.<sup>31,32</sup> Both complexes exhibit distorted tetrahedral geometries about the

Mg ion, with the phosphorus atoms showing pyramidal geometries in the solid state with Mg-P bond lengths of 2.487(2) Å (**10**) and 2.5023(19) Å (**11**).<sup>31,32</sup> Complex **11** exhibits a single resonance in the <sup>31</sup>P{<sup>1</sup>H} NMR spectrum at -294.7 ppm; the respective NMR data for **10** were not reported but a similar chemical shift would be expected.<sup>31,32</sup>

The dimeric heavy group 2 metal P'' complexes, [M(N'')(μ-P'')(sol)]<sub>2</sub> (**12-Ca**: M = Ca, sol = THF; **12-Ba**: M = Ba, sol = DME; Fig. 4) were synthesised by Westerhausen in 1995 and 1996, respectively, by the separate reactions of [Ca(N'')<sub>2</sub>(THF)]<sub>2</sub> or [Ba(N'')<sub>2</sub>] with HP'' in toluene or DME.<sup>33,34</sup> These bimetallic complexes, which represented the first mixed bis(trimethylsilyl)pnictide complexes at the time, consist of two four-coordinate (**12-Ca**) or five-coordinate (**12-Ba**) metal centres, each with a bridging P'', a terminal N'', and monodentate THF for **12-Ca** or bidentate DME for **12-Ba**.<sup>33,34</sup> The <sup>31</sup>P{<sup>1</sup>H} NMR spectrum of **12-Ca** indicates that a mixture of geometrical isomers is present in solution, with the *syn*-conformer resonance at (-229.96 ppm) and the *anti*-conformer at -254.82 ppm, whereas only one conformer is present in solution for **12-Ba** (-222.05 ppm).<sup>33,34</sup> For **12-Ca**, it was noted that the central Ca<sub>2</sub>P<sub>2</sub> ring is highly distorted, with Ca-P bond lengths of 2.927(2) and 3.005(2) Å, whereas the Ba<sub>2</sub>P<sub>2</sub> ring in **12-Ba** is essentially planar with Ba-P bond lengths of 3.289(2) and 3.340(2) Å; this is likely a consequence of the larger Ba ion better accommodating the bulky P'' ligands.<sup>33,34</sup>

The protonolysis reaction of **12-Ca** with HP'' in THF gave the monomeric solvated P'' complex *trans*-[Ca(P'')<sub>2</sub>(THF)]<sub>4</sub> (**13-Ca**, Fig. 4).<sup>33</sup> Similarly, the separate reactions of [M(N'')<sub>2</sub>(THF)]<sub>n</sub> (M = Sr, n = 0; M = Ba, n = 2) with two equivalents of HP'' in THF gave the heavier homologues, *trans*-[M(P'')<sub>2</sub>(THF)]<sub>4</sub> (**13-Ba**,<sup>36</sup> **13-Sr**, Fig. 4).<sup>35</sup> All three complexes exhibit distorted octahedral geometries with two axial P'' and four equatorial THF, with M-P distances of 2.911(2) and 2.924(2) Å (**13-Ca**),<sup>33</sup> 3.035(6) and 3.006(6) Å (**13-Sr**),<sup>35</sup> and 3.158(6) and 3.190(6) Å (**13-Ba**),<sup>36</sup> and near-linear P-M-P angles of 175.16(7)° (**13-Ca**),<sup>33</sup> 174.2(2)° (**13-Sr**)<sup>35</sup> and 174.9(1)° (**13-Ba**).<sup>36</sup> When **13-Sr** is heated under vacuum, the bimetallic complex, [Sr(P'')(μ-P'')<sub>3</sub>Sr(THF)]<sub>3</sub>



Fig. 4 Heavy group 2 P'' complexes [M(N'')(μ-P'')(sol)]<sub>2</sub> (**12-Ca**: M = Ca, sol = THF; **12-Ba**: M = Ba, sol = DME), *trans*-[M(P'')<sub>2</sub>(THF)]<sub>4</sub> (**13-M**; M = Ca, Sr, Ba), [Sr(P'')(μ-P'')<sub>3</sub>Sr(THF)]<sub>3</sub> (**14**), [Ca(P'')<sub>2</sub>(TMATA)]<sub>2</sub> (**15**, TMATA = 1,3,5-trimethyl-1,3,5-triazinane).



(**14**, Fig. 4) forms, which has one four-coordinate Sr ion with three bridging and one terminal P'', and one six-coordinate Sr ion with three bridging P'' and three bound THF.<sup>35</sup> Complex **14** is structurally analogous to that of the bimetallic Sm species discussed in Section 2.4.1.<sup>125</sup>

Finally, the reaction of  $[\text{Ca}(\text{N}'')(\mu\text{-N}'')]_2$  with four equivalents each of HP'' and 1,3,5-trimethyl-1,3,5-triazinane (TMTA) gave the eight-coordinate  $C_2$ -symmetric Ca complex,  $[\text{Ca}(\text{P}'')_2(\text{TMTA})_2]$  (**15**, Fig. 4).<sup>36</sup> This complex contains two P'' and two tridentate TMTA ligands in chair conformations, with two identical Ca–P bond lengths of 2.994(2) Å, and a wide range of Ca–N bond lengths between 2.575(4)–2.704(5) Å, which was attributed to steric buttressing.<sup>36</sup>

## 2.2 p-Block complexes

As stated previously p-block P'' compounds containing direct E–P bonds with E = non-metals and the metalloids B and Si are not within the scope of this review. Surprisingly no structurally authenticated examples of As, Sb, Bi or Te P'' complexes were found in the CSD;<sup>9</sup> we note that a synthetic route to a trimeric Bi(I) complex  $[\text{Bi}(\mu\text{-P}'')]_3$  has been reported and it has been claimed that the solid-state structure has been determined, but this dataset has not been deposited on the CSD.<sup>12e</sup> The absence of group 15 and 16 metal(loid) P'' complexes to date is noteworthy given that this chemistry has been developed for the lighter congener N''.<sup>7</sup> This section is therefore divided into group 13 and group 14 metal P'' complexes, and subdivided into homo- and hetero-metallic examples, with the heavier metalloid Ge included for comparisons with heavier homologous Sn and Pb complexes.

**2.2.1 Homometallic Group 13 complexes.** Group 13 P'' complexes have received interest as reactive precursors for further chemical transformations such as in the formation of N-heterocyclic rings and cages.<sup>37</sup> There are currently seven structurally characterised examples of mononuclear group 13

P'' complexes, which all contain bulky supporting ligands.<sup>37</sup> The Al(III) P'' complex  $[\text{Al}(\text{P}'')(\text{tmp})_2]$  (tmp = 2,2,6,6-tetramethyl-piperidino) (**16**, Fig. 5) was synthesised by Nöth and Paine in 2007 by the salt metathesis reaction of  $[\text{Al}(\text{tmp})_2(\text{Cl})]$  with  $\text{LiP}''$  in hexane.<sup>37</sup> The reactions of  $\text{CO}_2$ ,  $\text{CS}_2$  and  $\text{OCS}$  with **16** proceeded *via* insertion into the Al–P bond to yield bridged species.<sup>37</sup> From 2001–2006, Schulz reported synthetic routes to the four-coordinate M(III) complexes  $[\text{M}(\text{R})_2(\text{P}'')(\text{dmap})]$  (**17-MR**. **17-AlMe**: M = Al, R = Me; **17-GaMe**: M = Ga, R = Me; **17-Ga<sup>t</sup>Bu**: M = Ga, R = <sup>t</sup>Bu; dmap = 4-(dimethylamino)pyridine, Fig. 5) which involved the addition of a base (dmap) to the heterocyclic complexes  $[\text{M}(\text{R})_2(\text{P}'')]_x$  ( $x = 1-3$ ).<sup>38-40</sup> The M–P bond lengths of the methyl derivatives (**17-AlMe**: 2.375(1) Å; **17-GaMe**: 2.372(1) Å) are relatively short, whilst that of **17-Ga<sup>t</sup>Bu** is slightly longer (2.3948(6) Å) due to the additional steric effects of the <sup>t</sup>Bu groups.<sup>38-40</sup> The separate reactions of these complexes with various d-transition metal carbonyl complexes gave heterobimetallic complexes, which are discussed in Section 2.3.

In 2005, Jones and Murphy reported the synthesis of the Ga(III) P'' complexes,  $[\text{Ga}(\text{P}'')(i)\text{-}^t\text{Bu-DAB}]$  (**18**, Fig. 5) and  $[\text{Ga}(\text{P}'')_2(^t\text{Bu-DAB})]$  (**19**, Fig. 5) (<sup>t</sup>Bu-DAB =  $\{\text{N}(^t\text{Bu})\text{C}(\text{H})_2\}_2$ ), by the reaction of the Ga(II) precursors  $[\text{Ga}(^t\text{Bu-DAB})(i)]_2$  with either two or four equivalents of  $\text{MP}''$  (M = Li, Na) in  $\text{Et}_2\text{O}$  at  $-78^\circ\text{C}$ .<sup>41</sup> The mechanism of formation of **18** is unclear, but involves both salt elimination and Ga–Ga bond cleavage, with oxidation of Ga(II) to Ga(III). When only one equivalent of  $\text{MP}''$  was used the unreacted Ga(II) starting material was isolated, whilst the addition of a further equivalent of  $\text{MP}''$  to **18** gave **19** in moderate yield.<sup>41</sup> Both of these Ga(III) P'' complexes have distorted tetrahedral geometries, with the Ga–P bond lengths being longer for **19** (2.343(2) Å mean) than for **18** (2.300(2) Å) due to the increased steric bulk about the metal centre.<sup>41</sup> Complexes **18** and **19** did not give NMR data that could be interpreted due to their paramagnetism, but EPR spectroscopy showed that there was a small amount of unpaired spin density




Fig. 5 Mononuclear group 13 P'' complexes  $[\text{Al}(\text{P}'')(\text{tmp})_2]$  (tmp = 2,2,6,6-tetramethyl-piperidino) (**16**),  $[\text{M}(\text{R})_2(\text{P}'')(\text{dmap})]$  (**17-MR**. **17-AlMe**: M = Al, R = Me; **17-GaMe**: M = Ga, R = Me; **17-Ga<sup>t</sup>Bu**: M = Ga, R = <sup>t</sup>Bu; dmap = 4-(dimethylamino)pyridine),  $[\text{Ga}(\text{P}'')(i)\text{-}^t\text{Bu-DAB}]$ , (**18**, <sup>t</sup>Bu-DAB =  $\{\text{N}(^t\text{Bu})\text{C}(\text{H})_2\}_2$ ),  $[\text{Ga}(\text{P}'')_2(^t\text{Bu-DAB})]$  (**19**),  $[\text{M}(\text{Dipp})(\text{P}'')(\text{Cl})]$  (<sup>Dipp</sup>Nacnac =  $\{\text{HC}[\text{C}(\text{Me})\text{N}(\text{Dipp})_2]\}_2$ , Dipp =  $\text{C}_6\text{H}_5\text{Pr}_{2-2,6}$ ) (**20-M**; M = Ga, In).



at the P atoms due to the splitting of the signal from coupling to  $^{31}\text{P}$  nuclei. A smaller line width was seen for **19** due to increased spin delocalisation at the  $\text{P}''$  ligands; despite this giving increased spin density at gallium, the unpaired electrons are mostly localised on the diazabutadiene ligands.<sup>41</sup> In 2019, Scheer investigated a series of group 13 phosphido complexes for their potential applications in nanoparticles, semiconductors and optoelectronic layers.<sup>42</sup> The authors postulated that bulky co-ligands would be needed in order to isolate mononuclear parent phosphido complexes, as there is a tendency towards oligomerisation due to the Lewis acidity of the group 13 metal centres and Lewis basicity of  $\text{PH}_2$ .<sup>42</sup> The reaction of  $[\text{M}(\text{DippP}(\text{Nacnac}))(\text{Cl})_2]$  ( $\text{M} = \text{Ga}, \text{In}$ ;  $\text{DippP}(\text{Nacnac}) = \{\text{HC}[\text{C}(\text{Me})\text{N}(\text{Dipp})]\}_2$ ,  $\text{Dipp} = \text{C}_6\text{H}_3\text{Pr}_{2,6}$ ) with excess  $\text{LiP}''$  in toluene gave the monosubstituted complexes  $[\text{M}(\text{DippP}(\text{Nacnac}))(\text{P}'')(\text{Cl})]$  (**20-M**;  $\text{M} = \text{Ga}, \text{In}$ , Fig. 5); the disubstituted  $\text{P}''$  complexes could not be prepared due to the steric bulk of the  $\beta$ -diketiminato ligand.<sup>42</sup>  $^{31}\text{P}\{^1\text{H}\}$  NMR spectroscopy revealed singlet resonances at  $-255.0$  ppm (**20-Ga**) and  $-252.9$  ppm (**20-In**), whilst single crystal XRD showed that these complexes have heavily distorted tetrahedral geometries, with N–M–N angles of  $96.03(13)^\circ$  (**20-Ga**) and  $90.33(10)^\circ$  (**20-In**); the M–P bond length of  $2.3310(9)$  Å (**20-Ga**) is shorter than that found for **20-In** ( $2.4806(8)$  Å).<sup>42</sup>

There have been numerous published examples of symmetric dimeric group 13  $\text{P}''$  complexes,  $[\text{M}(\text{R})_2(\mu\text{-P}'')]_2$  (**21-MR**;  $\text{M} = \text{Al}, \text{Ga}, \text{In}, \text{Tl}$ ;  $\text{R} = \text{Me}, \text{Et}, \text{Ph}, \text{}^t\text{Bu}, \text{CH}_2\text{Ph}, \text{CH}_2\text{}^i\text{Pr}, \text{CH}_2\text{}^t\text{Bu}, \text{CH}_2\text{SiMe}_3$ , Fig. 6), as these have found interest as precursors for III–V semiconductor materials.<sup>43–57</sup> Each of these complexes consists of two four-coordinate distorted tetrahedral metal centres bridged by two  $\text{P}''$  ligands to give  $\text{M}_2\text{P}_2$  cores. In **21-ALR** the mean Al–P bond lengths increase as the steric influence of the R group increases: **21-AlMe**,  $2.457(2)$  Å; **21-AlEt**,  $2.458(1)$  Å; **21-AlCH}\_2\text{}^i\text{Pr}**,  $2.476(2)$  Å; **21-AlCH}\_2\text{SiMe}\_3,  $2.483(1)$  Å.<sup>43–46</sup> This trend**

is also seen for the **21-GaR** series: **21-GaMe**,  $2.450(1)$  Å; **21-GaEt**,  $2.4558(7)$  Å; **21-Ga}^t\text{Bu}**,  $2.4533(6)$  Å; **21-GaCH}\_2\text{}^t\text{Bu}**,  $2.517(3)$  Å; **21-GaCH}\_2\text{SiMe}\_3,  $2.4887(16)$  Å.<sup>40,47–51</sup> Finally, the In–P bond lengths in **21-InR** vary similarly: **21-InMe**,  $2.630(1)$  Å; **21-InEt**,  $2.645(1)$  Å; **21-InPh**,  $2.612(1)$  Å; **21-InCH}\_2\text{Ph}**,  $2.6123(6)$  Å; **21-InCH}\_2\text{SiMe}\_3,  $2.655(3)$  Å.<sup>52–55</sup> The variation of R groups in these complexes changes their functionalities; for example,  $[\text{In}(\text{}^t\text{Bu})_2(\mu\text{-P}'')]_2$  (**21-In}^t\text{Bu}**) was employed in the fabrication of nanowires of approximately 10–100 nm diameter.<sup>55</sup> In 2003 Schulz disclosed that the combination of **17-Al** with  $\text{GaMe}_3$  in the presence of dmap at  $-30$  °C yielded the adduct  $[\text{Al}(\text{Me})_2(\text{dmap})(\mu\text{-P}'')\text{Ga}(\text{Me})_3]$  (**22**, Fig. 6), as well as  $[\text{Ga}(\text{Me})_2(\mu\text{-P}'')]_2$  (**21-GaMe**)  $[\text{AlMe}_3(\text{dmap})]$ ; the latter products formed upon decomposition when warming to room temperature as a result of a methyl group transfer from Ga to Al with concomitant Al–P bond cleavage.<sup>58</sup> This rearrangement was monitored by VT  $^{31}\text{P}\{^1\text{H}\}$  NMR spectroscopy; from  $-50$  to  $-10$  °C there was only a single resonance corresponding to **22** ( $-262$  ppm), but above  $-10$  °C an additional signal appears at  $-219$  ppm due to the formation of **21-GaMe**.<sup>58</sup>****

In both 1992 and 1993, Wells reported the synthesis of three dimeric In(III)  $\text{P}''$  complexes,  $[\text{In}(\text{Me})(\text{R})(\mu\text{-P}'')]_2$  (**23-R**;  $\text{R} = \text{CH}_2\text{SiMe}_3, \text{CH}_2\text{}^t\text{Bu}$ , Fig. 6) and  $[\{\text{In}(\text{CH}_2\text{}^t\text{Bu})_2\}_2(\mu\text{-P}'')\{\mu\text{-PH}(\text{SiMe}_3)\}]$  (**24**, Fig. 6).<sup>59,60</sup> Complexes **23-R** exhibit puckered  $\text{In}_2\text{P}_2$  cores, with each In atom bound to two different alkyl groups, and mean In–P bond lengths of  $2.635(2)$  Å (**23-CH}\_2\text{SiMe}\_3**) and  $2.637(3)$  Å (**23-CH}\_2\text{}^t\text{Bu}**).<sup>59,60</sup> Complex **24** was isolated as a minor product from the reaction of  $[\text{In}(\text{CH}_2\text{}^t\text{Bu})_2(\text{Cl})]$  and  $\text{P}(\text{SiMe}_3)_3$ , and represented the first example of a dinuclear In phosphide complex in which the phosphorus centres were substituted asymmetrically; complex **24** exhibits a mean In–P bond length of  $2.650(3)$  Å and hydrocarbon solutions of this complex were found to be unstable at room temperature.<sup>60</sup>

There are numerous examples of dinuclear group 13 bridged  $\text{P}''$  complexes that contain either terminal or bridging halides as these are often used as precursors to functionalised complexes *via* salt metathesis reactions. For example,  $[\text{M}(\text{R})(\text{X})(\mu\text{-P}'')]_2$  (**25-MXR**;  $\text{M} = \text{Al}, \text{Ga}, \text{In}$ ;  $\text{X} = \text{Cl}, \text{Br}$ ;  $\text{R} = \text{Me}, \text{CH}_2\text{}^t\text{Bu}, \text{CH}_2\text{SiMe}_3, \eta^1\text{-Cp}^*$ ;  $\text{Cp}^* = \text{C}_5\text{Me}_5$ , Fig. 7) were synthesised from parent  $[\text{M}(\text{R})(\text{X})_2]$  with either  $\text{P}(\text{SiMe}_3)_3$  or  $\text{LiP}''$  between 1991–2004 by Schulz, Wells, and Theopold.<sup>50,59,61–63,126</sup> Each of these complexes has a  $\text{M}_2\text{P}_2$  core, with the four-coordinate group 13 metals having distorted tetrahedral geometries.<sup>50,59,61–63,126</sup>

Complex **25-AlBrCH}\_2\text{SiMe}\_3** exhibits a mean Al–P bond length of  $2.436(4)$  Å, whilst the Ga–P bond lengths of **25-GaXR** range from  $2.4106(10)$ – $2.424(3)$  Å, and the In–P distances of **25-InXR** vary from  $2.593(2)$ – $2.621(2)$  Å.<sup>50,59,61,63</sup> Wells reported several examples of group 13 metal  $\text{P}''$  complexes where a chloride has been incorporated into the central four membered ring,  $[\{\text{M}(\text{R})_2\}_2(\mu\text{-P}'')(\mu\text{-Cl})]$  (**26-MR**;  $\text{M} = \text{Ga}, \text{In}$ ;  $\text{R} = \text{Ph}, \text{CH}_2\text{}^t\text{Bu}, \text{CH}_2\text{SiMe}_3$ , Fig. 7).<sup>50,56,60,64</sup> For **26-GaCH}\_2\text{}^t\text{Bu}**, the central  $\text{Ga}_2\text{PX}$  core is planar, but **26-GaCH}\_2\text{SiMe}\_3** deviates from planarity by  $23^\circ$  and **26-GaCH}\_2\text{Ph}** deviates by  $6.4^\circ$  due to steric effects.<sup>50,64</sup>

In contrast, the In  $\text{P}''$  complexes **26-InCH}\_2\text{}^t\text{Bu}** and **26-InCH}\_2\text{SiMe}\_3** were found to be planar, due to the increased size of In vs. Ga allowing for the large steric bulk of both alkyl and silylphosphide ligands.<sup>56,60</sup> This increase in size from Ga to In



Fig. 6 Dinuclear group 13  $\text{P}''$  complexes  $[\text{M}(\text{R})_2(\mu\text{-P}'')]_2$  (**21-MR**;  $\text{M} = \text{Al}, \text{Ga}, \text{In}, \text{Tl}$ ;  $\text{R} = \text{Me}, \text{Et}, \text{Ph}, \text{}^t\text{Bu}, \text{CH}_2\text{Ph}, \text{CH}_2\text{}^i\text{Pr}, \text{CH}_2\text{}^t\text{Bu}, \text{CH}_2\text{SiMe}_3$ ),  $[\text{Al}(\text{Me})_2(\text{dmap})(\mu\text{-P}'')\text{Ga}(\text{Me})_3]$  (**22**),  $[\text{In}(\text{Me})(\text{R})(\mu\text{-P}'')]_2$  (**23-R**;  $\text{R} = \text{CH}_2\text{SiMe}_3, \text{CH}_2\text{}^t\text{Bu}$ ),  $[\{\text{In}(\text{CH}_2\text{}^t\text{Bu})_2\}_2(\mu\text{-P}'')\{\mu\text{-PH}(\text{SiMe}_3)\}]$  (**24**).





Fig. 7 Dimeric group 13 P'' complexes [M(R)(X)(μ-P'')] (25-MXR: M = Al, Ga, In; X = Cl, Br; R = Me, CH<sub>2</sub><sup>t</sup>Bu, CH<sub>2</sub>SiMe<sub>3</sub>, η<sup>1</sup>-Cp\*<sup>+</sup>; Cp\* = C<sub>5</sub>Me<sub>5</sub>), [(M(R))<sub>2</sub>(μ-P'')(μ-Cl)] (26-MR: M = Ga, In; R = Ph, CH<sub>2</sub><sup>t</sup>Bu, CH<sub>2</sub>SiMe<sub>3</sub>), [(M(R))<sub>2</sub>(μ-P'')(μ-As'')] (27-M: M = Al, R = Et, M = In, R = CH<sub>2</sub>SiMe<sub>3</sub>), [(M(R))<sub>2</sub>(μ-P'')(μ-As'')] (28-M: M = Al, R = Et, M = In, R = CH<sub>2</sub>SiMe<sub>3</sub>), [(Ga(Et))<sub>2</sub>(μ-P'')(μ-Sb'')] (29), [(Ga(X))<sub>2</sub>(μ-P'')(μ-E(SiMe<sub>3</sub>)<sub>2</sub>)] (30-ClP: X = Cl, Pn'' = P''; 30-BrP: X = Br, Pn'' = P''; 30-IP: X = I, Pn'' = P''; 30-IAs: X = I, Pn'' = As'').

is reflected in the respective mean M–P bond lengths; 26-GaR (2.391(4)–2.451(3) Å) and 26-InR (2.603(4)–2.622(5) Å).<sup>50,56,60,64</sup>

There are also examples of dinuclear group 13 P'' complexes that contain bridging {Pn(SiMe<sub>3</sub>)<sub>2</sub>} ligands; Pn = As (As'') or Sb (Sb''). In 1994 and 1995, Wells synthesised the respective Al and In P'' complexes [(M(R))<sub>2</sub>(μ-P'')(μ-As'')] (27-Al: M = Al, R = Et; 27-In: M = In, R = CH<sub>2</sub>SiMe<sub>3</sub>; Fig. 7) by the separate reactions of [M(R)<sub>2</sub>(P'')] with [M(R)<sub>2</sub>(As'')] (28-Al: M = Al, R = Et; 28-In: M = In, R = CH<sub>2</sub>SiMe<sub>3</sub>, Fig. 7),<sup>65,66</sup> whilst an analogous Ga complex, [(Ga(Et))<sub>2</sub>(μ-P'')(μ-Sb'')] (29, Fig. 7), was synthesised by Wells in 2000 *via* the combination of [Ga(Et)<sub>2</sub>(Cl)] with P(SiMe<sub>3</sub>)<sub>3</sub> and Sb(SiMe<sub>3</sub>)<sub>3</sub>; the As analogue of 29 was not structurally authenticated.<sup>48</sup> Complexes 27-M and 29 represent rare examples of two different group 15 elements bridging between Al, Ga or In; in the solid state planar central M<sub>2</sub>PE cores (M = Al, In, E = As; M = Ga, E = Sb) were observed in each case, with the metal centres displaying distorted tetrahedral geometries.<sup>48,65,66</sup> The mean M–P bond lengths increase with the size of the metal (27-Al: 2.497(1) Å; 29: 2.574(2) Å; 27-In: 2.691(3) Å).<sup>48,65,66</sup> Between 1993–1995, Wells reported several other examples of dimeric Ga(III) P'' halide complexes, namely [(Ga(X))<sub>2</sub>(μ-P'')(μ-Pn'')] (30-ClP: X = Cl, Pn'' = P''; 30-BrP: X = Br, Pn'' = P''; 30-IP: X =

I, Pn'' = P''; 30-IAs: X = I, Pn'' = As''); Fig. 7).<sup>67–69</sup> These complexes were isolated from the separate reactions of GaX<sub>3</sub> (X = Cl, Br, I) with either two equivalents of P(SiMe<sub>3</sub>)<sub>3</sub> (30-ClP, 30-BrP, 30-IP) or one equivalent each of P(SiMe<sub>3</sub>)<sub>3</sub> and As(SiMe<sub>3</sub>)<sub>3</sub> (30-IAs).<sup>67–69</sup> These complexes are the first examples of Ga pnictide dimers that contain only exocyclic halogen ligands, and in common with the previously discussed bridging pnictide complexes they all display a planar central Ga<sub>2</sub>PE core with the Ga atoms exhibiting distorted tetrahedral geometries. The mean Ga–P bond lengths range from 2.379(3)–2.398(4) Å, with the increase in size of the terminal halogen leading to longer bridging Ga–P bond distances.<sup>67–69</sup> The incorporation of the bridging As'' ligand in 30-IAs gives a longer Ga–P bond length (2.443(3) Å) than the mean Ga–P distance in 30-IP (2.398(4) Å).<sup>69</sup> The authors anticipated that these complexes would be viable starting materials for introducing further substitution at the Ga centres through dehalosilylation and salt elimination reactions.<sup>67–69</sup>

Although there are numerous reports of dimeric group 13 P'' complexes, there are just a handful of structurally characterised tri- or tetra-nuclear examples. The trinuclear complexes [M(H)<sub>2</sub>(μ-P'')] (M = Al, 31-Al; M = Ga, 31-Ga, Fig. 8), were



Fig. 8 Polynuclear 13 P'' complexes [M(H)<sub>2</sub>(μ-P'')] (31-M: M = Al, Ga), [(M(Me))<sub>2</sub>(μ-P'')(μ-P'')(μ-P'')] (32-M: M = Al, Ga), [(Ga(Me))<sub>2</sub>(μ-P'')(μ-P'')(μ-P'')] (33).



reported by Wells in 1997 and 1998, and  $[\{M(\text{Me})_2\}_3(\mu\text{-P}''\{\text{HP}(\text{SiMe}_3)_2\})_2]$  ( $M = \text{Al}$ , **32-Al**;  $M = \text{Ga}$ , **32-Ga**, Fig. 8), were reported by Robinson and Weidlein in 1994 and 1999, respectively.<sup>70–72</sup> These complexes were synthesised by the reaction of parent  $\text{H}_3\text{M-NMe}_3$  ( $M = \text{Al}$ ,  $\text{Ga}$ ) with  $\text{P}(\text{SiMe}_3)_3$  for **31-M**, or from parent  $[\text{M}(\text{Me})_2(\text{OME})]_3$  ( $M = \text{Al}$ ,  $\text{Ga}$ ) and  $\text{HP}''$  for **32-M**. Each complex features six-membered  $\text{M}_3\text{P}_3$  rings with three four-coordinate distorted tetrahedral  $\text{Al}/\text{Ga}$  atoms, with **31-M** exhibiting planar rings and **32-M** having twisted conformations in order to relieve steric congestion.<sup>70–72</sup> The mean  $\text{M-P}$  bond lengths increase from **31-M** to **32-M**: 2.398(4) Å (**31-Al**), 2.4287(13) Å (**32-Al**), 2.392(3) Å (**31-Ga**), and 2.415(2) Å (**32-Ga**).<sup>70–72</sup> The tetranuclear  $\text{Ga}(\text{III})$   $\text{P}''$  complex  $[\{\text{Ga}(\text{Me})_2(\mu\text{-P}'')\}_2\{\mu\text{-P}(\text{Me})\}_2]$  (**33**, Fig. 8), reported by Robinson in 1994, was prepared from the reaction of the Lewis acid–base adduct  $[\text{GaMe}_3\text{-PMe}_3]$  with  $\text{P}(\text{SiMe}_3)_3$ .<sup>73</sup> Complex **33** is bicyclic and features two fused  $\text{Ga}_2\text{P}_3$  five-membered rings, with distorted tetrahedral geometries of both  $\text{Ga}$  and  $\text{P}$  atoms, mean  $\text{Ga-P}$  bond lengths of 2.443(9) Å and a  $\text{P-P}$  single bond distance of 2.25(3) Å.<sup>73</sup> Remarkably, unlike the vast majority of  $\text{P}''$  complexes, **31-M**, **32-M** and **33** are all stable in air for a limited amount of time.<sup>70–73</sup>

**2.2.1.1 Heterometallic group 13 complexes.** There are a handful of structurally characterised examples of heterometallic mixed  $s$ - and  $p$ -block  $\text{P}''$  complexes, which are often used as precursors for complexes that possess bonds between group 13–15 elements.<sup>74</sup> The Li group 13 ‘ate’  $\text{P}''$  complexes,  $[\text{M}(\text{H})_2(\mu\text{-P}'')_2\text{Li}(\text{Et}_2\text{O})_2]$  (**34-M**;  $M = \text{Al}$ ,  $\text{Ga}$ ; Fig. 9) were synthesised by Wells in 1997 and 1998, respectively, *via* the reaction of  $\text{LiMH}_4$  with two equivalents of  $\text{P}(\text{SiMe}_3)_3$  ( $M = \text{Al}$ ,  $\text{Ga}$ ) in diethyl ether.<sup>74,75</sup> The  $\text{LiP}_2\text{M}$  ( $M = \text{Al}$ ,  $\text{Ga}$ ) cores of these complexes were confirmed to be planar by both NMR spectroscopy and single crystal XRD, and show mean  $\text{M-P}$  bond lengths of 2.4001(13) Å (**34-Al**) and 2.4122(12) Å (**34-Ga**).<sup>74,75</sup> Both the hydrogen and carbon atoms of the  $\text{SiMe}_3$  groups gave triplet resonances in the  $^1\text{H}$  and  $^{13}\text{C}\{^1\text{H}\}$  NMR spectra from second order coupling to two identical  $^{31}\text{P}$  nuclei, and signals at  $-282.0$  and  $-277.7$  ppm were seen in the  $^{31}\text{P}\{^1\text{H}\}$  NMR spectra of **34-Al** and **34-Ga**, respectively.<sup>74,75</sup> The  $\text{M-H}$  bonds were evidenced by stretching absorptions in IR spectra at  $1752\text{ cm}^{-1}$  (**34-Al**) and  $1838\text{ cm}^{-1}$  (**34-Ga**), together with broad resonances in  $^1\text{H}$  NMR spectra at 4.3 ppm (**34-Al**) and 4.7 ppm (**34-Ga**).<sup>74,75</sup> A structurally analogous  $\text{Al}$   $\text{P}''$  complex,  $[\text{Al}(\text{H})_2(\mu\text{-P}'')_2\text{Na}(\text{DME})_2]$  (**35**, Fig. 9), was synthesised by Hänisch in 2004 *via* the reaction of  $\text{NaAlH}_4$  with four equivalents of the primary phosphine,  $\text{H}_2\text{PSiMe}_3$  in DME.<sup>76</sup> Similarly to **34-M**, complex **35** has a planar



Fig. 9 Heterometallic group 13  $\text{P}''$  complexes  $[\text{M}(\text{H})_2(\mu\text{-P}'')_2\text{Li}(\text{Et}_2\text{O})_2]$  (**34-M**;  $M = \text{Al}$ ,  $\text{Ga}$ ),  $[\text{Al}(\text{H})_2(\mu\text{-P}'')_2\text{Na}(\text{DME})_2]$  (**35**).

four-membered  $\text{NaP}_2\text{Al}$  ring, mean  $\text{Al-P}$  bond lengths of 2.405(1) Å, triplet resonances in both its  $^1\text{H}$  and  $^{13}\text{C}\{^1\text{H}\}$  NMR spectra for the  $\text{SiMe}_3$  groups, and a single resonance at  $-283.0$  ppm in its  $^{31}\text{P}\{^1\text{H}\}$  NMR spectrum. Both symmetric and asymmetric stretches of the  $\text{Al-H}$  bonds were observed in the IR spectrum of **35** at 1762 and  $1728\text{ cm}^{-1}$ .<sup>76</sup>

**2.2.2 Homometallic group 14 complexes.** Group 14  $\text{P}''$  complexes are less common than group 13 examples, but have received particular interest as  $\sigma$ -donor ligands, and as frustrated Lewis pairs, due to their ability to mediate  $\text{E-H}$  bond activation.<sup>127–133</sup> The terphenyl  $\text{Ge}(\text{II})$   $\text{P}''$  complex  $[\text{Ge}(\text{P}'')(\text{Ar}^{\text{Mes}})]$ ,  $\text{Ar}^{\text{Mes}} = \text{C}_6\text{H}_3\text{Me}_2\text{-2,6}$  (**36**, Fig. 10), synthesised by Aldridge in 2016, features a two-coordinate  $\text{Ge}$  atom with a pyramidal phosphorus centre displaying a  $\text{C}_{\text{Ar}}\text{-Ge-P}$  angle of  $106.1(1)^\circ$  and a  $\text{Ge-P}$  bond length of 2.329(1) Å.<sup>77</sup> It was noted that this complex was unreactive towards  $\text{H}_2$  and  $\text{NH}_3$ , and as such is not well-suited for promoting small molecule activation reactions.<sup>77</sup> Complexes of the type  $[\text{M}(\text{DippNacnac})(\text{P}'')]$  ( $M = \text{Ge}$ , **37-Ge**;  $\text{Sn}$ , **37-Sn**;  $\text{Pb}$ , **37-Pb**; Fig. 10) have also been explored by several groups as  $\sigma$ -donor ligands.<sup>78–80</sup> These complexes exist as both *endo*- and *exo*-conformers in the solid state, with the *endo*-conformers exhibiting planar phosphorus centres (**37-Ge** and **37-Sn**) and the *exo*-conformers possessing pyramidal phosphorus centres.<sup>78–80</sup> Complexes **37-M** exhibit increasing  $\text{M-P}$  bond lengths with the size of the metal centre: **37-Ge**, 2.3912(8) Å; **37-Sn**, 2.5526(7) Å; **37-Pb**, 2.715(2) Å.<sup>78–80</sup> The  $^{31}\text{P}$  NMR spectra of the **37-M** triad respectively showed signals at  $-192.7$ ,  $-183.5$  ( $^1J_{\text{SnP}} = 2427\text{ Hz}$ ; assume mean coupling constants from 7.7% abundant  $^{117}\text{Sn}$  and 8.6% abundant  $^{119}\text{Sn}$  coupling constants, both  $I = \frac{1}{2}$ ) and  $-116.6$  ( $^1J_{\text{PbP}} = 1580\text{ Hz}$ , 22.1% abundant  $^{207}\text{Pb}$   $I = \frac{1}{2}$ ) ppm, with the  $\text{Sn}$  centres of **37-Sn** being planar and the  $\text{Pb}$  centres of **37-Pb** being pyramidal.<sup>78–80</sup> In 2017 Fulton reported that the addition of elemental  $\text{Se}$  to **37-Ge** gave the  $\text{Ge}(\text{IV})$  selenide complex,  $[\text{Ge}(\text{Se})(\text{DippNacnac})(\text{P}'')]$ , (**38**, Fig. 10), which displayed a  $\text{Ge-P}$  bond length (2.2976(7) Å) that is shorter than that of **37-Ge**.<sup>78,79,81</sup> A single resonance was observed in the  $^{31}\text{P}\{^1\text{H}\}$  NMR spectrum of **38** at  $-176.2$  ppm, with both selenium and silicon satellites ( $^1J_{\text{SeP}} = 52\text{ Hz}$ , 7.6% abundant  $^{77}\text{Se}$   $I = \frac{1}{2}$ , and  $^1J_{\text{PSe}} = 26\text{ Hz}$ ).<sup>81</sup> In 2010, Scheer reported the terphenyl  $\text{Ge}(\text{II})$  and  $\text{Sn}(\text{II})$   $\text{P}''$  complexes,  $[\text{M}(\text{Ph}^*)(\text{P}'')]$  ( $M = \text{Ge}$ , **39-Ge**;  $\text{Sn}$ , **39-Sn**;  $\text{Ph}^* = \text{C}_6\text{H}_3\text{Trip-2,6}$ ,  $\text{Trip} = \text{C}_6\text{H}_2\text{Pr-2,4,6}$ ; Fig. 10), which were synthesised by a redox reaction of  $[\text{M}(\text{Ph}^*)(\text{X})_3]$  ( $M = \text{Ge}$ ,  $\text{X} = \text{Cl}$ ;  $M = \text{Sn}$ ,  $\text{X} = \text{Cl}$ ,  $\text{Br}$ ) with  $\text{LiP}''$ .<sup>82</sup> Both complexes display a *trans*-bent geometry, as commonly seen for germylene/stannylenes complexes, with the phosphorus atom exhibiting a distorted trigonal pyramidal geometry and bond lengths of 2.291(4) Å (**39-Ge**) and 2.527(1) Å (**39-Sn**), signifying that the  $\text{P}$  lone pair is not donating to the group 14 centre.<sup>82</sup>

In 1993, Buhro reported the dimeric  $\text{Pb}(\text{II})$   $\text{P}''$  complex,  $[\text{Pb}(\text{P}'')(\mu\text{-P}'')_2]$  (**40**, Fig. 11) which was synthesised by the protonolysis reaction of  $[\text{Pb}(\text{N}'' )_2]$  with two equivalents of  $\text{HP}''$ ; the  $\text{Sn}(\text{II})$  analogue of **40** was synthesised by the same methods but was not structurally authenticated.<sup>83</sup> Complex **40** was reported to decompose to  $\text{Pb}$  and the diphosphine  $(\text{Me}_3\text{Si})_2\text{P-P}(\text{SiMe}_3)_2$  in refluxing benzene. Complex **40** exhibits a puckered central  $\text{M}_2\text{P}_2$  core and crystallises with the trigonal pyramidal terminal





**Fig. 10** Mononuclear group 14  $P''$  complexes  $[Ge(P'')(Ar^{Mes})]$  ( $Ar^{Mes} = C_6H_3Mes_{2-2,6}$ ) (**36**),  $[M(DiPPNacnac)(P'')]$  (**37-M**; M = Ge, Sn, Pb,  $DiPPNacnac = CH((CH_3)_2CN(Dipp))_2$ ;  $Dipp = C_6H_3^iPr_2-2,6$ ),  $[Ge(Se)(DiPPNacnac)(P'')]$  (**38**),  $[M(Ph^*)(P'')]$  (**39-M**; M = Ge, Sn,  $Ph^* = C_6H_3Trip-2,6$ ,  $Trip = C_6H_2^iPr_3-2,4,6$ ).



**Fig. 11** Complex  $[Pb(P'')(μ-P'')]_2$  (**40**).

$P''$  ligands mutually *syn*-, with mean Pb–P bond lengths of 2.77(1) Å (terminal) and 2.70(1) Å (bridging).<sup>83</sup> An equilibrium between the *syn*- and *anti*-conformers of **40** is seen in solution, with the  $^{31}P\{^1H\}$  NMR spectrum containing two sets of resonances in a 2 : 1 and 1 : 1 ratio (–217.3 ppm,  $^1J_{PbP} = 1264$  Hz, *anti*-terminal; –218.0 ppm,  $^1J_{PbP} = 1183$  Hz, *syn*-terminal; –281.4 ppm,  $^1J_{PbP} = 1598$  Hz, *syn*-bridging; –302.4 ppm,  $^1J_{PbP} = 1658$  Hz, *anti*-bridging).<sup>83</sup> Further assignment of these

conformers was enabled by  $^1H$  NMR spectroscopy as the  $SiMe_3$  groups on the bridging  $P''$  are inequivalent for the *syn*- and equivalent for the *anti*-conformer, thus three resonances are associated with the former and two for the latter conformer.<sup>83</sup> It was deduced from the intensities of the resonances at room temperature that solutions of **40** contained nearly equal amounts of *syn*- and *anti*-conformers.<sup>83</sup>

**2.2.2.1 Heterometallic group 14 complexes.** In 1994 Westershausen reported the synthesis of two Sn(II)  $P''$  complexes,  $[Sn(P'')(μ-P'')_2Ca(μ-P'')_2Ca(N'')]$  (**41**, Fig. 12) and  $[{Sn(P'')(μ_2-P'')}_2Ba]$  (**42**, Fig. 12), by the separate reactions of the respective M(II) group 2 bis-amide complexes,  $[M(N'')_2]$  (M = Ca, Ba) with  $[Sn(N'')_2]$  and five or six equivalents of  $HP''$  in toluene.<sup>34,84</sup> Both complexes exhibit central  $\{M(μ-P'')_4\}$  (M = Ca, Ba) cores, with **41** featuring one terminal  $CaN''$  and one terminal  $SnP''$  fragment,



**Fig. 12** Heterometallic group 14  $P''$  complexes  $[Sn(P'')(μ-P'')_2Ca(μ-P'')_2Ca(N'')]$  (**41**),  $[{Sn(P'')(μ_2-P'')}_2Ba]$  (**42**),  $[M(THF)_x\{Sn(μ_2-P'')_2(μ_3-P'')_2\}]$  (**43-M**; M = Ca, x = 2, M = Ba, x = 3, 4,  $P'' = \{P(SiMe_3)_2\}$ ).



and **42** having two terminal SnP'' moieties.<sup>34,84</sup> The central s-block atom of both complexes have distorted tetrahedral geometries, the terminal Ca atom in **41** is planar, and the terminal Sn atoms of both **41** and **42** are pyramidal. The bridging Sn-P bond distances in **41** are longer than that of the terminal Sn-P bond by approximately 0.09 Å, whereas the corresponding difference in **42** is *ca.* 0.11 Å.<sup>34,84</sup> The cluster complexes, [M(THF)<sub>x</sub>{Sn(μ<sub>2</sub>-P'')(μ<sub>3</sub>-P'')}<sub>2</sub>] (M = Ca, *x* = 2, **43-Ca**; M = Ba, *x* = 3, 4, **43-Ba**, P' = {P(SiMe<sub>3</sub>)<sub>2</sub>}; Fig. 12), were synthesized by Westerhausen in 1994 and 1995 using analogous methods but with THF as the reaction solvent. Both **43-M** feature two three-coordinate Sn atoms, each bound to one bridging P'' and two dianionic bridging silylphosphadiides, {P(SiMe<sub>3</sub>)<sub>2</sub>}, to the respective s-block metal; **43-Ca** and **43-Ba** have mean Sn-P bond lengths of 2.642(1) Å and 2.885(9) Å, respectively.<sup>33,34</sup>

### 2.3 d-Block P'' complexes

d-Transition metal P'' complexes are known to be versatile reagents for the formation of alkylidene-phosphides,<sup>134,135</sup> diphosphenyls<sup>136</sup> and cyclophosphines,<sup>137</sup> bulky supporting ligands are required in order to prevent oligomerisation to give monomeric complexes, and for homoleptic complexes oligomers tend to form. As expected and in common with the vast majority of other ligands,<sup>22,23</sup> first row d-transition metal P'' chemistry is more developed than for the heavier second and third row metals. It is noteworthy that no group 5 or 9 P'' complexes were found in the CSD search described above, and there are also no structurally authenticated Tc, Re, Ru or Os P'' complexes,<sup>9</sup> thus there are multiple opportunities here to expand d-block P'' chemistry rapidly.

**2.3.1 Group 3 complexes.** In 1998, Westerhausen synthesised the dimeric Y complex, [Y(P'')<sub>2</sub>]<sub>2</sub>(μ-P'')<sub>2</sub> (**44**, Fig. 13) with the crystal structure being reported in 2002, by the protonolysis reaction of [Y{CH(SiMe<sub>3</sub>)<sub>2</sub>}]<sub>3</sub> with three equivalents of HP''.<sup>86,87</sup> Both Y(III) cations in **44** are bound by two terminal and two bridging P'' in distorted tetrahedral geometries; the mean Y-P<sub>terminal</sub> bond lengths (2.678(4) Å) are shorter than the mean Y-P<sub>bridging</sub> bond lengths (2.849(4) Å) as expected.<sup>86,87</sup> These metrical parameters may be compared to those of a structurally analogous dimeric Y(III) amide complex, [Y{N(SiHMe<sub>2</sub>)<sub>2</sub>}]<sub>2</sub>(μ-N(SiHMe<sub>2</sub>)<sub>2</sub>)<sub>2</sub>, which has mean Y-N<sub>terminal</sub> bond lengths of 2.249(4) Å and mean Y-N<sub>bridging</sub> bond lengths of 2.479(4) Å.<sup>138</sup> The dimeric structure of **44** was shown to be maintained in solution by <sup>31</sup>P{<sup>1</sup>H} NMR spectroscopy, with a doublet of pentets observed at -107.8 ppm (<sup>1</sup>J<sub>YP</sub> = 56.7 Hz, <sup>2</sup>J<sub>PP</sub> = 5.0 Hz)

for bridging P'' and a doublet of triplets -104.8 ppm (<sup>1</sup>J<sub>YP</sub> = 122.4 Hz, <sup>2</sup>J<sub>PP</sub> = 5.0 Hz) for terminal P''.<sup>86,87</sup>

In 2015, as part of a study by Mézailles into how the anionic ligand *trans*- to a bound carbon atom in Sc(III) methanediide complexes influences complex geometry, the P'' complex, [Sc{C(PPh<sub>2</sub>)<sub>2</sub>}(P'')(py)<sub>2</sub>] (**45**, Fig. 13) was synthesised.<sup>88</sup> Complex **45** has a Sc-P bond length of 2.618(14) Å; it was found that the methanediide carbon atom is planar for all Sc(III) complexes explored in this study save the N'' analogue, [Sc{C(PPh<sub>2</sub>)<sub>2</sub>}(N'')(THF)], which has one THF bound to Sc rather than two pyridines. Density Functional Theory (DFT) calculations indicated that the coordinating solvent made little impact on the planarity of the methanediide. Natural Bond Orbital (NBO) analysis revealed that in **45** the donation from both the C and P atoms are stronger than the corresponding C and N atoms in [Sc{C(PPh<sub>2</sub>)<sub>2</sub>}(N'')(THF)], which leads to the difference in methanediide geometry.<sup>88</sup> In 2024, Mills and co-workers reported the synthesis of [La{P(SiMe<sub>3</sub>)<sub>2</sub>}<sub>3</sub>(THF)<sub>2</sub>] (**46-La**, Fig. 13) by a salt metathesis reaction of the respective [La(i<sub>3</sub>(THF))<sub>4</sub>] starting material with three equivalents of KP''.<sup>89</sup> Single crystal XRD showed that **46-La** to exhibit a distorted trigonal bipyramidal geometry, with three equatorial P'' and two axial THF molecules, with the O-La-O angle showing a small deviation from linearity: 175.14(8)° and a mean La-P bond length of 2.886(2) Å. Two of the equatorial P'' in **46-La** show pyramidal geometries about the phosphorus atom, whereas the third P'' exhibits a planar geometry. Solid state <sup>31</sup>P MAS NMR spectroscopy of **46-La** revealed two components in their spectra in a 2:1 ratio, with the major component assigned to pyramidal and the minor component to planar P environments. This is in contrast to the solution <sup>31</sup>P{<sup>1</sup>H} NMR spectrum of **46-La**, which exhibits one broad resonance due to dynamic processes and quadrupolar broadening due to the 99.9% abundant <sup>77</sup>La *I* = 7/2 nuclei.

**2.3.2 Group 4 complexes.** The first d-block P'' complex to the best of our knowledge, [Ti(NMe<sub>2</sub>)<sub>3</sub>(P'')], was reported in 1970 by Bürger, but this was not structurally authenticated.<sup>18</sup> In 1991, the paramagnetic bis-Cp Ti(III) P'' complex, [Ti(Cp)<sub>2</sub>(P'')] (**47**, Fig. 14), was synthesised by Fenske by the salt metathesis reaction of [Ti(Cp)(Cl)] with LiP''.<sup>90</sup> This complex features a planar phosphorus atom as a result of the π-interaction between Ti and P, and has a short Ti-P bond length of 2.468(1) Å, indicating some double bond character, although the sum of covalent radii of a Ti-P single bond (2.47 Å) matches that reported for **47**.<sup>139</sup> The <sup>31</sup>P{<sup>1</sup>H} NMR spectrum of **47**



Fig. 13 Group 3 P'' complexes [Y(P'')<sub>2</sub>]<sub>2</sub>(μ-P'')<sub>2</sub> (**44**), [Sc{C(PPh<sub>2</sub>)<sub>2</sub>}(P'')(py)<sub>2</sub>] (**45**), [La{P(SiMe<sub>3</sub>)<sub>2</sub>}<sub>3</sub>(THF)<sub>2</sub>] (**46-La**).





Fig. 14 Group 4  $P''$  complexes  $[Ti(Cp)_2(P'')]$  (**47**),  $[Hf(Cp)_2(P'')]_2$  (**48**),  $[Zr(Cp)_2(Cl)(P'')]$  (**49**),  $[Zr(Cp)_2(Me)(P'')]$  (**50**),  $[Zr(C_5H_4Me)_2(P'')]_2$  (**51**),  $[Zr(Cp)_2(Cl)(P(SiMe_3)_2C(H)C(Ph))]$  (**52**),  $[Zr(Cp)_2(R)(P(SiMe_3)_2C(H)C(Ph))]$  (**53-R**; R = Me,  $n$ Bu, C $\equiv$ CPh).

exhibits a single broad resonance at 251 ppm, and an isotropic signal was observed in the EPR spectrum with a  $g$ -value of 1.9779.<sup>90</sup> In 1987 Weber reported a series of heavy group 4 M(IV)  $P''$  complexes with two ancillary Cp ligands by salt metathesis methodologies.<sup>91</sup> The reaction of  $[Zr(Cp)_2(Cl)_2]$  with one or two equivalents of  $LiP''$  exclusively gave  $[Zr(Cp)_2(P'')]_2$  (**48-Zr**), whilst for the analogous reactions of the Hf analogue  $[Hf(Cp)_2(Cl)_2]$  with  $LiP''$  rapid substitution of the first chloride gave  $[Hf(Cp)_2(Cl)(P'')]$  (**49-Hf**), and the second substitution proceeds more slowly to give  $[Hf(Cp)_2(P'')]_2$  (**48-Hf**, Fig. 14). Complexes **48-Zr** and **49-Hf** were not structurally characterised in this study, but in 1988 Lappert reported the synthesis of  $[Zr(Cp)_2(Cl)(P'')]$  (**49-Zr**, Fig. 14) from the reaction of  $[Zr(Cp)_2(Cl)_2]$  with  $[Li(\mu-P'')(THF)_2]_2$ .<sup>91,92</sup> The reaction of  $[Zr(Cp)_2(Me)(Cl)]$  with  $LiP''$  gave the monophosphido complex,  $[Zr(Cp)_2(Me)(P'')]$  (**50**, Fig. 14).<sup>91</sup> All of these structurally characterised group 4 M(IV)  $P''$  complexes exhibit *pseudo*-tetrahedral geometries and additionally show some interesting structural features. For example, in **48-Hf** it was seen that one  $P''$  has a planar geometry with a Hf–P bond length of 2.553(1) Å, whilst the other  $P''$  is pyramidal with a longer Hf–P distance of 2.654(1) Å.<sup>91</sup> The shorter Hf–P bond length in the planar  $P''$  arises from its orthogonal lone pair being involved in  $\pi$ -bonding with Hf.<sup>91</sup> It was also noted in the  $^{31}P\{^1H\}$  NMR spectrum of **48-Hf** that this structural feature is maintained in solution, with two resonances seen at –69.90 and –98.83 ppm for the pyramidal and planar  $P''$ , respectively.<sup>91</sup> For complex **49-Zr**, a relatively short Zr–P bond length (2.547(6) Å) is seen for the near-planar  $P''$  ( $\Sigma$  angles = 344.4°), though this is slightly pyramidalized compared to the planar  $P''$  in **48-Hf** ( $\Sigma$  angles = 360°).<sup>92</sup> In 1992, Hey-Hawkins synthesised a derivative of **48-Zr**,  $[Zr(C_5H_4Me)_2(P'')]_2$  (**50**, Fig. 14), by the salt metathesis reaction of  $[Zr(C_5H_4Me)_2(Cl)_2]$  with  $LiP''$ .<sup>93</sup> Complex **50** shows a *pseudo*-tetrahedral geometry, where both phosphorus atoms are planar and the Zr–P bond lengths are longer than the corresponding distances in **49-Zr** (2.600(2) and 2.634(2) Å).<sup>93</sup> VT  $^{31}P\{^1H\}$  NMR spectra of **51** indicated that the planarity at phosphorus was retained in solution, with two equivalent P atoms evidenced by a single resonance at –75.3 ppm at 298 K, which shifts to –86.2 ppm at 173 K.<sup>93</sup>

Between 1992–1995, Hey-Hawkins synthesised zirconocene alkenyl complexes that feature  $P''$  incorporated into metallocycles.<sup>94,95</sup> The insertion reaction of  $[Zr(Cp)_2(P'')(Cl)]$  with phenylacetylene in toluene at reflux gave  $[Zr(Cp)_2(Cl)\{P(SiMe_3)_2C(H)C(Ph)\}]$  (**52**, Fig. 14), which can undergo subsequent salt metathesis reactions with RLi (R = Me,  $n$ Bu, C $\equiv$ CPh) to yield  $[Zr(Cp)_2(R)\{P(SiMe_3)_2C(H)C(Ph)\}]$  (R = Me,  $n$ Bu, C $\equiv$ CPh, **53-R**; Fig. 14); **53- $n$ Bu** was not structurally characterised.<sup>94,95</sup>  $^{31}P\{^1H\}$  NMR spectroscopy showed modest changes in chemical shifts when Cl is substituted for an R group (R = Cl, –179.5 ppm; Me, –179.3 ppm;  $n$ Bu, –176.6 ppm; C $\equiv$ CPh, –189.9 ppm).<sup>94,95</sup> For all of these complexes a doublet is observed in the  $^1H$  NMR spectra for the alkenyl proton, with  $^1J_{PH}$  coupling constants ranging from 8–12 Hz.<sup>94,95</sup> The alkenyl moieties exhibit the *Z*-configuration in the solid state, and the Zr–P distances (**52**: 2.855(4) Å; **53-Me**: 2.915(3) Å; **53-C $\equiv$ CPh**: 2.774(3) Å) indicate that Zr–P interactions persist in solution for these complexes.<sup>94,95</sup> The phosphorus atoms exhibit pyramidal geometries in **52** and **53-R**, due to steric effects causing wide Si–P–Si angles.<sup>94,95</sup>

**2.3.3 Group 6 complexes.** Dimeric mid-transition metal complexes have been investigated for their applications as molecular switches and molecular magnets due to the spin crossover and exchange coupling properties that these metals possess.<sup>83,85,96</sup> In 2013, Layfield and Scheer reported the synthesis of the dimeric Cr(II) Cp  $P''$  complex  $[Cr(Cp)(\mu-P'')]_2$  (**54-Cr**, Fig. 15) by the salt metathesis reaction of  $LiP''$  with  $[Cr(Cp)_2]$ .<sup>96</sup> In the solid state each Cr atom of **54-Cr** exhibits a *pseudo*-trigonal geometry, being bound to two phosphorus atoms and one  $\eta^5$ -Cp ligand; the Cr–P bond lengths of 2.3814(9) and 2.3864(9) Å fall within the known range for cyclic Cr phosphido complexes.<sup>9,85,96</sup> From magnetic susceptibility measurements, the high-temperature limit ( $\chi_M T$ ) of **54-Cr** was found to be 0.60 cm<sup>3</sup> K mol<sup>–1</sup>, which is lower than the spin-only value of 6.0 cm<sup>3</sup> K mol<sup>–1</sup> for two non-interacting high-spin Cr(II) ions and is indicative of strong antiferromagnetic coupling.<sup>96</sup> This is also observed in variable temperature susceptibility measurements, where there is a steady decrease of  $\chi_M T$  towards zero upon cooling to 2 K; for the diamagnetic spin ground state where  $S = 0$  the exchange coupling constant was calculated to be  $J = -166$  cm<sup>–1</sup>.<sup>96</sup>





Fig. 15 Group 6 P'' complexes  $[\text{Cr}(\text{Cp})(\mu\text{-P}'')]_2$  (**54-Cr**),  $[\text{Cr}(\text{Dippnacnac})(\text{P}'')]$  (**37-Cr**;  $\text{Dippnacnac} = \text{CH}(\text{CH}_3)_2\text{CN}(\text{Dipp})_2$ ;  $\text{Dipp} = \text{C}_6\text{H}_3\text{Pr}_2\text{-2,6}$ ),  $[\text{Li}(12\text{-crown-4})][\{\text{Mo}(\text{Cp})(\text{CO})_2\}_2(\mu\text{-P}'')]$  (**55**),  $[\text{1,2-W}_2(\text{P}'')(\text{NMe}_2)_4]$  (**56**),  $[\text{1,2-W}_2(\text{PCy}_2)(\text{P}'')(\text{NMe}_2)_4]$  (**57**).

In 2024, Weigend and Hänisch synthesised a Cr(II) P'' complex,  $[\text{Cr}(\text{Dippnacnac})(\text{P}'')]$  (**37-Cr**, Fig. 15) by the salt metathesis reaction of the Cr(II) precursor  $[\text{Cr}(\text{Dippnacnac})(\text{Cl})]$  with one equivalent of  $\text{LiP}''$  in  $\text{Et}_2\text{O}$  at  $-78^\circ\text{C}$ .<sup>140</sup> Complex **37-Cr** features a nearly planar P atom ( $\Sigma$  angles =  $356^\circ$ ), which allows its lone electron pair to act as a  $\pi$ -donor into a vacant Cr 3d orbital, leading to a short Cr–P bond length of  $2.3641(3) \text{ \AA}$ .<sup>140</sup>

In 2021, Scheer synthesised the dinuclear Mo(I) P'' complex,  $[\text{Li}(12\text{-crown-4})_2][\{\text{Mo}(\text{Cp})(\text{CO})_2\}_2(\mu\text{-P}'')]$  (**55**, Fig. 15) by the addition of  $\text{LiP}''$  to  $[\text{Mo}(\text{Cp})(\text{CO})_2]_2$  in THF; the addition of one equivalent of either  $\text{AsCl}_3$  or  $\text{SbCl}_3$  to **55** gives the mixed-dipnictogen complexes  $[\{\text{Mo}(\text{Cp})(\text{CO})_2\}_2(\mu\text{-PE})]$  ( $\text{E} = \text{As}, \text{Sb}$ ), releasing two equivalents of  $\text{SiMe}_3\text{Cl}$  in the process.<sup>97</sup> The Mo–P distances in **55** are  $2.4304(6) \text{ \AA}$ , which is between that expected for a single or double bond, and the phosphorus atom exhibits a distorted tetrahedral geometry. A long Mo–Mo bond length in **55** ( $3.1890 \text{ \AA}$ ) is observed compared to the starting material  $[\text{Mo}(\text{Cp})(\text{CO})_2]_2$  ( $2.4477(12) \text{ \AA}$ ), due to the degradation of the  $\text{Mo}\equiv\text{Mo}$  triple bond to a Mo–Mo single bond.<sup>97</sup>

In 1992, Chisholm synthesised two examples of dinuclear W(III) complexes that exhibit tungsten-tungsten triple bonds and incorporate both P'' and amide ligands.<sup>98</sup> The salt metathesis reaction of  $[\text{1,2-W}_2(\text{Cl})_2(\text{NMe}_2)_4]$  with two equivalents of  $\text{LiP}''$  gave  $[\text{1,2-W}_2(\text{P}'')_2(\text{NMe}_2)_4]$  (**56**), whilst combination of the mono-substituted precursor  $[\text{1,2-W}_2(\text{Cl})(\text{P}'')(\text{NMe}_2)_4]$  with  $\text{LiPCy}_2$  gave  $[\text{1,2-W}_2(\text{PCy}_2)(\text{P}'')(\text{NMe}_2)_4]$  (**57**, Fig. 15).<sup>98</sup> Complex **56** exhibits an *anti*-conformation in the solid-state across the triple bond, with trigonal planar nitrogen atoms and pyramidal

phosphorus atoms.<sup>98</sup> The  $\text{W}\equiv\text{W}$  triple bond lengths of  $2.2989(9) \text{ \AA}$  (**56**) and  $2.3016(10) \text{ \AA}$  (**57**) are similar to the mean values for  $\text{W}\equiv\text{W}$  triple bonds deposited on the CCDC ( $2.32(5) \text{ \AA}$ )<sup>9</sup> and the W–P distances are statistically equivalent at  $2.423(3) \text{ \AA}$  (**56**) and  $2.425(4) \text{ \AA}$  (**57**).<sup>98</sup> These complexes were characterised by both  $^1\text{H}$  and  $^{31}\text{P}\{^1\text{H}\}$  NMR spectroscopy, with the latter showing a dynamic equilibrium in solution between the *anti*- and *gauche*-conformations; the  $^{31}\text{P}$  chemical shifts are  $-106.7 \text{ ppm}$  (**56**) and  $-106.9 \text{ ppm}$  (**57**) for the former and  $-88.3 \text{ ppm}$  (**56**) and  $-96.3 \text{ ppm}$  (**57**) for the latter.<sup>98</sup> It was noted that upon lowering the temperature from 295 to 193 K, the  $^{31}\text{P}$  chemical shifts move to a higher field and a change of line shape is observed, which was attributed to inversion of the phosphorus centres.<sup>98</sup>

**2.3.4 Group 7 complexes.** In 2012 Layfield and Scheer reported the synthesis of the Mn(II) P'' complex  $[\{\text{Mn}(\text{Cp})(\mu\text{-P}'')\}_2]$  (**54-Mn**, Fig. 16) by the salt metathesis reaction of  $\text{LiP}''$  with  $[\text{Mn}(\text{Cp})_2]$ .<sup>85</sup> The solid state structure of **54-Mn** is analogous to the Cr(II) complex **54-Cr** discussed in the previous section, with the Mn–P bond lengths of **54-Mn** being longer at  $2.5075(5)$  and  $2.5123(5) \text{ \AA}$ .<sup>85</sup> Variable temperature magnetic measurements of **54-Mn** showed that  $\chi_{\text{M}}T$  decreases from  $5.71 \text{ cm}^3 \text{ K mol}^{-1}$  at 300 K to  $0.027 \text{ cm}^3 \text{ K mol}^{-1}$  at 2 K, with the former value being approximately 75% of the expected value ( $8.75 \text{ cm}^3 \text{ K mol}^{-1}$  at 300 K) for two  $S = 5/2$  Mn(II) ions. The small value of  $\chi_{\text{M}}T$  at 2 K was rationalised as the dimer having a diamagnetic ground state, likely due to antiferromagnetic exchange.<sup>85</sup>

In 1993, Buhro and co-workers reported the synthesis of the dinuclear Mn(II) P'' complex  $[\{\text{Mn}(\text{P}'')(\mu\text{-P}'')\}_2]\{\text{Mn}(\text{P}'')(\text{THF})\}$



Fig. 16 Group 7 P'' complexes  $[\text{Mn}(\text{Cp})(\mu\text{-P}'')]_2$  (**54-Mn**),  $[\{\text{Mn}(\text{P}'')(\mu\text{-P}'')\}_2]\{\text{Mn}(\text{P}'')(\text{THF})\}$  (**58**),  $[\text{Mn}(\text{Dippnacnac})(\text{P}'')(\text{THF})]$  (**37-Mn**;  $\text{Dippnacnac} = \text{CH}(\text{CH}_3)_2\text{CN}(\text{Dipp})_2$ ;  $\text{Dipp} = \text{C}_6\text{H}_3\text{Pr}_2\text{-2,6}$ ).



(**58**, Fig. 16) from the protonolysis reaction of  $[\text{Mn}(\text{N}'')_2(\text{THF})]$  with two equivalents of  $\text{HP}''$  in hexane.<sup>83</sup> In the solid state **58** consists of one trigonal planar three-coordinate Mn centre with two bridging and one terminal  $\text{P}''$ , and one four-coordinate distorted tetrahedral Mn centre with two bridging and one terminal  $\text{P}''$ , together with a coordinated THF. It was noted that the bound THF in **58** can be removed when this complex is exposed to a dynamic vacuum.<sup>83</sup> The central  $\text{Mn}_2\text{P}_2$  core in **58** is nearly planar and the phosphorus atoms of the terminal  $\text{P}''$  adopt pyramidal geometries (mean  $\Sigma$  angles  $334.2(3)^\circ$ ).<sup>83</sup> The bridging Mn–P distances range from 2.493(2)–2.565(2) Å, whereas the terminal Mn–P bond lengths are 2.417(3) Å for the three-coordinate Mn and 2.461(2) Å for the four-coordinate Mn centre.<sup>83</sup> Due to the paramagnetic nature of complex **58**, NMR spectroscopic data were intractable, but Evans method magnetic susceptibility measurements gave a  $\mu_{\text{eff}}$  value of  $3.33\mu_{\text{B}}$  per Mn atom; this is lower than the spin-only value for a high-spin system with  $S = 5/2$  ( $5.92\mu_{\text{B}}$ ) and was ascribed to the Mn centres being antiferromagnetically coupled.<sup>83</sup>

In 2024 Weigend and Hänisch synthesised the Mn(II)  $\text{P}''$  complex  $[\text{Mn}(\text{P}''\text{P}''\text{Nacnac})(\text{P}'')(\text{THF})]$  (**37-Mn**, Fig. 16), in an analogous manner to **37-Cr** discussed in the previous section.<sup>140</sup> The structure of **37-Mn** is similar to **37-Cr** but THF is additionally bound to Mn in the solid state; the Mn–P bond length of **37-Mn** (2.461(1) Å) is longer than the corresponding Cr–P distance seen in **37-Cr** and is similar to the mean Mn–P bond lengths found in **54-M** (2.5075(5) Å) and **58** (2.461(2) Å).<sup>83,85</sup>

**2.3.5 Group 8 complexes.** Although the syntheses of a number of group 8  $\text{P}''$  complexes have been reported, remarkably only one structurally authenticated Fe(II)  $\text{P}''$  complex has been reported to date.<sup>99</sup> The first group 8  $\text{P}''$  complex to be reported,  $[\text{Fe}(\text{Cp})(\text{CO})_2(\text{P}'')]$ , was prepared by Schäfer in 1980 by the salt metathesis reactions of  $[\text{Fe}(\text{Cp})(\text{CO})_2(\text{X})]$  ( $\text{X} = \text{Cl}, \text{Br}$ ) with  $\text{LiP}''$ .<sup>99a</sup> A reactivity study was reported in the same publication. The separate reactions of  $[\text{Fe}(\text{Cp})(\text{CO})_2(\text{P}'')]$  with  $[\text{Fe}_2(\text{CO})_9]$  and  $[\text{Ni}(\text{CO})_4]$  gave the  $\text{P}''$ -bridged complexes  $\{[\text{Fe}(\text{Cp})(\text{CO})_2(\mu\text{-P}'')\{\text{Fe}(\text{CO})_4\}]\}$  and  $\{[\text{Fe}(\text{Cp})(\text{CO})_2(\mu\text{-P}'')\{\text{Ni}(\text{CO})_3\}]\}$ , respectively. Photolysis of the former complex was proposed to give  $\{[\text{Fe}(\text{Cp})(\text{CO})_2(\mu\text{-P}'')(\mu\text{-CO})\{\text{Fe}(\text{CO})_3\}]\}$ , whilst the treatment of either complex with stoichiometric methanol gave respective  $\text{PH}_2$ -bridged products by cleavage of P–Si bonds. Between 1985 and 1987 Weber reported the synthesis of the group 8 M(II)  $\text{P}''$  complexes  $[\text{M}(\text{Cp}^*)(\text{CO})_2(\text{P}'')]$  ( $\text{M} = \text{Fe}, \text{Ru}, \text{Os}$ ) and  $[\text{Fe}(\text{Cp}^*)(\text{CO})(\text{PPh}_3)(\text{P}'')]$  by salt metathesis reactions,<sup>99b–e</sup> reactions of these complexes with  $\text{Mes}^*\text{PCl}_2$  gave M(II) diphosphene complexes by P–Si bond cleavage and loss of  $\text{ClSiMe}_3$ .<sup>99e</sup>

In 1992, Weber reported the synthesis of a series of Cp-substituted Fe(II)  $\text{P}''$  complexes  $[\text{Fe}(\text{C}_5\text{R}_5)(\text{CO})_2(\text{P}'')]$  (**59-R**;  $\text{C}_5\text{R}_5 = \text{C}_5\text{EtMe}_4$ , **59-EtMe<sub>4</sub>**;  $\text{C}_5^n\text{BuMe}_4$ , **59-<sup>n</sup>BuMe<sub>4</sub>**;  $\text{C}_5\text{H}_3^t\text{Bu}_2$ -1,3, **59-1,3-<sup>t</sup>Bu<sub>2</sub>H<sub>3</sub>**, Fig. 17) by the separate salt metathesis reactions of the parent monobromide complexes,  $[\text{Fe}(\text{C}_5\text{R}_5)(\text{CO})_2(\text{Br})]$  ( $\text{R}_5 = \text{EtMe}_4$ , <sup>n</sup>BuMe<sub>4</sub>, 1,3-<sup>t</sup>Bu<sub>2</sub>H<sub>3</sub>) with  $\text{LiP}''$ .<sup>99f</sup> The syntheses of three analogous Ru(II)  $\text{P}''$  complexes  $[\text{Ru}(\text{C}_5\text{R}_5)(\text{CO})_2(\text{P}'')]$  ( $\text{R}_5 = 1,2,4\text{-}^i\text{Pr}_3\text{H}_2$ , 1,3-<sup>t</sup>Bu<sub>2</sub>H<sub>3</sub>, 1,3-(SiMe<sub>3</sub>)<sub>2</sub>H<sub>3</sub>) were reported by Weber in 1994 but again these complexes were not structurally authenticated.<sup>99g</sup> We note that only **59-EtMe<sub>4</sub>** was structurally




Fig. 17 Group 8  $\text{P}''$  complexes  $[\text{Fe}(\text{C}_5\text{R}_5)(\text{CO})_2(\text{P}'')]$  (**59-R**;  $\text{C}_5\text{R}_5 = \text{C}_5\text{EtMe}_4$ , **59-EtMe<sub>4</sub>**;  $\text{C}_5^n\text{BuMe}_4$ , **59-<sup>n</sup>BuMe<sub>4</sub>**;  $\text{C}_5\text{H}_3^t\text{Bu}_2$ -1,3, **59-1,3-<sup>t</sup>Bu<sub>2</sub>H<sub>3</sub>**).

characterised by single crystal XRD in the former paper,<sup>99f</sup> thus comparative discussions of **59-R** focus on NMR and IR spectroscopy. The Fe–P distance in **59-EtMe<sub>4</sub>** is 2.359(3) Å and in the solid state the phosphorus atom has a pyramidal geometry ( $329.2(2)^\circ$ ), with the trimethylsilyl groups eclipsed with respect to the CO ligands. The  $^{31}\text{P}\{^1\text{H}\}$  NMR spectra of **59-R** exhibit singlets at  $-219.1$  ppm (**59-EtMe<sub>4</sub>**),  $-218.6$  ppm (**59-<sup>n</sup>BuMe<sub>4</sub>**) and  $-266.4$  ppm (**59-1,3-<sup>t</sup>Bu<sub>2</sub>H<sub>3</sub>**), whilst the IR spectra contain CO stretching modes at  $1986$  and  $1939$   $\text{cm}^{-1}$  (**59-EtMe<sub>4</sub>**),  $1988$  and  $1935$   $\text{cm}^{-1}$  (**59-<sup>n</sup>BuMe<sub>4</sub>**), and  $1995$  and  $1944$   $\text{cm}^{-1}$  (**59-1,3-<sup>t</sup>Bu<sub>2</sub>H<sub>3</sub>**). It was deduced that a reduced donor strength for the 1,3-bis-*tert*-butyl-substituted  $\text{C}_5\text{R}_5$  ligand gives rise to weaker M–C  $\pi$ -backbonding and therefore stronger CO bonding in **59-1,3-<sup>t</sup>Bu<sub>2</sub>H<sub>3</sub>**. Finally, Weber reported that the separate reactions of **59-R** or related Fe(II) and Ru(II)  $\text{P}''$  complexes with  $\text{Mes}^*\text{PCl}_2$  gave M(II) diphosphene complexes,<sup>99f,g</sup> analogously to the reactions described above.<sup>99e</sup>

**2.3.6 Group 10 complexes.** In 1987 Schäfer reported the synthesis of a family of monomeric Ni(II)  $\text{P}''$  complexes,  $[\text{Ni}(\kappa^2\text{-drpe})(\text{P}'')]$  (**60-R**;  $\text{drpe} = \text{R}_2\text{PCH}_2\text{CH}_2\text{PR}_2$ ,  $\text{R} = \text{Et}, \text{Cy}, \text{Ph}$ , Fig. 18), by the separate salt metathesis reactions of two equivalents of  $\text{LiP}''$  with  $[\text{Ni}(\kappa^2\text{-drpe})(\text{Cl})_2]$  ( $\text{R} = \text{Et}$ , **60-NiEt**;  $\text{Cy}$ , **60-NiCy**;  $\text{Ph}$ , **60-NiPh**); **60-NiCy** was the only complex in this series to be structurally characterised.<sup>100</sup> The intermediate mono-substituted



Fig. 18 Group 10  $\text{P}''$  complexes  $[\text{Ni}(\kappa^2\text{-drpe})(\text{P}'')]$  (**60-R**;  $\text{drpe} = \text{R}_2\text{PCH}_2\text{CH}_2\text{PR}_2$ ,  $\text{R} = \text{Et}, \text{Cy}, \text{Ph}$ ),  $[\text{M}(\text{X})(\text{R}_2\text{-bimz})_2(\text{P}'')]$  (**61-MR**,  $\text{M} = \text{Ni}$ ,  $\text{X} = \text{Br}$ ,  $\text{R} = ^n\text{Bu}$ , **61-Ni<sup>n</sup>Bu**;  $\text{M} = \text{Ni}$ ,  $\text{X} = \text{I}$ ,  $\text{R} = ^i\text{Pr}$ , **61-Ni<sup>i</sup>Pr**;  $\text{M} = \text{Pd}$ ,  $\text{X} = \text{I}$ ,  $\text{R} = ^n\text{Bu}$ , **61-Pd<sup>n</sup>Bu**;  $^i\text{Pr}$ , **61-Pd<sup>i</sup>Pr**;  $\text{bimz} = \text{benzimidazole-2-ylidene}$ ),  $[\text{Ni}(\text{PMe}_3)(\mu\text{-P}'')_2]$  (**62**),  $\{[\text{M}(\text{PPh}_3)_2(\text{P}'')]\{\text{Si}[\text{N}(^t\text{Bu})_2]\text{CPh}\}\}$  (**63-M**;  $\text{M} = \text{Pd}, \text{Pt}$ ).



complexes,  $[\text{Ni}(\kappa^2\text{-drpe})(\text{P}'')(\text{Cl})]$ , were observed by *in situ* NMR spectroscopy but could not be isolated.<sup>100</sup> The Ni atom in **60-NiCy** is three-coordinate and nearly planar, with the P'' phosphorus atom exhibiting a pyramidal geometry and a Ni–P bond length of 2.225(2) Å, which is slightly longer than the corresponding Ni–P distances of the coordinated diphosphine (2.192(2) and 2.202(2) Å).<sup>100</sup> In 2015 Corrigan synthesised the Ni(II) and Pd(II) complexes,  $[\text{M}(\text{X})(\text{R}_2\text{-bimy})_2(\text{P}'')]$  (**61-MR**, M = Ni, X = Br, R = <sup>n</sup>Bu, **61-Ni<sup>n</sup>Bu**; M = Ni, X = I, R = <sup>i</sup>Pr, **61-Ni<sup>i</sup>Pr**; M = Pd, X = I, R = <sup>n</sup>Bu, **61-Pd<sup>n</sup>Bu**; <sup>i</sup>Pr, **61-Pd<sup>i</sup>Pr**; bimy = benzimidazole-2-ylidene, Fig. 18) by the separate reactions of parent  $[\text{MX}_2(\text{R}_2\text{-bimy})_2]$  with LiP'' in THF at 60 °C; the high reaction temperature was required due to the poor solubility of the group 10 metal starting materials.<sup>101</sup> The <sup>i</sup>Pr-substituted complexes, **61-Ni<sup>i</sup>Pr** and **61-Pd<sup>i</sup>Pr**, were found to undergo P–Si bond cleavage in the presence of benzoyl chloride to give the respective complexes  $[\text{M}(\text{i})(\text{Pr}_2\text{-bimy})_2\text{-}\{\text{P}[\text{C}(\text{O})\text{Ph}]_2\}]$  (M = Ni, Pd).<sup>101</sup> Only the Pd analogues, **61-Pd<sup>n</sup>Bu** and **61-Pd<sup>i</sup>Pr**, were structurally characterised, and were both found to exhibit square-planar geometries at the d<sup>8</sup> Pd(II) centres as expected.<sup>101</sup> The NHC ligands in **61-Pd<sup>i</sup>Pr** cause a more distorted pyramidal geometry at the P atom (Si–P–Si: 98.53(6)°) compared to **61-Pd<sup>n</sup>Bu** (Si–P–Si: 100.68(9)°);<sup>101</sup> this difference was attributed to the larger steric requirements of <sup>i</sup>Pr vs. <sup>n</sup>Bu groups, as the latter are able to fold away to a greater extent to allow for the steric requirements of P''.<sup>101</sup> The Pd–P bond lengths vary slightly between the two complexes (2.3648(17) Å, **61-Pd<sup>n</sup>Bu**; 2.3442(12) Å, **61-Pd<sup>i</sup>Pr**) and there is a greater distortion from linearity of the C–Pd–C (171.41(13)°) and I–Pd–P (166.42(3)°) angles in **61-Pd<sup>i</sup>Pr** compared to **61-Pd<sup>n</sup>Bu** (176.6(2)° and 174.55(4)°, respectively).<sup>101</sup> It was noted that the two <sup>n</sup>Bu<sub>2</sub>-bimy ligands of **61-Pd<sup>n</sup>Bu** are twisted at 26.8° with respect to each other.<sup>101</sup>

In 1982, Schäfer reported the synthesis of the dimeric Ni(I) P'' complex  $[\text{Ni}(\text{PMe}_3)(\mu\text{-P}'')]_2$  (**62**, Fig. 18) by the decomposition of  $[\text{Ni}(\text{PMe}_3)_2(\text{P}'')]_2$  at room temperature to form both  $[\text{Ni}(\text{PMe}_3)_2\{\mu\text{-P}(\text{SiMe}_3)_2\}]$  and  $[\text{Ni}(\text{PMe}_3)(\mu\text{-P}'')]_2$ .<sup>102</sup> In the solid state **62** exhibits a planar central Ni<sub>2</sub>P<sub>2</sub> ring in which each Ni atom is three-coordinate and bound to two bridging P'' and one terminal trimethylphosphine.<sup>102</sup> The Ni–P bond lengths within the central core are 2.186(1) Å and the Ni–Ni separation is 2.381(1) Å.<sup>102</sup> In 2013 Inoue reported the synthesis of the dinuclear M(I) P'' complexes  $[\{\text{M}(\text{PPh}_3)_2(\text{P}'')\}\{\text{Si}[\text{N}(\text{tBu})_2\text{CPh}]\}]$  (**63-M**; M = Pd, Pt; Fig. 18); **63-Pt** was characterised by both NMR spectroscopy and single crystal XRD, whereas the formation of **63-Pd** was determined by NMR spectroscopy and high resolution mass (HRMS) spectrometry only.<sup>103</sup> The <sup>1</sup>H NMR spectra of **63-M** exhibit singlets (0.92 ppm, **63-Pt**; 0.86 ppm, **63-Pd**) for the <sup>t</sup>Bu groups, as well as a doublet (0.46 ppm, <sup>3</sup>J<sub>PH</sub> = 5.2 Hz, **63-Pt**; 0.54 ppm, <sup>3</sup>J<sub>PH</sub> = 5.2 Hz, **63-Pd**) for the trimethylsilyl groups. In the <sup>31</sup>P{<sup>1</sup>H} NMR spectrum of **63-Pt** a doublet at 56.0 ppm (<sup>1</sup>J<sub>PP</sub> = 5137 Hz and <sup>2</sup>J<sub>PP</sub> = 60.5 Hz, 33.8% abundant <sup>195</sup>Pt I =  $\frac{1}{2}$ ) and a triplet at –94.6 ppm (<sup>1</sup>J<sub>PP</sub> = 1633 Hz) are observed that respectively correspond to PPh<sub>3</sub> and P'', and these are seen at 41.2 and –163.4 ppm, respectively, for **63-Pd**.<sup>103</sup> The Pt–Pt bond length of **63-Pt** (2.6466(5) Å) falls within the reported range of Pt–Pt bonds (2.4059(6)–2.7183(10) Å),<sup>9</sup> whilst the P'' Pt–P bond lengths (2.360(2)

and 2.367(2) Å) are longer than the corresponding distances for the coordinated phosphines of 2.222(2) and 2.213(2) Å.<sup>103</sup>

**2.3.7 Group 11 complexes.** N-Heterocyclic carbene (NHC) ligands have been used as supporting ligands in d-transition metal P'' complexes in an effort to provide stable complexes that are suitable for long term storage.<sup>104</sup> Between 2015–2016 Corrigan reported the syntheses of three types of group 11 P'' complexes:  $[\text{Cu}(\text{Pr}_2\text{-bimy})_2(\text{P}'')]$  (**64**, <sup>i</sup>Pr<sub>2</sub>-bimy = 1,3-diisopropylbenzimidazole-2-ylidene) by the reaction of  $[\text{Cu}(\text{Pr}_2\text{-bimy})_2(\text{OAc})]$  with P(SiMe<sub>3</sub>)<sub>3</sub>,  $[\text{Au}(\text{Pr}_2\text{-bimy})(\text{P}'')]$  (**65**) by the reaction of  $[\text{Au}(\text{Pr}_2\text{-bimy})(\text{Cl})]$  with P(SiMe<sub>3</sub>)<sub>3</sub>, and  $[\text{M}(\text{IPr})(\text{P}'')]$  (**66-M**; M = Cu, Au, IPr = 1,3-bis(2,6-diisopropylphenyl)-2-ylidene) by the reaction of  $[\text{Cu}(\text{IPr})(\text{OAc})]$  or  $[\text{Au}(\text{IPr})(\text{Cl})]$  with P(SiMe<sub>3</sub>)<sub>3</sub> (Fig. 19); complex **65** was not structurally authenticated.<sup>104,105</sup> At the time of publication **64** and **66-M** provided some of the first examples of structurally authenticated terminal d-transition metal P'' complexes. The Cu centre in **64** exhibits a distorted trigonal planar geometry and the phosphorus atom is trigonal pyramidal; together with a long Cu–P bond length (2.291(7) Å); this demonstrates that the phosphorus lone pair does not contribute to the M–P bonding.<sup>104</sup> Complex **66-Cu** exhibits a shorter Cu–P bond length of 2.1913(15) Å and a near-linear geometry (C–Cu–P: 173.39(17)°), with the phosphorus atom being distorted trigonal pyramidal;<sup>104</sup> similarly, **66-Au** has a near-linear geometry (C–Au–P: 175.74(4)°) and a distorted trigonal pyramidal phosphorus atom, though the Au–P bond length in **66-Au** is longer (2.3174(10) Å).<sup>105</sup> The <sup>1</sup>H NMR spectrum of **64** confirms that the solid state structure is maintained in solution as there is a 2 : 1 ratio of <sup>i</sup>Pr<sub>2</sub>-bimy : P'', together with one resonance in the <sup>31</sup>P{<sup>1</sup>H} NMR spectrum at –261 ppm; the analogous signal for **65** is at δ<sub>p</sub> = –235 ppm.<sup>104,105</sup> The <sup>1</sup>H NMR spectrum of **66-Cu** showed that the solid-state structure was retained in solution due to the 1 : 1 ratio of IPr : P'' ligands. The <sup>31</sup>P{<sup>1</sup>H} NMR spectra of **66-M** each show one resonance, at –268 ppm for the Cu analogue and –235.7 ppm for the Au congener.<sup>104,105</sup> In 2023 Liu reported the synthesis of  $[\text{Au}(\text{CAAC}^{\text{MeEt}})(\text{P}'')]$  (**67**; CAAC<sup>MeEt</sup> = dimethyl-diethyl-cyclic(alkyl)(amino)carbene, Fig. 19) by the reaction of  $[\text{Au}(\text{CAAC}^{\text{MeEt}})\text{-}\{\text{P}(\text{Ph})_2\}]$  with KP''.<sup>106</sup> This synthetic route is remarkable as it involves Au–P bond cleavage occurring, despite the strong covalency between Au and P due to relativistic effects which result in Au having the highest electronegativity of any d-transition metal (Pauling χ = 2.54). As with **66-M**, **67** exhibits a near-linear geometry (C–Au–P: 174.45(5)°) with a distorted trigonal pyramidal geometry about the phosphorus atom and an Au–P bond length of 2.3253(5) Å. Complex **67** has one resonance in its <sup>31</sup>P{<sup>1</sup>H} NMR spectrum at –233.1 ppm.<sup>106</sup>

Group 11 metal P'' cluster complexes have been synthesised in order to explore their potential as precursors for semiconductor materials. The group 11 hexameric cyclic complexes  $[\text{M}(\mu\text{-P}'')]_6$  (**68-M**; M = Cu, Ag; Fig. 19) were synthesised in 2015 by Corrigan by the separate reactions of parent polymeric  $[\text{M}(\text{S}^t\text{Bu})_n]$  with equimolar P(SiMe<sub>3</sub>)<sub>3</sub> *via* the cleavage of P–Si and M–S bonds and the formation of <sup>t</sup>BuSSiMe<sub>3</sub> and **68-M**.<sup>104</sup> In the solid state the P atoms of **68-M** form the corners of a planar hexagonal structure, and the two-coordinate metal





Fig. 19 Group 11  $P''$  complexes  $[\text{Cu}(\text{iPr}_2\text{-bimy})_2(\text{P}'')]$  **64** ( $\text{iPr}_2\text{-bimy}$  = 1,3-di-isopropylbenzimidazole-2-ylidene),  $[\text{Au}(\text{iPr}_2\text{-bimy})(\text{P}'')]$  (**65**);  $[\text{M}(\text{iPr})(\text{P}'')]$  (**66-M**;  $\text{M} = \text{Cu}, \text{Au}$ ),  $\text{iPr}$  = (1,3-bis(2,6-diisopropylphenyl)-2-ylidene),  $[\text{Au}(\text{CAAC}^{\text{MeEt}})(\text{P}'')]$  (**67**,  $\text{CAAC}^{\text{MeEt}}$  = dimethyl-diethyl-cyclic(alkyl)(amino)carbene),  $[\text{M}(\mu\text{-P}'')]_6$  (**68-M**;  $\text{M} = \text{Cu}, \text{Ag}$ ).

centres form the sides.<sup>104</sup> The M–P bond lengths range from 2.2043(16)–2.2094(15) Å for **68-Cu** and 2.393(1)–2.411(2) Å for **68-Ag**, and the P–M–P angles deviate slightly from linearity in both complexes ( $\sim 178^\circ$ , **68-Cu**;  $\sim 177^\circ$ , **68-Ag**).<sup>104</sup> The phosphorus atoms in **68-M** exhibit distorted tetrahedral geometries such that the trimethylsilyl groups are equally arranged above and below the plane of the ring. It was noted that the voids above and below the centre of the ring are occupied by THF lattice solvent molecules, which is believed to be important for the formation of the cyclic structures, and this sensitivity is in line with the loss of solvent leading to discolouration of solutions of **68-M**.<sup>104</sup> NMR spectroscopy showed that the symmetrical structures are retained in solution for both complexes,

with only one resonance in each  $^1\text{H}$  NMR spectrum at 0.50 ppm (**68-Cu**) and 0.48 ppm (**68-Ag**), and each  $^{31}\text{P}\{^1\text{H}\}$  NMR spectrum at  $-149$  ppm (**68-Cu**) and  $-236$  ppm (**68-Ag**).<sup>104</sup>

**2.3.8 Group 12 complexes.** In 2024, Weigend and Hänisch synthesised the monomeric  $\text{Zn}(\text{II})$  complex  $[\text{Zn}(\text{DippNacnac})(\text{P}'')]$  (**37-Zn**, Fig. 20) in an analogous manner to that discussed earlier for **37-M** ( $\text{M} = \text{Cr}, \text{Mn}$ ).<sup>140</sup> The Zn centre of **37-Zn** has a distorted trigonal pyramidal geometry with a Zn–P bond length of 2.2728(3) Å, which compares well to other literature compounds such as  $[\text{Zn}(\text{P}'')(\mu\text{-P}'')]_2$  (mean Zn–P<sub>terminal</sub>: 2.295(1) Å).<sup>83,140</sup> The  $^{31}\text{P}\{^1\text{H}\}$  NMR spectrum of **37-Zn** has one resonance at  $-288.85$  ppm, with satellites observed due to coupling to  $^{29}\text{Si}$  nuclei ( $J_{\text{P-Si}} = 28.5$  Hz).<sup>140</sup> In 1993 Buhro




Fig. 20 Monomeric and dimeric group 12  $P''$  complexes  $[\text{Zn}(\text{DippNacnac})(\text{P}'')]$  (**37-Zn**;  $\text{DippNacnac}$  =  $\text{CH}((\text{CH}_3)_2\text{CN}(\text{Dipp}))_2$ ;  $\text{Dipp}$  =  $\text{C}_6\text{H}_3\text{iPr}_2\text{-2,6}$ ),  $[\text{Hg}(\text{P}'')]_2$  (**69**),  $[\text{M}(\text{P}'')(\mu\text{-P}'')]_2$  (**70-M**,  $\text{M} = \text{Zn}, \text{Cd}$ ),  $[\text{Zn}(\text{R})(\mu\text{-P}'')]_2$  (**71-R**,  $\text{R} = \text{tBu}, \text{iPr}, \text{CH}_2\text{SiMe}_3$ ),  $[\text{M}(\text{P}'')\text{Pr}_3(\text{X})(\mu\text{-P}'')]_2$  (**72-Zn**:  $\text{M} = \text{Zn}$ ,  $\text{X} = \text{Cl}$ ; **72-Cd**:  $\text{M} = \text{Cd}$ ,  $\text{X} = \text{Br}$ ).



synthesised a monomeric Hg(II) P'' complex [Hg(P'')] (69, Fig. 20) by the reaction of [Hg(N'')] with HP''.<sup>83</sup> Complex 69 displays a near-linear geometry in the solid state (P–Hg–P = 175.8°), and the phosphorus atoms have pyramidal geometries.<sup>83</sup> The monomers of 69 are oriented in the lattice such that long-range dimers are formed with two bridging and two terminal P''. However, the resultant bridging Hg...P interactions are only slightly shorter (3.246(1) Å) than the sum of the van der Waals radii of Hg and P (3.35 Å). This interaction can be seen as a weak dative bond, which leads to the Hg–P bond lengths being nearly equal to one another (2.402(1) and 2.410(1) Å).<sup>83</sup> The NMR spectra of 69 indicate that a monomeric form dominates in solution, with one resonance seen in both <sup>1</sup>H (0.36 ppm) and <sup>31</sup>P{<sup>1</sup>H} (–162.0 ppm) NMR spectra.<sup>83</sup>

Many dimeric group 12 complexes have been reported with bridging P'', with varying terminal ligands such as alkyls, aryls, halides, phosphines and phosphides. In 1993 Buhro synthesised the homoleptic M(II) P'' complexes [M(P'')(μ-P'')] (70-M, M = Zn, Cd, Fig. 20) by the protonolysis reaction of [M(N'')] (M = Zn, Cd) with HP''.<sup>83</sup> Single crystal XRD studies of 70-M showed that each metal in the dimer has a distorted trigonal planar geometry and is bound by two bridging and one terminal P'', with the terminal phosphorus atom being pyramidal; the distorted rectangular M<sub>2</sub>P<sub>2</sub> cores in 70-M have bridging M–P–M angles which are closer to 90° rather than the usual 120°. The M–P<sub>bridging</sub> (2.419(1) and 2.421(1) Å for 70-Zn; 2.575(1) and 2.612(1) Å for 70-Cd) and M–P<sub>terminal</sub> (2.295(1) Å, 70-Zn; 2.459(1) Å, 70-Cd) bond lengths also lie within the expected range for bridging and terminal Zn–P and Cd–P bonds, respectively (CSD: Zn–P (2.212(3)–2.9159(7) Å), Cd–P (2.2379(8)–2.800(6) Å)).<sup>9,83</sup> Both complexes exhibit fluxional properties in solution, with low temperature NMR data (230 K) consistent with the presence of the dimer, as evidenced by two resonances in the <sup>31</sup>P{<sup>1</sup>H} NMR spectra for bridging and terminal phosphides (–183.0 ppm, P<sub>bridging</sub>, –237.3 ppm P<sub>terminal</sub>, 70-Zn; –180.1 ppm, P<sub>bridging</sub>, –229.5 ppm, P<sub>terminal</sub>, 70-Cd).<sup>83</sup> Upon conducting high temperature NMR experiments, line broadening and coalescence of these two resonances is seen for both complexes, which was assigned to bridging-to-terminal site exchange with a calculated barrier of ΔG<sub>360</sub><sup>‡</sup> = 14.3(2) kcal mol<sup>–1</sup> (70-Zn) and ΔG<sub>321</sub><sup>‡</sup> = 12.7(6) kcal mol<sup>–1</sup> (70-Cd).<sup>83</sup> The mechanism of this exchange was proposed to be dissociative where the dimers dissociate into the respective monomers and

reassociate, or that one of the M–P<sub>bridging</sub> bonds is broken, the remaining M–P<sub>bridging</sub> bond rotates, and then the cleaved M–P<sub>bridging</sub> bond reforms to give the exchanged dimer.<sup>83</sup>

The synthesis of the dimeric Zn(II) P'' complexes [Zn(R)(μ-P'')] (71-R; R = <sup>t</sup>Bu, <sup>i</sup>Pr, CH<sub>2</sub>SiMe<sub>3</sub>; Fig. 20) was reported in 1995 by Westerhausen by the reaction of parent ZnR<sub>2</sub> (R = <sup>t</sup>Bu, <sup>i</sup>Pr, CH<sub>2</sub>SiMe<sub>3</sub>) with HP''; however 71-<sup>t</sup>Bu was not structurally characterised.<sup>107</sup> Complexes 71-<sup>i</sup>Pr and 71-CH<sub>2</sub>SiMe<sub>3</sub> are dimers that consist of two trigonal planar Zn centres with bridging P'' and terminal alkyl groups, with nearly planar Zn<sub>2</sub>P<sub>2</sub> cores. The Zn–P bond lengths are 2.405(3) and 2.416(3) Å for 71-<sup>i</sup>Pr and 2.394(1) and 2.436(1) Å for 71-CH<sub>2</sub>SiMe<sub>3</sub>.<sup>107</sup> <sup>31</sup>P{<sup>1</sup>H} NMR spectroscopy showed one resonance for each complex, at –216.3 ppm (71-<sup>i</sup>Pr), –205.8 ppm (71-CH<sub>2</sub>SiMe<sub>3</sub>) and –216.7 ppm (71-<sup>t</sup>Bu).<sup>107</sup> Dimeric group 12 metal P'' complexes [M(P''Pr<sub>3</sub>)(X)(μ-P'')] (72-Zn: M = Zn, X = Cl; 72-Cd: M = Cd, X = Br; Fig. 20) featuring both a terminal phosphine and halide were synthesised in 1996 by Fenske via the separate reactions of MX<sub>2</sub> (M = Zn, X = Cl; M = Cd, X = Br) with P(SiMe<sub>3</sub>)<sub>3</sub> in the presence of P''Pr<sub>3</sub>.<sup>108</sup> Both complexes exhibit distorted tetrahedral metal centres in the solid state, with the central M<sub>2</sub>P<sub>2</sub> rings showing planar geometries with typical M–P<sub>bridging</sub> (2.426(1) and 2.447(1) Å for 72-Zn; 2.571(1) and 2.609(1) Å for 72-Cd) and M–P<sub>terminal</sub> (2.408(1) and 2.430(1) Å for 72-Zn; 2.584(1) and 2.571(1) Å for 72-Cd) bond distances.<sup>108</sup>

In 1996, Fenske reported the synthesis of a Zn(II) P'' cluster complex, [(Zn(Cl)(MeCN)(μ-P''))<sub>2</sub>{Zn(μ-Cl)}<sub>2</sub>] (73, Fig. 21), by the reaction of ZnCl<sub>2</sub> with equimolar P(SiMe<sub>3</sub>)<sub>3</sub> in MeCN.<sup>108</sup> Single crystal XRD revealed that the four Zn and four P atoms form a Zn<sub>4</sub>P<sub>4</sub> eight membered ring, with two chlorides bridging two of the Zn centres to give a Zn<sub>2</sub>Cl<sub>2</sub> core. The two Zn atoms that are not involved in the central four-membered ring are distorted tetrahedral and coordinated by two bridging P'' (mean Zn–P: 2.405(2) Å), a terminal chloride and MeCN, whereas the coordination spheres of the other two Zn atoms are completed by bridging P'', with a shorter mean Zn–P bond length (2.343(2) Å) giving much more strongly distorted tetrahedra (P–Zn–P: 153.88(4)°). Solutions of 73 decompose upon warming to room temperature, precluding the collection of NMR spectroscopic data that could be confidently assigned.<sup>108</sup> In the same report a Cd(II) cluster complex, [N''Bu<sub>4</sub>]<sub>2</sub>[Cd<sub>4</sub>(I)<sub>8</sub>(P'')] (74, Fig. 21), was formed from the reaction of [Cd(I)<sub>2</sub>{P(SiMe<sub>3</sub>)<sub>3</sub>}<sub>2</sub>] with N''Bu<sub>4</sub>I, with ISiMe<sub>3</sub> eliminated as a byproduct.<sup>108</sup> Complex 74 consists of four C atoms with four bridging iodides and two bridging P''




Fig. 21 Group 12 P'' cluster complexes [(Zn(Cl)(MeCN)(μ-P''))<sub>2</sub>{Zn(μ-Cl)}<sub>2</sub>] (73), [N''Bu<sub>4</sub>]<sub>2</sub>[Cd<sub>4</sub>(I)<sub>8</sub>(P'')] (74), [Zn(R)(P'')] (75-R; R = Me, Et, <sup>i</sup>Pr, <sup>n</sup>Bu).



to yield an adamantane-like skeleton; each distorted tetrahedral Cd atom also bears one terminal iodide.<sup>108</sup> The mean Cd–P bond distance is 2.507(7) Å, which is shorter than those found for other Cd–P<sub>bridging</sub> bonds in **70-Cd** (2.612(1) Å) and **72-Cd** (2.609(1) Å).<sup>83,108</sup> The trimeric Zn(II) P'' complexes [Zn(R)(P'')]<sub>3</sub> (**75-R**; R = Me, Et, <sup>1</sup>Pr, <sup>n</sup>Bu, Fig. 21) were reported in 1994 by Westerhausen, *via* the addition of ZnR<sub>2</sub> to HP''. Single crystal XRD showed that these complexes contain six-membered Zn<sub>3</sub>P<sub>3</sub> rings in twisted boat conformations, with the three Zn atoms having near trigonal planar geometries with terminal alkyl groups, and the phosphorus atoms showing distorted tetrahedral geometries; the solid-state structure of **75-Et** was not reported.<sup>107</sup> In solution this series of complexes were found to exhibit equilibria with the dimeric complexes discussed earlier in this section (**71-R**), with the larger alkyl groups favouring the formation of dimers.<sup>107</sup> For complex **75-Me**, two of the phosphorus atoms are situated 0.525 Å and 0.857 Å above and below the plane, respectively, whereas in **75-<sup>n</sup>Bu** these two phosphorus atoms are located more symmetrically at ±0.640 Å.<sup>107</sup> The mean Zn–P bond lengths of 2.390(3) Å (**75-Me**), 2.408(6) Å (**75-<sup>1</sup>Pr**) and 2.388(4) Å (**75-<sup>n</sup>Bu**) fall within the expected range for bridging Zn–P bonds (2.245(4)–2.8049(7) Å).<sup>9,107</sup> Chemical shifts in the <sup>31</sup>P{<sup>1</sup>H} NMR spectra for the trimeric complexes of –246.8 ppm (**75-Me**), –246.9 ppm (**75-Et**), –243.6 ppm (**75-<sup>1</sup>Pr**), –246.0 ppm (**75-<sup>n</sup>Bu**) are shifted to lower chemical shifts than the related dimers, which are seen at –216.3 ppm (**71-<sup>1</sup>Pr**) and –216.7 ppm (**71-<sup>t</sup>Bu**).<sup>107</sup>

**2.3.9 Heterometallic complexes.** To date, all examples of heterometallic d-block P'' complexes are also bound by carbon monoxide. From 2001–2002, Schulz reported the separate reactions of [Al(Me)<sub>2</sub>(P'')(dmap)] (**17-Al**, see Section 2.2.1) with the d-transition metal carbonyl complexes [Cr(NMe<sub>3</sub>)(CO)<sub>5</sub>], [Fe<sub>3</sub>(CO)<sub>12</sub>] and [Ni(CO)<sub>4</sub>] to give the respective M(0) P''-bridged complexes, [M(CO)<sub>n</sub>(μ-P'')<sub>2</sub>{Al(dmap)(Me)<sub>2</sub>}] (M = Cr, n = 5, **76-Cr**; M = Fe, n = 4, **76-Fe**; M = Ni, n = 3, **76-Ni**, Fig. 22).<sup>39,109</sup> Complexes **76-M** exhibit staggered conformations, with N–Al–P–M torsion angles of 36.9(1)° for **76-Cr** and 34.1(1)° for **76-Fe**; **76-Ni** shows a range of N–Al–P–Ni torsion angles from 21.60(13)–88.05(13)° as there are six independent molecules in the unit cell.<sup>39,109</sup> The M–P bond lengths are 2.528(1) Å (**76-Cr**), 2.377(1) Å (**76-Fe**) and 2.315(2) Å (**76-Ni**), with the former two being the longest reported examples of such bonds in

P–M(CO)<sub>n</sub> moieties for these metals at the time of publication.<sup>39,109</sup> The long Al–P bonds and short Al–N bonds in **76-M**, combined with their long P–M(CO)<sub>n</sub> bonds, indicate that there is some weak M → P π-backbonding within these systems.<sup>39,109</sup> Dimeric heterobimetallic P'' complexes were separately reported by Hey-Hawkins and Eisen in order to study the cooperative reactivity between an early and late d-transition metal for their potential in the heterolytic bond cleavage of polar substrates.<sup>111</sup> In 1995–1996, the separate reactions of the Zr(IV) complex, [Zr(Cp)<sub>2</sub>(P'')<sub>2</sub>] (**77-M**, Fig. 22), with [Ni(CO)<sub>4</sub>] or [Mo(CO)<sub>4</sub>(NBD)] (NBD = norbornadiene) gave [Zr(Cp)<sub>2</sub>(μ-P'')<sub>2</sub>M(CO)<sub>n</sub>] (M = Mo, n = 4, **77-Mo**; M = Ni, n = 2, **77-Ni**),<sup>110,111</sup> a similar complex [Zr(C<sub>5</sub>H<sub>4</sub>Me)<sub>2</sub>(μ-P'')<sub>2</sub>Cr(CO)<sub>n</sub>] (**77-Cr**) has also been synthesised *via* analogous methods from [Cr(CO)<sub>4</sub>(NBD)] and **51**.<sup>110</sup> The central ZrP<sub>2</sub>M rings in these complexes are slightly puckered, with equivalent Zr–P bond lengths of 2.654(4) and 2.657(4) Å (**77-Cr**), 2.6711(9) Å and 2.6585(7) Å (**77-Mo**) and 2.655(1), 2.652(1) Å (**77-Ni**) as well as equivalent M–P bond lengths of 2.513(4) and 2.502(4) Å (**77-Cr**), 2.6263(7) and 2.6311(10) Å (**77-Mo**) and 2.264(1) and 2.266(1) Å (**77-Ni**).<sup>110,111</sup> There are no Zr–M bonds in these complexes, with Zr...M distances of 3.414(3) Å (**77-Cr**), 3.459(3) Å (**77-Mo**) and 3.038(1) Å (**77-Ni**).<sup>110,111</sup> The <sup>31</sup>P{<sup>1</sup>H} NMR spectra each exhibit a single resonance at –38.4 ppm (**77-Cr**), –57.9 ppm (**77-Mo**), –42.1 ppm (**77-Ni**), which are all at higher frequencies than the starting material (–71.1 ppm, **48-Zr**).<sup>110,111</sup> Complexes **77-Ni** and **77-Mo** did not demonstrate any catalytic activity towards the polymerisation of ethylene. However, when these complexes were activated by a Lewis acid such as methylaluminoxane (MAO), polymerisation of ethylene occurred and was found to outperform cationic group 4 metallocene systems.<sup>110,111</sup>

## 2.4 f-Block P'' complexes

**2.4.1 Lanthanides.** Rabe reported the first Ln P'' complexes, [Ln(P'')<sub>3</sub>(THF)<sub>2</sub>] (**46-Ln**, Fig. 23) (Ln = Nd, Tm) in 1995, which at the time were the first structurally characterised Ln tris-phosphide complexes.<sup>112,113</sup> Later, in 2024, Mills and co-workers reported further examples in this series for Ln = La, Ce, Pr and Sm.<sup>89</sup> All complexes were prepared by salt metathesis reactions of the respective [Ln(III)<sub>3</sub>(THF)<sub>3,5</sub>] starting material with three equivalents of KP''. Single crystal XRD showed that **46-Ln** have distorted trigonal bipyramidal geometries, with three equatorial P'' and two axial THF molecules, with the O–Ln–O angles showing a small deviation from linearity: 175.6(6) (**46-Ce**), 175.58(13) (**46-Pr**), 175.68(9) (**46-Nd**), 175.36(13) (**46-Sm**) and 172.06(14)° (**46-Tm**).<sup>89,112,113</sup> Two of the equatorial P'' in **46-Ln** show pyramidal geometries about the phosphorus atom, whereas the third P'' exhibits a planar geometry.<sup>89,112,113</sup> The mean Ln–P bond lengths (2.849(3) (**46-Ce**), 2.837(3) (**46-Pr**), 2.818(2) (**46-Nd**), 2.789(3) (**46-Sm**) and 2.707(16) (**46-Tm**) Å) can be compared to the mean Ln–N distances in the Ln(III) N'' complexes [Ln(N'')<sub>3</sub>] (Ln = Nd, 2.244(16); Yb, 2.149(12) Å);<sup>141–143</sup> the structure of [Tm(N'')<sub>3</sub>] has not been reported, but the six-coordinate ionic radius of Yb(III) (0.868 Å) is similar to that of Tm(III) (0.880 Å).<sup>123</sup> The differences in solvation between **46-Ln** and [Ln(N'')<sub>3</sub>] can be attributed to the Ln–P bonds in the former being longer than the Ln–N bonds in the latter leading to a



Fig. 22 Heterometallic P'' complexes [M(CO)<sub>n</sub>(μ-P'')<sub>2</sub>{Al(dmap)(Me)<sub>2</sub>}] (**76-M**; M = Cr, n = 5, **76-Cr**; M = Fe, n = 4, **76-Fe**; M = Ni, n = 3, **76-Ni**), [Zr(Cp)<sub>2</sub>(μ-P'')<sub>2</sub>M(CO)<sub>n</sub>] (**77-M**; M = Mo, n = 4, **77-Mo**; M = Ni, n = 2, **77-Ni**).



Fig. 23 Ln P'' complexes [Ln(P'')<sub>3</sub>(THF)<sub>2</sub>] (**46-Ln**; Ln = Ce, Pr, Nd, Sm, Tm), [Sm(P'')(μ-P'')<sub>3</sub>Sm(THF)<sub>3</sub>] (**78**), [(Ln(P'')<sub>3</sub>(THF))<sub>2</sub>(μ-I)K<sub>3</sub>(THF)] (**79-Ln**; Ln = Sm, Eu), [KYb(P'')<sub>3</sub>(μ-K(P''))<sub>2</sub>]<sub>∞</sub> (**80**), *trans*-[Ln(P'')<sub>2</sub>(py)<sub>4</sub>] (**81-Ln**; Ln = Sm, Eu, Yb), [Ln(P'')<sub>2</sub>(18-crown-6)] (**82-Ln**; Ln = Sm, Eu, Yb).

reduced steric saturation of the Ln coordination sphere.<sup>112,113</sup>

Comparisons may also be made between **46-Ln** and the structurally analogous solvated Ln(III) tris-amide complexes [Ln{N(SiMe<sub>2</sub>H)<sub>2</sub>}<sub>3</sub>(THF)<sub>2</sub>] (Ln = Nd, Nd–N = 2.344(9) Å, O–Nd–O = 163.18(15)°; Ln = Yb, Yb–N = 2.224(8) Å, O–Yb–O = 160.50(14)°).<sup>114,144</sup> The solution <sup>31</sup>P{<sup>1</sup>H} NMR spectra of **46-Ln** (Ln = Ce, Pr, Nd, Sm) exhibited broad and paramagnetically shifted singlets at 616.7 (**46-Ce**), 1894.2 (**46-Pr**), 2570.3 (**46-Nd**) and –259.2 (**46-Sm**) ppm with no resonances being observed in <sup>13</sup>C{<sup>1</sup>H} and <sup>31</sup>P{<sup>1</sup>H} NMR spectra that could be assigned to **46-Tm**.<sup>89,112,113</sup> Solid state <sup>31</sup>P MAS NMR spectroscopy of **46-Ln** (Ln = Ce–Sm) revealed two components in their spectra in a 2 : 1 ratio, with the major component assigned to pyramidal and the minor component to planar P environments.<sup>89</sup> This is in contrast to the one resonance observed in the solution <sup>31</sup>P{<sup>1</sup>H} NMR spectra of **46-Ln** due to dynamic processes and broad resonances.<sup>89</sup>

In 1996, Rabe reported the synthesis of the bimetallic Sm(II) complex [Sm(P'')(μ-P'')<sub>3</sub>Sm(THF)<sub>3</sub>] (**78**, Fig. 23) by the salt metathesis reaction of two equivalents of KP'' with SmI<sub>2</sub>(THF)<sub>2</sub> in THF at room temperature, analogously to the preparation of **14** discussed in Section 2.1.1.<sup>125</sup> Single crystal XRD showed that **78** features one four-coordinate Sm ion with one terminal and three bridging P'', and one six-coordinate Sm ion with three bridging P'' and three bound THF. The N'' analogue of **78**, [Sm(N'')<sub>2</sub>(THF)<sub>2</sub>],<sup>145</sup> is monometallic, following a general trend that higher nuclearities tend to be observed for P'' complexes compared to their N'' counterparts.<sup>146</sup> The solid state structure of **78** differs from those of the six-coordinate monometallic Ln(II) bis-(diphenyl)phosphide complexes, [Ln(PPh<sub>2</sub>)<sub>2</sub>(THF)<sub>4</sub>] (Ln = Sm, Yb) and [Sm(PPh<sub>2</sub>)<sub>2</sub>(*N*-methylimidazole)<sub>4</sub>],<sup>147</sup> and bears more resemblance to the solvent-free Eu(II) N'' 'ate'

complex, [Eu(N'')(μ-N'')<sub>2</sub>Na].<sup>148</sup> The terminal Sm–P bond length of **78** (3.027(3) Å) is similar to the mean bridging Sm–P distances (3.039(5) Å), and all are shorter than those reported for six-coordinate [Sm(η<sup>1</sup>-dibenzophospholy)<sub>2</sub>(THF)<sub>4</sub>] (3.1908(6) Å),<sup>148</sup> [Sm(PPh<sub>2</sub>)<sub>2</sub>(*N*-methylimidazole)<sub>4</sub>] (3.139(3) Å),<sup>147</sup> and [Sm(η<sup>5</sup>-C<sub>8</sub>H<sub>4</sub>PMe<sub>2</sub>-2,3)<sub>2</sub>(THF)<sub>2</sub>] (3.0775(1) Å).<sup>149</sup> VT <sup>1</sup>H NMR spectra of **78** were collected in both *d*<sub>8</sub>-toluene and *d*<sub>8</sub>-THF in order to determine its structure in solution, but these studies were inconclusive. In *d*<sub>8</sub>-THF at 293 K there is a singlet at 0.42 ppm (*ν*<sub>1/2</sub> = 10 Hz), and at 173 K this becomes two singlets at 3.94 ppm (*ν*<sub>1/2</sub> = 16 Hz) and –0.01 ppm (*ν*<sub>1/2</sub> = 8 Hz). In *d*<sub>8</sub>-toluene at 293 K there is one singlet at 0.99 ppm (*ν*<sub>1/2</sub> = 20 Hz), but at 193 K seven broad resonances were observed at 13.60, 8.50, 6.10, 3.80, 3.10, –0.54 and –2.93 ppm.<sup>125</sup>

In 2024, Mills and co-workers showed that the addition of two or three equivalents of KP'' to [LnI<sub>2</sub>(THF)<sub>2</sub>] (Ln = Sm, Eu, Yb) in Et<sub>2</sub>O at –78 °C gave the dimeric Ln(II) P'' complexes [(Ln(P'')<sub>3</sub>(THF))<sub>2</sub>(μ-I)K<sub>3</sub>(THF)] (**79-Ln**; Ln = Sm, Eu) and polymeric [KYb(P'')<sub>3</sub>(μ-K(P''))<sub>2</sub>]<sub>∞</sub> (**80**), Fig. 23.<sup>115</sup> The addition of pyridine to either **79-Ln** or **80** at room temperature gave the monomeric pyridine-solvated Ln(II) P'' complexes *trans*-[Ln(P'')<sub>2</sub>(py)<sub>4</sub>] (**81-Ln**; Ln = Sm, Eu, Yb), whereas the addition of 18-crown-6 to **79-Ln** and **80** at room temperature afforded [Ln(P'')<sub>2</sub>(18-crown-6)] (**82-Ln**; Ln = Sm, Eu, Yb), Fig. 23.<sup>115</sup> The dimeric Ln(II) 'ate' complexes **79-Ln** consist of two {Ln(P'')<sub>3</sub>(THF)} fragments with KI encapsulated between them. The Ln centres exhibit distorted trigonal bipyramidal geometries with three equatorial P'', one axial THF and one axial iodide, which bridges the two fragments.<sup>115</sup> The mean Ln–P bond lengths in **79-Ln** (Ln = Sm, 3.033(7) Å; Eu, 3.038(7) Å) are shorter than the bridging Sm–P'' bonds in **78** but are statistically equivalent to the terminal Sm–P'' bond length of **78**.<sup>115,125</sup> Due to the





further reactivity, including them showing no reaction with  $KP''$ .<sup>116</sup>

In 2018, Liddle, Scheer and co-workers reported the synthesis of a series of An(IV) pnictogen complexes containing bulky triamidoamine ancillary ligands, including the  $P''$  complexes  $[An(Tren^{DMBS})(P'')]$  (**86-An**) and  $[An(Tren^{TIPS})(P'')]$  (**87-An**) (An = Th, U) ( $Tren^{DMBS} = \{N(CH_2CH_2NSiMe_2^tBu)_3\}$ ,  $Tren^{TIPS} = \{N(CH_2CH_2NSi^iPr_3)_3\}$ , Fig. 24) by the reaction of the respective triamidoamine precursor complexes  $[An(Tren^R)(L)][BPh_4]$  (An = Th, L = DME; U, L = THF) with one equivalent of  $KP''$ .<sup>117</sup> It was noted that for **86-Th** the Th–P bond length (2.9406(11) Å) was 0.08 Å longer than the U–P bond in **86-U** (2.8646(14) Å) despite the single bond covalent radius of Th only being 0.05 Å larger than U.<sup>90</sup> Conversely, the U–P bond distance in **87-U** (2.8391(9) Å) is shorter than that seen for **87-U**, and the geometry about the P atom is more planar in the former complex (359.94(5)°) than the latter (354.06(8)°); this was determined by the sum of the angles about the phosphorus atom and the deviation of this value from 360°, which can be attributed to the more sterically demanding  $Tren^{TIPS}$  ligand.<sup>117</sup> As part of a study to probe the propensity of An–E (An = Th, U; E = P, As, Sb, Bi) bonds to undergo either homolytic cleavage or acid–base/dehydrocoupling to occur, **86-U** and **87-U** were heated to 80 °C and exposed to a 125 W UV lamp for 2 h. It was noted that both complexes are remarkably robust, and undergo <5% thermal and photolytic decomposition under these conditions, likely by homolytic cleavage, whereas acid–base/dehydrocoupling is the most likely decomposition route for more redox-robust Th(IV)–E bonds.

### 3. Bis(triisopropylsilyl)phosphide ( $P^{\dagger\dagger}$ ) complexes

In accord with the paucity of structurally authenticated  $P''$  complexes compared to  $N''$  across the periodic table, there are even fewer examples of complexes containing bulkier bis(trialkylsilyl)phosphides.<sup>146,152–160</sup> As bulkier bis(trialkylsilyl)-amides such as bis(triisopropylsilyl)amide ( $\{N(Si^iPr_3)_2\}$ ,  $N^{\dagger\dagger}$ ) are

now starting to grow in popularity as ligands in low-coordinate s-block and f-block chemistry,<sup>5,161</sup> we also cover bis(triisopropylsilyl)phosphide ( $\{P(Si^iPr_3)_2\}$ ,  $P^{\dagger\dagger}$ ) chemistry here; only seven structurally authenticated  $P^{\dagger\dagger}$  complexes have been reported to date.

In 1996, Driess reported the synthesis of  $[\{Li(\mu-P^{\dagger\dagger})\}_3\{Li[\mu-P(Si^iPr_3)H]\}]$  (**88**) by the lithiation of  $HP(Si^iPr_3)_2$  with  $nBuLi$  in toluene.<sup>155</sup> Complex **88** is an eight-membered ring consisting of four Li centres, three bridging  $P^{\dagger\dagger}$  and one bridging  $HP(Si^iPr_3)$  ligand.<sup>155</sup> Each Li atom in **88** is two-coordinate and the mean Li–P bond length is 2.44(5) Å which is shorter than those seen for the ladder-like  $Li-P''$  complexes **3** and **4**; bulky  $P^{\dagger\dagger}$  suppresses the rearrangement to a ladder-like conformation. The  $^{31}P$  NMR spectrum of a solution of **88** in  $d_8$ -toluene at  $-70$  °C exhibited two singlets at  $-338$  and  $-370$  ppm and a broad doublet at  $-351$  ppm ( $^1J_{PH} = 170$  Hz). This provided evidence that in solution **88** is a mixed aggregate made up of various  $[Li\{P(Si^iPr_3)_2\}(H)]$  and  $[Li(P^{\dagger\dagger})]$  building blocks.<sup>155</sup> In 2004, von Hae-nisch published the solid-state structure of  $[Li(\mu-P^{\dagger\dagger})(THF)]_2$  (**89**), Fig. 25, *via* a CSD communication.<sup>162</sup> Later, in 2005, Westerhausen also reported the synthesis of **89** by analogous methods to that of **88**, but with THF used as the solvent.<sup>163</sup> Complex **89** is a dimeric complex consisting of two three-coordinate Li centres bridged by two  $P^{\dagger\dagger}$ , with each Li bound by one THF.<sup>163</sup> The Li–P bond length in **89** is 2.533(6) Å, which falls within the range observed for  $Li-P''$  complexes discussed in this review. It is noted however that although this bond length is neither longer nor shorter than for  $Li-P''$  complexes, the coordination number of Li in **89** is only three, compared to four for **1**.<sup>163</sup> The  $^{29}Si\{^1H\}$  NMR spectrum of **89** shows a single resonance at 21.6 ppm for the  $Si^iPr_3$  groups, which exhibits coupling to the  $^{31}P$  nuclei ( $^1J_{PSi} = 49.7$  Hz), and one resonance was observed in the  $^{31}P\{^1H\}$  NMR spectrum at  $-374.7$  ppm.<sup>163</sup> In 2024, Kays and Mills separately reported the synthesis of  $NaP^{\dagger\dagger}$  by the addition of two or three equivalents of  $Si^iPr_3Cl$  to a refluxing DME solution of red phosphorus, Na metal and naphthalene to yield crystals of either  $[Na(P^{\dagger\dagger})(DME)_2]$  (**90**), or  $[Na(\mu-P^{\dagger\dagger})(THF)]_2$  (**91**) following treatment with THF, Fig. 25.<sup>164,165</sup>




Fig. 25  $P^{\dagger\dagger}$  complexes  $[\{Li(\mu-P^{\dagger\dagger})\}_3\{Li[\mu-P(Si^iPr_3)H]\}]$  (**88**),  $[Li(\mu-P^{\dagger\dagger})(THF)]_2$  (**89**),  $[Na(P^{\dagger\dagger})(DME)_2]$  (**90**),  $[Na(\mu-P^{\dagger\dagger})(THF)]_2$  (**91**),  $[W(P^{\dagger\dagger})(CO)_3][Li(THF)_4]$  (**92**),  $[M(P^{\dagger\dagger})_2]$  (**93-M**; M = Zn, Cd, Hg),  $[Ln(P^{\dagger\dagger})_2(THF)_3]$  (**94-Ln**; Ln = Sm, Eu),  $[Yb(P^{\dagger\dagger})_2(THF)_2]$  (**94-Yb**).



In contrast to  $P''$  chemistry, where Na salts are yet to be structurally authenticated (see above), the solid-state structures of both **90** and **91** were determined by single crystal XRD. Complex **90** consists of a five-coordinate Na centre with one  $P^{††}$  and two bound DME solvent molecules,<sup>164</sup> whilst **91** is dimeric with a  $Na_2P_2$  core and bridging  $P^{††}$ , with the Na coordination spheres completed by a single THF each.<sup>165</sup> The mean Na–P bond length in **90** is 2.824(3) Å,<sup>164</sup> compared to 2.805(2) Å for **91**,<sup>165</sup> which fall within the mean bond lengths discussed for both Li– $P''$  and K– $P''$  complexes. The geometry of the phosphorus centre in **90** is trigonal pyramidal, with the sum of angles about the phosphorus centre adding to 332.53(10)°. <sup>29</sup>Si DEPT90 NMR spectrum of **90** or **91** in  $d_8$ -THF contains a doublet at 20.65 ppm with  $^1J_{PSi} = 60.2$  Hz, whilst a singlet is observed in the  $^{31}P\{^1H\}$  NMR spectrum at –384.26 ppm with satellites observed with  $^1J_{PSi} = 59.8$  Hz.<sup>164,165</sup>

In 2005, Westerhausen reported the synthesis of a W(0)  $P^{††}$  complex by addition of a THF solution of  $[W(CO)_5(THF)]$  to **89** to yield  $[W(P^{††})(CO)_5][Li(THF)_4]$  (**92**), Fig. 25.<sup>163</sup> Complex **92** exhibits a distorted octahedral geometry with five coordinated CO and one  $P^{††}$ .<sup>163</sup> The W–P bond length of **92** is 2.6665(7) Å and the geometry about the phosphorus centre is trigonal pyramidal (350.38(6)°), which was noted to reflect the shift in the *trans*-CO stretching vibration to smaller wavenumbers (1904  $cm^{-1}$ ).<sup>163</sup> As phosphides are strong  $\sigma$ -donors but weak  $\pi$ -acceptors, the  $\pi$ -backbonding from the W to the *trans*-CO ligand is increased, which weakens this W–C bond.<sup>163</sup> The  $^{29}Si\{^1H\}$  NMR spectrum of **92** exhibits a triplet at 21.1 ppm with coupling to  $^{31}P$  of  $^1J_{PSi} = 40.8$  Hz, and a singlet resonance at –409.2 ppm is observed in its  $^{31}P\{^1H\}$  NMR spectrum.<sup>163</sup> In 2024, Kays reported the synthesis of the group 12 M(II) complexes  $[M(P^{††})_2]$  (**93-M**; M = Zn, Cd, Hg) by the salt metathesis reactions of parent  $ZnCl_2$ ,  $CdI_2$  or  $HgBr_2$  with two equivalents of **90**.<sup>165</sup> The average M–P bond lengths of 2.2263(5) (**93-Zn**), 2.4215(10) (**93-Cd**) and 2.3938(7) (**93-Hg**) Å are consistent with those seen for the terminal  $P''$  ligands in **68** and **69-M**.<sup>83,165</sup> In addition, complexes **93-M** exhibit an increasing P–M–P angle from Zn to Hg (**93-Zn**; 168.747(12)°, **93-Cd**; 169.215(19)°, **93-Hg**; 170.086(16)°).<sup>165</sup> The  $^{31}P\{^1H\}$  NMR spectra for **93-M** exhibit singlets at –287.8, –284.0 and –209.2 for the Zn, Cd and Hg complexes respectively.<sup>165</sup> For **93-Cd**, the  $^{113}Cd$  NMR spectrum exhibited a triplet resonance at 137.68 ppm ( $^1J_{CdP} = 350$  Hz, 12.2% abundant  $^{113}Cd$   $I = \frac{1}{2}$ ) whereas the  $^{199}Hg$  NMR spectrum of **93-Hg** contains a triplet at 13.2 ppm ( $^1J_{HgP} = 407.7$  Hz, 16.9% abundant  $^{199}Hg$   $I = \frac{1}{2}$ ).<sup>165</sup> Interestingly, the  $^{29}Si$  NMR spectra for **93-Zn** and **93-Cd** display apparent doublets, whereas for **93-Hg** the  $^{29}Si$  NMR spectrum is consistent with an AA'XX' spin system with virtual coupling. This spectrum was accurately simulated using  $^1J_{PSi} = 50.6$  Hz,  $^3J_{P'Si} = 0.0$  Hz and  $^2J_{PP'} = 19.0$  Hz.<sup>165</sup>

In 2024, Mills and co-workers reported the synthesis of a series of three Ln(II)  $P^{††}$  complexes by the addition of two equivalents of **90** to  $[LnI_2(THF)_2]$  (Ln = Sm, Eu, Yb) to yield  $[Ln(P^{††})_2(THF)_x]$  (Ln = Sm,  $x = 3$ , **94-Sm**; Eu,  $x = 3$ , **94-Eu**; Yb,  $x = 2$ , **94-Yb**), Fig. 25.<sup>164</sup> Complexes **94-Sm** and **94-Eu** exhibit five-coordinate Ln centres which are bound by two  $P^{††}$  and three THF, whereas **94-Yb** features four-coordinate Yb bound by two  $P^{††}$  and two THF.<sup>164</sup> The mean Ln–P bond lengths of 3.0336(13) (**94-Sm**),

3.0237(18) (**94-Eu**) and 2.8065(13) (**94-Yb**) Å are shorter than those seen for the Ln(II)  $P''$  complexes **79-M**, **81-M** and **82-M** despite the increased bulk of the  $P^{††}$  ligand, though the coordination numbers of **94-Ln** are lower.<sup>164</sup> Complexes **94-M** exhibit bent geometries, as evidenced by the P–Ln–P angles (156.11(3)°, **94-Sm**; 156.10(3)°, **94-Eu**; 133.48(3)°, **94-Yb**).<sup>164</sup> For the diamagnetic complex **94-Yb**, a virtual triplet is observed in the  $^{29}Si$  DEPT90 NMR spectrum at 24.30 ppm, with coupling to  $^{31}P$  nuclei  $^1J_{PSi} = 15.8$  Hz. The  $^{31}P\{^1H\}$  NMR spectrum contains a singlet at –301.10 ppm which shows satellites from coupling to  $^{171}Yb$ , with  $^1J_{YbP} = 1382.1$  Hz, as well as to  $^{29}Si$ ,  $^1J_{PSi} = 18.1$  Hz.<sup>164</sup> The  $^{171}Yb\{^1H\}$  NMR spectrum of **94-Yb** contains a triplet resonance at 682 ppm due to coupling to two equivalent  $^{31}P$  nuclei ( $^1J_{YbP} = 1382.9$  Hz).

## 4. Conclusions

We have shown that metal  $P''$  coordination chemistry lags far behind that of its lighter congener  $N''$ , and that there are therefore a huge number of avenues to be explored and exploited. Although the CSD search performed above gave 210  $P''$  vs. 4062  $N''$  metal complexes, we note that an analogous search for the other heavy group 15 congeners  $As''$  (101),  $Sb''$  (58) and  $Bi''$  (22) returns progressively fewer hits upon descent of the group.<sup>9</sup> Perhaps the most surprising omissions in s-block metal  $P''$  chemistry are that there are no structurally characterised examples of Na or Be  $P''$  complexes to date.<sup>9</sup> The former of these absences is quite remarkable considering that synthetic routes to Na  $P''$  salts are well-known,<sup>12</sup> and that solid-state structures of solvated Na  $P^{††}$  complexes are already known despite the chemistry of this bulkier bis(silyl)phosphide ligand being in its relative infancy.<sup>164,165</sup> We have found there to be relatively few p-block metal  $P''$  complexes, including no structurally authenticated group 15 and 16 metal(loid)  $P''$  complexes to date.<sup>9</sup> This is quite astonishing considering the potential of such molecular precursors in the synthesis of solid-state III–V semiconductors;<sup>42–57</sup> we noted that a Bi  $P''$  complex has been reported without a solid-state structure,<sup>12e</sup> and this is also the case for Sb.<sup>166</sup> We have identified multiple gaps in d-block metal  $P''$  chemistry that can be exploited, mainly for the second and third row d-transition metals. There are ten d-transition metal M–P bonds that have yet to be structurally authenticated for this ligand, surprisingly including the entirety of groups 5 and 9.<sup>9</sup> We have discussed that numerous 18  $e^-$  Fe(II), Ru(II) and Os(II)  $P''$  complexes have been synthesised but not yet structurally characterised,<sup>99</sup> similarly, there are numerous other 18  $e^-$  d-transition metal  $P''$  complexes that are known but lack solid-state structural data, including Mn(I)<sup>167</sup> and Re(I)<sup>168</sup> examples. The synthesis of group 7 M(I) and group 8 M(II)  $P''$  complexes are particularly promising avenues to pursue as these are diamagnetic and follow the 18  $e^-$  rule, allowing reactions to be easily monitored by  $^{31}P$  NMR spectroscopy and in principle providing relatively stable products.<sup>22,23</sup> Finally, there is also a notable paucity of f-block metal  $P''$  complexes, considering how crucial the  $N''$  ligand has been to the development of molecular f-block chemistry,<sup>5</sup> and this has only started to be addressed relatively recently.<sup>115,116,164</sup>





- 23 *Comprehensive Coordination Chemistry III*, ed. E. C. Constable and L. Que Jr., Elsevier B. V., Amsterdam, 2007.
- 24 E. Hey, P. B. Hitchcock, M. F. Lappert and A. K. Rai, *J. Organomet. Chem.*, 1987, **325**, 1–12.
- 25 G. Becker, H.-M. Hartmann and W. Schwarz, *Z. Anorg. Allg. Chem.*, 1989, **577**, 9–22.
- 26 E. Hey-Hawkins and E. Sattler, *J. Chem. Soc., Chem. Commun.*, 1992, **24**, 775–776.
- 27 U. Englich, K. Hassler, K. Ruhlandt-Senge and F. Uhlig, *Inorg. Chem.*, 1998, **37**, 3532–3537.
- 28 A. S. Ionkin, W. J. Marshall, B. M. Fish, A. A. Marchione, L. A. Howe, F. Davidson and C. N. McEwen, *Eur. J. Inorg. Chem.*, 2008, 2386–2390.
- 29 M. Westerhausen, M. H. Digeser, B. Wieneke, H. Nöth and J. Knizek, *Eur. J. Inorg. Chem.*, 1998, 517–521.
- 30 M. Westerhausen, M. H. Digeser, H. Nöth, T. Seifert and A. Pfitzner, *J. Am. Chem. Soc.*, 1998, **120**, 6722–6725.
- 31 M. Westerhausen and W. Schwarz, *Z. Anorg. Allg. Chem.*, 1994, **620**, 304–308.
- 32 M. Westerhausen and A. Pfitzner, *J. Organomet. Chem.*, 1995, **487**, 187–195.
- 33 M. Westerhausen and W. Schwarz, *Z. Anorg. Allg. Chem.*, 1996, **622**, 903–913.
- 34 M. Westerhausen, H.-D. Hausen and W. Schwarz, *Z. Anorg. Allg. Chem.*, 1995, **621**, 877–888.
- 35 M. Westerhausen, *J. Organomet. Chem.*, 1994, **479**, 141–151.
- 36 M. Westerhausen and W. Schwarz, *J. Organomet. Chem.*, 1993, **463**, 51–63.
- 37 T. Habereeder, H. Nöth and R. T. Paine, *Eur. J. Inorg. Chem.*, 2007, 4298–4305.
- 38 F. Thomas, S. Schulz and M. Nieger, *Eur. J. Inorg. Chem.*, 2001, 161–166.
- 39 F. Thomas, S. Schulz, M. Nieger and K. Nättinen, *Chem. – Eur. J.*, 2002, **8**, 1915.
- 40 M. Matar, S. Schulz and U. Flörke, *Z. Anorg. Allg. Chem.*, 2007, **633**, 162–165.
- 41 K. L. Antcliff, R. J. Baker, C. Jones, D. M. Murphy and R. P. Rose, *Inorg. Chem.*, 2005, **44**, 2098–2105.
- 42 B. Li, S. Bauer, M. Seidl, A. Y. Timoshkin and M. Scheer, *Chem. – Eur. J.*, 2019, **25**, 13714–13718.
- 43 L. K. Krannich, C. L. Watkins, S. J. Schauer and C. H. Lake, *Organometallics*, 1996, **15**, 3980–3984.
- 44 E. Hey-Hawkins, M. F. Lappert, J. L. Atwood and S. G. Bott, *J. Chem. Soc., Dalton Trans.*, 1991, 939–948.
- 45 R. L. Wells, A. T. McPhail, M. F. Self and J. A. Laske, *Organometallics*, 1993, **12**, 3333–3339.
- 46 M. Westerhausen, C. Birg, H. Nöth, J. Knizek and T. Seifert, *Eur. J. Inorg. Chem.*, 1999, 2209–2214.
- 47 D. Wiedmann, H.-D. Hausen and J. Weidlein, *Z. Anorg. Allg. Chem.*, 1995, **621**, 1351–1357.
- 48 R. J. Jouet, R. L. Wells, A. L. Rheingold and C. D. Incarvito, *J. Organomet. Chem.*, 2000, **601**, 191–198.
- 49 S. T. Barry, S. Belhumeur and D. S. Richeson, *Organometallics*, 1997, **16**, 3588–3592.
- 50 R. L. Wells, R. A. Baldwin, P. S. White, W. T. Pennington, A. L. Rheingold and G. P. A. Yap, *Organometallics*, 1996, **15**, 91–97.
- 51 R. L. Wells, E. E. Foos, R. A. Baldwin, A. L. Rheingold and G. P. A. Yap, *Heteroat. Chem.*, 1998, **9**, 147–150.
- 52 F. Thomas, S. Schulz and M. Nieger, *Z. Anorg. Allg. Chem.*, 2002, **628**, 235–242.
- 53 C. von Hänisch, *Z. Anorg. Allg. Chem.*, 2001, **627**, 68–72.
- 54 R. L. Wells, A. T. McPhail, L. J. Jones and M. F. Self, *Polyhedron*, 1993, **12**, 141–147.
- 55 C. Banerjee, D. L. Hughes, M. Bochmann and T. Nann, *Dalton Trans.*, 2012, **41**, 7244.
- 56 R. L. Wells, A. T. McPhail and M. F. Self, *Organometallics*, 1992, **11**, 221–225.
- 57 F. Thomas, S. Schulz, H. Mansikkamäki and M. Nieger, *Angew. Chem., Int. Ed.*, 2003, **42**, 5641–5644.
- 58 F. Thomas, S. Schulz, H. Mansikkamäki and M. Nieger, *Organometallics*, 2003, **22**, 3471–3477.
- 59 R. L. Wells, A. T. McPhail, L. J. Jones III and M. F. Self, *J. Organomet. Chem.*, 1993, **449**, 85–94.
- 60 R. L. Wells, A. T. McPhail and M. F. Self, *Organometallics*, 1993, **12**, 3363–3367.
- 61 R. L. Wells, E. E. Foos, A. L. Rheingold, G. P. A. Yap, L. M. Liable-Sands and P. S. White, *Organometallics*, 1998, **17**, 2869–2875.
- 62 M. Nieger, F. Thomas and S. Schulz, CCDC 244687: Experimental Crystal Structure Determination, 2004.
- 63 T. Douglas and K. H. Theopold, *Inorg. Chem.*, 1991, **30**, 594–596.
- 64 R. L. Wells, S. R. Aubuchon, M. F. Self, J. P. Jasinski, R. C. Woudenberg and R. J. Butcher, *Organometallics*, 1992, **11**, 3370–3375.
- 65 J. A. L. Cooke, R. L. Wells and P. S. White, *Organometallics*, 1995, **14**, 3562–3565.
- 66 L. J. I. Jones, A. T. McPhail and R. L. Wells, *Organometallics*, 1994, **13**, 2504–2507.
- 67 R. L. Wells, M. F. Self, A. T. McPhail, S. R. Aubuchon, R. C. Woudenberg and J. P. Jasinski, *Organometallics*, 1993, **12**, 2832–2834.
- 68 S. R. Aubuchon, A. T. McPhail, R. L. Wells, J. A. Giambra and J. R. Bowser, *Chem. Mater.*, 1994, **6**, 82–86.
- 69 R. L. Wells, S. R. Aubuchon, M. S. Lube and P. S. White, *Main Group Chem.*, 1995, **1**, 81–88.
- 70 J. F. Janik, R. L. Wells and P. S. White, *Inorg. Chem.*, 1998, **37**, 3561–3566.
- 71 J. F. Janik, R. L. Wells, V. G. Young, A. L. Rheingold and I. A. Guzei, *J. Am. Chem. Soc.*, 1998, **120**, 532–537.
- 72 A. Schaller, H.-D. Hausen, J. Weidlein and P. Fischer, *Z. Anorg. Allg. Chem.*, 2000, **626**, 616–618.
- 73 J. A. Burns, W. T. Pennington and G. H. Robinson, *Organometallics*, 1995, **14**, 1533–1535.
- 74 J. F. Janik, R. L. Wells and P. S. White, *Organometallics*, 1998, **17**, 2361–2365.
- 75 J. F. Janik, R. L. Wells, V. G. Young and J. A. Halfen, *Organometallics*, 1997, **16**, 3022–3026.
- 76 C. von Hänisch and B. Rolli, *Z. Anorg. Allg. Chem.*, 2004, **630**, 1987–1990.
- 77 M. Usher, A. V. Protchenko, A. Rit, J. Campos, E. L. Kolychev, R. Tirfoin and S. Aldridge, *Chem. – Eur. J.*, 2016, **22**, 11685–11698.



- 78 E. C. Y. Tam, N. A. Maynard, D. C. Apperley, J. D. Smith, M. P. Coles and J. R. Fulton, *Inorg. Chem.*, 2012, **51**, 9403–9415.
- 79 S. Yao, M. Brym, K. Merz and M. Driess, *Organometallics*, 2008, **27**, 3601–3607.
- 80 S. Yao, S. Block, M. Brym and M. Driess, *Chem. Commun.*, 2007, 3844–3846.
- 81 L. M. Harris, E. C. Y. Tam, S. J. W. Cummins, M. P. Coles and J. R. Fulton, *Inorg. Chem.*, 2017, **56**, 3087–3094.
- 82 B. P. Johnson, S. Almstätter, F. Dielmann, M. Bodensteiner and M. Scheer, *Z. Anorg. Allg. Chem.*, 2010, **636**, 1275–1285.
- 83 S. C. Goel, M. Y. Chiang, D. J. Rauscher and W. E. Buhro, *J. Am. Chem. Soc.*, 1993, **115**, 160–169.
- 84 M. Westerhausen, M. M. Enzelberger and W. Schwarz, *J. Organomet. Chem.*, 1995, **491**, 83–90.
- 85 S. Scheuermayer, F. Tuna, M. Bodensteiner, M. Scheer and R. A. Layfield, *Chem. Commun.*, 2012, **48**, 8087–8089.
- 86 M. Westerhausen, S. Schneiderbauer, M. Hartmann, M. Warchhold and H. Nöth, *Z. Anorg. Allg. Chem.*, 2002, **628**, 330–332.
- 87 M. Westerhausen, M. Hartmann and W. Schwarz, *Inorg. Chim. Acta*, 1998, **269**, 91–100.
- 88 M. Fustier, X.-F. Le Goff, M. Lutz, J. C. Slootweg and N. Mézailles, *Organometallics*, 2015, **34**, 63–72.
- 89 J. Baldwin, K. L. Bonham, T. R. C. Thompson, G. K. Gransbury, G. F. S. Whitehead, I. J. Vitorica-Yrezabal, D. Lee, N. F. Chilton and D. P. Mills, *JACS Au*, 2025, DOI: [10.1021/jacsau.4c01018](https://doi.org/10.1021/jacsau.4c01018).
- 90 D. Fenske, A. Grissinger, E. M. Hey-Hawkins and J. Magull, *Z. Anorg. Allg. Chem.*, 1991, **595**, 57–66.
- 91 L. Weber, G. Meine, R. Boese and N. Augart, *Organometallics*, 1987, **6**, 2484–2488.
- 92 E. Hey, M. F. Lappert, J. L. Atwood and S. G. Bott, *Polyhedron*, 1988, **7**, 2083–2086.
- 93 F. Lindenberg and E. Hey-Hawkins, *J. Organomet. Chem.*, 1992, **435**, 291–297.
- 94 E. Hey-Hawkins and F. Lindenberg, *Chem. Ber.*, 1992, **125**, 1815–1819.
- 95 F. Lindenberg and E. Hey-Hawkins, *Z. Anorg. Allg. Chem.*, 1995, **621**, 1531–1534.
- 96 S. Scheuermayer, F. Tuna, E. M. Pineda, M. Bodensteiner, M. Scheer and R. A. Layfield, *Inorg. Chem.*, 2013, **52**, 3878–3883.
- 97 L. Dütsch, C. Riesinger, G. Balázs and M. Scheer, *Chem. – Eur. J.*, 2021, **27**, 8804–8810.
- 98 W. E. Buhro, M. H. Chisholm, K. Folting, J. C. Huffman, J. D. Martin and W. E. Streib, *J. Am. Chem. Soc.*, 1992, **114**, 557–570.
- 99 (a) H. Schäfer, *Z. Anorg. Allg. Chem.*, 1980, **467**, 105–122; (b) L. Weber, K. Reizig and R. Boese, *Chem. Ber.*, 1985, **118**, 1193–1203; (c) L. Weber, K. Reizig and R. Boese, *Organometallics*, 1985, **4**, 2097–2101; (d) L. Weber and D. Bungardt, *J. Organomet. Chem.*, 1986, **311**, 269–280; (e) L. Weber, K. Reizig, D. Bungardt and R. Boese, *Organometallics*, 1987, **6**, 110–114; (f) L. Weber, I. Schumann, H.-G. Stammmler and B. Neumann, *Z. Naturforsch.*, 1992, **B47**, 1134–1140; (g) L. Weber, H. Misiak, S. Buchwald, H.-G. Stammmler and B. Neumann, *Organometallics*, 1994, **13**, 2194–2204.
- 100 H. Schäfer, D. Binder, B. Deppisch and G. Mattern, *Z. Anorg. Allg. Chem.*, 1987, **546**, 79–98.
- 101 M. Madadi, B. Khalili Najafabadi, M. A. Fard and J. F. Corrigan, *Eur. J. Inorg. Chem.*, 2015, 3094–3101.
- 102 B. Deppisch and H. Schäfer, *Z. Anorg. Allg. Chem.*, 1982, **490**, 129–135.
- 103 N. C. Breit, T. Szilvási, T. Suzuki, D. Gallego and S. Inoue, *J. Am. Chem. Soc.*, 2013, **135**, 17958–17968.
- 104 B. Khalili Najafabadi and J. F. Corrigan, *Dalton Trans.*, 2015, **44**, 14235–14241.
- 105 B. Khalili Najafabadi and J. F. Corrigan, *Can. J. Chem.*, 2016, **94**, 593–598.
- 106 R. Wei, X. Wang and L. Leo Liu, *Z. Anorg. Allg. Chem.*, 2023, **649**, e202300005.
- 107 B. Rademacher, W. Schwarz and M. Westerhausen, *Z. Anorg. Allg. Chem.*, 1995, **621**, 287–300.
- 108 O. Fuhr and D. Fenske, *Z. Anorg. Allg. Chem.*, 1999, **625**, 1229–1236.
- 109 F. Thomas, S. Schulz and M. Nieger, *Organometallics*, 2001, **20**, 2405–2408.
- 110 F. Lindenberg, T. Gelbrich and E. Hey-Hawkins, *Z. Anorg. Allg. Chem.*, 1995, **621**, 771–778.
- 111 F. Lindenberg, T. Shribman, J. Sieler, E. Hey-Hawkins and M. S. Eisen, *J. Organomet. Chem.*, 1996, **515**, 19–25.
- 112 G. W. Rabe and J. W. Ziller, *Inorg. Chem.*, 1995, **34**, 5378–5379.
- 113 G. W. Rabe, J. Riede and A. Schier, *J. Chem. Soc., Chem. Commun.*, 1995, **5**, 577–578.
- 114 W. A. Herrmann, R. Anwander, F. C. Munck, W. Scherer, V. Dufaud, N. W. Huber and G. R. J. Artus, *Z. Naturforsch., B*, 1994, **49**, 1789–1797.
- 115 J. Baldwin, A. Brookfield, G. F. S. Whitehead, L. S. Natrajan, E. J. L. McInnes, M. S. Oakley and D. P. Mills, *Inorg. Chem.*, 2024, **63**, 18120–18136.
- 116 S. W. Hall, J. C. Huffman, M. M. Miller, L. R. Avens, C. J. Burns, A. P. Sattelberger, D. S. J. Arney and A. F. England, *Organometallics*, 1993, **12**, 752–758.
- 117 T. M. Rookes, E. P. Wildman, G. Balázs, B. M. Gardner, A. J. Wooles, M. Gregson, F. Tuna, M. Scheer and S. T. Liddle, *Angew. Chem., Int. Ed.*, 2018, **57**, 1332–1336.
- 118 G. Fritz and W. Hölderich, *Z. Anorg. Allg. Chem.*, 1976, **422**, 104–114.
- 119 W. N. Setzer and P. von R. Schleyer, *Adv. Organomet. Chem.*, 1985, **24**, 353–451.
- 120 R. D. Rogers, J. L. Atwood and R. Grüning, *J. Organomet. Chem.*, 1978, **157**, 229–237.
- 121 D. Mootz, A. Zinnius and B. Böttcher, *Angew. Chem., Int. Ed. Engl.*, 1969, **8**, 378–379.
- 122 L. M. Engelhardt, A. S. May, C. L. Raston and A. H. White, *J. Chem. Soc., Dalton Trans.*, 1983, **8**, 1671–1673.
- 123 R. D. Shannon, *Acta Crystallogr., Sect. A*, 1976, **A32**, 751–767.
- 124 R. E. Mulvey and S. D. Robertson, Modern Developments in Magnesium Reagent Chemistry for Synthesis, in



- Alkaline-Earth Metal Compounds: Oddities and Applications*, ed. S. Harder, Springer Berlin, Heidelberg, 2013.
- 125 G. W. Rabe, J. Riede and A. Schier, *Organometallics*, 1996, **15**, 439–441.
- 126 A. Kuczkowski, S. Schulz and M. Nieger, *Appl. Organomet. Chem.*, 2004, **18**, 244–251.
- 127 D. Martin, M. Soleilhavoup and G. Bertrand, *Chem. Sci.*, 2011, **2**, 389–399.
- 128 Y. Mizuhata, T. Sasamori and N. Tokitoh, *Chem. Rev.*, 2009, **109**, 3479–3511.
- 129 D. Bourissou, O. Guerret, F. P. Gabbaï and G. Bertrand, *Chem. Rev.*, 2000, **100**, 39–92.
- 130 D. W. Stephan and G. Erker, *Angew. Chem., Int. Ed.*, 2015, **54**, 6400–6441.
- 131 P. P. Power, *Acc. Chem. Res.*, 2011, **44**, 627–637.
- 132 R. C. Fischer and P. P. Power, *Chem. Rev.*, 2010, **110**, 3877–3923.
- 133 P. P. Power, *Nature*, 2010, **463**, 171–177.
- 134 L. Weber, K. Reizig and M. Frebel, *Chem. Ber.*, 1986, **119**, 1857–1867.
- 135 L. Weber and D. Bungardt, *J. Organomet. Chem.*, 1986, **311**, 269–280.
- 136 L. Weber, K. Reizig, D. Bungardt and R. Boese, *Organometallics*, 1987, **6**, 110–114.
- 137 L. Weber and G. Meine, *Chem. Ber.*, 1987, **120**, 457–459.
- 138 H. M. Dietrich, C. Meermann, K. W. Törnroos and R. Anwänder, *Organometallics*, 2006, **25**, 4316–4321.
- 139 P. Pykkö, *J. Phys. Chem. A*, 2015, **119**, 2326–2337.
- 140 C. von Hänisch, K. Dollberg, J. Moritz, K. Mészáros, C. Jakobi, R.-M. Richter and F. Weigend, *Z. Anorg. Allg. Chem.*, 2024, e202400023.
- 141 R. A. Andersen, D. H. Templeton and A. Zalkin, *Inorg. Chem.*, 1978, **17**, 2317–2319.
- 142 M. Niemeyer, *Z. Anorg. Allg. Chem.*, 2002, **628**, 647–657.
- 143 D. C. Bradley, J. S. Ghotra and F. A. Hart, *J. Chem. Soc., Dalton Trans.*, 1973, **10**, 1021–1023.
- 144 G. Occhipinti, C. Meermann, H. M. Dietrich, R. Litlabø, F. Auras, K. W. Törnroos, C. Maichle-Mössner, V. R. Jensen and R. Anwänder, *J. Am. Chem. Soc.*, 2011, **133**, 6323–6337.
- 145 W. J. Evans, D. K. Drummond, H. Zhang and J. L. Atwood, *Inorg. Chem.*, 1988, **27**, 575–579.
- 146 M. A. Matchett, M. Y. Chiang and W. E. Buhro, *Inorg. Chem.*, 1994, **33**, 1109–1114.
- 147 G. W. Rabe, G. P. A. Yap and A. L. Rheingold, *Inorg. Chem.*, 1995, **34**, 4521–4522.
- 148 T. D. Tilley, R. A. Andersen and A. Zalkin, *Inorg. Chem.*, 1984, **23**, 2271–2276.
- 149 F. Nief and L. Ricard, *J. Organomet. Chem.*, 1994, **464**, 149–154.
- 150 J. M. Manriquez, P. J. Fagan, T. J. Marks, C. S. Day and V. W. Day, *J. Am. Chem. Soc.*, 1978, **100**, 7112–7114.
- 151 S. J. Simpson, H. W. Turner and R. A. Andersen, *Inorg. Chem.*, 1981, **20**, 2991–2995.
- 152 M. Westerhausen, M. H. Digeser, H. Nöth and J. Knizek, *Z. Anorg. Allg. Chem.*, 1999, **2**, 215–220.
- 153 M. Driess, U. Winkler, W. Imhof, L. Zsolnai and G. Huttner, *Chem. Ber.*, 1994, **6**, 1031–1035.
- 154 M. Westerhausen, G. Lang and W. Schwarz, *Chem. Ber.*, 1996, **9**, 1035–1040.
- 155 M. Driess and H. Pritzkow, *Z. Anorg. Allg. Chem.*, 1996, **9**, 1524–1530.
- 156 A. S. Ionkin and W. J. Marshall, *Organometallics*, 2003, **22**, 4136–4144.
- 157 A. Lorbach, S. Breitung, I. Sängler, F. Schödel, M. Bolte, M. Wagner and H.-W. Lerner, *Inorg. Chim. Acta*, 2011, **1**, 1–9.
- 158 M. Petrie and P. P. Power, *J. Chem. Soc., Dalton Trans.*, 1993, **11**, 1737–1745.
- 159 C. von Hänisch and D. Nikolova, *Eur. J. Inorg. Chem.*, 2006, 4770–4773.
- 160 M. Driess, R. Janoschek, H. Pritzkow, S. Rell and U. Winkler, *Angew. Chem., Int. Ed. Engl.*, 1995, **15**, 1614–1616.
- 161 F. Ortu and D. P. Mills, Low Coordinate Rare Earth and Actinide Complexes, in *Handbook on the Physics and Chemistry of Rare Earths*, ed. J.-C. G. Bünzli and V. K. Pecharsky, Elsevier B. V., Amsterdam, 2019, ch. 306, 55, pp. 1–87.
- 162 C. von Hänisch, CCDC 154239: Experimental Crystal Structure Determination, 2004.
- 163 M. Westerhausen, T. Rotter, H. Górls, C. Birg, M. Warchhold and H. Noth, *Z. Naturforsch., B*, 2005, **60**, 766–770.
- 164 J. Baldwin, A. Brookfield, G. F. S. Whitehead, L. S. Natrajan, E. J. L. McInnes, M. S. Oakley and D. P. Mills, *Inorg. Chem.*, 2024, **63**, 20295–20306.
- 165 O. P. Churchill, A. Dase, L. J. Taylor, S. P. Argent, N. T. Coles, G. S. Walker and D. L. Kays, *Inorg. Chem.*, 2024, **63**, 20286–20294.
- 166 B. Ross, J. Belz and M. Nieger, *Chem. Ber.*, 1990, **123**, 975–978.
- 167 L. Weber and G. Meine, *Chem. Ber.*, 1987, **120**, 457–459.
- 168 L. Weber, G. Meine, R. Boese and D. Bläser, *Chem. Ber.*, 1988, **122**, 853–857.
- 169 G. Lacorre and H. Grützmacher, *Dalton Trans.*, 2022, **51**, 3497–3501.

