



Cite this: *Chem. Commun.*, 2020, 56, 8553

Advancing the fundamental understanding and practical applications of photo-bioelectrocatalysis

Matteo Grattieri,  Kevin Beaver,  Erin M. Gaffney,  Fangyuan Dong  and Shelley D. Minteer *

Photo-bioelectrocatalysis combines the natural and highly sophisticated process of photosynthesis in biological entities with an abiotic electrode surface, to perform semi-artificial photosynthesis. However, challenges must be overcome, from the establishment and understanding of the photoexcited electron harvesting process at the electrode to the electrochemical characterization of these biotic/abiotic systems, and their subsequent tuning for enhancing energy generation (chemical and/or electrical). This Feature Article discusses the various approaches utilized to tackle these challenges, particularly focusing on powerful multi-disciplinary approaches for understanding and improving photo-bioelectrocatalysis. Among them is the combination of experimental evidence and quantum mechanical calculations, the use of bioinformatics to understand photo-bioelectrocatalysis at a metabolic level, or bioengineering to improve and facilitate photo-bioelectrocatalysis. Key aspects for the future development of photo-bioelectrocatalysis are presented alongside future research needs and promising applications of semi-artificial photosynthesis.

Received 13th April 2020,
Accepted 17th June 2020

DOI: 10.1039/d0cc02672g

rsc.li/chemcomm

1. Introduction

Over billions of years of evolution, nature developed an elegant approach to convert solar energy into chemical energy, while

generating oxygen, by means of photosynthesis.^{1,2} Inspired by nature, scientists developed synthetic strategies to artificially perform the photosynthetic process, utilizing sunlight to drive the water-splitting reaction and to reduce an abundant electron acceptor (*i.e.*, CO₂) into valuable compounds (*i.e.*, CO, formaldehyde, methanol). This process is defined as “artificial photosynthesis”,^{3–6}

Department of Chemistry, University of Utah, 315 S 1400 E Rm 2020, Salt Lake City, UT 84112, USA. E-mail: minteer@chem.utah.edu



Matteo Grattieri

Matteo is currently a Postdoctoral Research Associate at the University of Utah, working in the group of Shelley D. Minteer. He focuses on interfacing biological entities with electrodes, and his research interests span the fields of extracellular electron transfer in photosynthetic and halotolerant bacteria, artificial redox mediating systems, and biosensors development. Prior to join the University of Utah, he studied Chemistry in Milano (Università degli Studi di Milano and Politecnico di Milano), and he was a visiting researcher at the University of New Mexico with Plamen Atanassov and at the University of Buenos Aires with Ernesto J. Calvo.



Kevin Beaver

Kevin is a first-year graduate student pursuing his PhD at the University of Utah, under the mentorship of Shelley D. Minteer and Aaron Puri. With dual BS degrees in Biochemistry & Molecular Biology and Environmental Science, his research combines bioengineering and microbial electrochemistry for environmental applications, including renewable energy and bioremediation. Kevin began his bioelectrochemistry research with Michelle Rasmussen at Lebanon Valley College (Annville, PA). He also was mentored by Joseph Kieber (University of North Carolina at Chapel Hill) and Shelley D. Minteer as part of the National Science Foundation Research Experience for Undergraduate (NSF-REU) program.

and provides the opportunity to fabricate robust catalysts,⁷ as well as the use of ligand design to tune them for selectivity. Furthermore, artificial light absorbers can be synthesized to cover a broad portion of the solar spectrum, increasing their light-to-product conversion efficiency. However, the outstanding performance of artificial photosynthesis often comes at the cost of high-purity catalysts. Additionally, several molecular catalysts performing artificial photosynthesis under aqueous conditions require noble metals (*i.e.*, ruthenium), sacrificial reagents, and present challenges for their heterogenization that limits the performance of molecular-catalyst-based electrodes. We refer the reader to a recent review discussing advantages and challenges of artificial photosynthesis.⁸ Focusing on the natural photosynthesis, chlorophyll-based light harvesting at the basis of photosynthesis in plants, algae, and bacteria, covers only a specific part of the solar spectrum that varies depending on the photosynthetic entity utilized, thus significantly decreasing the final efficiency of the process. However, the natural photosynthetic apparatuses of biological entities

offer the unique advantage of having sophisticated self-repair mechanisms,⁹ as well as mechanisms to protect against excessive illumination.¹⁰ Another critical advantage of biological entities is the ability to utilize sunlight to drive complex reactions performed by highly sophisticated “enzymatic machinery”, which is not yet feasible with synthetic approaches, particularly under mild aqueous conditions. In order to take advantage of the benefits of both natural and artificial photosynthesis, while minimizing their drawbacks, there has been increasing interest in combining abiotic systems with photosynthetic biological entities to perform semi-artificial photosynthesis.¹¹ Accordingly, the goal of semi-artificial photosynthesis is to merge the strengths of natural photosynthesis (*i.e.*, selectivity towards complex products and regeneration features), and artificial photosynthesis (*i.e.*, improved light absorption, efficiency towards a desired pathway, facilitated charge transfer). Furthermore, it should be considered that plants, algae, purple bacteria, and cyanobacteria constitute an abundant, diverse, and broadly available and sustainable source of bio-photocatalysts, paving the way for industrialization of photo-bioelectrochemical systems. The recent progressions in synthetic biology could further expand the synthetic diversity enabled by biohybrid systems, making semi-artificial photosynthesis even more appealing with exciting new research possibilities. As discussed in detail below, a variety of biological entities have been coupled to an electrode surface, having various advantages and strengths.^{12–14} In this Feature Article, the basics of photo-bioelectrocatalysis are first introduced, exploring intriguing possibilities provided by biohybrid systems for the renewable energy field. Thereafter, the specific challenges for the application of biological entities in photo-bioelectrochemical systems are presented. The other sections of the article focus on discussing recent works performed to address the presented challenges, posing particular focus on the importance of multidisciplinary approaches, which combine the knowledge of chemists, biologists, computational scientists, material scientists and engineers. We strongly believe that only by gaining a better understanding of the fundamental aspects of



Erin M. Gaffney

Erin is currently a third-year graduate student pursuing a PhD in Analytical Chemistry at the University of Utah, working in the group of Shelley D. Minteer. Her research focuses on microbial electrochemical system development. She aims to elucidate extracellular electron transfer mechanisms in halotolerant bacteria through a combination of bioinformatics and electrochemical characterization. Previously she has obtained her bachelor's degree in Biochemistry from Fort Lewis College under the research supervision of Prof. Kathryn D. Mouzakis.

Erin is currently a third-year graduate student pursuing a PhD in Analytical Chemistry at the University of Utah, working in the group of Shelley D. Minteer. Her research focuses on microbial electrochemical system development. She aims to elucidate extracellular electron transfer mechanisms in halotolerant bacteria through a combination of bioinformatics and electrochemical characterization. Previously she has obtained her bachelor's degree in Biochemistry from Fort Lewis College under the research supervision of Prof. Kathryn D. Mouzakis.



Fangyuan Dong

Fangyuan is currently a graduate student at the University of Utah, working in the group of Shelley D. Minteer. She focuses on bioelectrocatalysis and bioelectrosynthesis system development, and her research interests span the fields of biphasic asymmetric bioelectrocatalysis, microbial bioelectrosynthesis through bioengineering, and cofactor regeneration. Prior to join the University of Utah, she studied Chemistry in Shanghai at East China Normal University under the research supervision of Prof. Yang Tian.

Fangyuan is currently a graduate student at the University of Utah, working in the group of Shelley D. Minteer. She focuses on bioelectrocatalysis and bioelectrosynthesis system development, and her research interests span the fields of biphasic asymmetric bioelectrocatalysis, microbial bioelectrosynthesis through bioengineering, and cofactor regeneration. Prior to join the University of Utah, she studied Chemistry in Shanghai at East China Normal University under the research supervision of Prof. Yang Tian.



Shelley D. Minteer

Shelley is the Dale and Susan Poulter Endowed Chair in Biological Chemistry and the Associate Chair of the Department of Chemistry at University of Utah. She received her PhD in Analytical Chemistry at University of Iowa, followed by positions as Assistant Professor, Associate Professor, Professor, and CAS Endowed Professor of Chemistry at Saint Louis University before moving to University of Utah in 2011. Her research interests are in electrocatalysis and bioelectrocatalysis for sensing, energy, and synthesis applications.

Shelley is the Dale and Susan Poulter Endowed Chair in Biological Chemistry and the Associate Chair of the Department of Chemistry at University of Utah. She received her PhD in Analytical Chemistry at University of Iowa, followed by positions as Assistant Professor, Associate Professor, Professor, and CAS Endowed Professor of Chemistry at Saint Louis University before moving to University of Utah in 2011. Her research interests are in electrocatalysis and bioelectrocatalysis for sensing, energy, and synthesis applications.

photo-bioelectrocatalysis, will it be possible to pave the way for the employment of semi-artificial photosynthesis in the field of renewable energy.

Photo-bioelectrocatalysis and the future of the renewable energy field

Perhaps, when analyzing the relationship between photo-bioelectrocatalysis and renewable energy, the first aspect that should be considered is that Earth receives solar energy in the range of 120 000–178 000 TW (where 1 TW corresponds to 10^{12} W).^{15–17} For comparison, the worldwide primary energy consumption in 2018 was in the range of 635×10^{18} J, for an average energy consumption rate of approximately 20 TW. This value is projected to rise to about 30 TW by 2050.^{18,19} Thus, it is clear that the world energy requirement would be largely exceeded if sustainable approaches to efficiently harvest solar energy became available. Silicon-based photovoltaics have reached impressive efficiencies for solar to electrical energy conversion and widespread worldwide application.²⁰ However, in the effort to increase the amount of usable solar energy, multiple approaches should be employed, expanding beyond the sunlight to electricity conversion. Semi-artificial photosynthesis provides several pathways for the conversion of solar energy and sustainable alternatives to common energy-intensive processes, including the sunlight-driven synthesis of valuable compounds from broadly available substrates and monitoring water decontamination with sunlight-powered biosensors.

Basics of photo-bioelectrocatalysis

Photo-bioelectrocatalysis covers a broad research field, as it refers to the employment of biological entities capable of bridging sunlight energy and electrical energy, while catalyzing a chemical reaction at an electrode surface. Three main components are at play during photo-bioelectrocatalysis (Fig. 1): (i) light-harvesting complexes (LHCs), which harvest light energy (photons); (ii) protein complexes (green ellipse) that convert light into a charge-separated state after receiving the excitation energy (generation of an electron–hole pair at high energy); and (iii) an electrode acting as the terminal electron acceptor (or primary electron donor for a photocathode) for the redox reactions enabled by the charge-separated state. The high-energy electron obtained after the charge-separation event undergoes a series of electron transfer processes in the metabolisms of photosynthetic microorganisms and plants. At the same time, the high-energy hole allows the oxidation of various substrates, which span from water to organic carbon depending on the photosynthetic entity, as discussed in Section 3. By diverting the photosynthetically generated electrons to the electrode (or from the electrode to the photosynthetic electron transfer chain), the process of photo-bioelectrocatalysis can be selectively tuned toward facilitating specific reactions, opening opportunities for several energy-related applications.

Practical applications of photo-bioelectrocatalysis

Below, some reports of practical applications of photo-bioelectrocatalysis are presented to provide a broad overview




Fig. 1 Scheme of the main components for photo-bioelectrocatalysis. Light harvesting complexes (LHCs), protein complexes enabling the charge-separation event (green ellipse), electrode acting as electron acceptor in a biophotoanodes (for electrical energy generation, black arrows), or as an electron donor in a biophotocathode (for chemical energy generation, blue dashed arrows).

of the recent trends and opportunities offered by this research field. Both isolated photosynthetic apparatuses and intact organisms have been employed for the sunlight-driven online sensing of various toxic compounds, avoiding the requirement of a power source, and providing an easily measured signal.²¹ In some previous work from our group, the photo-bioelectrocatalytic features of both intact cyanobacteria or isolated thylakoids membranes were employed for the online detection of herbicides.²² A self-powered system obtained by coupling a thylakoid-based photoanode with an air-breathing cathode enabled detection of three commercial herbicides at concentrations below $1 \mu\text{g L}^{-1}$.²³ On-going experiments in our group are focused on expanding the range of photosynthetic entities that can be utilized to develop self-powered system for the detection of herbicides, focusing on the sustainability of the entire photo-bioelectrochemical system. Recent reports from other groups showed the possibility to employ both isolated reaction centers or intact organisms for the detection of herbicides and other toxic compounds,²⁴ as well as a self-powered system with microalgae.²⁵

Moving to the possibility to utilize photobioelectrocatalysis for sustainable power generation, considerable improvements have been reported employing cyanobacterial mutants in biophotovoltaic systems, which allowed the achievement of a power density of 0.5 W m^{-2} .^{26,27}

Another extremely interesting application of photo-bioelectrocatalysis is to accomplish the bioelectrosynthesis of valuable products with high energy efficiency compared to the synthetic approaches commonly utilized. Leddy and co-workers initially reported the possibility to electrochemically stimulate the cyanobacterium *Anabaena variabilis* SA-1 (*A. variabilis*) immobilized on an electrode to enhance ammonia production.²⁸ As discussed in detail below, we recently studied this photo-bioelectrocatalytic process to unveil the redox intermediates

allowing sunlight driven ammonia production.²⁹ Other photosynthetic entities were also utilized by other groups for product synthesis, such as purple bacteria in a bioelectrochemical system allowing sustainable H₂ production and CO₂ reduction utilizing wastewater as substrate.³⁰ Self-assembly of photosynthetic entities in protocells (artificial cell models) enabled obtaining highly oriented photosynthetic reaction centers from *Rhodobacter sphaeroides* (*R. sphaeroides*) in giant unilamellar vesicles. The orientation of the reaction centers allowed the generation of a proton gradient that could be used for ATP synthesis in a biohybrid photoelectrochemical system.^{31,32}

2. Challenges of semi-artificial photosynthesis

The opportunities offered by photo-bioelectrocatalysis for converting sunlight energy come with several critical challenges to be overcome. Fig. 2 presents some of the challenges, which will be specifically discussed in this Feature Article. The first and, perhaps, most important challenge of photo-bioelectrocatalysis is gaining a deeper understanding of the photoexcited electron harvesting process at an electrode surface. While this is, relatively, easier when isolated photosystems are employed, extremely little information is available for the case of more complex apparatuses or intact organisms. Gaining a comprehensive understanding of the process would allow specifically tackling the various bottlenecks, thus optimizing the process. Another fundamental challenge is to understand how environmental parameters (*i.e.*, temperature, pH, pollutants, salinity, *etc.*)

influence photo-bioelectrocatalysis. This is critical for the future applications of semi-artificial photosynthesis in industrial scenarios. Microbial photo-bioelectrocatalysis could be employed for energy recovery, decontamination, and monitoring of industrial wastewater that encounter fluctuations in salinity, temperature, pH, and contaminant concentrations. Additionally, for enzymatic photo-bioelectrocatalysis systems, changes in solvent polarity, ionic strength, and temperature can challenge the applications.

Expanding to the more applicative field of photo-bioelectrocatalysis, several other challenges can be identified. First, the photosynthetic apparatus of biological entities has been optimized by nature to utilize a specific wavelength of the incoming photons. However, from an efficiency point of view, maximizing the range of usable light is critical.

As we consider maximizing the efficiency of the process, another critical aspect is to artificially facilitate the photoexcited electron transfer (PEET) to the electrode surface. In fact, biological entities are not designed to exchange electrons with an external solid electron acceptor (*i.e.*, the electrode), and for the case of intact microorganisms, to funnel charge transfer to the desired pathway rather than to biomass accumulation. When focusing on product synthesis, an important challenge is defined by the need to tune the semi-artificial photosynthesis process in order to selectively obtain a desired product, which would greatly facilitate its purification and isolation. Finally, to make photo-bioelectrochemical systems appealing to the industrial community, it will be critical to enhance their stability to enable long-term application, and to develop cost-effective systems (considering both their assembly and operation).

As discussed below, there is no single approach that would allow solving all of the presented challenges. However, the implementation of multi-disciplinary approaches has been recently reported in literature, providing critical insights toward exciting future research directions.

3. Understanding the photo-excited electron harvesting process and the effect of environmental parameters

Isolating photosynthetic apparatuses constitutes a model system for the study of the PEET with an electrode surface, thanks to their simpler structure and the possibility to directly deposit them on the electrode surface. Various photosynthetic entities are initially presented, starting from the most up to date understanding of their electrical communication with an electrode surface. The case of intact photosynthetic microorganisms is then presented in a detailed section, which initially describes anoxygenic (purple bacteria) and oxygenic (cyanobacteria) microorganisms, and concludes with the case of eukaryotic algae. It is important to note that when utilizing intact bacterial cells or algae, the challenge of accomplishing the PEET with an electrode surface is magnified by the presence of various membranes separating the photosynthetic apparatus and the electrode, as well as by the metabolism of the microorganisms requiring




Fig. 2 Challenges of semi-artificial photosynthesis. (top) Understanding the photoexcited electron transfer and the effects of environmental parameters, as well as the redox intermediate at play to shuttle the photoexcited electrons between the photosynthetic apparatus (both isolated or *in vivo*) and the electrode surface. (bottom) Developing approaches to artificially tune the photoexcited electron transfer to the electrode surface and the efficiency of the light conversion process.

electrons for their own sustenance. As a result, the fundamental aspects of the PEET process for this category of photosynthetic entities remain largely unexplored.

The approaches utilized for the detailed study of photo-bioelectrocatalysis discussed below include: (i) bioengineering for selective inhibition (or removal) of redox intermediates of the photosynthetic electron transfer chain to study their involvement in the PEET process; (ii) theoretical models to predict the concentrations of redox species in the biohybrid photoelectrochemical systems; (iii) localized electrochemical techniques (*i.e.*, scanning (photo)electrochemical microscopy); (iv) quantum mechanical calculations that enable studying structures and states of the various components involved in photo-bioelectrocatalysis (*i.e.*, redox mediators, electrodes, substrates, and products) to probe reaction mechanisms; (v) bioinformatic tools to unveil gene and transcript expression in organisms exposed to various stimuli/conditions to study their influence on the PEET process.

3.1 Thylakoids

Among the model systems for investigating photo-bioelectrocatalysts are thylakoids, pigmented membranous compartments localized within the chloroplasts of plant and algal cells, and situated free in the cytosol of cyanobacteria. As shown in Fig. 3, the thylakoid membrane consists of a series of two photosystems functioning in tandem: photosystem II and I (PSII and PSI, respectively). The oxidation of water to oxygen in oxygenic photosynthesis can be accomplished at PSII thanks to three main components: (i) a chlorophyll dimer referred to as P680, which has an extremely high midpoint redox potential of 1.1–1.4 V *vs.* SHE; (ii) a tyrosine residue; and (iii) the tetranuclear Mn cluster that stores four-oxidative equivalents and reduces the tyrosine residue.³³ PSII has both light-harvesting and charge-separation functions, utilizing ambient light to catalyze the oxidation of water. After the charge-separation

event, the photoexcited electrons are shuttled down an electron transport chain to PSI. From there, a second electron transport chain shuttles the electrons from PSI to ferredoxin, a small protein containing a single 2Fe–2S center, to catalyze the reduction of nicotinamide adenine dinucleotide phosphate (NADP⁺) to NADPH, which powers the metabolisms of plants, algae, and cyanobacteria. This photosynthetic electron transfer is commonly referred to as the “Z scheme”.

Thylakoids can be isolated from a diversity of organisms. Accordingly, our group investigated the photo-bioelectrocatalytic properties of thylakoids obtained from different sources (including chard and beets),³⁴ showing that spinach thylakoids yielded the highest current density. Following, the efforts from our group were focused on unveiling the detailed electron transfer process of thylakoid membranes with an abiotic carbon paper electrode.³⁵ A multidisciplinary approach was utilized, combining electrochemical studies with the bioengineering of the thylakoids membranes by removal, activation, and inhibition of several components involved in light-dependent reactions (plastoquinone, ferredoxin, or ferredoxin–NADP⁺ reductase). Interestingly, it was possible to show that direct electron transfer (DET) can be obtained, where both PSII and PSI contribute to the biophotocurrent, along with a significant contribution from plastoquinone, the cytochrome *b₆f* complex, and plastocyanin. For the sake of clarity, we define DET between a biological entity and an electrode as a condition where no exogenous redox mediator is added to the investigated system, and the redox molecule responsible for the establishment of the electron transfer is not an endogenous redox-active compound secreted outside the cellular membrane of a living organism (*i.e.*, *Shewanella oneidensis* that secretes riboflavin and uses it as a redox shuttle between the membrane-bound cytochromes and an electrode surface³⁶). Conversely, mediated electron transfer (MET) is obtained in the presence of exogenous redox mediators artificially added to the system (*i.e.*, diffusible redox mediators such as methyl viologen, or redox polymers), or




Fig. 3 Scheme of the electron transfer chain in thylakoid membrane: first PSII catalyses the photo-driven water splitting reaction to generate protons, oxygen and electrons. After a series of electron transfer steps, the water-derived electrons are transferred to the plastoquinone pool (PQ and PQH₂ for the oxidized and reduced forms, respectively) and downstream to cytochrome *b₆f* complex. The electrons thereby flow to PSI via the electron carrier plastocyanin (PC). Following the electrons are passed to ferredoxin (Fd). In the noncyclic pathway, the electrons reduce NADP⁺ with ferredoxin–NADP⁺ reductase (Z-scheme photosynthetic electron transfer chain). In the cyclic pathway, the reduced Fd transfers the electrons to PQ, completing the cycle. The proton gradient created by water photolysis and electron movements from the overall process powers ATP synthase to regenerate ATP.

endogenous redox-active compounds secreted outside the cellular membrane of a living organism.

3.2 Isolated reaction centers and photosystems

3.2.1 Isolated reaction centers from purple bacteria. The bacterial reaction center (RC) can be considered the photochemical core of photosynthesis.³⁷ It is a transmembrane pigment-protein complex, which was first isolated in 1968 from the purple bacterium *R. sphaeroides*.³⁸ A scheme of the RC is reported in Fig. 4A (purple ellipse). Thanks to the relatively easy isolation, simplicity, and good stability, the RC has been utilized as a model to study the photosynthetic process.^{39–45} The bacterial reaction center and photosystem II of plants, algae, and cyanobacteria show various similarities, and the RC served as a structural and functional model for the study of photosystem II. The discussion of the differences and similarities between these two complexes has been reviewed in previous literature, and we refer the reader to the work of Allen and co-workers for an extensive discussion of these aspects.⁴⁶ In brief, the RC is found in anoxygenic organisms, while PSII is found in oxygenic organisms. As previously introduced in Section 3.1, the chlorophyll dimer in photosystem II, P680, has an extremely high midpoint redox potential of 1.1–1.4 V vs. SHE,

while its counterpart in the RC, P870, has a considerably lower midpoint redox potential of 0.5 V vs. SHE. Furthermore, compared to the tyrosine residue that reduces P680 in PSII, a cytochrome docks to the RC and leaves after transferring one electron to P870. Due to these differences, water splitting cannot be accomplished in the bacterial reaction center, and only the oxidation of molecules with lower redox potentials can be performed (*i.e.*, malic acid). Another difference is that PSII is a considerably larger complex than the RC. In fact, PSII has both light-harvesting and charge-separation functions, while two light-harvesting complexes are necessary to capture and direct light to the RC.

The redox processes taking place between the RC and diffusible quinone redox mediators were investigated in detail by scanning electrochemical microscopy (SECM).⁴⁷ The study allowed measuring kinetic constants for the interaction between RC and mediators, which highlighted the importance of the chemical environment in which the RC is embedded. In a recent study from the same group, the photo-bioelectrocatalytic process for solubilized reaction centers was described by means of a theoretical model, unveiling kinetic parameters and physicochemical factors affecting the light conversion process.⁴⁸ The importance of achieving preferential orientation of the RC on indium tin oxide electrodes modified with cytochrome *c* for the



Fig. 4 (A) Purple bacteria photosynthetic electron transfer chain. Light harvesting complexes (LHC); the reaction center (purple ellipse, RC) where a “special pair”, consisting of two bacteriochlorophyll molecules, absorbs light at 870 nm (P870); an excited state of the “special pair” (P870*); monomeric bacteriochlorophyll (BChl); bacteriopheophytin (BPh); a primary quinone acceptor (Q_A); a secondary quinone acceptor (Q_B); the quinone pool (Q_P); cytochrome *bc1* (Cyt *bc1*); cytochrome *c2* (Cyt *c2*). From the LHC, the excitation energy is transferred to the “special pair” of bacteriochlorophyll in the RC, leading to the charge separation event that is then followed by electron transfer steps, ultimately resulting in the reduction of quinone molecules in the quinone pool. A reduced quinone molecule transfers an e^- and a proton to the Cyt *bc1* complex that generates a proton gradient for ATP synthesis and a reduction of Cyt *c2*, thereby regenerating the cycle. Purple bacteria have alternative electron sinks to divert excess of electrons, through N_2 fixation, CO_2 fixation or dimethyl sulfoxide (DMSO) reduction. Adapted with permission from E. M. Gaffney, M. Grattieri, K. Beaver, J. Pham, C. McCartney and S. D. Minter, Unveiling salinity effects on photo-bioelectrocatalysis through combination of bioinformatics and electrochemistry, *Electrochim. Acta*, 2020, **337**, 135731. Copyright 2020 Elsevier.⁶³ (B) The photosynthetic apparatus of purple bacteria is embedded in the inner membrane of the bacterial cell, requiring a mediator to cross both the outer and inner membranes in order to harvest the photoexcited electrons at an electrode surface. Adapted from M. Grattieri, Purple bacteria photo-bioelectrochemistry: enthralling challenges and opportunities, *Photochem. Photobiol. Sci.*, 2020, **19**, 424–435¹² with permission from the European Society for Photobiology, the European Photochemistry Association, and The Royal Society of Chemistry.

generation of a photovoltage has also been discussed in a recent work.⁴⁹

3.2.2 Isolated photosystem II. The light-activated PSII protein is the benchmark natural photocatalyst for the oxidation of water, allowing the development of biophotoanodes operating at low potential (close to 0 V vs. SHE).⁵⁰ For comparison, the only other enzyme reported to accomplish the water-splitting reaction, laccase, requires an electron acceptor at very high potentials (about 0.95 V vs. SHE).⁵¹ Isolated photosystems have been reported to enable facile PEET with an electrode surface, a great advantage compared to thylakoids or intact photosynthetic organisms, where the photoreactive centers are buried under membranes and other cell components. With the aim to clarify how the isolation of PSII facilitates photo-bioelectrocatalysis and affects stability, a very interesting study reported a comparison between the photo-bioelectrocatalytic response of PSII *in vitro* versus *in vivo*, where cyanobacteria biofilms were deposited on the electrode surface.⁵² While the PSII *in vitro* enabled photocurrent about 30-fold higher compared to PSII *in vivo*, the latter showed high stability and an increasing photocurrent generation over time. A recent study employed scanning photoelectrochemical microscopy (SPECM) to unveil the capability of redox polymers utilizing osmium redox moieties to effectively mediate electron transfer from both the Q_A and Q_B site to the electrode surface.⁵³

3.2.3 Isolated photosystem I. Conversely from PSII, which is utilized in biophotoanodes, PSI can be employed also for developing biophotocathodes. As previously introduced, PSI allows the photo-excitation of an electron that is necessary for the reduction of the energy-carrying cofactor NADP⁺. The direct electron transfer between isolated PSI and an electrode surface was studied in detail by Ciobanu *et al.* utilizing modified gold electrodes.⁵⁴ Later, the same group studied the mediated electron transfer between PSI and an electrode surface, exploring both metal-based and organic redox mediators, and clarifying the influence of electrochemical and physico-chemical properties of the mediators on photocurrent production.⁵⁵ It was shown that utilizing redox mediators that absorb light in the same regions of PSI significantly lowered the photocurrent response, particularly for those absorbing in the red region (650–750 nm) corresponding to the Q_y transition band of chlorophyll a. Furthermore, the photocurrent from PSI-modified gold electrodes could be correlated to the relative redox potentials of the PSI redox centers and the formal redox potential ($E^{0'}$) of the soluble redox mediators utilized, with a favorable trend for the redox mediators with more positive $E^{0'}$. Finally, metal-based mediators outperformed organic compounds, possibly due to a faster electron transfer rate between the metal-based mediators and PSI cofactors. Accordingly, the study provided critical findings for the development of PSI-based biohybrid photoelectrochemical systems.

An outstanding 3-D architecture was reported by Ciornii *et al.* where PSI is combined with a redox enzyme and cytochrome *c* resulting in a well-defined interprotein electron transfer allowing different pathways to switch on and off based on experimental conditions.⁵⁶ Recently, Robinson *et al.* reported an electrochemical reaction–diffusion model for PSI

multilayer film atop a metal electrode that allowed shedding light on the electrochemical, physico-chemical, and transport processes taking place at the photoelectrode.⁵⁷ The reader is also referred to a recent review discussing the recent trends in PSI-based biophotovoltaics.⁵⁸

3.3 Intact photosynthetic microorganisms

Advantageous to bioelectrochemical systems is the use of whole-cells on an electrode surface due to the enhanced long term stability, owing to the ability of microorganisms to self-replicate and repair. Whole-bacterial cells can easily be deposited on an electrode surface and form bacterial communities known as biofilms. The biofilm is composed of the whole-cells and self-secreted extracellular polymeric substances that function as structural support and protection from the external environment, providing enhanced resistance to inhibiting agents.⁵⁹ Electroactive biofilms arise from having electroactive microorganisms present in the biofilm, which establish the extracellular electron transfer with the electrode surface.^{60,61}

3.3.1 Purple bacteria. Purple bacteria are anoxygenic photosynthetic organisms that utilize sunlight as an energy source to catalyze the oxidation of various organic and inorganic substrates thanks to the RC (previously introduced in Section 3.2.1). These bacteria are characterized by an outer and an inner membrane, with a periplasm layer in between, and their photosynthetic apparatus is embedded in the inner membrane of the bacterial cells, as shown in Fig. 4B. As a result, the electron transfer to the electrode surface is considerably hindered, and only limited photocurrents have been reported in the absence of redox mediators, which is attributed to the limited interaction between the ubiquinol and the electrode surface.⁶²

Accordingly, particular interest has been posed on unveiling the capability of diffusible redox mediators to cross the membranes and reach the photosynthetic apparatus. A pioneering study was performed by Cai *et al.* in 2002, where SECM was used to clarify the properties of redox molecules capable of crossing both the outer and inner membrane.⁶⁴ A class of reduced hydrophobic redox mediators crossing the inner cellular membrane to react with redox components of the photosynthetic electron transfer chain was defined. Furthermore, the study unveiled the potential of the redox cofactor interacting with the diffusible redox mediators being in a range of -110 ± 90 mV vs. Ag|AgCl. Another interesting aspect was that, utilizing the formal redox potential of the selected redox mediators, a correlation between the heterogeneous rate constant and the redox potential could not be obtained, pointing to the presence of additional limitations in the PEET process. Recently, our group focused on combining electrochemical experimental evidence and quantum mechanical calculations, specifically density functional theory (DFT), to clarify the rate-determining step in the PEET process.⁶⁵ First, the electronic and structural parameters of various quinone forms were obtained by DFT calculations to investigate their redox processes. Then, the computationally-derived electrochemical properties of the mediators were utilized to fit the

experimentally measured current density obtained from purple bacteria employing the various redox mediators during malic acid oxidation. It was possible to identify a model correlating to the PEET rates, unveiling that the rate determining step is the one-electron, proton-decoupled electron transfer taking place in the lipophilic membrane of purple bacteria. The result also suggests that the mediators interact, accepting one electron, from the active site Q_A in the photosynthetic electron transfer chain, rather than the quinone pool, a finding that is supported also from the SECM results of Cai *et al.*⁶⁴ In view of future studies, modeling the binding between biological entities and electrodes by means of quantum mechanical calculations would provide the possibility to gain an atomic-level understanding of the biotic/abiotic interface, which has been reported for redox proteins on carbon-based electrodes.^{66,67}

The extreme versatile metabolism of purple bacteria allows for their employment also in contaminated environments,⁶⁸ paving the way for the use of photo-bioelectrocatalysis for the light-driven degradation of various wastewater pollutants. However, determining the effects of environmental parameters—such as the presence of contaminants, or variations in pH, salinity, and temperature—on photo-bioelectrocatalysis from a fundamental point of view is critical to elucidate the PEET taking place under environmentally relevant conditions. Extreme variations in pH, temperature, and salinity are harmful to non-tolerant bacteria; however, small variations can have drastic effects on the metabolisms of bacteria. Additionally, tolerant bacteria such as extremophilic organisms, provide a useful platform to study and overcome these conditions.⁶⁹ Although gene expression and metabolic changes vary widely among species, temperature and pH changes generally result in a variation of the lipid membrane composition.⁷⁰ Few studies elucidated these effects in photosynthetic microorganisms, with the effects of salinity studied at the genomic level. Salinity is a critical parameter affecting the metabolism of biological entities. As a result, studies have been focused on quantifying gene expression in purple bacteria exposed to saline conditions to unveil their adaptation mechanisms. *R. sphaeroides* was shown to increase the expression of glycine–betaine uptake systems, due to glycine–betaine acting as an osmotic protectant.⁷¹ Although the electrochemical performance of *R. sphaeroides* has not been evaluated in high salinity, closely related purple bacterium, *Rhodobacter capsulatus* (*R. capsulatus*), has been shown to have interesting electrochemical performance resulting from saline adaptation. In a study from our group, it was shown that *R. capsulatus* adaptation to salinity, and the resulting increased photo-bioelectrocatalysis, occurred only with gradual increases in salinity achieved during several growth cycles.⁷² Following, bioinformatic analysis was performed and coupled to electrochemical evidence to analyze differential gene expression for understanding the adaptation process from a metabolic point of view.⁶³ Several key metabolic groups showed differences in their expression values between the adapted and non-adapted strains. The genes important to photo-bioelectrocatalysis included those involved in the photosynthetic electron transfer chain and the nitrogen

metabolisms of this bacterium. With saline adaptation, there was a decrease in expression of key genes in the photosynthetic electron transport chain, including cytochrome *c2*, cytochrome *c550*, and phytoene synthase. Additionally, decreased expression of several genes involved in nitrogen metabolism regulations and many genes in the actual nitrogen metabolism revealed a key link between these systems. The nitrogen metabolism is a known electron sink of the photosynthetic electron transfer chain, as shown in Fig. 4A. This justifies its over-expression during the hindered photosynthetic electron transfer inferred from the decreased photocurrent produced by the saline-adapted strain. Interestingly, the overexpression of an osmoprotectant system offered the ability to artificially tune for rapid saline tolerance of this bacterium and an instant increase in photo-bioelectrocatalysis when compared to the non-adapted strain.

Transcriptomic approaches were utilized also by Volpicella *et al.* to clarify the mechanism of adaptation for *R. sphaeroides* to high concentrations of cobalt ions at a metabolic level.⁷³ It was possible to reveal the involvement of energetic metabolisms on the capability of the bacterial cells to cope with high cobalt concentrations. Such a detailed understanding of the effects of contaminants on the metabolisms of photosynthetic organisms could pave the way for the development of biohybrid systems where the interaction between bacterial cells and the electrode surface could be used to tune the metabolism in a fashioned way.

Expanding the study of the effects of environmental parameters on the PEET process, the simultaneous presence of contaminants and redox mediators in a solution containing *R. capsulatus* cells was recently investigated by Borghese *et al.*⁷⁴ Interestingly, the use of a diffusible naphthoquinone mediator allowed *R. capsulatus* cells to extracellularly reduce tellurite, which also enabled the production of tellurium nanoparticles.

3.3.2 Cyanobacteria. Cyanobacteria, conversely from purple bacteria, are oxygenic photosynthetic organisms that account for 20–30% of worldwide primary photosynthetic productivity,⁷⁵ converting solar energy at the rate of ~ 450 TW.¹⁶ These bacteria have evolved with impressive light-harvesting systems and charge-generation capabilities, making them an outstanding candidate for solar energy conversion. Light-dependent electrogenic activity has been observed within various genera of cyanobacteria, including biofilm-forming and pelagic strains. Cyanobacteria are characterized by their unique thylakoid membranes, containing protein complexes for both the photosynthetic electron transport chain and respiratory electron transport chain (Fig. 3). Previous studies have shown that electrons can flow between the photosynthetic and respiratory systems *via* the shared plastoquinone pool.⁷⁶ This allows the surplus light energy quenched by PSII to be transferred to the quinol oxidases of the respiratory electron transport chain on thylakoid membranes, alleviating photo-damage of the photosynthetic proteins. Therefore, both a better stability and a longer lifetime on electrode surfaces are achieved compared to isolated photosystems of higher plants, where photosynthesis takes place in the chloroplast and respiration in the mitochondria. However, the entangled respiratory chain and photosynthetic chain on the cyanobacteria thylakoid membrane

also add complexity to the cellular electron transfer pathway. Groundbreaking studies have been done to elucidate the electrogenic pathway in cyanobacteria using site-specific inhibitors and mutations that target components of photosynthetic electron transport chain,^{77–79} and electron paramagnetic resonance.⁸⁰ Another interesting aspect is that under intense light, the rate of electron production at PSII will surpass consumption by the Calvin cycle, and consequently, the plastoquinone-carried electrons will build up in the thylakoid membrane. Those overflow electrons can be directly transmitted to cytochrome *bd*-quinol oxidase and possibly flow into a route that contributes to the photocurrent generation of cyanobacteria.⁸¹

As many cyanobacteria exhibit the natural capability of electrochemical communication with an electrode surface, our group explored the possibility to utilize electrochemical stimulation to enhance ammonia generation from a mutant strain of *A. variabilis* when immobilized on an indium tin oxide (ITO) electrode.²⁹ The ammonium assimilation step by glutamine synthetase in the *A. variabilis* mutant strain SA-1 is blocked, hence ammonium will be constantly produced instead of regulated by cellular requirements. A bioelectrocatalytic signal was observed in the presence of light on the *A. variabilis* SA-1 modified ITO electrode, and nitrogenase was determined as the enzyme that most of the redox moiety is communicating within *A. variabilis* SA-1. The intriguing part is that nitrogenase is not positioned to be capable of direct electron transfer with the electrode, indicating that an internal mediation system is at play. To clarify the electron transfer process, the electrolyte obtained after the electrochemical characterization of *A. variabilis* SA-1 in the presence of N₂ and light was collected. UV-vis absorption spectra of the concentrated electrolyte closely matched the spectrum of commercial ferredoxin (Fig. 5), which supported the electrochemical evidence of a redox couple attributed to ferredoxin. Accordingly, the electrostimulated ammonia production by *A. variabilis* SA-1 could be due to endogenously produced ferredoxin exiting cells to mediate electron transfer between the ITO electrode and nitrogenase.

DET from cyanobacteria to the electrode through membrane bound cytochromes has also been reported, a process similar to that of metal reducing microorganisms.^{77,82} A recent study investigated the role of PSII, PSI, and the respiratory metabolism of *Synechocystis* cells on the obtained photocurrent.⁸³ It was shown that the respiratory metabolism could supply electrons to the PSI which could then photo-reduce the anode, generating the photocurrent. Additionally, bacterial conductive nanowires have also been discovered to facilitate the EET process in the cyanobacterium *Synechocystis* sp. PCC-6803 using scanning tunnelling microscopy.⁸⁴

As discussed above for purple bacteria, environmental stress effects need to be elucidated for the successful deployment of photo bioelectrochemical systems, and genome analysis is an excellent platform to study. In a study by RübSam *et al.*, the mRNA levels of *Synechocystis* sp. PCC-6803 were analysed in response to saline and iron depletion. It was shown that this strain accumulates the compatible solute glucosylglycerol to overcome saline stress, and an iron-stress activated RNA1 that




Fig. 5 Comparison of UV-vis absorption spectra for concentrated spent electrolyte collected after electrochemical characterization of an ITO electrode modified with *A. variabilis* SA-1 in N₂ purged buffer in the presence of light (solid black line, absorbance plotted on left) and commercial ferredoxin (green dashed line, absorbance plotted on right). Reproduced with permission from K. L. Knoche, E. Aoyama, K. Hasan and S. D. Minteer, Role of Nitrogenase and Ferredoxin in the Mechanism of Bioelectrocatalytic Nitrogen Fixation by the Cyanobacteria *A. variabilis* SA-1 Mutant Immobilized on Indium Tin Oxide (ITO) Electrodes, *Electrochim. Acta*, 2017, **232**, 396–403. Copyright 2017 Elsevier.²⁹

is directly involved with the photosynthetic electron transport chain.⁸⁵ Although the resulting electrochemical performance of *Synechocystis* sp. PCC-6803 has not been evaluated with these environmental stress factors, it is likely that due to the difference in the mRNA expression, both saline and iron could play a role in its photo-bioelectrochemical performance.

3.3.3 Eukaryotic algae. Unlike cyanobacteria, eukaryotic algae contain membrane-bound organelles, including a chloroplast that houses stacks of thylakoids called grana, like in plant cells. Hence, in these systems, PEET to an electrode surface can be mainly hindered due to the extra and intracellular membranes to be crossed, in addition to the aqueous compartments. Accordingly, research has been focused on determining the capability of exogenous redox mediators to harvest the photoexcited electrons.^{86,87} The group of Frédéric Lemaître showed that diffusible 2,6-dichlorobenzoquinone allowed effective harvesting of photoexcited electrons from a suspension of a mutant green algae *Chlamydomonas reinhardtii*.⁸⁸ The mutant allowed electron harvesting from only PSII in the presence of the exogenous mediator, achieving biophotocurrents up to 60 $\mu\text{A cm}^{-2}$. In a later study, the same group investigated photo-bioelectrocatalysis of the same mutant algae suspension over time, to unveil photoinhibition and cytotoxicity effects.⁸⁹ Recently, they reported an interesting approach for studying the interaction between algae and diffusible redox mediators combining electrochemistry and pulse amplitude modulation fluorescence.⁹⁰ The approach unveiled the importance of quenching properties of the redox mediator, distinguishing the photoexcited electron harvesting from some side effects due to quinones in real time. Finally, the possibility of direct PEET between algae and electrodes has also been

recently investigated. By employing the genus *Choricystis* spp. and performing additions of boric acid, current densities on the order of $15 \mu\text{A cm}^{-2}$ could be obtained.⁹¹ The DET was attributed to the electron carriers plastoquinone, ferredoxin, and plastocyanin.

4. Artificial approaches for upgraded photo-bioelectrocatalysis

In view of utilizing photo-bioelectrocatalysis for practical applications, it is critical to maximize the efficiency of the process, both in terms of current density and desired product synthesis. Current densities tend to be lower in whole-cell systems than cell-component systems, due to resistances of charge transport and mass transport from additional membranes. Furthermore, sunlight energy conversion efficiencies are also lower, due to respiration competing with the photosynthetic electron transfer pathways and nutrients requirement for growth to continue. However, an advantage of using intact cells is exploiting their regulatory and replicative machinery to increase the overall lifetime and stability of the bioanode (or biocathode). Accordingly, research efforts have been focused on developing artificial approaches to facilitate the PEET process and to drive the electron flux towards desired pathways. A scheme comparing the performance of the various photosynthetic biological entities in biohybrid systems is reported in Fig. 6.

4.1 Artificial redox mediating systems

The group of Lo Gorton performed pioneering studies on wiring purple bacteria with an electrode surface utilizing redox polymers. The artificial systems enabling electron-shuttling employed redox moieties constituted by osmium functionalities. Interestingly, it was shown that by modifying the chemical structure of these redox moieties, the photoexcited electron harvesting could be tuned, achieving biophotocurrents in the range of $10 \mu\text{A cm}^{-2}$.^{62,92}

Our group focused particular efforts on developing artificial mediating systems for diverting the photoexcited electrons to an abiotic electrode. Specifically, inspired by the use of quinone redox moieties in biological processes, we developed a redox polymer containing naphthoquinone redox moieties immobilized on a linear polyethyleneimine backbone.⁹³ The redox polymer was first efficiently employed to enhance the electron transfer between an electrode surface and various enzymes.⁹⁴ Following, the capability of the polymer to mediate electron transfer from intact bacterial cells was demonstrated as well.^{95,96} Inspired by these results, we explored the possibility to utilize the redox polymer for harvesting photoexcited electrons.⁹⁷ Finely controlling the ratio of bacterial cells and redox moieties in a 3-D redox hydrogel geometry allowed establishing PEET with a low overpotential, achieving $\sim 3 \mu\text{A cm}^{-2}$.

We also expanded the applicability of this naphthoquinone-based redox polymer to oxygenic photosynthetic entities, specifically for wiring intact chloroplasts (Fig. 7A).⁹⁸ Electrical wiring of chloroplasts with the redox polymer allowed a five-fold increase in




Fig. 6 Scheme of efficiencies, stabilities, and current generations for isolated bacterial reaction centers (RC), photosystem II and I (PSII and PSI, respectively), thylakoid membranes (TM), purple bacteria (PB), cyanobacteria (CB), and algae. Red circles are indicative and not to scale.

photocurrent generation compared to the current obtained under DET conditions. Furthermore, a mixed redox polymer-diffusible mediator system was explored, by adding dichlorobenzoquinone to the system. The new artificial redox mediation resulted in an outstanding increase of the photocurrent, enabling an additional five-fold increase for the harvested current, as shown in Fig. 7B.

Expanding to polymers employing different redox moieties, osmium redox polymers have been recently utilized to achieve the electrical wiring of both PSI (used at the biophotocathode) and PSII (used at the biophotoanode) obtaining a fully light-driven semi-artificial electrochemical system that performed water splitting at the anode while obtaining hydrogen evolution at the cathode.⁹⁹ The combination of an artificial redox mediating system with osmium moieties and an engineered hierarchically-structured indium tin oxide electrode surface was reported by Sokol *et al.*¹⁰⁰ The efficient wiring by the redox polymer and the higher loading enabled by the engineered electrode allowed achieving the outstanding biophotocurrent of $410 \mu\text{A cm}^{-2}$.

Regarding cyanobacteria photo-bioelectrocatalysis, major interest has been focused on facilitating cross-membranes electron transportation. Accordingly, various studies on photosynthetic microbial fuel cells employed exogenous electron mediators, using both lipid-insoluble (*i.e.*, ferricyanide), and lipid-soluble (*i.e.*, cytochrome, 2,6-dichloro-1,4-benzoquinone) molecule to harvest photoexcited electrons from cyanobacteria.^{83,101,102} However, lipid-soluble mediators can disrupt electron transfer processes used for important cellular events within the organism, and eventually cause cell death. Conversely, lipid-insoluble mediators, like ferricyanide ion, accepting electrons from the cell surface offered a more bio-compatible choice over long durations.¹⁰² The combination of an osmium-based redox polymer and soluble ferricyanide allowed achieving a biophotocurrent of about $48 \mu\text{A cm}^{-2}$ from cyanobacteria *Lyptolyngbia* sp. (CYN82).¹⁰³ Finally, for the case of isolated RC-LHC complexes, the combined used of diffusible ubiquinones and cytochrome *c* as redox mediators allowed achieving photocurrents in the range of $45 \mu\text{A cm}^{-2}$.




Fig. 7 (A) Artificial PEET process through the naphthoquinone-based redox polymer that harvests photoexcited electrons from intact chloroplast and shuttles them to the electrode surface. (B) Comparison of photocurrent generation from chloroplasts in direct electron transfer condition (TP + Chl, black), in the presence of the naphthoquinone redox polymer (TP + NQ-LPEI + Chl, red), and in the presence of both the redox polymer and diffusible dichloro benzoquinone (TP + NQ-LPEI + DCBQ + Chl, green). Adapted with permission from K. Hasan, R. D. Milton, M. Grattieri, T. Wang, M. Stephanz and S. D. Minteer, Photobioelectrocatalysis of Intact Chloroplasts for Solar Energy Conversion, *ACS Catal.*, 2017, 7, 2257–2265. Copyright 2017 American Chemical Society.⁹⁸

4.2 Engineering of the biotic/abiotic interface

Crucial to the success of photo-bioelectrochemical systems is improving the performance of the abiotic electrode and its biocompatibility for the photosynthetic biological entity utilized. For each division of biophotocatalysts, pairing of nanoscale materials with a biological entity provides the opportunity for overall current and system enhancement. The use of a hierarchically-structured inverse opal indium tin oxide as support for PSII hybrid electrodes with Os-based redox polymer has been previously introduced.¹⁰⁰ Recently, the use of mesoporous NiO surface as a p-type semiconductor in conjunction with PSI has been reported to obtain a biophotocathode capable of tandem operation with PSI-TiO₂ biophotoanode, for a biophotovoltaic cell with open-circuit potential of 0.7 V.¹⁰⁴

To enhance the current density produced by thylakoid photoanodes, multi-walled carbon nanotubes (CNTs) are often mixed in a suspension with thylakoids before depositing on the surface of the electrode. The structure of CNTs increases the electrode surface area, while their hydrophobicity helps to immobilize the thylakoids on the surface, and their combined employment with both diffusible mediators and redox polymer has been reported. Exogenous quinone-based mediators and CNTs-modified thylakoids electrodes were explored by Takeuchi *et al.*, testing a variety of quinone-based and pyridine-based mediators, greatly enhancing the obtained photocurrent.¹⁰⁵ The research group of Lo Gorton has employed CNTs in their thylakoid-modified electrodes, where rather than using diffusible quinone-based mediators, an osmium-based redox polymer was used to immobilize thylakoids on graphite electrodes and facilitate mediated electron transfer.¹⁰⁶ This quinone-free system allowed biophotocurrents densities similar to those reported with diffusible mediators (on the order of 100 $\mu\text{A cm}^{-2}$). We also refer the reader to a recent review highlighting the advancements in thylakoid membranes photobioelectrochemical systems.¹⁰⁷ The use of carbon nanotubes to enhance photocurrent production was

also explored for cyanobacteria photoanodes. Specifically, carbon nanotube modified electrodes served as a tool to bridge the PEET from membranes to the electrode surface.^{108,109} Additionally, cyanobacterium *Leptolyngbia* sp. was paired with a Au electrode modified with a conducting polymer and Au nanoparticles to increase current performance in a photo-biofuel cell.¹¹⁰ Recently, Wolfe *et al.* explored the effect of different three-dimensional electrode morphologies on the photocurrent production from PSL.¹¹¹ Interestingly, macroporous electrodes (5 μm porous) outperformed mesoporous electrodes (20–100 nm porous) when in the presence of diffusible mediators, achieving biophotocurrents in the range of 40 $\mu\text{A cm}^{-2}$.

When pairing whole-cell entities to abiotic electrodes, it is of interest to pair the microorganisms as closely to the electrode surface as possible to minimize ohmic losses. Nanoscale engineering of the abiotic electrode surface offers one approach toward this goal. In a recent work by Cornejo *et al.*, the combination was achieved on the nanoscale with the incorporation of a tin oxide inorganic catalyst layer, and a silica oxide layer embedded with molecular wires interfaced with the outer membrane cytochromes of *Shewanella oneidensis* MR-1, as represented in Fig. 8.¹¹² Although this work was not with photosynthetic microorganisms, the approach was a great advancement in wiring bacterial cells with the electrode surface, which could be extended for applications in biophotovoltaics. Recently, work has also been done to improve the solar-powered CO₂ reduction at the cathode of a biophotovoltaic cell for acetate production. Interfacing silicon nanowires with *Sporomusa ovata* cells allowed for development of a close-packed nanowire-bacteria cathode that, along with optimization of electrolyte buffering capacity and pH, exhibited a 7-day efficiency of 3.6% for solar to acetate conversion.¹¹³

4.3 Bioengineering of biological entities for enhancing PEET

Due to the advances in genomic editing tools, the possibility for engineering strains to provide electroactivity by inserting



Fig. 8 Wiring of *Shewanella oneidensis* MR-1 to an electrode surface. The pairing of outer membrane cytochromes, a SnO₂ membrane with -CN and -NO₂ functionalized *p*-oligo(phenylene vinylene) molecular wires, and a SnO₂ coated Pt electrode (a) and the corresponding potentials of each (b). Reproduced with permission from J. Cornejo, H. Sheng, E. Edri, C. Ajo-Franklin, and H. Frei, Nanoscale membranes that chemically isolate and electronically wire up the abiotic/biotic interface, *Nat. Commun.*, 2018, **9**, 2263. Copyright 2018 Springer Nature with a Creative Commons Attribution 4.0 International License (<http://creativecommons.org/licenses/by/4.0/>).

critical genes has attracted particular interest.¹¹⁴ The most well-studied extracellular electron transfer pathways from model electrogenic microorganisms are the mtrCAB pathway from *Shewanella oneidensis* MR-1,¹¹⁵ and outer membrane cytochrome (omc) genes from *Geobacter sulfurreducens*, specifically omcS.¹¹⁶ These metabolic pathways can then be engineered into microorganisms, transforming an easily-grown but non-electrogenic strain, such as *Escherichia coli*, into a strain capable of communicating with and wiring to an electrode. In this section, a brief overview of the recent synthetic biology approaches for purple bacteria and cyanobacteria will be given, and we refer the reader to a recent review describing the bioengineering of photovoltaics in detail.¹¹⁷ For non-photosynthetic microorganisms, electricity generation has been achieved from the pairing of the oxidation of organic substrates to the current generation at an electrode. Specifically, the model genetically modified microorganism, *Escherichia coli*.¹¹⁸ Similar approaches are being explored to pair the photosynthetic process with an electrode through bioengineering of these electrogenic pathways into photosynthetic microorganisms. As discussed above, the large limitation for purple bacteria and cyanobacteria is the transfer of electrons through the thick cellular membrane to an electrode surface. The mediation approaches discussed face

challenges of non-specific electron extraction due to their reactivity with any redox intermediate present with a thermodynamically-favored potential.¹¹⁷ For specific and selective electron harvesting from the photosynthetic electron chain, bioengineering extracellular electron transfer pathways into these microorganisms offer a favorable approach. Genetic bioengineering for enhanced extracellular electron transfer has been achieved in two strains of photosynthetic microorganisms *Synechocystis* sp. PCC 6803, and *Synechococcus elongatus* PCC 7942. In a study by Bradley *et al.*,⁷⁶ the terminal oxidases, *bd*-quinol oxidase, cytochrome *c* oxidase, and an “alternative respiratory terminal oxidase”, were deleted in *Synechocystis* sp. PCC 6803. Without these terminal oxidases, electrons could not be utilized for O₂ reduction and would be available for use in the photosynthetic electron transport chain. The deletion resulted in a 10% increase in power generation in dark conditions. In *Synechococcus elongatus* PCC 7942, Sekar *et al.* engineered the strain to contain the omcS gene from *Geobacter sulfurreducens* in order to express an outer membrane cytochrome protein, OmcS.¹¹⁹ The engineered strain enabled a nine-fold increase in photocurrent (reaching about 15 μA cm⁻²) compared to the non-engineered strain. In a subsequent work by the same group, OmcS was expressed while also deleting the terminal oxidases in *Synechococcus elongatus* PCC 7942, resulting in a increased current when compared to both the wildtype strain and the deleted terminal oxidases mutant strain.¹²⁰

Bioengineering of photosynthetic microorganisms has multiple challenges, including the difficulty of synthetic biology techniques, and unknown metabolic effects when compared to model microorganisms such as *Escherichia coli*.¹¹⁷ Despite these challenges, the advances in understanding these extracellular electron transfer pathways in the model organisms allow methods to be applied to photosynthetic bacteria, offering increased applications in bioelectrochemical systems.

4.4 Tuning product selectivity

The metabolic pathways and enzyme networks that characterize intact microorganisms provide the possibility to perform complex chemistry, channeling substrates along reaction cascades. Accordingly, selectively driving a specific metabolism to obtain a desired product in a biohybrid system with intact microorganisms involves modifying the flux of redox carriers and ions between, and within, the electrode, the electrolyte, and the microorganisms. Both genetic and bioelectrochemical tools can play a critical role in achieving these modifications.¹²¹ Genetic tools for metabolic engineering have been extensively developed for model organisms, such as *Escherichia coli*,^{122,123} showing the selective bioelectrochemical production of succinate from CO₂.¹²⁴ Conversely, limited toolsets are available for non-model strains, such as photosynthetic microorganisms, and future efforts to develop synthetic biology approaches for these entities will be required to tune photo-bioelectrocatalysis for product synthesis.¹²⁵

The use of bioelectrochemical approaches to finely tune product synthesis has been investigated to a greater degree than genetic approaches. We have previously discussed how electrostimulation

allowed the enhancement of ammonia generation from a mutant strain of *A. variabilis*.²⁹ Recently, the use of electron-regulated flux by means of polarized electrodes has been reported also to increase CH₄ formation and CO₂ utilization in a bioelectrochemical system for electromethanogenesis.¹²⁶ Additionally, active modification of electron fluxes through electrochemical polarization was recently utilized to maximize H₂ production and CO₂ fixation in a photo-microbial electrochemical system with purple bacteria.³⁰

4.5 Artificial light-harvesting complexes and biohybrid systems

Enhancing the usable portion of light, by expanding the absorption region of photosynthetic entities, could enable higher efficiency for the energy conversion process, further facilitating the implementation of semi-artificial photosynthesis. In biological photosynthetic entities, light is collected and funneled to the photosystems by pigment-protein complexes (*i.e.*, the light-harvesting complexes), which are also referred to as “antennas”. A significant amount of research has been focused on developing artificial antennas to develop hybrid systems with tuned light-harvesting efficiency. In a pioneering study, Nabiev *et al.* reported the coupling of photoluminescent quantum dots to transfer the energy from an excited electron to the photosynthetic reaction center of the purple bacterium *R. sphaeroides* by Förster resonance energy transfer (FRET).¹²⁷ Following, Milano *et al.* reported a tailored molecular organic dye, absorbing at 450 nm and emitting at 602 nm, where the RC has an absorption peak (Fig. 9B). A succinimidyl ester group enables selective covalent binding of the synthesized fluorophore to the lysines of the photosynthetic reaction center of *R. sphaeroides* R26, as shown in Fig. 9A.¹²⁸

Such molecular antennas expanded the tuning possibility for the hybrid system, thanks to the diversity of molecular compounds and their shape and flexibility, reducing the impact of the artificial antennas on the structure and function of the photosynthetic reaction center. The same group further extended the capability to utilize artificial molecular antennas absorbing light in the visible region (400–700 nm) and emitting in correspondence of the near infra-red absorption peak of the reaction center (800 nm) in the following works.^{129,130} Another study utilized commercially available fluorescent dyes molecules with a genetically modified purple bacteria reaction center.¹³¹ The use of organic antennas to increase light absorption was extended to the PSI of the thermophilic cyanobacterium *Thermosynechococcus elongates* targeting the “green gap” (*i.e.*, the lack of light absorption in the range 450–600 nm),¹³² as well as for the light-harvesting complex LH2, obtained from a strain of *Rhodospseudomonas acidophila*, expanding light-harvesting to the 620–750 nm range.¹³³ Artificial antennas have also been coupled to intact organisms. Specifically, the cyanobacterium *Synechococcus* sp. was combined with quantum dots to enhance the current generation under low light conditions through FRET. A nanocomposite of CdTe quantum dots (broadening the excitation spectrum to 350 nm), graphene nanoplatelets, and silk-fibroin cast on a graphite electrode enabled a 5.7 fold increase in biophotocurrent.¹³⁴ Recently, Grayson *et al.* reported that artificial tailor-made light-harvesting antennas can be created *in vivo*, by



Fig. 9 (A) Crystallographic structure of the *Rhodobacter sphaeroides* R26 reaction center. The ubiquinone-10 molecules (Q_A, Q_B), two bacterio-pherophytins, and four bacteriochlorophylls are in pink. Lysines are van der Waals green spheres. The terminal N_ϵ lysine targets of bioconjugation are in blue. The detergent toroid surrounding the hydrophobic portion of the protein is shown as a grey cylinder, including the detergent molecules (not to scale). The structure of the activated organic fluorophore antenna is shown in the upper left corner (C, cyan; N, blue; O, red; and S, yellow). The antenna and the RC structures are to scale. (B) Absorption spectra of the RC (black) and the artificial antenna (red). Fluorescence spectrum of the antenna (blue). Adapted with permission from F. Milano, R. R. Tangorra, O. Hassan Omar, R. Ragni, A. Operamolla, A. Agostiano, G. M. Farinola and M. Trotta, Enhancing the Light Harvesting Capability of a Photosynthetic Reaction Center by a Tailored Molecular Fluorophore, *Angew. Chem., Int. Ed.*, 2012, **51**, 11019–11023. Copyright 2012 Wiley-VCH Verlag GmbH & Co. KGaA.

genetically modifying the photosystem of *R. sphaeroides* to incorporate the yellow fluorescent protein.¹³⁵

In two recent works, Liu *et al.* utilized a very interesting approach to combine the features of oxygenic photosynthesis in plants with the anoxygenic photosynthesis of purple bacteria, augmenting the near-infrared solar energy conversion of the latter to the visible light spectrum.^{136,137}

5. Conclusions and future outlook for research in photo-bioelectrocatalysis

Employing multidisciplinary approaches has provided unique tools to critically expand our understanding of photo-bioelectrocatalysis. The combination of quantum mechanical

calculations and electrochemical experimental evidence enabled us to probe reaction mechanisms taking place in biohybrid systems composed of intact bacterial cells, exogenous redox mediators, and abiotic electrodes, which are otherwise particularly challenging to be studied. While the combination of quantum mechanical methods and microbial electrochemical systems can be considered in its infancy, it could provide critical insights for future developments of the field, both in terms of shedding light on biotic/abiotic materials interactions and reaction mechanisms between biological entities and exogenous molecules. On a similar note, the insights provided by bioinformatics studies allow the understanding of photo-bioelectrocatalysis at a metabolic level. As shown for purple bacteria adaptation to salinity, understanding the metabolic changes resulting from the adaptation process allows artificially tuning the adaptation. Expanding to other environmentally relevant parameters, the knowledge provided by bioinformatic studies can be used to enable rational tuning of detailed metabolisms to drive the biohybrid systems towards desired directions. It is critical to underline that in order to accomplish this goal, bioengineering of the biological entities will play a critical role. The possibility to selectively insert heterologous electron transfer pathways, or to modify the natural metabolic pathways of photosynthetic entities provides unprecedented research possibilities in order to perform multi-step reactions in a fashioned way.

Accurately combining synthetic biology with engineering of the biotic/abiotic interface and artificial redox mediating systems, as well as implementing artificial light-harvesting complexes, will provide powerful tools to improve the efficiency of the sunlight conversion process. Such results will critically promote the implementation of semi-artificial photosynthesis for practical applications.

In conclusion, a critical aspect to be underlined is that no single approach will enable the successful development and implementation of photo-bioelectrocatalysis, but rather the combination of the multidisciplinary approaches presented will provide the unique features to accomplish this task. A strong collaboration of the community working in the photo-bioelectrochemical field, including (bio)electrochemists, biologists, engineers, organic chemists, computational scientists, material scientists, and biophysicist is vital to provide the required tools to fully explore photo-bioelectrocatalysis of biological entities and enable the development of biohybrid photosynthetic systems with unprecedented capabilities.

Conflicts of interest

There are no conflicts to declare.

Acknowledgements

The authors would like to acknowledge the National Science Foundation Grant # 1561427.

Notes and references

- 1 D. E. Canfield, *Annu. Rev. Earth Planet. Sci.*, 2005, **33**, 1–36.
- 2 S. A. Crowe, L. N. Dossing, N. J. Beukes, M. Bau, S. J. Kruger, R. Frei and D. E. Canfield, *Nature*, 2013, **501**, 535–538.
- 3 S. Y. Reece, J. A. Hamel, K. Sung, T. D. Jarvi, A. J. Esswein, J. J. H. Pijpers and D. G. Nocera, *Science*, 2011, **334**, 645–648.
- 4 D. G. Nocera, *Acc. Chem. Res.*, 2012, **45**, 767–776.
- 5 A. Pannwitz and O. S. Wenger, *Chem. Commun.*, 2019, **55**, 4004–4014.
- 6 K. E. Dalle, J. Warnan, J. J. Leung, B. Reuillard, I. S. Karmel and E. Reisner, *Chem. Rev.*, 2019, **119**, 2752–2875.
- 7 A. Sartorel, M. Carraro, F. M. Toma, M. Prato and M. Bonchio, *Energy Environ. Sci.*, 2012, **5**, 5592–5603.
- 8 B. Zhang and L. Sun, *Chem. Soc. Rev.*, 2019, **48**, 2216–2264.
- 9 S. Jarvi, M. Suorsa and E. M. Aro, *Biochim. Biophys. Acta*, 2015, **1847**, 900–909.
- 10 C. Ota, K. Sugihara, Y. Kinoshita, Y. Kashiyama, Y. Nagasawa and H. Tamiaki, *Photochem. Photobiol. Sci.*, 2019, **18**, 64–70.
- 11 N. Kornienko, J. Z. Zhang, K. K. Sakimoto, P. Yang and E. Reisner, *Nat. Nanotechnol.*, 2018, **13**, 890–899.
- 12 M. Grattieri, *Photochem. Photobiol. Sci.*, 2020, **19**, 424–435.
- 13 K. Hasan, V. Grippo, E. Sperling, M. A. Packer, D. Leech and L. Gorton, *ChemElectroChem*, 2017, **4**, 412–417.
- 14 F. Milano, A. Punzi, R. Ragni, M. Trotta and G. M. Farinola, *Adv. Funct. Mater.*, 2019, **29**, 1805521.
- 15 R. E. Blankenship, D. M. Tiede, J. Barber, G. W. Brudvig, G. Fleming, M. Ghirardi, M. R. Gunner, W. Junge, D. M. Kramer, A. Melis, T. A. Moore, C. C. Moser, D. G. Nocera, A. J. Nozik, D. R. Ort, W. W. Parson, R. C. Prince and R. T. Sayre, *Science*, 2011, **332**, 805–809.
- 16 J. M. Pisciotta, Y. Zou and I. V. Baskakov, *PLoS One*, 2010, **5**, e10821.
- 17 O. Kruse, J. Rupprecht, J. H. Mussgnug, G. C. Dismukes and B. Hankamer, *Photochem. Photobiol. Sci.*, 2005, **4**, 957–970.
- 18 U.S. Energy Information Administration, U.S. Department of Energy, Washington, DC, 2019. <https://www.eia.gov/outlooks/ieo/pdf/ieo2019.pdf>.
- 19 N. S. Lewis and D. G. Nocera, *Proc. Natl. Acad. Sci. U. S. A.*, 2006, **103**, 15729–15735.
- 20 K. Yoshikawa, H. Kawasaki, W. Yoshida, T. Irie, K. Konishi, K. Nakano, T. Uto, D. Adachi, M. Kanematsu, H. Uzu and K. Yamamoto, *Nat. Energy*, 2017, **2**, 17032.
- 21 M. Grattieri and S. D. Minter, *ACS Sens.*, 2018, **3**, 44–53.
- 22 M. Tucci, M. Grattieri, A. Schievano, P. Cristiani and S. D. Minter, *Electrochim. Acta*, 2019, **302**, 102–108.
- 23 M. Rasmussen and S. D. Minter, *Anal. Methods*, 2013, **5**, 1140–1144.
- 24 M. Chatzipetrou, F. Milano, L. Giotta, D. Chirizzi, M. Trotta, M. Massaouti, M. R. Guascito and I. Zergioti, *Electrochem. Commun.*, 2016, **64**, 46–50.
- 25 J. Chouler, M. D. Monti, W. J. Morgan, P. J. Cameron and M. Di Lorenzo, *Electrochim. Acta*, 2019, **309**, 392–401.
- 26 K. L. Saar, P. Bombelli, D. J. Lea-Smith, T. Call, E.-M. Aro, T. Müller, C. J. Howe and T. P. J. Knowles, *Nat. Energy*, 2018, **3**, 75–81.
- 27 M. Grattieri and S. D. Minter, *Nat. Energy*, 2018, **3**, 8–9.
- 28 J. Leddy and T. M. Pachekewitz, *US Pat.*, US 9506085 B2, University of Iowa Research Foundation, United States, 2016.
- 29 K. L. Knoche, E. Aoyama, K. Hasan and S. D. Minter, *Electrochim. Acta*, 2017, **232**, 396–403.
- 30 I. A. Vasiliadou, A. Berná, C. Manchon, J. A. Melero, F. Martinez, A. Esteve-Núñez and D. Puyol, *Front. Energy Res.*, 2018, **6**, 107.
- 31 E. Altamura, F. Milano, R. R. Tangorra, M. Trotta, O. H. Omar, P. Stano and F. Mavelli, *Proc. Natl. Acad. Sci. U. S. A.*, 2017, **114**, 3837–3842.
- 32 J. P. Allen, *Proc. Natl. Acad. Sci. U. S. A.*, 2017, **114**, 3790–3791.
- 33 F. Rappaport, M. Guergova-Kuras, P. J. Nixon, B. A. Diner and J. Lavergne, *Biochemistry*, 2002, **41**, 8518–8527.
- 34 M. Rasmussen, A. Wingersky and S. D. Minter, *Electrochim. Acta*, 2014, **140**, 304–308.
- 35 M. Rasmussen and S. D. Minter, *Electrochim. Acta*, 2014, **126**, 68–73.
- 36 E. Marsili, D. B. Baron, I. D. Shikhare, D. Coursolle, J. A. Gralnick and D. R. Bond, *Proc. Natl. Acad. Sci. U. S. A.*, 2008, **105**, 3968–3973.
- 37 A. Operamolla, R. Ragni, F. Milano, R. Roberto Tangorra, A. Antonucci, A. Agostiano, M. Trotta and G. Farinola, *J. Mater. Chem. C*, 2015, **3**, 6471–6478.

- 38 D. W. Reed and R. K. Clayton, *Biochem. Biophys. Res. Commun.*, 1968, **30**, 471–475.
- 39 J. P. Allen, G. Feher, T. O. Yeates, H. Komiya and D. C. Rees, *Proc. Natl. Acad. Sci. U. S. A.*, 1988, **85**, 8487–8491.
- 40 G. Feher, J. P. Allen, M. Y. Okamura and D. C. Rees, *Nature*, 1989, **339**, 111–116.
- 41 T. O. Yeates, H. Komiya, A. Chirino, D. C. Rees, J. P. Allen and G. Feher, *Proc. Natl. Acad. Sci. U. S. A.*, 1988, **85**, 7993–7997.
- 42 R. A. Isaacson, F. Lendzian, E. C. Abresch, W. Lubitz and G. Feher, *Biophys. J.*, 1995, **69**, 311–322.
- 43 J. P. Allen and J. C. Williams, *FEBS Lett.*, 1998, **438**, 5–9.
- 44 F. Milano, A. Agostiano, F. Mavelli and M. Trotta, *Eur. J. Biochem.*, 2003, **270**, 4595–4605.
- 45 F. Ciriaco, R. R. Tangorra, A. Antonucci, L. Giotta, A. Agostiano, M. Trotta and F. Milano, *Eur. Biophys. J.*, 2015, **44**, 183–192.
- 46 L. Kalman, J. C. Williams and J. P. Allen, *Photosynth. Res.*, 2008, **98**, 643–655.
- 47 F. Longobardi, P. Cosma, F. Milano, A. Agostiano, J. Mauzeroll and A. J. Bard, *Anal. Chem.*, 2006, **78**, 5046–5051.
- 48 F. Milano, F. Ciriaco, M. Trotta, D. Chirizzi, V. De Leo, A. Agostiano, L. Valli, L. Giotta and M. R. Guascito, *Electrochim. Acta*, 2019, **293**, 105–115.
- 49 M. Di Lauro, G. Buscemi, M. Bianchi, A. De Salvo, M. Berto, S. Carli, G. M. Farinola, L. Fadiga, F. Biscarini and M. Trotta, *MRS Adv.*, 2020, **5**, 985–990.
- 50 M. Kato, J. Z. Zhang, N. Paul and E. Reisner, *Chem. Soc. Rev.*, 2014, **43**, 6485–6497.
- 51 M. Pita, D. M. Mate, D. Gonzalez-Perez, S. Shlevv, V. M. Fernandez, M. Alcalde and A. L. De Lacey, *J. Am. Chem. Soc.*, 2014, **136**, 5892–5895.
- 52 J. Z. Zhang, P. Bombelli, K. P. Sokol, A. Fantuzzi, A. W. Rutherford, C. J. Howe and E. Reisner, *J. Am. Chem. Soc.*, 2018, **140**, 6–9.
- 53 F. Zhao, V. Hartmann, A. Ruff, M. M. Nowaczyk, M. Rögner, W. Schuhmann and F. Conzuelo, *Electrochim. Acta*, 2018, **290**, 451–456.
- 54 M. Ciobanu, H. A. Kincaid, V. Lo, A. D. Dukes, G. K. Jennings and D. E. Cliffel, *J. Electroanal. Chem.*, 2007, **599**, 72–78.
- 55 G. Chen, G. LeBlanc, G. K. Jennings and D. E. Cliffel, *J. Electrochem. Soc.*, 2013, **160**, H315–H320.
- 56 D. Ciornii, M. Riedel, K. R. Stieger, S. C. Feifel, M. Hejazi, H. Lokstein, A. Zouni and F. Lisdat, *J. Am. Chem. Soc.*, 2017, **139**, 16478–16481.
- 57 M. T. Robinson, D. E. Cliffel and G. K. Jennings, *J. Phys. Chem. B*, 2018, **122**, 117–125.
- 58 M. T. Robinson, E. A. Gizzie, F. Mwambutsa, D. E. Cliffel and G. K. Jennings, *Curr. Opin. Electrochem.*, 2017, **5**, 211–217.
- 59 Y. Xiao and F. Zhao, *Curr. Opin. Electrochem.*, 2017, **4**, 206–211.
- 60 D. R. Bond and D. R. Lovley, *Appl. Environ. Microbiol.*, 2003, **69**, 1548–1555.
- 61 P. Cristiani, M. L. Carvalho, E. Guerrini, M. Daghigh, C. Santoro and B. Li, *Bioelectrochemistry*, 2013, **92**, 6–13.
- 62 K. Hasan, K. V. R. Reddy, V. Eßmann, K. Górecki, P. Ó. Conghaile, W. Schuhmann, D. Leech, C. Hägerhäll and L. Gorton, *Electroanalysis*, 2015, **27**, 118–127.
- 63 E. M. Gaffney, M. Grattieri, K. Beaver, J. Pham, C. McCartney and S. D. Minteer, *Electrochim. Acta*, 2020, **337**, 135731.
- 64 C. Cai, B. Liu, M. V. Mirkin, H. A. Frank and J. F. Rusling, *Anal. Chem.*, 2002, **74**, 114–119.
- 65 M. Grattieri, Z. Rhodes, D. P. Hickey, K. Beaver and S. D. Minteer, *ACS Catal.*, 2019, **9**, 867–873.
- 66 S. Babanova, I. Matanovic, M. S. Chavez and P. Atanassov, *J. Am. Chem. Soc.*, 2015, **137**, 7754–7762.
- 67 I. Matanovic, S. Babanova, M. S. Chavez and P. Atanassov, *J. Phys. Chem. B*, 2016, **120**, 3634–3641.
- 68 R. Blasco and F. Castillo, *Appl. Environ. Microbiol.*, 1992, **58**, 690–695.
- 69 B. E. Logan, R. Rossi, A. Ragab and P. E. Saikaly, *Nat. Rev. Microbiol.*, 2019, **17**, 307–319.
- 70 M. F. Siliakus, J. van der Oost and S. W. M. Kengen, *Extremophiles*, 2017, **21**, 651–670.
- 71 M. Tsuzuki, O. V. Moskvina, M. Kuribayashi, K. Sato, S. Retamal, M. Abo, J. Zeilstra-Ryalls and M. Gomelsky, *Appl. Environ. Microbiol.*, 2011, **77**, 7551–7559.
- 72 M. Grattieri, K. Beaver, E. Gaffney and S. D. Minteer, *Faraday Discuss.*, 2019, **215**, 15–25.
- 73 M. Volpicella, A. Costanza, O. Palumbo, F. Italiano, L. Claudia, A. Placido, E. Picardi, M. Carella, M. Trotta and L. R. Ceci, *FEMS Microbiol. Ecol.*, 2014, **88**, 345–357.
- 74 R. Borghese, M. Malferrari, M. Brucale, L. Ortolani, M. Franchini, S. Rapino, F. Borsetti and D. Zannoni, *Bioelectrochemistry*, 2020, **133**, 107456.
- 75 M. Paumann, G. Regelsberger, C. Obinger and G. A. Peschek, *Biochim. Biophys. Acta*, 2005, **1707**, 231–253.
- 76 R. W. Bradley, P. Bombelli, D. J. Lea-Smith and C. J. Howe, *Phys. Chem. Chem. Phys.*, 2013, **15**, 13611–13618.
- 77 J. M. Pisciotta, Y. Zou and I. V. Baskakov, *Appl. Microbiol. Biotechnol.*, 2011, **91**, 377–385.
- 78 A. G. Roberts, M. K. Bowman and D. M. Kramer, *Biochemistry*, 2004, **43**, 7707–7716.
- 79 J. Yan, G. Kurisu and W. A. Cramer, *Proc. Natl. Acad. Sci. U. S. A.*, 2006, **103**, 69–74.
- 80 B. V. Trubitsin, V. V. Ptushenko, O. A. Koksharova, M. D. Mamedov, L. A. Vitukhnovskaya, I. A. Grigor'ev, A. Y. Semenov and A. N. Tikhonov, *Biochim. Biophys. Acta*, 2005, **1708**, 238–249.
- 81 D. Pils and G. Schmetterer, *FEMS Microbiol. Lett.*, 2001, **203**, 217–222.
- 82 A. J. McCormick, P. Bombelli, A. M. Scott, A. J. Phillips, A. G. Smith, A. C. Fisher and C. J. Howe, *Energy Environ. Sci.*, 2011, **4**, 4699–4709.
- 83 G. Saper, D. Kallmann, F. Conzuelo, F. Zhao, T. N. Toth, V. Liveanu, S. Meir, J. Szymanski, A. Aharoni, W. Schuhmann, A. Rothschild, G. Schuster and N. Adir, *Nat. Commun.*, 2018, **9**, 2168.
- 84 Y. A. Gorbly, S. Yanina, J. S. McLean, K. M. Rosso, D. Moyles, A. Dohnalkova, T. J. Beveridge, I. S. Chang, B. H. Kim, K. S. Kim, D. E. Cullley, S. B. Reed, M. F. Romine, D. A. Saffarini, E. A. Hill, L. Shi, D. A. Elias, D. W. Kennedy, G. Pinchuk, K. Watanabe, S. Ishii, B. Logan, K. H. Nealson and J. K. Fredrickson, *Proc. Natl. Acad. Sci. U. S. A.*, 2006, **103**, 11358–11363.
- 85 H. Rubsam, F. Kirsch, V. Reimann, A. Erban, J. Kopka, M. Hagemann, W. R. Hess and S. Klahn, *Environ. Microbiol.*, 2018, **20**, 2757–2768.
- 86 G. Longatte, H. Y. Fu, O. Buriez, E. Labbe, F. A. Wollman, C. Amatore, F. Rappaport, M. Guille-Collignon and F. Lemaître, *Biophys. Chem.*, 2015, **205**, 1–8.
- 87 G. Longatte, F. Rappaport, F. A. Wollman, M. Guille-Collignon and F. Lemaître, *Photochem. Photobiol. Sci.*, 2016, **15**, 969–979.
- 88 G. Longatte, F. Rappaport, F. A. Wollman, M. Guille-Collignon and F. Lemaître, *Electrochim. Acta*, 2017, **236**, 337–342.
- 89 G. Longatte, A. Sayegh, J. Delacotte, F. Rappaport, F. A. Wollman, M. Guille-Collignon and F. Lemaître, *Chem. Sci.*, 2018, **9**, 8271–8281.
- 90 L. Beauzamy, J. Delacotte, B. Bailleul, K. Tanaka, F. A. Wollman and F. Lemaître, *Anal. Chem.*, 2020, **92**, 7532–7539.
- 91 E. Cevik, H. Tombuloglu, I. Anil, M. Senel, H. Sabit, S. Abdulazeez, J. F. Borgio and M. Barghouthi, *Int. J. Hydrogen Energy*, 2020, **45**, 11330–11340.
- 92 K. Hasan, S. A. Patil, K. Gorecki, D. Leech, C. Hagerhall and L. Gorton, *Bioelectrochemistry*, 2013, **93**, 30–36.
- 93 R. D. Milton, D. P. Hickey, S. Abdellaoui, K. Lim, F. Wu, B. Tan and S. D. Minteer, *Chem. Sci.*, 2015, **6**, 4867–4875.
- 94 S. Abdellaoui, R. D. Milton, T. Quah and S. D. Minteer, *Chem. Commun.*, 2016, **52**, 1147–1150.
- 95 K. Hasan, M. Grattieri, T. Wang, R. D. Milton and S. D. Minteer, *ACS Energy Lett.*, 2017, **2**, 1947–1951.
- 96 G. Pankratova, D. Pankratov, R. D. Milton, S. D. Minteer and L. Gorton, *Adv. Energy Mater.*, 2019, **9**, 1900215.
- 97 M. Grattieri, S. Patterson, J. Copeland, K. Klunder and S. D. Minteer, *ChemSusChem*, 2020, **13**, 230–237.
- 98 K. Hasan, R. D. Milton, M. Grattieri, T. Wang, M. Stephanz and S. D. Minteer, *ACS Catal.*, 2017, **7**, 2257–2265.
- 99 F. Zhao, P. Wang, A. Ruff, V. Hartmann, S. Zacarias, I. A. C. Pereira, M. M. Nowaczyk, M. Rögner, F. Conzuelo and W. Schuhmann, *Energy Environ. Sci.*, 2019, **12**, 3133–3143.
- 100 K. P. Sokol, D. Mersch, V. Hartmann, J. Z. Zhang, M. M. Nowaczyk, M. Rögner, A. Ruff, W. Schuhmann, N. Plumeré and E. Reisner, *Energy Environ. Sci.*, 2016, **9**, 3698–3709.
- 101 J. Tschortner, B. Lai and J. O. Kromer, *Front. Microbiol.*, 2019, **10**, 866.
- 102 A. C. Gonzalez-Aravena, K. Yunus, L. Zhang, B. Norling and A. C. Fisher, *RSC Adv.*, 2018, **8**, 20263–20274.
- 103 K. Hasan, H. Bekir Yildiz, E. Sperling, P. O. Conghaile, M. A. Packer, D. Leech, C. Hagerhall and L. Gorton, *Phys. Chem. Chem. Phys.*, 2014, **16**, 24676–24680.

- 104 Y. Takekuma, N. Ikeda, K. Kawakami, N. Kamiya, M. Nango and M. Nagata, *RSC Adv.*, 2020, **10**, 15734–15739.
- 105 R. Takeuchi, A. Suzuki, K. Sakai, Y. Kitazumi, O. Shirai and K. Kano, *Bioelectrochemistry*, 2018, **122**, 158–163.
- 106 D. Pankratov, J. Zhao, M. A. Nur, F. Shen, D. Leech, Q. Chi, G. Pankratova and L. Gorton, *Electrochim. Acta*, 2019, **310**, 20–25.
- 107 D. Pankratov, G. Pankratova and L. Gorton, *Curr. Opin. Electrochem.*, 2020, **19**, 49–54.
- 108 M. Sawa, A. Fantuzzi, P. Bombelli, C. J. Howe, K. Hellgardt and P. J. Nixon, *Nat. Commun.*, 2017, **8**, 1327.
- 109 N. Sekar, Y. Umasankar and R. P. Ramasamy, *Phys. Chem. Chem. Phys.*, 2014, **16**, 7862–7871.
- 110 E. Cevik, M. Buyukharman and H. B. Yildiz, *Biotechnol. Bioeng.*, 2019, **116**, 757–768.
- 111 K. D. Wolfe, D. Dervishogullari, C. D. Stachurski, J. M. Passantino, G. K. Jennings and D. E. Cliffel, *ChemElectroChem*, 2019, **7**, 596–603.
- 112 J. A. Cornejo, H. Sheng, E. Edri, C. M. Ajo-Franklin and H. Frei, *Nat. Commun.*, 2018, **9**, 2263.
- 113 Y. Su, S. Cestellos-Blanco, J. M. Kim, Y.-X. Shen, Q. Kong, D. Lu, C. Liu, H. Zhang, Y. Cao and P. Yang, *Joule*, 2020, **4**, 800–811.
- 114 F. Li, L. Wang, C. Liu, D. Wu and H. Song, *Curr. Opin. Electrochem.*, 2018, **10**, 37–45.
- 115 L. Shi, K. M. Rosso, T. A. Clarke, D. J. Richardson, J. M. Zachara and J. K. Fredrickson, *Front. Microbiol.*, 2012, **3**, 50.
- 116 M. Akhujkar, J. Krushkal, G. DiBartolo, A. Lapidus, M. L. Land and D. R. Lovley, *BMC Microbiol.*, 2009, **9**, 109.
- 117 N. Schuergers, C. Werlang, C. M. Ajo-Franklin and A. A. Boghossian, *Energy Environ. Sci.*, 2017, **10**, 1102–1115.
- 118 H. Jensen, M. Ter Avest, M. Kokish and C. M. Ajo-Franklin, *ACS Synth. Biol.*, 2016, **5**, 679–688.
- 119 N. Sekar, R. Jain, Y. Yan and R. P. Ramasamy, *Biotechnol. Bioeng.*, 2016, **113**, 675–679.
- 120 N. Sekar, J. Wang, Y. Zhou, Y. Fang, Y. Yan and R. P. Ramasamy, *Biotechnol. Bioeng.*, 2018, **115**, 1361–1366.
- 121 L. Su and C. M. Ajo-Franklin, *Curr. Opin. Biotechnol.*, 2019, **57**, 66–72.
- 122 M. A. TerAvest and C. M. Ajo-Franklin, *Biotechnol. Bioeng.*, 2016, **113**, 687–697.
- 123 M. A. TerAvest, T. J. Zajdel and C. M. Ajo-Franklin, *ChemElectroChem*, 2014, **1**, 1874–1879.
- 124 Z. Wu, J. Wang, J. Liu, Y. Wang, C. Bi and X. Zhang, *Microb. Cell Fact.*, 2019, **18**, 15.
- 125 F. Kracke, B. Lai, S. Yu and J. O. Kromer, *Metab. Eng.*, 2018, **45**, 109–120.
- 126 J. S. Sravan, O. Sarkar and S. V. Mohan, *Sustainable Energy Fuels*, 2020, **4**, 700–712.
- 127 I. Nabiev, A. Rakovich, A. Sukhanova, E. Lukashev, V. Zagidullin, V. Pachenko, Y. P. Rakovich, J. F. Donegan, A. B. Rubin and A. O. Govorov, *Angew. Chem., Int. Ed.*, 2010, **49**, 7217–7221.
- 128 F. Milano, R. R. Tangorra, O. Hassan Omar, R. Ragni, A. Operamolla, A. Agostiano, G. M. Farinola and M. Trotta, *Angew. Chem., Int. Ed.*, 2012, **51**, 11019–11023.
- 129 O. Hassan Omar, S. la Gatta, R. R. Tangorra, F. Milano, R. Ragni, A. Operamolla, R. Argazzi, C. Chiorboli, A. Agostiano, M. Trotta and G. M. Farinola, *Bioconjugate Chem.*, 2016, **27**, 1614–1623.
- 130 R. Ragni, G. Leone, S. la Gatta, G. Rizzo, M. Lo Presti, V. De Leo, F. Milano, M. Trotta and G. M. Farinola, *MRS Adv.*, 2018, **4**, 1143–1148.
- 131 P. K. Dutta, S. Lin, A. Loskutov, S. Levenberg, D. Jun, R. Saer, J. T. Beatty, Y. Liu, H. Yan and N. W. Woodbury, *J. Am. Chem. Soc.*, 2014, **136**, 4599–4604.
- 132 P. I. Gordiichuk, D. Rimmerman, A. Paul, D. A. Gautier, A. Gruszka, M. Saller, J. W. de Vries, G. J. Wetzelaer, M. Manca, W. Gomulya, M. Matmor, E. Gloukhikh, M. Loznik, N. Ashkenasy, P. W. Blom, M. Rogner, M. A. Loi, S. Richter and A. Herrmann, *Bioconjugate Chem.*, 2016, **27**, 36–41.
- 133 Y. Yoneda, T. Noji, T. Katayama, N. Mizutani, D. Komori, M. Nango, H. Miyasaka, S. Itoh, Y. Nagasawa and T. Dewa, *J. Am. Chem. Soc.*, 2015, **137**, 13121–13129.
- 134 S. Kaushik, M. K. Sarma and P. Goswami, *J. Mater. Chem. A*, 2017, **5**, 7885–7895.
- 135 K. J. Grayson, K. M. Faries, X. Huang, P. Qian, P. Dilbeck, E. C. Martin, A. Hitchcock, C. Vasilev, J. M. Yuen, D. M. Niedzwiedzki, G. J. Leggett, D. Holten, C. Kirmaier and C. Neil Hunter, *Nat. Commun.*, 2017, **8**, 13972.
- 136 J. Liu, V. M. Friebe, R. N. Frese and M. R. Jones, *Nat. Commun.*, 2020, **11**, 1542.
- 137 J. Liu, J. Mantell and M. R. Jones, *ACS Nano*, 2020, **14**, 4536–4549.