



Cite this: *Environ. Sci.: Processes Impacts*, 2025, 27, 2212

Risk assessment of tire wear in the environment – a literature review†

Kathrin Müller, ^a Kenny Unice,^b Julie Panko,^c Benoît J. D. Ferrari, ^d
Florian Breider ^e and Stephan Wagner *^{af}

Tread wear emission inventories, uncertainty about the future development of the emissions and observed adverse effects of tire constituents in the environment have raised the need for an environmental risk assessment of tire wear emissions. While progress has been made in exposure and hazard assessment of tire wear emissions in the environment, the complexity of tire wear emissions creates some challenges which are not yet overcome. For instance, there is no universal agreed risk assessment framework for tire wear emissions. It was proposed that existing frameworks, for example for microplastics, be adapted to tire wear emissions because there are similarities between particulate tire wear emissions and microplastics, e.g. particulate material with a polymer backbone. The review discusses whether these are applicable for tire wear emissions and proposes adaptations. It provides a comprehensive assessment of exposure and hazard data for tire wear emission and reveals needs and data gaps for environmental risk assessment of tire wear. Based on the available exposure and hazard data sets a low risk prioritization of particulate tire wear emissions in aquatic and terrestrial environments was estimated. Risk prioritization of leachables from tire emissions is not yet possible due to inconsistent hazard data sets. It was found that for environmental risk assessment, insufficient consistent exposure and hazard data is available. It is suggested to develop clear harmonization guidelines how exposure and hazard studies should be designed. Such guidelines should be developed between all relevant stakeholders covering the entire product life cycle.

Received 3rd February 2025
Accepted 20th May 2025

DOI: 10.1039/d5em00088b

rsc.li/espi

Environmental significance

This structured literature review concludes with a risk prioritization of environmental compartment with respect to particulate tire wear emissions and leachables from tire wear. The review reveals also data and knowledge gaps that must be addressed in order to facilitate a comprehensive risk assessment of tire wear emissions. These needs include data and knowledge regarding exposure and hazard of tire wear emissions for marine environments and more specifically exposure and hazard data of leachable tire chemicals and their transformation products. Furthermore, it proposes a series of possible adaptations of microplastics risk assessment frameworks that could be employed in the context of tire wear emissions.

1 Introduction

As reported by the International Organization of Motor Vehicle Manufacturers, in 2024 global vehicle production reached approximately 92.5 million units.¹ The average lifespan of a tire

is approximately four years, or 50 000 kilometers. During this period, the tire undergoes a loss of approximately 10% of its mass. This equates to a reduction of between 1 and 1.5 kg for a tire weighing between 7 and 12 kg, or an average wear per vehicle of approximately 120 g per 1000 km.^{2,3} For example, it is estimated that approximately 500 000 tons of tire wear is generated in the EU on an annual basis.⁴ How these mass generation rates will evolve during the ongoing electrification of the vehicle fleet is not yet known. Tread wear emission inventories, uncertainty about the future development of the emissions and observed adverse effects of tire constituents in the environment have raised the need of an environmental risk of tire wear emissions.^{5–7}

The characterization of environmental risk is a process for estimating the probability of an adverse outcome or event due to pressures or changes in environmental conditions resulting from human activities. Risk is determined by incorporating

^aHochschule Fresenius, Institute for Analytical Research, Idstein, Germany. E-mail: stephan.wagner@hs-fresenius.de

^bStantec Consulting Services, Inc., Pittsburgh, PA, USA

^cToxStrategies LLC, Pittsburgh, PA, USA

^dEcotox Centre – Ecole Polytechnique Fédérale de Lausanne – EPFL, IIE, ENAC, Station 2, Switzerland

^eEcole Polytechnique Fédérale de Lausanne – EPFL, Central Environmental Laboratory, IIE, ENAC, Station 2, Lausanne, Switzerland

^fKompetenzzentrum Wasser Hessen, Max-von-Laue-Straße 13, 60438 Frankfurt am Main, Germany

† Electronic supplementary information (ESI) available. See DOI: <https://doi.org/10.1039/d5em00088b>



knowledge of both the hazard and exposure potential and involves several steps, including (1) problem formulation, (2) exposure assessment, (3) hazard assessment and dose-response, and (4) risk characterization. Problem formulation defines the scope of environmental risk characterization by specifying the affected environmental compartment and potential harmful effects. The risk is determined based on a comparison between the hazard and the exposure assessment.^{8,9} There is no universal agreed risk assessment framework for tire wear emissions. It has been suggested to adapt existing frameworks for instance for microplastics^{10–13} to tire wear emissions because there are similarities between particulate tire wear emissions and microplastics, *e.g.* particulate material with a polymer backbone. Risk characterization of tire wear emissions may include particulate but also volatile and dissolved emissions so called leachables.¹⁴ Therefore, there is

a need to discuss the applicability of the existing microplastic risk assessment framework and consider the possibility of making adaptations.

This structured literature review summarizes available risk assessment concepts for microplastics and assesses whether these are applicable for tire wear emissions and suggests adaptations of these concepts toward tire wear emissions. Subsequently a comprehensive assessment of exposure and hazard data for tire wear emission is carried out and the needs and data gaps for environmental risk assessment of tire wear are identified. Based on the available exposure and hazard data a risk prioritization for freshwater, terrestrial and marine environments is done. The structured literature search was done from 1980 to 2024. Ultimately, the present study explores the question of whether environmental risk assessment of tire



Kathrin Müller

Kathrin Müller received her Bachelor of Science degree in Applied Chemistry in 2018 and her Master of Science degree in Bioanalytical Chemistry and Pharmaceutical Analysis in 2019 at Hochschule Fresenius gem. Trägergesellschaft mbH. During her studies she worked at SGS Institut Fresenius GmbH and after her master studies, she started working at the Institute for Analytical Research (IFAR). Since, 2020 she is a PhD student

at Hochschule Fresenius gem. Trägergesellschaft mbH and her research focuses on environmental and analytical chemistry in the context of tire wear emissions.



Benoît J. D. Ferrari

Dr Benoît J. D. Ferrari is a researcher in the field of ecotoxicology. He is the Director of the Swiss Centre for Applied Ecotoxicology (Ecotox Centre) and Head of the Soil and Sediment Ecotoxicology group. His main areas of interest are the ecodynamic of contaminants and their effects at different levels of biological organisation. In particular, he is involved in the development of exposure- and effect-based tools, the

development of integrated approaches to assess the hazard and risk of chemicals on ecosystems, and the transfer of such tools and approaches to end-users.



Kenny Unice

Kenny Unice is a Principal Health Scientist at Stantec with 25 years of quantitative human health risk, environmental modeling experience and product stewardship experience. His expertise is in environmental fate and transport models and the application of risk assessment principles in occupational, community, consumer product and environmental health evaluations. He has co-authored more than 70

peer-reviewed articles on environment, toxicokinetics, bio-monitoring, pathogen risk assessment, exposure reconstruction, and subsurface bacterial transport since 2004. He has served as an invited peer-reviewer for several journals, U.S. EPA draft modeling approaches, and of California Sea Grant microplastic research proposals.



Florian Breider

Dr Florian Breider is the head of the Central Environmental Laboratory at Ecole Polytechnique Fédérale de Lausanne - EPFL. He studied environmental chemistry and physics at the University of Lausanne and earned his PhD in isotope biogeochemistry from the University of Neuchâtel. He then worked as a research associate at Tokyo Institute of Technology. His research focuses on the fate and effects of micropollutants

and micro/nanoplastics in ecosystems, and their implications for human health. Since 2020, he has been investigating the environmental impact of tire and road wear particles and their associated chemicals in both aquatic and terrestrial systems.



wear emissions is already possible and formulates most pressing research needs in this field.

2 Adaptation of the risk assessment concepts for microplastics to tire wear emissions

2.1 Risk assessment concepts for microplastics

In the past decade, ecological assessments of microplastics have consisted of particle mass or number exposure concentration measurement studies aimed at determining nature and extent in aquatic, terrestrial and airborne exposure settings. While progress has been made on the problem formulation and the general scheme (Table 1), a global consensus has not been agreed on the critical elements of a robust regulatory risk assessment framework.¹¹ Nonetheless, the proposed frameworks include (1) consideration of problem formulation, (2) exposure assessment, hazard assessment and (3) the integration of hazard and exposure to characterize the risk (Table 1). Differences exist in the terminology and approach in frequently cited frameworks that have been proposed for regulatory risk assessment (Table 2). The four main concepts proposed for microplastic risk assessment to date are based on this stepwise assessment requiring exposure and hazard data but differ with respect to information inputs.

The International Council of Chemical Associations (ICCA) proposed an environmental risk assessment framework (ERA) which would identify species and compartments of concern using exposure, fate, and hazard studies for a wide range of polymer types with diverse physicochemical properties¹⁵ resulting in a high number of tests. The concept by Koelmans *et al.* is based on polymer-specific probability density functions (PDF) which reflect the diverse microplastic properties in the environment^{10,16} and the identification of most sensitive species

towards the heterogeneous polymer mixtures. This approach may not allow systemic identification of drivers for adverse effects. A third approach, the microplastics management framework, categorizes microplastics concentrations and applies tiers for mitigation actions.¹⁷ A fourth concept uses a relative level of concern based on summed categorical hazard such as none, low, moderate, elevated, or high which are based on a harmonized exposure and hazard testing protocols.¹⁸ All concepts have in common that they focus on the particulate nature of the microplastics, but chemical additives are not included despite some of them are for instance endocrine disruptors, *e.g.* bisphenol A or phthalates and nonylphenols.¹⁹

2.2 An extended risk assessment framework for tire wear emissions needs to cover particulate tire wear emissions and leachables

Among the proposed frameworks in Table 2 none can be directly applied to tire wear emissions because (i) physicochemical and morphological differences exist between particulate tire wear emissions and microplastics and (ii) both particulate and chemical hazard are relevant for tire wear emissions while microplastic concepts focus only on assessment of particles. The PDF approach by Koelmans *et al.*¹⁰ represents the most flexible approach because it accounts for the heterogeneity of the physical and chemical properties of these materials. It is also comprehensive and consistent with modern ecological risk assessment principles incl. quantitation of uncertainty (Tables 1 and 2). Similar to microplastics, the uncertainty of exposure and hazard data from tire wear emissions remains high and needs to be reduced to draw reliable conclusions. For a tire wear risk assessment framework, the existing microplastic framework may be complemented by specific aspects of chemical compounds, so called tire leachables and their transformation products, as tire wear particles may act as a source for these chemicals. As revealed by this



Julie Panko

Ms Panko is a Principal Scientist at ToxStrategies LLC. She has >30 years of experience conducting and managing various risk assessments. She is a Certified Industrial Hygienist (CIH). Her work focuses on evaluating chemical risks from industrial, commercial, and consumer products within the context of various regulatory programs and voluntary initiatives. Ms. Panko's expertise include quantitative exposure assessment

with retrospective analysis of occupational and community-based exposures to industrial emissions; human health risk assessment for contaminated sites; industrial hygiene program development and implementation; indoor air quality related to industrial, commercial, and residential environments; and product stewardship and sustainability.



Stephan Wagner

Stephan Wagner is the head of the Institute for Analytical Research at the Hochschule Fresenius University of Applied Sciences in Idstein, Germany and a professor for environmental analysis. His research focuses on detection and fate of anthropogenic materials in the aquatic environment and water treatment systems. He graduated from Technische Universität Dresden, Germany in waste and contaminated site

management and received his PhD in mine water treatment in 2011 from Brandenburgische Technische Universität Cottbus, Germany. Before joining Hochschule Fresenius in July 2022, he worked at several research institutions in Germany and Austria as Postdoc and as research group leader.



Table 1 Overview of the multidimensional regulatory risk assessment scheme and emphasis on ecologically relevant metrics (ERM) proposed by ref. 10 and 20

| Scheme element | Step 1: Problem formulation | Step 2: Exposure and hazard | Step 3: Risk assessment |
|----------------|--|---|--|
| Question | What is the protection objective? | What is the effect mechanism? | What is the risk? |
| Outcome | Preliminary hypotheses about past or future ecological effects | Selection of exposure (A) and dose–response metrics (B) | Risk for each assessed toxicological and/or ecological effect mechanism |
| Concepts | ● Exposure pathways | ● Exposure metric and profile | ● Continuum (e.g. distribution of sizes) |
| Considerations | ● Ecological or toxicological effect ● Harmonization ● Quality assurance and control | ● Effect metric and profile ● Mass and surface area ● Volume and aspect ratio | ● Acceptable dose distributions ● Bioavailable fraction ● Probability density function |

literature review, leachables from tire wear emissions are most susceptible to adverse effects. Therefore, an extension of the risk assessment of microplastics towards methodologies for leachables including transformation products is required.

Because a risk assessment is conducted for a specific geographical scale and mostly for a specific site, it precludes the

drawing of general conclusions regarding the risk of tire wear emissions. However, a discussion on available data and information regarding the three steps of the regulatory risk assessment scheme depicted in Table 1 for tire wear emissions is possible.

Table 2 Comparison of four proposed environmental risk assessment frameworks

| Framework | Contributors | Key first tier inputs | First tier outputs | Higher tier assessment | Data needs | Reference |
|--|---|--|--|--|--|---------------|
| International Council of Chemical Associations (ICCA) symposium summary of a proposed conceptual model | Scientists from academia, industry and government attending ICCA symposium | Fate; physio-chemical properties; harmonized effect studies (physical and cellular effects); exposure and dose–response modeling | Compartments and species of concern; screening-level risk characterization; identification of data needs to address uncertainty | Refined risk characterization based on mesocosm, field studies, mixture toxicity, and/or species sensitivity distributions | Research on method development and use to strengthen data quality | 15 |
| Development of default polymer-specific probability density functions (PDF) | Lead by scientists at Aquatic Ecology and Water Quality Management Group, Wageningen University & Research ^a | PDFs for particle width, width to length ratio, area, specific surface area, volume and mass distributions | Screening-level alignment of monodisperse effect threshold with polydisperse environmental exposures for risk characterization | Authors suggest conceptual workflow be used refine the generalized PDFs and exponent values | Mechanism of adverse effects for freshwater, marine and terrestrial toxicological or ecological endpoints | 10, 16 and 21 |
| Microplastics management framework (MPP) developed by Microplastics Health Effects Workshop experts | International scientists participating in Southern California Coastal Water Research Project (SCCWRP) workshop | Particles per liter or mass (mg per liter) in water scaled operationally defined as particles sized 1 to 5000 µm | Management tier or level of concern based on concentration for food dilution and tissue translocation Tier 1 (no concern); Tier 2 (low; monitoring); Tier 3 (moderate; study sources); Tier 4 (elevated; mitigation); Tier 5 (pollution control) | Freshwater and aquatic species were merged to derived tiered thresholds of concern; data for taxa specific to geography and habitat, if available, would refine the assessment | Evaluation of the assumption that mechanisms influence toxicity more than habitat; thresholds for sediment | 17 |
| Bucci and Rochman (BR) | Scientists at Department of Ecology and Evolutionary Biology, University of Toronto | Categorical dimensions of shape, polymer type, size and ecosystem water quality | Relative level of concern based on summed categorical hazard such as none, low, moderate, elevated or high | Metaanalyses of literature or future toxicity tests to inform risk management in line with objectives and risk tolerance | Harmonized exposure measurements and ecotoxicity testing | 18 |

^a Researchers from Alfred Wegener Institute Helmholtz Centre for Polar and Marine Research (Germany) and Department of The Built Environment, Aalborg University (Denmark) contributed to the development of particle-specific PDFs.



3 Risk assessment of tire wear emissions in aquatic and terrestrial environments

3.1 Step 1: Problem formulation

In step 1 the protection objective is defined and preliminary hypotheses about past or future ecological effects are set. This includes the geographical scale of the risk assessment; does it assess a local site or a region, continent or even globally. Endpoints of adverse effects and test organism are to be defined as well; classics are growth or mortality or other end points which may be related to chronic effects or effects on cellular level, metabolisms or on population scale.

With the compilation of the release and distribution pathways of tire wear emissions during vehicle use, relevant exposure pathways and affected environmental compartments are identified. These pathways have been described earlier.^{22,23} Tire wear emissions during vehicle use comprises particulate tire wear emissions, leachable constituents and volatile or semi-volatile organic chemicals. While the emission pathways and exposure routes of particulate tire emissions and leachables have been studied in more detail, there are substantial knowledge and data gaps for the emission of volatiles.²⁴ The main distribution pathways of leachables and particulate tire emissions is *via* terrestrial and aquatic compartments and thus, considered as relevant environmental exposure routes. While air is considered as a transport pathway of airborne particulate tire wear emission eventually being deposited in terrestrial and aquatic compartments, it is only considered as relevant exposure route for human health assessment but not ecological risk assessment.²²

Many microplastic researchers have acknowledged that the largest testing uncertainty is the lack of representativeness of the test material because most microplastic testing has been conducted with particles of the pristine polymers rather than present in the environment. Hence, the pristine microplastics lack many of the plastic additives. Compared to tire wear emissions microplastics are expected to contain an even higher diversity of additives due to the very broad range of applications and the required material properties while the mass concentration of additives in rubber is comparably high; up to 60%.²⁵ Representative test materials for tire wear emissions may include particulate materials mimicking tire and road wear particles (TRWPs) and dissolved compounds representing typical leachable composition. Ideally the test material for hazard and fate studies should mimic the heterogeneity of the environmental tire wear emissions including physical and chemical properties *e.g.* size, density, and chemical composition, including transformation products.²⁶ For instance, it was observed that aging of particulate tire wear results in changes of the toxicity of the leachates; leachates from pristine tire wear particles were more toxic compared to leachates from aged tire wear particles.²⁷ Although the tire wear emission researchers use some type of surrogate for TRWP, the test material is highly variable including particles from laboratory generated TRWP,^{28,29} manual abrasion from the tire surface,^{30–32}

commercially produced recycled crumb rubber,³³ both of which may be further ground to a specific size range, and more recently, a standardized cryogenic milled tire tread (CMTT) mix.³⁴ Additionally, there is an appreciable difference in test system designs with respect to particle size, concentration of tire particles used either directly or from which leachates are prepared (100 to >100 000 mg L⁻¹), duration of leaching time, temperature at which the leachate is created, and aging of the particles.^{35–38} Test material representing a tread compound mix of tire leachables, including their transformation products, needs to be defined for the exposure assessment of the entirety of tire leachables. As for now leachates are produced mostly from pristine tire materials or laboratory aged tires either using organic extraction agents or water-based extraction protocols while environmental material rich in TRWP has not been used to generate representative leachable composition.³⁹

3.2 Step 2A: Exposure assessment in relevant environmental compartments

A structured literature search was conducted of relevant scientific publications using Web of Science and PubMed as research database to identify exposure and hazard measurements of particulate tire emissions or leachables in terrestrial and aquatic environmental compartments. In total 327 publications were reviewed, and 130 publications were found stating exposure or hazard concentration data for tire wear emissions (Table 3). Exposure data for freshwater environments (38 items) was reported more than four times as much as for marine (11 items) and terrestrial environments (10 items). There were also more literature hits for leachables compared to particulate tire wear emissions. Additionally, data for road dust, treatment systems, storm water and snow (other) were compiled as they may act as source and entrance pathways for tire emissions into natural environments.

3.2.1 Concentrations of particulate tire emissions are overall consistent despite high uncertainties in the quantitation methods. Most data were found for road dust, one of the initial receiver media of particulate tire wear emission, and the different road dust origins can be rank from lowest to highest concentration as follows: rural < industrial = urban < tunnel (Table 4). The mean concentration ranges between several thousand to tens of thousands mg kg⁻¹, while maximum concentrations of over 100 000 mg kg⁻¹ are reported for urban and tunnel road dust.^{40,41} For roadside soil as primary environmental receiving media, a mean concentration of several thousand mg kg⁻¹ is reported but can even reach up to 52 720 mg kg⁻¹⁴² and thus, are in the same range as for road dust. For the aquatic compartment as receiving media, storm water is considered as main distribution pathway, indicated by the mean concentration of particulate tire wear emission in the two-digit mg L⁻¹ range.⁴³ Due to dilution and further transformation and removal processes in the receiving water compartments two order of magnitude lower concentrations are reported for the river water column and sea water column.² For example, the particulate tire emission is eventually deposited in the sediment (secondary environmental receiving media) after



Table 3 Number publications found during the structured literature review for exposure and hazard data of particulate tire wear emissions and tire leachables

| Topic | Total | Freshwater | Marine | Terrestrial | Other |
|--|-------|------------|--------|-------------|-------|
| <i>Overall</i> | 130 | 68 | 27 | 21 | 43 |
| (i) Exposure data of particulate tire wear emission | 31 | 10 | 3 | 5 | 20 |
| (ii) Exposure data of tire leachables | 49 | 28 | 8 | 5 | 21 |
| (iii) Hazard assessment data of particulate tire wear emission | 29 | 16 | 6 | 8 | 1 |
| (iv) Hazard assessment data of tire leachables | 32 | 23 | 11 | 4 | 1 |

reaching the aquatic compartment.²⁹ The reported river sediment concentrations range from several hundred to thousand mg kg⁻¹, with exception of the results of Goßmann *et al.*²⁵ which are significantly lower reporting only 0.05 mg kg⁻¹ on average. This uncertainty is usually explained with different sample locations, sampling methods, as well as analyses methods.^{3,44} This may also be true for the reported high concentration in lake sediment employing Zn as marker.⁴⁵ Barber *et al.*⁴⁶ adapted the Zn-based tire wear detection method and yielded more comparable results for the same samples analyzed once employing Zn as marker and once employing styrene-butadiene rubber (SBR), butadiene rubber (BR) and natural rubber (NR) pyrolysis products as marker. For sea sediment the authors reported concentrations in the low hundred mg kg⁻¹ range, while again the results of Goßmann *et al.*²⁵ for sea sediment are significantly lower reporting the overall lowest concentration in any compartment. As this concentration is one order of magnitude lower than the authors reported river sediment concentrations, the concentration of particulate tire wear emission in sea sediments is assumed to be the lowest. Yet, further studies of lake and sea sediments are needed to verify these results.

The observed high concentration variations within one sampling compartment may be attributed, among other things, to the conditions at the different sample locations of each study. The formation of particulate tire wear emissions depends on several influencing parameters such as vehicle speed, road profile/section, road surface, tire types and more.⁴⁷⁻⁵² Thus, urban road dust samples from Beijing, China,⁵³ from Seoul, Republic of Korea⁵⁴ and from Leipzig, Germany⁴⁵ are difficult to directly compare. Furthermore, differences in the analysis method increase the variation as eleven different markers were used to determine the particulate tire wear concentration using average content of marker compounds in tire tread. Interestingly, the mean concentrations in urban road dust are one order of magnitude higher if organic marker compounds are employed compared to polymer markers (compare Table 4). While these differences indicate a potential bias, no general conclusion can be drawn about the marker suitability and if concentrations are overestimated or underestimated, respectively.

3.2.2 Exposure data of tire leachables is dominated by a few tire chemicals. Over 200 potential leachables emitted from tires or particulate tire emissions have been identified^{30,76} but

only for a few selected compounds such as *N*-(1,3-dimethylbutyl)-*N'*-phenyl-*p*-phenyldiamine (6PPD) and its transformation product *N*-(1,3-dimethylbutyl)-*N'*-phenyl-*p*-phenyldiamine quinone (6PPD-Q), 1,3-diphenylguanidine (DPG), hexa(methoxymethyl)melamine (HMMM), benzotriazole and benzothiazole derivatives has their environmental occurrence and distribution been reported (Tables 5 and S1†). Among the assessed compounds 6PPD-Q is studied the most (38 publications) followed by DPG (20 publications) which is detected most frequently and is reported with the overall highest concentration of 0.3 mg L⁻¹ in storm water.⁷⁷ The highest concentrations in environmental media are reported in general for each compound in storm water, ranging from ng L⁻¹ to µg L⁻¹, as it acts as transportation media for leachables.⁷⁷⁻⁸¹ Additionally, for roadside snow, high concentrations are detected due to the accumulation of particulate tire emissions in the snow and the resulting continuous leaching of the particles.^{77,82} Thus, snow melt water must also be considered as relevant mobilization and transportation media for leachables. The same is true for tunnel wash water. While it does not represent an environmental compartment, it acts as a potential source and entrance pathway, reporting concentrations of several µg L⁻¹.^{83,84}

In the receiving compartments, freshwater concentrations in the ng L⁻¹ range are reported and lowest concentrations are reported for marine water due to the dilution effect and other fate processes *e.g.* transformation, breakdown and possibly sorption processes. Thus, also higher concentrations for freshwater sediment are reported reaching up to 801 µg kg⁻¹ for HMMM and 619 µg kg⁻¹ BTR,^{85,86} while for marine sediment one-digit µg kg⁻¹ concentrations are reported. Yet, data on sediment concentrations for each compound are limited as mostly only one study assessed those (Tables 5 and S1†). Soil as receiving media is studied the least and concentrations of up to mg kg⁻¹ are reported with the highest concentration of 4.1 mg kg⁻¹ for DPG.⁸⁷ Especially for the terrestrial receiving media and sediments a more comprehensive data set is needed for a reliable exposure estimation and thus a risk characterization of leachables.

Another consideration in the exposure and thus risk characterization is the uncertainty about the known and unknown leachables. While the presented exposure data are needed for a risk characterization of each compound, no defined exposure measure for the risk assessment of tire leachables exists.



Table 4 Reported concentrations of tire wear particles in various environmental and anthropogenic matrices using different chemical markers. The data refers to the tire tread component of TRWP particles

| Sampling compartment | Marker | Mean | Min | Max | Unit | References |
|----------------------------|---------------|--------|--------|---------|---------------------|---------------------|
| Terrestrial | | | | | | |
| Roadside soil | SBR | 3347 | 179 | 18 700 | mg kg ⁻¹ | 55 and 56 |
| | SBR/BR | 3900 | 200 | 20 000 | mg kg ⁻¹ | 29 |
| | SBR/SBS/BR | 9381 | 3700 | 20 067 | mg kg ⁻¹ | 42 and 57 |
| | Zn | 8400 | 475 | 52 720 | mg kg ⁻¹ | 55 |
| <i>Road dust</i> | | | | | | |
| Urban | SBR | 5683 | 4050 | 7000 | mg kg ⁻¹ | 45 |
| | SBR/BR | 3580 | 2270 | 4950 | mg kg ⁻¹ | 58 |
| | SBR/NR | 1294 | 105 | 6813 | mg kg ⁻¹ | 59 and 60 |
| | SBR/BR/NR | 2666 | 1500 | 4500 | mg kg ⁻¹ | 54 |
| | BT | 35 400 | | | mg kg ⁻¹ | 61 |
| | Bonded sulfur | 23 400 | 10 400 | 48 600 | mg kg ⁻¹ | 53 |
| | 24MoBT | 17 000 | 5000 | 124 000 | mg kg ⁻¹ | 40 and 41 |
| | NCBA | 34 000 | 2000 | 78 000 | mg kg ⁻¹ | 40 and 41 |
| | Oleamide | 28 958 | 18 250 | 42 750 | mg kg ⁻¹ | 62 |
| | Zn | 4017 | 3350 | 4700 | mg kg ⁻¹ | 45 |
| | Rural | SBR/BR | 510 | | | mg kg ⁻¹ |
| Industrial | SBR/NR | 7645 | 860 | 20 660 | mg kg ⁻¹ | 59 and 63 |
| | SBR/NR | 59 748 | 38 037 | 87 472 | mg kg ⁻¹ | 60 |
| Tunnel | SBR/SBS/BR | 62 900 | 8650 | 186 500 | mg kg ⁻¹ | 64 |
| | 24MoBT | 70 600 | 7725 | 210 700 | mg kg ⁻¹ | 41 |
| | NCBA | 67 100 | 14 404 | 203 600 | mg kg ⁻¹ | 41 |
| | Zn | 57 500 | 55 000 | 60 000 | mg kg ⁻¹ | 65 |
| Tunnel wash water | SBR/SBS/BR | 16.7 | 4.9 | 34.6 | mg L ⁻¹ | 64 |
| Aquatic | | | | | | |
| <i>River</i> | | | | | | |
| Sediment | SBR/BR | 1036 | 13 | 31 508 | mg kg ⁻¹ | 29 and 66 |
| | SBR/BR/NR | 250 | 70 | 1150 | mg kg ⁻¹ | 67 |
| | SBR/SBS/BR | 0.05 | 0.01 | 0.07 | mg kg ⁻¹ | 25 |
| | 24MoBT | 900 | 429 | 3900 | mg kg ⁻¹ | 40 and 41 |
| | NCBA | 3500 | 530 | 7300 | mg kg ⁻¹ | 40 and 41 |
| | Zn | 611 | 0.9 | 2560 | mg kg ⁻¹ | 67 |
| Water column | SBR/BR | 0.38 | 0.004 | 9.2 | mg L ⁻¹ | 66, 68 and 69 |
| | SBR/BR/NR | 0.01 | 0.004 | 0.03 | mg L ⁻¹ | 67 |
| <i>Lake</i> | | | | | | |
| Sediment | SBR | 52.1 | 0.075 | 172.9 | mg kg ⁻¹ | 45 and 70 |
| | Zn | 20 000 | | | mg kg ⁻¹ | 45 |
| <i>Sea</i> | | | | | | |
| Sediment | SBR/BR/NR | 202 | 103 | 361 | mg kg ⁻¹ | 46 |
| | SBR/SBS/BR | 0.002 | | | mg kg ⁻¹ | 25 |
| | Zn | 179 | 73.5 | 522 | mg kg ⁻¹ | 46 |
| Water column | SBR/NR | 0.0008 | | | mg L ⁻¹ | 46 |
| | SBR/SBS/BR | 0.002 | 0.0006 | 0.007 | mg L ⁻¹ | 71 |
| Storm water | SBR/BR/NR | 4.06 | 0.01 | 12.2 | mg L ⁻¹ | 72 |
| | 24MoBT | 43.9 | 0.64 | 164.8 | mg L ⁻¹ | 40 and 41 |
| | NCBA | 34.0 | 2.23 | 171.7 | mg L ⁻¹ | 40 and 41 |
| Roadside snow | SBR/SBS/BR | 2090 | 0.11 | 14 500 | mg L ⁻¹ | 57 and 73 |
| Treatment system | | | | | | |
| <i>Sedimentation basin</i> | | | | | | |
| Sediment | SBR | 34 300 | 950 | 95 000 | mg kg ⁻¹ | 45 and 74 |
| | SBR/BR | 24 400 | 2611 | 374 300 | mg kg ⁻¹ | 66 |
| | Zn | 53 900 | 190 | 240 000 | mg kg ⁻¹ | 31 and 45 |
| Water column | SBR/BR | 1 | 0.20 | 15.1 | mg L ⁻¹ | 66 |
| <i>gully pot</i> | | | | | | |
| Sediment | SBR/BR | 6700 | 305 | 188 700 | mg kg ⁻¹ | 66 |
| | SBR/SBS/BR | 26 800 | 5900 | 76 800 | mg kg ⁻¹ | 57 and 64 |
| | STA + FTIR | 43 100 | 800 | 149 600 | mg kg ⁻¹ | 75 |
| Water column | SBR/BR | 1.9 | 0.44 | 27.8 | mg L ⁻¹ | 66 |



Table 5 Reported concentrations of TRWP related leachable compounds in various environmental matrices. For concentration data of further TRWP related leachable compounds refer to Table S1. 6PPD *N*-(1,3-Dimethylbutyl)-*N'*-phenyl-*p*-phenylenediamine, 6PPD-Q *N*-(1,3-dimethylbutyl)-*N'*-phenyl-*p*-phenylenediamine quinone, HMMM hexa(methoxymethyl)melamine, DPG 1,3-diphenylguanidine

| Analyte | Sample | Mean | Min | Max | Unit | Reference |
|---------|---------------------|--------|-------|---------|---------------------|--|
| 6PPD | Freshwater | 28.0 | 0.27 | 163 | ng L ⁻¹ | 89–92 |
| 6PPD | Marine water | 2.65 | 0.02 | 6.37 | ng L ⁻¹ | 93–95 |
| 6PPD | Storm water | 36.7 | <0.04 | 907 | ng L ⁻¹ | 78, 81, 91, 92, 96 and 97 |
| 6PPD | Snow | 329 | 65.6 | 784 | ng L ⁻¹ | 82 |
| 6PPD | Freshwater sediment | 51.4 | 0.14 | 468 | µg kg ⁻¹ | 85, 90, 98 and 99 |
| 6PPD | Marine sediment | 2.87 | <0.02 | 11.1 | µg kg ⁻¹ | 93, 94 and 98 |
| 6PPD | Soil | 182 | 3.03 | 1290 | µg kg ⁻¹ | 97 and 100 |
| 6PPD | WWTP influent | 10.6 | 1.10 | 59.0 | ng L ⁻¹ | 101 and 102 |
| 6PPD | WWTP effluent | 5.94 | 0.30 | 15.0 | ng L ⁻¹ | 101 and 102 |
| 6PPD | Tunnel wash water | 21 554 | 1743 | 91 200 | ng L ⁻¹ | 83 |
| 6PPD-Q | Freshwater | 76.6 | 0.05 | 2850 | ng L ⁻¹ | 68, 89–92 and 103–115 |
| 6PPD-Q | Marine water | 3.47 | 0.02 | 16.3 | ng L ⁻¹ | 80, 93–95 and 109 |
| 6PPD-Q | Storm water | 733 | 0.002 | 5580 | ng L ⁻¹ | 77, 78, 80, 81, 91, 92, 96, 97, 103–105, 109 and 115 |
| 6PPD-Q | Snow | 217 | 4.43 | 756 | ng L ⁻¹ | 77, 82, 96 and 107 |
| 6PPD-Q | Freshwater sediment | 8.72 | <0.30 | 46.0 | µg kg ⁻¹ | 85, 90, 98 and 116 |
| 6PPD-Q | Marine sediment | 2.10 | 0.02 | 4.88 | µg kg ⁻¹ | 93, 94, 98 and 116 |
| 6PPD-Q | Soil | 206 | 0.85 | 939 | µg kg ⁻¹ | 97, 100 and 117 |
| 6PPD-Q | WWTP influent | 28.3 | 1.90 | 470 | ng L ⁻¹ | 91, 101, 102 and 118 |
| 6PPD-Q | WWTP effluent | 8.78 | 1.10 | 37.0 | ng L ⁻¹ | 84, 91, 101 and 102 |
| 6PPD-Q | Tunnel wash water | 1048 | 49.5 | 3440 | ng L ⁻¹ | 83 and 84 |
| HMMM | Freshwater | 365 | 0.30 | 2260 | ng L ⁻¹ | 68, 89, 110, 111, 114 and 119–122 |
| HMMM | Marine water | 11.5 | <0.22 | 57.0 | ng L ⁻¹ | 80, 94 and 115 |
| HMMM | Storm water | 1681 | 0.2 | 11 548 | ng L ⁻¹ | 78, 80, 81 and 96 |
| HMMM | Snow | 3311 | 1110 | 8419 | ng L ⁻¹ | 82 and 96 |
| HMMM | Freshwater sediment | 6.40 | <0.03 | 801 | µg kg ⁻¹ | 85 |
| HMMM | Marine sediment | 0.27 | 0.09 | 0.57 | µg kg ⁻¹ | 94 |
| HMMM | WWTP influent | 7.87 | <1.60 | 23.1 | ng L ⁻¹ | 123 |
| HMMM | WWTP effluent | 39.4 | <1.60 | 83.2 | ng L ⁻¹ | 123 |
| HMMM | Tunnel wash water | 13 129 | 2214 | 18 629 | ng L ⁻¹ | 83 |
| DPG | Freshwater | 139 | 0.50 | 1894 | ng L ⁻¹ | 68, 77, 91, 110, 111, 113, 114 and 119–121 |
| DPG | Marine water | 7.65 | 0.28 | 93.0 | ng L ⁻¹ | 80, 93, 94 and 115 |
| DPG | Storm water | 34 098 | 0.29 | 364 250 | ng L ⁻¹ | 77, 78, 80, 81 and 91 |
| DPG | Snow | 5912 | 15.8 | 14 196 | ng L ⁻¹ | 77 and 82 |
| DPG | Marine sediment | 0.15 | <0.12 | 2.07 | µg kg ⁻¹ | 93 and 94 |
| DPG | Soil | 1100 | 15.0 | 7600 | µg kg ⁻¹ | 87 |
| DPG | WWTP influent | 1086 | 73.0 | 3570 | ng L ⁻¹ | 91 and 101 |
| DPG | WWTP effluent | 123 | 58.2 | 296 | ng L ⁻¹ | 91 and 101 |
| DPG | Tunnel wash water | 29 561 | 2674 | 63 086 | ng L ⁻¹ | 83 |

Furthermore, several transformation products have been identified for the leachables and some still potentially being unknown, which need to be considered. Therefore, it is necessary to define a representative compound mix of tire leachables, including their transformation products, for the exposure assessment of the entire range of tire leachables. To generate such a representative compound mix there is a need for focused data on the assessment of vulcanization and decomposition products of additives, such as potential environmental weathering products of unknown and variable composition of functionally important classes of chemicals, such as amine antioxidants. An illustration of a previously unknown weathering product is 6PPD-Q, which has been identified as a decomposition of 6PPD formed during its intended function of protecting tire tread from oxidative attack.⁷⁹ Subsequently, a broad collection of studies necessary to inform risk

management of this environmentally formed product have been initiated⁸⁸ and it is currently most studied tire leachable.

3.3 Step 2B: Hazard assessment of tire wear emissions

3.3.1 What are relevant endpoints for hazard assessment of tire wear emissions? There are no agreed upon methods for testing the ecotoxicity of particulate tire emissions, however like microplastics, researchers have investigated effects from both direct exposure of organisms to the particles and exposure to leachates created from the particles. Many microplastics researchers have concluded that the relevant endpoints for direct particle exposure is food dilution, wherein part of the organism diet is replaced by the microplastics and creates a false sense of satiation and reduce nutrient ingestion, as well as translocation of microplastics across epithelial barriers which may induce tissue inflammation and oxidative stress.¹⁷ A wide range of particle size is considered for food dilution (1–



5000 μm), but a more limited particle size is considered for tissue translocation (1–83 μm).¹²⁴ For these impacts, test systems have included acute and chronic exposures, and the endpoints evaluated included those at the organism (e.g. mortality, body mass, body length, fecundity), tissue (e.g. Hepatic and gonad somatic index), cell (e.g. cell density and morphology) and sub-cell (e.g. metabolism indicators including glutathione *S*-transferase and superoxide dismutase activity and endocrine indicators like estradiol to testosterone ratio) levels.¹⁷

For exposure to microplastics leachates, no common relevant endpoints have been proposed, which is also the same for particulate tire emission leachates. Endpoints measured for tire particle leachate studies include mortality, growth, reproduction, population increase or decrease, hematological changes, and gene expression. The leachate research studies indicated that the effects from particle leachate are associated with the chemical composition of the particles themselves and are highly variable depending on the test material, age, and leaching conditions.

Mixture toxicity for chemicals associated with tire emissions is assessed when using particulate tire emissions and its surrogates because particulate tire emissions is itself a mixture comprised of the tread compound (a mixture of polymers and additives from different tires) and roadway particles (asphalt, minerals), as well as vehicle-related particles (brake wear particles, tailpipe emission particles).¹²⁵ As such, studies of particulate tire emissions rather than its base polymers provide the most realistic test material for investigation of the ecological hazard properties because the chemical mixture is inherently assessed. Given the difficulty in producing particulate tire emissions in the laboratory, cryogenic milled tire tread (CMTT) particles are the next best surrogate test material at the moment. If environmental materials rich in TRWP such as tunnel road dust may be a better surrogate test material has not been evaluated yet.²⁶

3.3.2 Hazard data exists predominantly for freshwater and terrestrial organisms. The ecological hazard of particulate tire wear emission has been assessed directly through exposure of organisms to the particles directly or indirectly through exposure to the leachates prepared from the particles. Particle exposure data have been presented in 29 studies (Table 3) to a total of 27 different terrestrial, aquatic and sediment dwelling organisms, involving many different test designs. Acute and chronic exposure for freshwater, marine water and terrestrial dwelling organisms have been investigated, while for freshwater sediment and marine sediment dwelling organisms only chronic exposure tests were performed (see Fig. 1). The most common endpoints investigated were those associated with organism level effects including mortality (67% of studies), growth/development (10% of studies), and reproduction (23% of studies). Only two studies were identified wherein sub-cellular effects were evaluated for acute freshwater exposure and include various gene expressions (midge *Chironomus riparius*¹²⁶) and antioxidant enzyme levels (water flea *Daphnia magna*¹²⁷).

Most aquatic toxicity tests of particulate tire wear emission were carried out for acute and chronic exposures of freshwater

organisms, with mortality and reproductive endpoints of *Daphnia magna* being the most studied. Other acute exposure studies have assessed mortality of invertebrates (*H. azteca*, *A. aquaticus*, *C. riparius*), growth for algae (*P. subcapitata*), mortality of amphibian (*S. tropicalis*), and mortality and reproduction in fish (*P. promelas*, *D. rerio*). All the of the available LC50 or EC50 values exceed 10 mg L^{-1} except for the amphibian *S. tropicalis*.¹²⁸ For chronic studies of the freshwater organisms, the data are more scattered with some hazard values less than 10 mg L^{-1} . In general, studies for *D. magna* report no effect to mortality after chronic exposure to particulate tire emission but effects to reproduction at concentrations between approximately 1 to 10 mg L^{-1} .¹²⁹ Growth was impacted in green algae (*P. subcapitata*) at $\sim 31 \text{ mg L}^{-1}$ (ref. 129) and mortality was observed in the amphibian *S. tropicalis* and on zebrafish (*D. rerio*) after chronic exposure to particulate tire emissions.^{128,129}

The toxicity of particulate tire wear emission in freshwater sediment organisms has also been well studied under chronic exposure conditions including mortality for amphibian (*R. sylvatica*), as well as seven invertebrate species (*G. pulex*, *Tubifex*, *C. riparius*, *C. dilutus*, *H. azteca*, *L. variegatus*, *A. aquaticus*) and reproduction for *L. variegatus*. All of the no observed effect concentrations (NOECs) for mortality or reproduction in these organisms are greater than 1000 mg kg^{-1} . There are no studies identified on the acute toxicity of particular tire wear emission in freshwater sediment organisms.

In addition, one study evaluated freshwater fish *in vitro* toxicity of tire wear particulate emissions (as CMTT) in rainbow trout gill (RTgill-W1) and gut (RTgutGC) cell lines.¹³⁰ Cell metabolic activity, membrane integrity, and lysosome integrity were assessed after 24 hour exposure to CMTT or thermooxidized CMTT particles to assess the effect of particle aging on toxicity. In both cell lines, metabolic activity was the most sensitive endpoint. Further, aged CMTT was less toxic in both cell lines. The EC50s for metabolic activity were 2023 mg L^{-1} RTgill-W1 and 4651 mg L^{-1} for RTgutGC.

Terrestrial toxicity studies of particulate tire wear emission have also been conducted and include a limited number of acute mortality studies on soil invertebrates (*E. crypticus*, *F. candida*, *E. andrei*). However, there are 10 studies of terrestrial invertebrates for chronic endpoints of mortality (*E. fetida*, *P. scaber*, *F. candida*, *E. andrei*, *E. crypticus*), and reproduction (*F. Candida*, *E. andrei*, *E. crypticus*). Only one study of a plant species (*V. radiata*) was reported in the literature and no effect was observed at the highest test concentrations.¹³¹ All datasets have hazard values (e.g. LC50, EC50, NOEC) exceeding 1000 mg kg^{-1} , except for one of the studies for *E. crypticus*.¹³²


The toxicity to particulate tire wear emission among marine aquatic and sediment dwelling species has been studied the least, compared to freshwater and terrestrial environments. For marine water organisms, mortality was evaluated across five species, however, no mortality was observed at the highest test concentrations greater than 1 mg L^{-1} for four of the organisms (*F. heteroclitus*, *M. beryllina*, *A. bahia*, *C. virginica*). Only one acute study of the marine copepod, *T. japonicus*, reported a definitive LC50 for mortality.¹³³ There is only one study on the chronic toxicity to particulate tire wear emissions for marine



A) Freshwater



B) Freshwater sediment



C) Marine water



D) Marine sediment



E) Terrestrial



Fig. 1 Environmental concentrations, L(E)C50 and NOEC of direct particulate tire wear exposure: Acute L(E)C50 and chronic NOEC values by species for the compartments (A) freshwater, (B) freshwater sediment, (C) marine water, (D) marine sediment and (E) terrestrial for the endpoint's mortality, reproduction, and growth. Figure contains point estimates, minima, maxima, and ranges. Multiple ranges and estimates are provided per species track and open patterned shapes indicate nondefinitive values (dashed pattern: the actual value is greater than reported value; dotted pattern: the actual value is lower than reported value). Peak shapes refer to the investigated endpoints. X axis is log transformed. Each species is represented by a different color. Taxonomic groups are abbreviated: Al – algae, Am – amphibian, F – fish, In – invertebrate, Pl – plant. For references see Table S2.†



aquatic organisms.¹³⁴ In this study, no effect to mortality was observed after chronic exposure to the highest test concentration.

Similarly, acute exposure to particulate tire wear emission in marine sediment did not result in mortality at the highest test concentrations for two species studied (*S. plana* and *H. diversicolor*). However, sub-lethal endpoints were examined in the sediment organisms in a chronic test where the endpoints included behavioral (feeding and burial rates), physiological (condition index) and biochemical (energy reserves and oxidative stress) markers; although differences between controls and exposed groups were only observed at the highest test concentration of 50 000 mg kg⁻¹.¹³⁵ Additional studies of marine sediment organisms are necessary to support hazard assessment for the marine sediment compartment.

Many research studies have investigated the toxicity of tire particle leachates to aquatic organisms (Table S3†). Thirty studies have investigated mortality, reproduction, growth, and other endpoints across the main trophic levels of fish, invertebrate, amphibian, algae/plant, and various microbes. In all, 46 species have been evaluated. Given the lack of a standardized leachate testing framework, however, it is extremely difficult to compare across the studies to characterize the potential adverse effect levels. The studies varied significantly in the leachate preparation with respect to the solid to liquid ratio which ranged from 0.1 to >100 g L⁻¹, percent concentration of leachate for the dose groups, duration of the leaching, and temperature of the solution. Additionally, the test material was quite different from study to study. Nevertheless, findings from these studies show effects may occur in some species after exposure to leachate with preparation concentrations equal to or greater than 0.1 g tire particle per L.

In addition to the *in vivo* toxicity tests described above for tire particulate emissions, Bergmann *et al.*¹³⁶ applied three high-performance thin-layer chromatography (HPTLC) bioassays to evaluate the estrogenicity, genotoxicity, and bacterial luminescence inhibition of leachable chemicals from CMTT. Aqueous leachates of CMTT were prepared at 100 g L⁻¹ under different conditions (*e.g.*, with or without sediment). The authors reported that estrogenic and luminescence-inhibiting chemicals leached from CMTT into aqueous media. No genotoxicity was detected in CMTT leachates. None of the other leachables studies directly evaluated the potential for endocrine disruption. However, several studies identified chemicals in the leachate solutions that have been identified as endocrine disruptors including bis-phenol A,^{137,138} and DiBP, DBP, BBP and DEHP.³⁷ Recent guidance from the European Chemicals Agency (ECHA) regarding the classification of endocrine disrupting properties states that the classification criteria require evidence on three elements, including adverse effect(s), endocrine activity, and a biologically plausible link between the endocrine activity and the adverse effect(s).¹³⁹ Overall, the acute toxicity of tire emissions as particle or leachate to aquatic species has been well studied, however, it may be helpful to have tire particle toxicity data on other fish species. Additionally, there are numerous chronic studies for freshwater aquatic and sediment organisms as well as terrestrial organisms,

however chronic data for the marine sediment species are lacking. Lastly, the potential for endocrine disruption from the tire emission leachables has not been well studied, and as such additional characterization *via* bioassays using environmentally relevant leachate solutions are necessary to accurately characterize this hazard endpoint. Thus, additional studies of species across the trophic levels and expansion of hazard data for the marine environment and to characterize the potential for endocrine disruption are necessary for a risk assessment.

3.3.3 Trophic transfer of tire chemicals from leachate is more relevant as for tire particles. Several of the tire particle emission studies showed that laboratory organisms can ingest tire particle emissions when present in their habitat, however, once removed from the exposure scenario the particles were egested.^{140–143} The uptake of chemical constituents from the tire particles was demonstrated in three laboratory studies to investigate the potential for bioaccumulation. Hägg *et al.*¹⁴³ found no significant difference in metal concentrations between the controls and the exposed fish, but for targeted organic chemicals there were significant differences, demonstrating the bioaccessibility of phenylenediamine substances. Bioaccumulation of tire-related chemicals was also demonstrated in earthworms *E. andrei*.¹⁴⁴ In an *in vitro* study, Masset *et al.*¹⁴⁵ demonstrated the low to moderate bioaccessibility of various tire related chemicals including, their transformation products and PAHs. One study was identified that investigated a natural estuary for the presence of various tire additive chemicals and their transformation products.⁹⁵ The authors evaluated biota samples in the food web of the estuary (snails, shrimp, sea cucumbers and fish) and reported bioaccumulation for two phenylenediamine substances and through Monte Carlo analysis predicted biomagnification trends. Overall, studies related to trophic transfer of tire related chemicals are limited and warrant further research.

3.4 Step 3: Risk characterization of tire wear emissions in the environment

Environmental risk characterization of tire wear emissions utilizes exposure concentration and hazard data. For a risk characterization of tire wear emissions, both particulate tire emissions such as TRWP and dissolved tire leachables are of interest. In risk characterization the exposure concentrations are compared to the effect concentrations. In this work we derived relevant exposure pathways,²² assessed the exposure concentrations and hazard data based on the structure literature review. Although the available exposure concentration data lacks comparability, as does the hazard data because of differences in the applied methodologies, it offers the opportunity to at least develop an idea on potentially sensitive species and affected environmental compartments. As a screening level risk prioritization, we identify environmental compartments and species of interest and comparing the exposure data with the hazard data for different environmental compartments and test organisms (Fig. 1).

3.4.1 Data reveals low risk prioritization of particulate tire wear emissions. The concentrations of tire wear particulates are



shown in the first row of Fig. 1 for each of the environmental compartments. For freshwater organisms, there are only 4 datasets available to which we can compare the hazard values. Nevertheless, all species have acute hazard values that are orders of magnitude above the median and 75th percentile measured concentrations, except for the *S. tropicalis*,¹²⁸ where the hazard values are within the range of the measured concentrations. A comparison of the chronic hazard values is similar to that of the acute, where most are an order of magnitude higher than the median measured water concentration, except for *S. tropicalis*.¹²⁸ Nevertheless, there is an appreciable lack of environmental measurements in the freshwater compartment and the available concentrations span 4 orders of magnitude (Table 4) suggesting highly location specific influences, such as traffic levels, extent of impervious surfaces, stormwater discharge, and sampling conditions (*i.e.*, wet or dry). As such, additional freshwater measurement data is crucial to drawing a firm conclusion regarding risk prioritization.

For freshwater sediment, there are eight datasets available to which we can compare the hazard values. All of the tested species have hazard values that are well above the median and 75th percentile measured concentrations, some by more than 2 orders of magnitude. This screening level comparison indicates a low prioritization of risk, although similar to the water measurements, the concentrations span many orders of magnitude and are likely influenced by the same factors.

The marine environment including water and sediment are not well characterized, with only one measured dataset for each compartment. Although quite limited in terms of species, where hazard values are available, they are many orders of magnitude higher than the measured sediment concentrations. Nevertheless, the sparse environmental concentration data for the marine environment, coupled with the limited hazard dataset do not allow for confidence in conclusions regarding prioritization for risk to marine organisms.

For terrestrial organisms, there are three soil concentration data sets available to which we can compare the hazard values. Although quite limited in terms of terrestrial species, the acute hazard values indicated no effects at concentrations within the range of the reported soil concentrations. More species hazard data is available for the chronic exposures, but similar to the acute hazard values, no effects were reported at concentrations within the range of the measured soil concentrations, except for *E. crypticus*.¹³² For this species, three studies were identified and one of the three showed adverse effects at concentrations lower than the range of soil measurements and 2–3 orders of magnitude lower than the other studies of this organism. Overall, the screening level comparison indicates a low potential for risk to terrestrial organisms, however, additional investigation of *E. crypticus* may be helpful to understand the marked difference in hazard values for the same endpoints in this species.

3.4.2 Risk prioritization of leachables from tire emissions is not yet possible due to inconsistent hazard data sets. The available hazard data of leachate from particulate tire wear allows for an understanding of the risk prioritization from tire

emissions that are not related to direct exposure to the particles. These data are most relevant for risk assessment in the road run-off exposure scenario where precipitation initiates leaching of deposited particulate tire emissions allowing chemical components of the particle to migrate into the water phase and subsequently transported to the aquatic and terrestrial environment *via* direct discharge to surface waters, wastewater treatment plants or roadside soil. The study of coho salmon urban runoff mortality syndrome is a case where observations of acute mortality in freshwater streams were associated with rainfall events and chemical analysis of the stormwater lead to implications for chemicals associated with tires.⁷⁹ A toxic identification evaluation study from tire particle leachate revealed that the mortality may be associated with 6PPD-Q, rather than the overall mixture of the leachate. Subsequent laboratory studies showed similarity of effects when 6PPD-Q was specifically studied and additional studies demonstrated coho salmon as a highly sensitive species.^{79,146,147} Given that the effects seen in the tire emission leachate studies are likely associated with the chemicals in the leachate, the chemical composition resulting from the leachate preparation is important to characterize; recognizing of course that the chemicals may include parent chemicals present in the tread rubber, as well as transformation products resulting from interactions with the ambient environmental conditions and weathering over time.

4 Risk characterization for tire wear emissions is not only challenged by exposure and hazard data

4.1 From laboratory to reality: the key gaps for tire wear risk assessment

Risk assessment requires a thorough definition of the problem addressing its scope and purpose. The problem formulation usually reflects potential harms, the pathways to those harms, and the information needed to assess the risk. The necessary particulate tire wear emission and non-tire microplastic hazard and exposure studies have not been completed to fully inform the problem formulation.^{7,11,22,39,79,148–150} The tire wear specific data gap identified consistently in the three stages (problem, hazard & exposure and risk) is the assessment of the unknown and variable composition components (UVCC) and weathering products of tread additives which may either be related to particulate contaminants or leachables. Tread additives must already comply with international regulatory environmental safety requirements, often including chemical-specific toxicity testing inclusive of unknown or variable composition of the chemical as formulated.¹⁵¹ These assessments include quantitative measures of risk, such as the risk quotient derived from predicted environmental concentrations and predicted no-effect concentrations. These data must be comparable and uncertainties quantifiable. It is therefore necessary to establish and apply harmonized analytical methods (sampling, identification, and quantification) for tire wear emissions for the relevant environmental compartments and hazard tests. While



there is already a number of data points for the aquatic environment available, the review revealed that exposure data of particulate tire wear emissions in terrestrial environments and for leachables and UVCC substances for various environmental compartments *e.g.* groundwater is required for more comprehensive risk characterization.

Risk assessment also relates to representative and harmonized test materials for tire wear emissions bridging the gap between laboratory and reality. Hazard testing is often challenged by the complexity of the real conditions making the identification of effect drivers hardly possible. Therefore, it is proposed to implement a tiered approach for hazard assessment starting with effect testing of tire tread leachates and particulate tire wear making use of suitable and harmonized test systems and test materials. In a second tier, if positive findings occur, further tests under more realistic conditions *e.g.* presence of matrices should be conducted. In a third tier, depending whether local or global assessments are performed, data on *e.g.* site specific environmental concentrations of tire wear particles and tire leachable concentrations including transformation products; reliable emission data, global distribution data incl. uncertainties and most affected environments for particles and leachables, fate data (transformation, breakdown, half-life), information on sensitive species, are needed. Such a stepwise approach combined with harmonized and robust methods for sampling and sample analysis as well as hazard testing would reduce the uncertainty of the data and thus enhance its interpretability.

4.2 Building on progress: globally harmonized risk assessment of tire wear emissions

The implementation of an extended risk assessment framework requires interdisciplinary action from various stakeholders including tire and automotive industry, road management, and regulation. There are already existing frameworks which may help the development of such an extension *e.g.* the safe and sustainable-by-design framework.¹⁵² Tire wear emissions do not conform to the International Organization for Standardization definition of plastic as shaped by flow.^{153–155} Thus, appreciable physiochemical and morphological differences exist between particulate tire wear emissions such as TRWP, which contain tread elastomer as well as road materials and thermoset plastics or thermoplastics-derived microplastics. As such, it is useful to examine the essential components of the PDF as described by Koelmans *et al.* (2023)¹¹ and assess the suitability or adaptation of each element for the study of TRWP (Table S4†). Overall, the PDF framework for MPs is suitable for the risk assessment and communication of environmental tire wear emission risks. Much of the information and tools necessary to perform the screening and first tiers of an environmental risk assessment for tire wear emissions have already been generated as it was demonstrated in Sections 2 and 3. However, there are still knowledge and data gaps with respect to higher tiers. Therefore, research on tire wear emission would benefit from a clear harmonization guideline how exposure and hazard studies should be designed. Such guidelines should be developed

between all relevant stakeholder covering the entire product life cycle.

5 The way forward: suggestions for future tire wear emission research

While progress has been made in exposure and hazard assessment of tire wear emissions in the environment, the complexity of tire wear emissions creates challenges which are not yet overcome. The following challenges, needs and data gaps for environmental risk assessment of tire wear emissions were derived:

- While there is existing aquatic exposure data, the review highlights the necessity for additional data on tire wear emissions in terrestrial environments and for leachables and substances of unknown or variable composition, including transformation products in environmental media, such as groundwater, freshwater and marine environments.
- A key data gap identified across all stages of assessment (problem, hazard, and exposure) is the evaluation of unknown and variable composition components and weathering products from tread additives, which are related to both particulate contaminants and leachables.
- There is a recommendation for further research into the trophic transfer of tire-related chemicals, including studies on bioaccumulation in various biota and the potential for biomagnification in food webs.
- For a risk assessment, insufficient consistent exposure and hazard data are available. Quantitative risk measures, like risk quotients derived from predicted environmental and no-effect concentrations, must be comparable and have quantifiable uncertainties. Consequently, there is a need to establish harmonized analytical methods for the sampling, identification, and quantification of tire wear emissions across relevant environmental media and hazard tests. These guidelines should be collaboratively established by all relevant stakeholders involved throughout the product life cycle and aligned with existing regulatory practices.

Data availability

The data supporting this article have been included as part of the ESI.†

Conflicts of interest

The authors declare the following facts which may be considered as potential conflict of interest: Kathrin Müller, Julie Panko, Kenny Unice and Stephan Wagner report that partial financial support was provided for the research by the Tire Industry Project, which is a part of the World Business Council for Sustainable Development, Switzerland.

Acknowledgements

The authors acknowledge Stephanie Kennedy for her technical support in the hazard data visualization. The authors



acknowledge John D. Spengler (Harvard T.H. Chan School of Public Health) and Ng How Yong (Beijing Normal University Zhuhai) for their scientific discussion on the topic of tire emissions which helped structure the manuscript and present the findings of the literature study. The work presented here was funded by the Tire Industry Project (TIP), which operates under the umbrella of the World Business Council for Sustainable Development (WBCSD), with the study design, execution, interpretation, and manuscript preparation independently conducted by the authors. Formed in 2005, the Tire Industry Project (TIP) is a voluntary CEO-driven initiative with a mission to anticipate, understand, and address global environmental, social, and governance (ESG) issues relevant to the tire industry and its value chain. TIP acts by commissioning independent research of the highest standards, collaborating on sectoral solutions, and engaging with external stakeholders. TIP currently brings together 10 leading tire companies that represent more than 60% of the world's tire manufacturing capacity.

References

- OICA, *World Motor Vehicle Production*, OICA Correspondents Survey, <https://www.oica.net/wp-content/uploads/By-country-region-2024.pdf>, accessed 2025-05-08.
- B. Baensch-Baltruschat, B. Kocher, C. Kochleus, F. Stock and G. Reifferscheid, Tyre and Road Wear Particles - A Calculation of Generation, Transport and Release to Water and Soil with Special Regard to German Roads, *Sci. Total Environ.*, 2021, 752, 141939, DOI: [10.1016/j.scitotenv.2020.141939](https://doi.org/10.1016/j.scitotenv.2020.141939).
- P. M. Mayer, K. D. Moran, E. L. Miller, S. M. Brander, S. Harper, M. Garcia-Jaramillo, V. Carrasco-Navarro, K. T. Ho, R. M. Burgess, L. M. Thornton Hampton, E. F. Granek, M. McCauley, J. K. McIntyre, E. P. Kolodziej, X. Hu, A. J. Williams, B. A. Beckingham, M. E. Jackson, R. D. Sanders-Smith, C. L. Fender, G. A. King, M. Bollman, S. S. Kaushal, B. E. Cunningham, S. J. Hutton, J. Lang, H. V. Goss, S. Siddiqui, R. Sutton, D. Lin and M. Mendez, Where the Rubber Meets the Road: Emerging Environmental Impacts of Tire Wear Particles and Their Chemical Cocktails, *Sci. Total Environ.*, 2024, 927, 171153, DOI: [10.1016/J.SCITOTENV.2024.171153](https://doi.org/10.1016/J.SCITOTENV.2024.171153).
- S. Hann, C. Sherrington, O. Jamieson, M. Hickman, P. Kershaw, A. Bapasola and G. Cole, *Investigating Options for Reducing Releases in the Aquatic Environment of Microplastics Emitted by (But Not Intentionally Added in) Products Interim Report*, London, 2018, <http://www.eumicroplastics.com/eumpwp/wp-content/uploads/investigating-options-eunomia-draft-report-v4-main-report-public.pdf>.
- OECD, *Non-Exhaust Particulate Emissions from Road Transport: An Ignored Environmental Policy Challenge*, OECD Publishing, Paris, 2020, DOI: [10.1787/4a4dc6ca-en](https://doi.org/10.1787/4a4dc6ca-en).
- K. McCarty, H. R. Mian, G. Chhipi-Shrestha, K. Hewage and R. Sadiq, Ecological Risk Assessment of Tire and Road Wear Particles: A Preliminary Screening for Freshwater Sources in Canada, *Environ. Pollut.*, 2023, 325, 121354, DOI: [10.1016/j.envpol.2023.121354](https://doi.org/10.1016/j.envpol.2023.121354).
- A. Wik and G. Dave, Occurrence and Effects of Tire Wear Particles in the Environment – A Critical Review and an Initial Risk Assessment, *Environ. Pollut.*, 2009, 157(1), 1–11, DOI: [10.1016/j.envpol.2008.09.028](https://doi.org/10.1016/j.envpol.2008.09.028).
- I. Lerche and W. Glaesser, *Environmental Risk Assessment*, Springer Berlin Heidelberg, Berlin, Heidelberg, 2006, DOI: [10.1007/3-540-29709-X](https://doi.org/10.1007/3-540-29709-X).
- I. V. Sukhorukova and N. A. Chistyakova, Mathematical Model for Assessing Environmental Risk, *IOP Conf. Ser.: Mater. Sci. Eng.*, 2020, 828(1), 012027, DOI: [10.1088/1757-899X/828/1/012027](https://doi.org/10.1088/1757-899X/828/1/012027).
- A. A. Koelmans, P. E. Redondo-Hasselerharm, N. H. M. Nor, V. N. de Ruijter, S. M. Mintenig and M. Kooi, Risk Assessment of Microplastic Particles, *Nat. Rev. Mater.*, 2022, 7(2), 138–152, DOI: [10.1038/s41578-021-00411-y](https://doi.org/10.1038/s41578-021-00411-y).
- A. A. Koelmans, B. M. Gebreyohanes Belay, S. M. Mintenig, N. H. Mohamed Nor, P. E. Redondo-Hasselerharm and V. N. de Ruijter, Towards a Rational and Efficient Risk Assessment for Microplastics, *TrAC, Trends Anal. Chem.*, 2023, 165, 117142, DOI: [10.1016/j.trac.2023.117142](https://doi.org/10.1016/j.trac.2023.117142).
- V. Adam, T. Yang and B. Nowack, Toward an Ecotoxicological Risk Assessment of Microplastics: Comparison of Available Hazard and Exposure Data in Freshwaters, *Environ. Toxicol. Chem.*, 2019, 38(2), 436–447, DOI: [10.1002/etc.4323](https://doi.org/10.1002/etc.4323).
- C. Schmidt, R. Kumar, S. Yang and O. Büttner, Microplastic Particle Emission from Wastewater Treatment Plant Effluents into River Networks in Germany: Loads, Spatial Patterns of Concentrations and Potential Toxicity, *Sci. Total Environ.*, 2020, 737, 139544, DOI: [10.1016/J.SCITOTENV.2020.139544](https://doi.org/10.1016/J.SCITOTENV.2020.139544).
- F. R. Khan, E. S. Rødland, P. J. Kole, F. G. A. J. Van Belleghem, A. Jaén-Gil, S. F. Hansen and A. Gomiero, An Overview of the Key Topics Related to the Study of Tire Particles and Their Chemical Leachates: From Problems to Solutions, *TrAC, Trends Anal. Chem.*, 2024, 172, 117563, DOI: [10.1016/J.TRAC.2024.117563](https://doi.org/10.1016/J.TRAC.2024.117563).
- T. Gouin, R. A. Becker, A. G. Collot, J. W. Davis, B. Howard, K. Inawaka, M. Lampi, B. S. Ramon, J. Shi and P. W. Hopp, Toward the Development and Application of an Environmental Risk Assessment Framework for Microplastic, *Environmental Toxicology and Chemistry*, Wiley Blackwell, 2019, pp. 2087–2100, DOI: [10.1002/etc.4529](https://doi.org/10.1002/etc.4529).
- A. A. Koelmans, P. E. Redondo-Hasselerharm, N. H. Mohamed Nor and M. Kooi, Solving the Nonalignment of Methods and Approaches Used in Microplastic Research to Consistently Characterize Risk, *Environ. Sci. Technol.*, 2020, 54(19), 12307–12315, DOI: [10.1021/acs.est.0c02982](https://doi.org/10.1021/acs.est.0c02982).
- A. C. Mehinto, S. Coffin, A. A. Koelmans, S. M. Brander, M. Wagner, L. M. Thornton Hampton, A. G. Burton, E. Miller, T. Gouin, S. B. Weisberg and C. M. Rochman, Risk-Based Management Framework for Microplastics in



- Aquatic Ecosystems, *Microplast. Nanoplast.*, 2022, 2(17), DOI: [10.1186/s43591-022-00033-3](https://doi.org/10.1186/s43591-022-00033-3).
- 18 K. Bucci and C. M. Rochman, Microplastics: A Multidimensional Contaminant Requires a Multidimensional Framework for Assessing Risk, *Microplast. Nanoplast.*, 2022, 2(7), DOI: [10.1186/s43591-022-00028-0](https://doi.org/10.1186/s43591-022-00028-0).
- 19 P. Chakraborty, G. K. Bharat, O. Gaonkar, M. Mukhopadhyay, S. Chandra, E. H. Steindal and L. Nizzetto, Endocrine-Disrupting Chemicals Used as Common Plastic Additives: Levels, Profiles, and Human Dietary Exposure from the Indian Food Basket, *Sci. Total Environ.*, 2022, **810**, 152200, DOI: [10.1016/J.SCITOTENV.2021.152200](https://doi.org/10.1016/j.scitotenv.2021.152200).
- 20 A. A. Koelmans, E. Besseling, E. Foekema, M. Kooi, S. Mintenig, B. C. Ossendorp, P. E. Redondo-Hasselerharm, A. Verschoor, A. P. Van Wezel and M. Scheffer, Risks of Plastic Debris: Unravelling Fact, Opinion, Perception, and Belief, *Environ. Sci. Technol.*, 2017, **51**(20), 11513–11519, DOI: [10.1021/acs.est.7b02219](https://doi.org/10.1021/acs.est.7b02219).
- 21 M. Kooi, S. Primpke, S. M. Mintenig, C. Lorenz, G. Gerdt and A. A. Koelmans, Characterizing the Multidimensionality of Microplastics across Environmental Compartments, *Water Res.*, 2021, **202**, 117429, DOI: [10.1016/j.watres.2021.117429](https://doi.org/10.1016/j.watres.2021.117429).
- 22 B. Baensch-Baltruschat, B. Bänsch-Baltruschat, B. Kocher, F. Stock, F. Stock, F. Stock and G. Reifferscheid, Tyre and Road Wear Particles (TRWP) - A Review of Generation, Properties, Emissions, Human Health Risk, Ecotoxicity, and Fate in the Environment, *Sci. Total Environ.*, 2020, **733**, 137823, DOI: [10.1016/j.scitotenv.2020.137823](https://doi.org/10.1016/j.scitotenv.2020.137823).
- 23 J. E. Tamis, A. A. Koelmans, R. Dröge, N. H. B. M. Kaag, M. C. Keur, P. C. Tromp and R. H. Jongbloed, Environmental Risks of Car Tire Microplastic Particles and Other Road Runoff Pollutants, *Microplast. Nanoplast.*, 2021, **1**(1), 1–17, DOI: [10.1186/S43591-021-00008-W/METRICS](https://doi.org/10.1186/S43591-021-00008-W/METRICS).
- 24 K. Müller, K. Unice, J. Panko and S. Wagner, Tire Emissions during the Use Phase of Tires - Today and Future Trends, *Environ. Sci.: Adv.*, 2025, DOI: [10.1039/D4VA00407H](https://doi.org/10.1039/D4VA00407H).
- 25 I. Gofsmann, M. Halbach and B. M. Scholz-Böttcher, Car and Truck Tire Wear Particles in Complex Environmental Samples – A Quantitative Comparison with “Traditional” Microplastic Polymer Mass Loads, *Sci. Total Environ.*, 2021, **773**, 145667, DOI: [10.1016/j.scitotenv.2021.145667](https://doi.org/10.1016/j.scitotenv.2021.145667).
- 26 S. Wagner, P. Klöckner and T. Reemtsma, Aging of Tire and Road Wear Particles in Terrestrial and Freshwater Environments - A Review on Processes, Testing, Analysis and Impact, *Chemosphere*, 2022, **288**, 132467, DOI: [10.1016/j.chemosphere.2021.132467](https://doi.org/10.1016/j.chemosphere.2021.132467).
- 27 L. L. Halle, A. Palmqvist, K. Kampmann, A. Jensen, T. Hansen and F. R. Khan, Tire Wear Particle and Leachate Exposures from a Pristine and Road-Worn Tire to *Hyalella Azteca*: Comparison of Chemical Content and Biological Effects, *Aquat. Toxicol.*, 2021, **232**, 105769, DOI: [10.1016/j.aquatox.2021.105769](https://doi.org/10.1016/j.aquatox.2021.105769).
- 28 C. A. Marwood, B. L. McAtee, M. L. Kreider, R. S. Ogle, B. L. Finley, L. Sweet and J. M. Panko, Acute Aquatic Toxicity of Tire and Road Wear Particles to Alga, Daphnid, and Fish, *Ecotoxicology*, 2011, **20**, 2079–2089, DOI: [10.1007/s10646-011-0750-x](https://doi.org/10.1007/s10646-011-0750-x).
- 29 K. M. Unice, M. L. Kreider and J. M. Panko, Comparison of Tire and Road Wear Particle Concentrations in Sediment for Watersheds in France, Japan, and the United States by Quantitative Pyrolysis GC/MS Analysis, *Environ. Sci. Technol.*, 2013, **47**, 8183–8147, DOI: [10.1021/es400871j](https://doi.org/10.1021/es400871j).
- 30 K. Müller, D. Hübner, S. Huppertsberg, T. P. Knepper and D. Zahn, Probing the Chemical Complexity of Tires: Identification of Potential Tire-Borne Water Contaminants with High-Resolution Mass Spectrometry, *Sci. Total Environ.*, 2022, **802**, 149799, DOI: [10.1016/j.scitotenv.2021.149799](https://doi.org/10.1016/j.scitotenv.2021.149799).
- 31 P. Klöckner, T. Reemtsma, P. Eisentraut, U. Braun, A. S. Ruhl and S. Wagner, Tire and Road Wear Particles in Road Environment – Quantification and Assessment of Particle Dynamics by Zn Determination after Density Separation, *Chemosphere*, 2019, **222**, 714–721, DOI: [10.1016/j.chemosphere.2019.01.176](https://doi.org/10.1016/j.chemosphere.2019.01.176).
- 32 E. S. Rødland, G. Binda, D. Spanu, S. Carnati, L. R. Bjerke and L. Nizzetto, Are Eco-Friendly “Green” Tires Also Chemically Green? Comparing Metals, Rubbers and Selected Organic Compounds in Green and Conventional Tires, *J. Hazard. Mater.*, 2024, **476**, 135042, DOI: [10.1016/J.JHAZMAT.2024.135042](https://doi.org/10.1016/J.JHAZMAT.2024.135042).
- 33 S. Weyrauch, B. Seiwert, M. Voll, S. Wagner and T. Reemtsma, Accelerated Aging of Tire and Road Wear Particles by Elevated Temperature, Artificial Sunlight and Mechanical Stress — A Laboratory Study on Particle Properties, Extractables and Leachables, *Sci. Total Environ.*, 2023, **904**, 166679, DOI: [10.1016/j.scitotenv.2023.166679](https://doi.org/10.1016/j.scitotenv.2023.166679).
- 34 USTMA, U.S. Tire Manufacturers Association to Produce Standardized Tire Tread Particles to Advance Scientific Research, <https://www.ustires.org/newsroom/us-tire-manufacturers-association-produce-standardized-tire-tread-particles-advance>, accessed 2025-02-02.
- 35 A. Foscari, N. Schmidt, B. Seiwert, D. Herzke, R. Sempéré and T. Reemtsma, Leaching of Chemicals and DOC from Tire Particles under Simulated Marine Conditions, *Front. Environ. Sci.*, 2023, **11**, 1206449, DOI: [10.3389/fenvs.2023.1206449](https://doi.org/10.3389/fenvs.2023.1206449).
- 36 Y. Jeong, S. Lee and S.-H. Woo, Chemical Leaching from Tire Wear Particles with Various Treadwear Ratings, *Int. J. Environ. Res. Public Health*, 2022, **19**(10), 6006, DOI: [10.3390/ijerph19106006](https://doi.org/10.3390/ijerph19106006).
- 37 C. Halsband, L. Sørensen, A. M. Booth and D. Herzke, Car Tire Crumb Rubber: Does Leaching Produce a Toxic Chemical Cocktail in Coastal Marine Systems?, *Front. Environ. Sci.*, 2020, **8**(125), DOI: [10.3389/fenvs.2020.00125](https://doi.org/10.3389/fenvs.2020.00125).
- 38 K. M. Unice, J. L. Bare, M. L. Kreider and J. M. Panko, Experimental Methodology for Assessing the Environmental Fate of Organic Chemicals in Polymer Matrices Using Column Leaching Studies and OECD 308



- Water/Sediment Systems: Application to Tire and Road Wear Particles, *Sci. Total Environ.*, 2015, **533**, 476–487, DOI: [10.1016/j.scitotenv.2015.06.053](https://doi.org/10.1016/j.scitotenv.2015.06.053).
- 39 S. Wagner, T. Hüffer, P. Klöckner, M. Wehrhahn, T. Hofmann and T. Reemtsma, Tire Wear Particles in the Aquatic Environment - A Review on Generation, Analysis, Occurrence, Fate and Effects, *Water Res.*, 2018, **139**, 83–100, DOI: [10.1016/j.watres.2018.03.051](https://doi.org/10.1016/j.watres.2018.03.051).
- 40 H. Kumata, J. Yamada, K. Masuda, H. Takada, Y. Sato, T. Sakurai and K. Fujiwara, Benzothiazolamines as Tire-Derived Molecular Markers: Sorptive Behavior in Street Runoff and Application to Source Apportioning, *Environ. Sci. Technol.*, 2002, **36**(4), 702–708, DOI: [10.1021/es0155229](https://doi.org/10.1021/es0155229).
- 41 H. Kumata, Y. Sanada, H. Takada and T. Ueno, Historical Trends of N-Cyclohexyl-2-Benzothiazolamine, 2-(4-Morpholinyl)Benzothiazole, and Other Anthropogenic Contaminants in the Urban Reservoir Sediment Core, *Environ. Sci. Technol.*, 2000, **34**(2), 246–253, DOI: [10.1021/es990738k](https://doi.org/10.1021/es990738k).
- 42 E. S. Rødland, L. S. Heier, O. C. Lind and S. Meland, High Levels of Tire Wear Particles in Soils along Low Traffic Roads, *Sci. Total Environ.*, 2023, **903**, 166470, DOI: [10.1016/j.scitotenv.2023.166470](https://doi.org/10.1016/j.scitotenv.2023.166470).
- 43 F. N. F. Parker-Jurd, I. E. Napper, G. D. Abbott, S. Hann and R. C. Thompson, Quantifying the Release of Tyre Wear Particles to the Marine Environment via Multiple Pathways, *Mar. Pollut. Bull.*, 2021, **172**, 112897, DOI: [10.1016/j.marpolbul.2021.112897](https://doi.org/10.1016/j.marpolbul.2021.112897).
- 44 E. Rødland and Y. Lin, Actions Are Needed to Deal with the High Uncertainties in Tire Wear Particle Analyses, *Environ. Sci. Technol.*, 2023, **57**(23), 8461–8462, DOI: [10.1021/ACS.EST.3C02393/ASSET/IMAGES/LARGE/ES3C02393_0002.JPEG](https://doi.org/10.1021/ACS.EST.3C02393/ASSET/IMAGES/LARGE/ES3C02393_0002.JPEG).
- 45 P. Klöckner, B. Seiwert, P. Eisentraut, U. Braun, T. Reemtsma and S. Wagner, Characterization of Tire and Road Wear Particles from Road Runoff Indicates Highly Dynamic Particle Properties, *Water Res.*, 2020, **185**, 116262, DOI: [10.1016/j.watres.2020.116262](https://doi.org/10.1016/j.watres.2020.116262).
- 46 T. R. Barber, F. Ribeiro, S. Claes, Y. Kawamura, J. Yeung, H. A. Byrne, S. Weyrauch, T. Reemtsma and K. M. Unice, The Identification and Quantification of Tire and Road Wear Particles in Osaka Bay, Japan, by Two Analytical Methods, *Mar. Pollut. Bull.*, 2025, **211**, 117363, DOI: [10.1016/j.marpolbul.2024.117363](https://doi.org/10.1016/j.marpolbul.2024.117363).
- 47 T. Feißel, M. Kunze, D. Hesse, V. Ivanov, K. Augsburg and S. Gramstat, On-Road Vehicle Measurement of Tire Wear Particle Emissions and Approach for Emission Prediction, *Tire Sci. Technol.*, 2024, **52**(1), 2–14, DOI: [10.2346/tire.22.21024](https://doi.org/10.2346/tire.22.21024).
- 48 D. Hesse, T. Feißel, M. Kunze, E. Bachmann, T. Bachmann and S. Gramstat, Comparison of Methods for Sampling Particulate Emissions from Tires under Different Test Environments, *Atmosphere*, 2022, **13**(8), 1262, DOI: [10.3390/atmos13081262](https://doi.org/10.3390/atmos13081262).
- 49 Y. Li, S. Zuo, L. Lei, Y. Xianwu, X. Yang and X. Wu, Analysis of Impact Factors of Tire Wear, *J. Vib. Control*, 2012, **18**(6), 833–840, DOI: [10.1177/1077546311411756](https://doi.org/10.1177/1077546311411756).
- 50 S. Schläfle, H.-J. Unrau and F. Gauterin, Influence of Load Condition, Tire Type, and Ambient Temperature on the Emission of Tire–Road Particulate Matter, *Atmosphere*, 2023, **14**, 1095, DOI: [10.3390/atmos14071095](https://doi.org/10.3390/atmos14071095).
- 51 S. Schläfle, M. Zhang, H. J. Unrau and F. Gauterin, Influence of Vertical Load, Inflation Pressure, and Driving Speed on the Emission of Tire–Road Particulate Matter and Its Size Distribution, *Atmosphere*, 2024, **15**, 502, DOI: [10.3390/atmos15040502](https://doi.org/10.3390/atmos15040502).
- 52 T. De Oliveira, B. Muresan, S. Ricordel, L. Lumière, X.-T. Truong, L. Poirier and J. Gasperi, Realistic Assessment of Tire and Road Wear Particle Emissions and Their Influencing Factors on Different Types of Roads, *J. Hazard. Mater.*, 2024, **465**, 133301, DOI: [10.1016/j.jhazmat.2023.133301](https://doi.org/10.1016/j.jhazmat.2023.133301).
- 53 Y. Ren, W. Li, Q. Jia, Y. Zhao, C. Qu, L. Liu, J. Liu and C. Wu, Separation and Quantification of Tire and Road Wear Particles in Road Dust Samples: Bonded-Sulfur as a Novel Marker, *J. Hazard. Mater.*, 2024, **465**, 133089, DOI: [10.1016/j.jhazmat.2023.133089](https://doi.org/10.1016/j.jhazmat.2023.133089).
- 54 E. Chae, U. Jung and S. S. Choi, Types and Concentrations of Tire Wear Particles (TWPs) in Road Dust Generated in Slow Lanes, *Environ. Pollut.*, 2024, **346**, 123670, DOI: [10.1016/j.envpol.2024.123670](https://doi.org/10.1016/j.envpol.2024.123670).
- 55 A. Müller, B. Kocher, K. Altmann and U. Braun, Determination of Tire Wear Markers in Soil Samples and Their Distribution in a Roadside Soil, *Chemosphere*, 2022, **294**, 133653, DOI: [10.1016/j.chemosphere.2022.133653](https://doi.org/10.1016/j.chemosphere.2022.133653).
- 56 J. Thomas, T. Cutright, C. Pugh and M. D. Soucek, Quantitative Assessment of Additive Leachates in Abiotic Weathered Tire Cryogrinds and Its Application to Tire Wear Particles in Roadside Soil Samples, *Chemosphere*, 2023, **311**, 137132, DOI: [10.1016/j.chemosphere.2022.137132](https://doi.org/10.1016/j.chemosphere.2022.137132).
- 57 E. S. Rødland, S. Samanipour, C. Rauert, E. D. Okoffo, M. J. Reid, L. S. Heier, O. C. Lind, K. V. Thomas and S. Meland, A Novel Method for the Quantification of Tire and Polymer-Modified Bitumen Particles in Environmental Samples by Pyrolysis Gas Chromatography Mass Spectroscopy, *J. Hazard. Mater.*, 2022, **423**, 127092, DOI: [10.1016/j.jhazmat.2021.127092](https://doi.org/10.1016/j.jhazmat.2021.127092).
- 58 T. De Oliveira, B. Muresan, S. Ricordel, L. Lumière, X. T. Truong, L. Poirier and J. Gasperi, Realistic Assessment of Tire and Road Wear Particle Emissions and Their Influencing Factors on Different Types of Roads, *J. Hazard. Mater.*, 2024, **465**, 133301, DOI: [10.1016/j.jhazmat.2023.133301](https://doi.org/10.1016/j.jhazmat.2023.133301).
- 59 E. Wi, E. Park, H. Shin, J. Hong, S. Jeong, J.-T. Kwon, H. Lee, J. Lee and Y. Kim, Overall Distribution of Tire-Wear Particles, Nano-carbon Black, and Heavy Metals in Size-Fractionated Road Dust Collected from Steel Industrial Complexes, *Sci. Total Environ.*, 2023, **884**, 163878, DOI: [10.1016/j.scitotenv.2023.163878](https://doi.org/10.1016/j.scitotenv.2023.163878).
- 60 J. Kim, E. Wi, H. Moon, H. Son, J. Hong, E. Park, J.-T. Kwon, D. Y. Seo, H. Lee and Y. Kim, Quantitative Analysis of the Concentration of Nano-carbon Black Originating from



- Tire-Wear Particles in the Road Dust, *Sci. Total Environ.*, 2022, **842**, 156830, DOI: [10.1016/j.scitotenv.2022.156830](https://doi.org/10.1016/j.scitotenv.2022.156830).
- 61 W. F. Rogge, P. M. Medeiros and B. R. T. Simoneit, Organic Compounds in Dust from Rural and Urban Paved and Unpaved Roads Taken During the San Joaquin Valley Fugitive Dust Characterization Study, *Environ. Eng. Sci.*, 2012, **29**(1), 1–13, DOI: [10.1089/ees.2010.0124](https://doi.org/10.1089/ees.2010.0124).
- 62 E. Chae, U. Jung and S.-S. Choi, Quantification of Tire Tread Wear Particles in Microparticles Produced on the Road Using Oleamide as a Novel Marker, *Environ. Pollut.*, 2021, **288**, 117811, DOI: [10.1016/j.envpol.2021.117811](https://doi.org/10.1016/j.envpol.2021.117811).
- 63 S. Jeong, H. Ryu, H. Shin, M. G. Lee, J. Hong, H. Kim, J. T. Kwon, J. Lee and Y. Kim, Quantification of Tire Wear Particles in Road Dust Based on Synthetic/Natural Rubber Ratio Using Pyrolysis-Gas Chromatography–Mass Spectrometry across Diverse Tire Types, *Sci. Total Environ.*, 2024, **942**, 173796, DOI: [10.1016/j.scitotenv.2024.173796](https://doi.org/10.1016/j.scitotenv.2024.173796).
- 64 E. S. Rødland, O. C. Lind, M. Reid, L. S. Heier, E. Skogsberg, B. Snilsberg, D. Gryteselv and S. Meland, Characterization of Tire and Road Wear Microplastic Particle Contamination in a Road Tunnel: From Surface to Release, *J. Hazard. Mater.*, 2022, **435**, 129032, DOI: [10.1016/j.jhazmat.2022.129032](https://doi.org/10.1016/j.jhazmat.2022.129032).
- 65 P. Klöckner, B. Seiwert, S. Weyrauch, B. I. Escher, T. Reemtsma and S. Wagner, Comprehensive Characterization of Tire and Road Wear Particles in Highway Tunnel Road Dust by Use of Size and Density Fractionation, *Chemosphere*, 2021, **279**, 130530, DOI: [10.1016/j.chemosphere.2021.130530](https://doi.org/10.1016/j.chemosphere.2021.130530).
- 66 E. L. Gaggini, M. Polukarova, M. Bondelind, E. Rødland, A. M. Strömvall, Y. Andersson-Sköld and E. Sokolova, Assessment of Fine and Coarse Tyre Wear Particles along a Highway Stormwater System and in Receiving Waters: Occurrence and Transport, *J. Environ. Manage.*, 2024, **367**, 121989, DOI: [10.1016/j.jenvman.2024.121989](https://doi.org/10.1016/j.jenvman.2024.121989).
- 67 T. R. Barber, S. Claes, F. Ribeiro, A. E. Dillon, S. L. More, S. Thornton, K. M. Unice, S. Weyrauch and T. Reemtsma, Abundance and Distribution of Tire and Road Wear Particles in the Seine River, France, *Sci. Total Environ.*, 2024, **913**, 169633, DOI: [10.1016/j.scitotenv.2023.169633](https://doi.org/10.1016/j.scitotenv.2023.169633).
- 68 C. Rauert, N. Charlton, E. D. Okoffo, R. S. Stanton, A. R. Agua, M. C. Pirrung and K. V. Thomas, Concentrations of Tire Additive Chemicals and Tire Road Wear Particles in an Australian Urban Tributary, *Environ. Sci. Technol.*, 2022, **56**(4), 2421–2431, DOI: [10.1021/acs.est.1c07451](https://doi.org/10.1021/acs.est.1c07451).
- 69 C. Rauert, S. Vardy, B. Daniell, N. Charlton and K. V. Thomas, Tyre Additive Chemicals, Tyre Road Wear Particles and High Production Polymers in Surface Water at 5 Urban Centres in Queensland, Australia, *Sci. Total Environ.*, 2022, **852**, 158468, DOI: [10.1016/j.scitotenv.2022.158468](https://doi.org/10.1016/j.scitotenv.2022.158468).
- 70 M. S. Molazadeh, F. Liu, J. Lykkemark, L. Iordachescu, A. H. Nielsen and J. Vollertsen, What Is Hiding below the Surface – MPs Including TWP in an Urban Lake, *Environ. Int.*, 2023, **182**, 108282, DOI: [10.1016/j.envint.2023.108282](https://doi.org/10.1016/j.envint.2023.108282).
- 71 I. Gofsmann, K. Mattsson, M. Hassellöv, C. Crazzolara, A. Held, T.-B. Robinson, O. Wurl and B. M. Scholz-Böttcher, Unraveling the Marine Microplastic Cycle: The First Simultaneous Data Set for Air, Sea Surface Microlayer, and Underlying Water, *Environ. Sci. Technol.*, 2023, **57**, 16541–16551, DOI: [10.1021/acs.est.3c05002](https://doi.org/10.1021/acs.est.3c05002).
- 72 M. Vercauteren, I. Semmouri, E. V. Acker, E. Pequeur, L. V. Esch, I. Uljee, J. Asselman and C. R. Janssen, Assessment of Road Run-off and Domestic Wastewater Contribution to Microplastic Pollution in a Densely Populated Area (Flanders, Belgium), *Environ. Pollut.*, 2023, **333**, 122090, DOI: [10.1016/j.envpol.2023.122090](https://doi.org/10.1016/j.envpol.2023.122090).
- 73 E. S. Rødland, O. C. Lind, M. J. Reid, L. S. Heier, E. D. Okoffo, C. Rauert, K. V. Thomas and S. Meland, Occurrence of Tire and Road Wear Particles in Urban and Peri-Urban Snowbanks, and Their Potential Environmental Implications, *Sci. Total Environ.*, 2022, **824**, 153785, DOI: [10.1016/j.scitotenv.2022.153785](https://doi.org/10.1016/j.scitotenv.2022.153785).
- 74 P. Eisentraut, E. Dümichen, A. S. Ruhl, A. S. Ruhl, M. Jekel, M. Albrecht, M. Albrecht, M. Gehde and U. Braun, Two Birds with One Stone—Fast and Simultaneous Analysis of Microplastics: Microparticles Derived from Thermoplastics and Tire Wear, *Environ. Sci. Technol. Lett.*, 2018, **5**, 608–613, DOI: [10.1021/acs.estlett.8b00446](https://doi.org/10.1021/acs.estlett.8b00446).
- 75 D. Mengistu, A. Heistad and C. Coutris, Tire Wear Particles Concentrations in Gully Pot Sediments, *Sci. Total Environ.*, 2021, **769**, 144785, DOI: [10.1016/j.scitotenv.2020.144785](https://doi.org/10.1016/j.scitotenv.2020.144785).
- 76 B. Seiwert, P. Klöckner, S. Wagner and T. Reemtsma, Source-Related Smart Suspect Screening in the Aqueous Environment: Search for Tire-Derived Persistent and Mobile Trace Organic Contaminants in Surface Waters, *Anal. Bioanal. Chem.*, 2020, **412**(20), 4909–4919, DOI: [10.1007/s00216-020-02653-1](https://doi.org/10.1007/s00216-020-02653-1).
- 77 J. K. Challis, H. Popick, S. Prajapati, P. Harder, J. P. Giesy, K. McPhedran and M. Brinkmann, Occurrences of Tire Rubber-Derived Contaminants in Cold-Climatic Urban Runoff, *Environ. Sci. Technol. Lett.*, 2021, **8**(11), 961–967, DOI: [10.1021/acs.estlett.1c00682](https://doi.org/10.1021/acs.estlett.1c00682).
- 78 Y.-H. Liu, Y.-X. Mei, X.-N. Liang, Z.-Y. Ge, Z. Huang, H.-Y. Zhang, J.-L. Zhao, A. Liu, C. Shi and G.-G. Ying, Small-Intensity Rainfall Triggers Greater Contamination of Rubber-Derived Chemicals in Road Stormwater Runoff from Various Functional Areas in Megalopolis Cities, *Environ. Sci. Technol.*, 2024, **58**(29), 13056–13064, DOI: [10.1021/acs.est.3c10737](https://doi.org/10.1021/acs.est.3c10737).
- 79 Z. Tian, H. Zhao, K. T. Peter, M. Gonzalez, J. Wetzel, C. Wu, X. Hu, J. Prat, E. Mudrock, R. Hettlinger, A. E. Cortina, R. G. Biswas, F. V. C. Kock, R. Soong, A. Jenne, B. Du, F. Hou, H. He, R. Lundeen, A. Gilbreath, R. Sutton, N. L. Scholz, J. W. Davis, M. C. Dodd, A. Simpson, J. K. McIntyre and E. P. Kolodziej, A Ubiquitous Tire Rubber-Derived Chemical Induces Acute Mortality in Coho Salmon, *Science*, 2021, **371**(6525), 185–189, DOI: [10.1126/science.abd6951](https://doi.org/10.1126/science.abd6951).
- 80 K. T. Peter, A. Gilbreath, M. Gonzalez, Z. Tian, A. Wong, D. Yee, E. L. Miller, P. M. Avellaneda, D. Chen,



- A. Patterson, N. Fitzgerald, C. P. Higgins, E. P. Kolodziej and R. Sutton, Storms Mobilize Organophosphate Esters, Bisphenols, PFASs, and Vehicle-Derived Contaminants to San Francisco Bay Watersheds, *Environ. Sci.: Processes Impacts*, 2024, **26**, 1760, DOI: [10.1039/d4em00117f](https://doi.org/10.1039/d4em00117f).
- 81 Y.-H. Liu, Y.-X. Mei, J.-Y. Wang, S.-S. Chen, J.-L. Chen, N. Li, W.-R. Liu, J.-L. Zhao, Q.-Q. Zhang and G.-G. Ying, Precipitation Contributes to Alleviating Pollution of Rubber-Derived Chemicals in Receiving Watersheds: Combining Confluent Stormwater Runoff from Different Functional Areas, *Water Res.*, 2024, **264**, 122240, DOI: [10.1016/j.watres.2024.122240](https://doi.org/10.1016/j.watres.2024.122240).
- 82 L. Maurer, E. Carmona, O. Machate, T. Schulze, M. Krauß and W. Brack, Contamination Pattern and Risk Assessment of Polar Compounds in Snow Melt: An Integrative Proxy of Road Runoffs, *Environ. Sci. Technol.*, 2023, **57**, 4143–4152, DOI: [10.1021/acs.est.2c05784](https://doi.org/10.1021/acs.est.2c05784).
- 83 S. Meland, G. M. Granheim, J. T. Rundberget and E. Rødland, Screening of Tire-Derived Chemicals and Tire Wear Particles in a Road Tunnel Wash Water Treatment Basin, *Environ. Sci. Technol. Lett.*, 2024, **11**(1), 35–40, DOI: [10.1021/acs.estlett.3c00811](https://doi.org/10.1021/acs.estlett.3c00811).
- 84 F. Kryuchkov, A. Foldvik, R. Sandodden and S. Uhlig, Presence of 6PPD-Quinone in Runoff Water Samples from Norway Using a New LC-MS/MS Method, *Front. Environ. Chem.*, 2023, **4**, 1194664, DOI: [10.3389/fenvc.2023.1194664](https://doi.org/10.3389/fenvc.2023.1194664).
- 85 L.-J. Zhou, S. Liu, M. Wang, N.-N. Wu, R. Xu, L.-N. Wei, X.-R. Xu, J.-L. Zhao, P. Xing, H. Li, J. Zeng and Q. L. Wu, Nationwide Occurrence and Prioritization of Tire Additives and Their Transformation Products in Lake Sediments of China, *Environ. Int.*, 2024, **193**, 109139, DOI: [10.1016/j.envint.2024.109139](https://doi.org/10.1016/j.envint.2024.109139).
- 86 L. X. Hu, Y. X. Cheng, D. Wu, L. Fan, J. H. Zhao, Q. Xiong, Q. L. Chen, Y. S. Liu and G. G. Ying, Continuous Input of Organic Ultraviolet Filters and Benzothiazoles Threatens the Surface Water and Sediment of Two Major Rivers in the Pearl River Basin, *Sci. Total Environ.*, 2021, **798**, 149299, DOI: [10.1016/j.scitotenv.2021.149299](https://doi.org/10.1016/j.scitotenv.2021.149299).
- 87 Z.-M. Li, V. K. Pal, P. Kannan, W. Li and K. Kannan, 1,3-Diphenylguanidine, Benzothiazole, Benzotriazole, and Their Derivatives in Soils Collected from Northeastern United States, *Sci. Total Environ.*, 2023, **887**, 164110, DOI: [10.1016/j.scitotenv.2023.164110](https://doi.org/10.1016/j.scitotenv.2023.164110).
- 88 S. C. Ihenetu, Q. Xu, Z. H. Khan, S. S. U. H. Kazmi, J. Ding, Q. Sun and G. Li, Environmental Fate of Tire-Rubber Related Pollutants 6PPD and 6PPD-Q: A Review, *Environ. Res.*, 2024, 119492, DOI: [10.1016/j.envres.2024.119492](https://doi.org/10.1016/j.envres.2024.119492).
- 89 C. Johannessen, P. Helm and C. D. Metcalfe, Detection of Selected Tire Wear Compounds in Urban Receiving Waters, *Environ. Pollut.*, 2021, **287**, 117659, DOI: [10.1016/j.envpol.2021.117659](https://doi.org/10.1016/j.envpol.2021.117659).
- 90 J. Zhu, R. Guo, F. Ren, S. Jiang and H. Jin, Occurrence and Partitioning of P-Phenylenediamine Antioxidants and Their Quinone Derivatives in Water and Sediment, *Sci. Total Environ.*, 2024, **914**, 170046, DOI: [10.1016/j.scitotenv.2024.170046](https://doi.org/10.1016/j.scitotenv.2024.170046).
- 91 H. Zhang, Z. Huang, Y. Liu, L. Hu, L.-Y. He, Y.-S. Liu, J. Zhao and G. Ying, Occurrence and Risks of 23 Tire Additives and Their Transformation Products in an Urban Water System, *Environ. Int.*, 2023, **171**, 107715, DOI: [10.1016/j.envint.2022.107715](https://doi.org/10.1016/j.envint.2022.107715).
- 92 H. Zhao, X. Hu, Z. Tian, M. Gonzalez, C. A. Rideout, K. T. Peter, M. C. Dodd and E. P. Kolodziej, Transformation Products of Tire Rubber Antioxidant 6PPD in Heterogeneous Gas-Phase Ozonation: Identification and Environmental Occurrence, *Environ. Sci. Technol.*, 2023, **57**(14), 5621–5632, DOI: [10.1021/acs.est.2c08690](https://doi.org/10.1021/acs.est.2c08690).
- 93 R. Jin, B. Li, Y. Wu, Y. Li, X. Du, C. Xia, H. Zhao and M. Liu, Unpuzzling Spatio-Vertical and Multi-Media Patterns of Aniline Accelerators/Antioxidants in an Urban Estuary, *Water Res.*, 2024, **266**, 122427, DOI: [10.1016/j.watres.2024.122427](https://doi.org/10.1016/j.watres.2024.122427).
- 94 H.-Y. Zhang, Y.-H. Liu, L.-N. Wei, R.-Q. Zhu, J.-L. Zhao, S. Liu, X.-R. Xu and G.-G. Ying, Unveiling Spatiotemporal Distribution, Partitioning, and Transport Mechanisms of Tire Additives and Their Transformation Products in a Highly Urbanized Estuarine Region, *Sci. Total Environ.*, 2024, **954**, 176804, DOI: [10.1016/j.scitotenv.2024.176804](https://doi.org/10.1016/j.scitotenv.2024.176804).
- 95 L. N. Wei, N. N. Wu, R. Xu, S. Liu, H. X. Li, L. Lin, R. Hou, X. R. Xu, J. L. Zhao and G. G. Ying, First Evidence of the Bioaccumulation and Trophic Transfer of Tire Additives and Their Transformation Products in an Estuarine Food Web, *Environ. Sci. Technol.*, 2024, **58**(14), 6370–6380, DOI: [10.1021/acs.est.3c10248](https://doi.org/10.1021/acs.est.3c10248).
- 96 L. Hanssen, S. Schmidt and V. Nikiforov, Screening of Compounds in Tire Wear Road Run Off, *TemaNord*, 2024, 525, DOI: [10.6027/temanord2024-525](https://doi.org/10.6027/temanord2024-525).
- 97 G. Cao, W. Wang, J. Zhang, P. Wu, X. Zhao, Z. Yang, D. Hu and Z. Cai, New Evidence of Rubber-Derived Quinones in Water, Air, and Soil, *Environ. Sci. Technol.*, 2022, **56**(7), 4142–4150, DOI: [10.1021/acs.est.1c07376](https://doi.org/10.1021/acs.est.1c07376).
- 98 L. Zeng, Y. Li, Y. Sun, L.-Y. Liu, M. Shen and B. Du, Widespread Occurrence and Transport of P-Phenylenediamines and Their Quinones in Sediments across Urban Rivers, Estuaries, Coasts, and Deep-Sea Regions, *Environ. Sci. Technol.*, 2023, **57**(6), 2393–2403, DOI: [10.1021/acs.est.2c07652](https://doi.org/10.1021/acs.est.2c07652).
- 99 Y. Wu, M. Venier and R. A. Hites, Broad Exposure of the North American Environment to Phenolic and Amino Antioxidants and to Ultraviolet Filters, *Environ. Sci. Technol.*, 2020, **54**(15), 9345–9355, DOI: [10.1021/acs.est.0c04114](https://doi.org/10.1021/acs.est.0c04114).
- 100 W. Wang, G. Cao, J. Zhang, W. Chang, Y. Sang and Z. Cai, Fragmentation Pattern-Based Screening Strategy Combining Diagnostic Ion and Neutral Loss Uncovered Novel Para-Phenylenediamine Quinone Contaminants in the Environment, *Environ. Sci. Technol.*, 2024, **58**(13), 5921–5931, DOI: [10.1021/acs.est.4c00027](https://doi.org/10.1021/acs.est.4c00027).
- 101 Z. M. Li and K. Kannan, Mass Loading, Removal, and Emission of 1,3-Diphenylguanidine, Benzotriazole, Benzothiazole, N-(1,3-Dimethylbutyl)-N'-Phenyl-p-Phenylenediamine, and Their Derivatives in a Wastewater



- Treatment Plant in New York State, USA, *ACS ES&T Water*, 2024, **4**(6), 2721–2730, DOI: [10.1021/acsestwater.4c00221](https://doi.org/10.1021/acsestwater.4c00221).
- 102 G. Cao, W. Wang, J. Zhang, P. Wu, H. Qiao, H. Li, G. Huang, Z. Yang and Z. Cai, Occurrence and Fate of Substituted *p*-Phenylenediamine-Derived Quinones in Hong Kong Wastewater Treatment Plants, *Environ. Sci. Technol.*, 2023, **57**(41), 15635–15643, DOI: [10.1021/acs.est.3c03758](https://doi.org/10.1021/acs.est.3c03758).
- 103 R. F. Lane, K. L. Smalling, P. M. Bradley, J. B. Greer, S. E. Gordon, J. D. Hansen, D. W. Kolpin, A. R. Spanjer and J. R. Masoner, Tire-Derived Contaminants 6PPD and 6PPD-Q: Analysis, Sample Handling, and Reconnaissance of United States Stream Exposures, *Chemosphere*, 2024, **363**, 142830, DOI: [10.1016/j.chemosphere.2024.142830](https://doi.org/10.1016/j.chemosphere.2024.142830).
- 104 R. Zhang, S. Zhao, X. Liu, L. Tian, Y. Mo, X. Yi, S. Liu, J. Liu, J. Li and G. Zhang, Aquatic Environmental Fates and Risks of Benzotriazoles, Benzothiazoles, and *p*-Phenylenediamines in a Catchment Providing Water to a Megacity of China, *Environ. Res.*, 2023, **216**, 114721, DOI: [10.1016/j.envres.2022.114721](https://doi.org/10.1016/j.envres.2022.114721).
- 105 J. Monaghan, A. Jaeger, A. R. Agua, R. S. Stanton, M. Pirrung, C. G. Gill and E. T. Krogh, A Direct Mass Spectrometry Method for the Rapid Analysis of Ubiquitous Tire-Derived Toxin *N*-(1,3-Dimethylbutyl)-*N'*-Phenyl-*p*-Phenylenediamine Quinone (6-PPDQ), *Environ. Sci. Technol. Lett.*, 2021, **8**, 1051–1056, DOI: [10.1021/acs.estlett.1c00794](https://doi.org/10.1021/acs.estlett.1c00794).
- 106 J. Monaghan, A. Jaeger, J. K. Jai, H. Tomlin, J. Atkinson, T. M. Brown, C. G. Gill and E. T. Krogh, Automated, High-Throughput Analysis of Tire-Derived *p*-Phenylenediamine Quinones (PPDQs) in Water by Online Membrane Sampling Coupled to MS/MS, *ACS ES&T Water*, 2023, **3**(10), 3293–3304, DOI: [10.1021/acsestwater.3c00275](https://doi.org/10.1021/acsestwater.3c00275).
- 107 S. Ren, Y. Xia, X. Wang, Y. Zou, Z. Li, M. Man, Q. Yang, M. Lv, J. Ding and L. Chen, Development and Application of Diffusive Gradients in Thin-Films for in-Situ Monitoring of 6PPD-Quinone in Urban Waters, *Water Res.*, 2024, **266**, 122408, DOI: [10.1016/j.watres.2024.122408](https://doi.org/10.1016/j.watres.2024.122408).
- 108 J. J. Halama, R. B. McKane, B. L. Barnhart, P. P. Pettus, A. F. Brookes, A. K. Adams, C. K. Gockel, K. S. Djang, V. Phan, S. M. Chokshi, J. J. Graham, Z. Tian, K. T. Peter and E. P. Kolodziej, Watershed Analysis of Urban Stormwater Contaminant 6PPD-Quinone Hotspots and Stream Concentrations Using a Process-Based Ecohydrological Model, *Front. Environ. Sci.*, 2024, **12**, 1364673, DOI: [10.3389/fenvs.2024.1364673](https://doi.org/10.3389/fenvs.2024.1364673).
- 109 P. Ross, S. Scott, M. Noel and N. Klasios, *Green/Cheakamus Watershed: Water Quality Report for the 2023 Dry Season*, 2024, DOI: [10.70766/955.423](https://doi.org/10.70766/955.423).
- 110 I. C. Tanui, F. Kandie, M. Krauss, A. Piotrowska, A. Kiprop, N. Shahid, M. Liess and W. Brack, Seasonal Hot Spots of Pollution and Risks in Western Kenya: A Spatial-Temporal Analysis of Almost 800 Organic Micropollutants, *Sci. Total Environ.*, 2024, **949**, 175036, DOI: [10.1016/j.scitotenv.2024.175036](https://doi.org/10.1016/j.scitotenv.2024.175036).
- 111 C. Johannessen, P. A. Helm, J. Mark Parnis, S. Kleywegt and C. D. Metcalfe, Targeted Screening of Passive Samplers as an “Early Warning” of Novel Contaminants in the Great Lakes Basin, *J. Great Lakes Res.*, 2024, **50**(2), 102298, DOI: [10.1016/j.jglr.2024.102298](https://doi.org/10.1016/j.jglr.2024.102298).
- 112 P. A. Helm, M. Raby, S. Kleywegt, R. J. Sorichetti, G. Arabian, D. Smith, E. T. Howell and J. Thibeau, Assessment of Tire-Additive Transformation Product 6PPD-Quinone in Urban-Impacted Watersheds, *ACS ES&T Water*, 2024, **4**(4), 1422–1432, DOI: [10.1021/acsestwater.3c00589](https://doi.org/10.1021/acsestwater.3c00589).
- 113 C. Johannessen, P. Helm, B. Lashuk, V. Yargeau and C. D. Metcalfe, The Tire Wear Compounds 6PPD-Quinone and 1,3-Diphenylguanidine in an Urban Watershed, *Arch. Environ. Contam. Toxicol.*, 2022, **82**(2), 171–179, DOI: [10.1007/s00244-021-00878-4](https://doi.org/10.1007/s00244-021-00878-4).
- 114 C. Rauert, S. Vardy, B. Daniell, N. Charlton and K. V. Thomas, Tyre Additive Chemicals, Tyre Road Wear Particles and High Production Polymers in Surface Water at 5 Urban Centres in Queensland, Australia, *Sci. Total Environ.*, 2022, **852**, 158468, DOI: [10.1016/j.scitotenv.2022.158468](https://doi.org/10.1016/j.scitotenv.2022.158468).
- 115 Z. Tian, K. T. Peter, A. D. Gipe, H. Zhao, F. Hou, D. A. Wark, T. Khangaonkar, E. P. Kolodziej and C. A. James, Suspect and Nontarget Screening for Contaminants of Emerging Concern in an Urban Estuary, *Environ. Sci. Technol.*, 2020, **54**(2), 889–901, DOI: [10.1021/acs.est.9b06126](https://doi.org/10.1021/acs.est.9b06126).
- 116 T.-T. Xu, J.-L. Guo, J.-F. Yi, S.-N. Huang, L.-Y. Liu and Y. Guo, Occurrence and Risk Assessment of *p*-Phenylenediamines and Their Quinones in Aquatic Environment: From City Wastewater to Deep Sea, *Sci. Total Environ.*, 2024, **954**, 176123, DOI: [10.1016/j.scitotenv.2024.176123](https://doi.org/10.1016/j.scitotenv.2024.176123).
- 117 W. Wu, Q. Xu, J. Li, Z. Wang and G. Li, The Spatio-Temporal Accumulation of 6 PPD-Q in Greenbelt Soils and Its Effects on Soil Microbial Communities, *Environ. Pollut.*, 2024, **358**, 124477, DOI: [10.1016/j.envpol.2024.124477](https://doi.org/10.1016/j.envpol.2024.124477).
- 118 B. Seiwert, M. Nihemaiti, M. Troussier, S. Weyrauch and T. Reemtsma, Abiotic Oxidative Transformation of 6-PPD and 6-PPD Quinone from Tires and Occurrence of Their Products in Snow from Urban Roads and in Municipal Wastewater, *Water Res.*, 2022, **212**, 118122, DOI: [10.1016/j.watres.2022.118122](https://doi.org/10.1016/j.watres.2022.118122).
- 119 K. T. Peter, F. Hou, Z. Tian, C. Wu, M. Goehring, F. Liu and E. P. Kolodziej, More Than a First Flush: Urban Creek Storm Hydrographs Demonstrate Broad Contaminant Pollutographs, *Environ. Sci. Technol.*, 2020, **54**(10), 6152–6165, DOI: [10.1021/acs.est.0c00872](https://doi.org/10.1021/acs.est.0c00872).
- 120 C. Johannessen, P. Helm and C. D. Metcalfe, Runoff of the Tire-Wear Compound, Hexamethoxymethyl-Melamine into Urban Watersheds, *Arch. Environ. Contam. Toxicol.*, 2022, **82**(2), 162–170, DOI: [10.1007/s00244-021-00815-5](https://doi.org/10.1007/s00244-021-00815-5).
- 121 F. Hou, Z. Tian, K. T. Peter, C. Wu, A. D. Gipe, H. Zhao, E. A. Alegria, F. Liu and E. P. Kolodziej, Quantification of Organic Contaminants in Urban Stormwater by Isotope Dilution and Liquid Chromatography-Tandem Mass Spectrometry, *Anal. Bioanal. Chem.*, 2019, **411**(29), 7791–7806, DOI: [10.1007/s00216-019-02177-3](https://doi.org/10.1007/s00216-019-02177-3).



- 122 C. Rauert, S. L. Kaserzon, C. Veal, R. Y. Yeh, J. F. Mueller and K. V. Thomas, The First Environmental Assessment of Hexa(Methoxymethyl)Melamine and Co-Occurring Cyclic Amines in Australian Waterways, *Sci. Total Environ.*, 2020, **743**, 140834, DOI: [10.1016/j.scitotenv.2020.140834](https://doi.org/10.1016/j.scitotenv.2020.140834).
- 123 C. Johannessen and C. D. Metcalfe, The Occurrence of Tire Wear Compounds and Their Transformation Products in Municipal Wastewater and Drinking Water Treatment Plants, *Environ. Monit. Assess.*, 2022, **194**(10), 731, DOI: [10.1007/s10661-022-10450-9](https://doi.org/10.1007/s10661-022-10450-9).
- 124 E. K. Hataley, H. K. McIlwraith, D. Roy and C. M. Rochman, Towards a Management Strategy for Microplastic Pollution in the Laurentian Great Lakes—Ecological Risk Assessment and Management (Part 2), *Can. J. Fish. Aquat. Sci.*, 2023, **80**(10), 1669–1678, DOI: [10.1139/cjfas-2023-0023](https://doi.org/10.1139/cjfas-2023-0023).
- 125 F. Sommer, V. Dietze, A. Baum, J. Sauer, S. Gilge, C. Maschowski and R. Gieré, Tire Abrasion as a Major Source of Microplastics in the Environment, *Aerosol Air Qual. Res.*, 2018, **18**(8), 2014–2028, DOI: [10.4209/aaqr.2018.03.0099](https://doi.org/10.4209/aaqr.2018.03.0099).
- 126 V. Carrasco-Navarro, A. B. Muñoz-González, J. Sorvari and J. L. Martínez-Guitarte, Altered Gene Expression in Chironomus Riparius (Insecta) in Response to Tire Rubber and Polystyrene Microplastics, *Environ. Pollut.*, 2021, **285**, 117462, DOI: [10.1016/j.envpol.2021.117462](https://doi.org/10.1016/j.envpol.2021.117462).
- 127 J. Liu, Q. Feng, H. Yang, X. Fan, Y. Jiang and T. Wu, Acute Toxicity of Tire Wear Particles and Leachate to Daphnia Magna, *Comp. Biochem. Physiol., Part C: Toxicol. Pharmacol.*, 2023, **272**, 109713, DOI: [10.1016/j.cbpc.2023.109713](https://doi.org/10.1016/j.cbpc.2023.109713).
- 128 R. S. Cheong, E. R. Dumont, P. E. Thomson, D. C. Castañeda-Cortés, L. M. Hernandez, X. Gao, J. Zheng, A. Baesu, J. Macairan, A. J. Smith, H.-L. Bui, H. C. E. Larsson, S. Ghoshal, S. Bayen, V. S. Langlois, S. A. Robinson and N. Tufenkji, Nanoparticle-Specific and Chemical-Specific Effects of Tire Wear Particle Leachate on Amphibian Early Life Stages, *J. Hazard. Mater. Adv.*, 2023, **12**, 100357, DOI: [10.1016/j.hazadv.2023.100357](https://doi.org/10.1016/j.hazadv.2023.100357).
- 129 S. Magni, E. Tediosi, D. Maggioni, R. Sbarberi, F. Noé, F. Rossetti, D. Fornai, V. Persici and M. C. Neri, Ecological Impact of End-of-Life-Tire (ELT)-Derived Rubbers: Acute and Chronic Effects at Organism and Population Levels, *Toxics*, 2022, **10**(5), 201, DOI: [10.3390/toxics10050201](https://doi.org/10.3390/toxics10050201).
- 130 W. Dufefoi, B. J. D. Ferrari, F. Breider, T. Masset, G. Leger, E. Vermeirssen, A. J. Bergmann and K. Schirmer, Evaluation of Tire Tread Particle Toxicity to Fish Using Rainbow Trout Cell Lines, *Sci. Total Environ.*, 2024, **912**, 168933, DOI: [10.1016/j.scitotenv.2023.168933](https://doi.org/10.1016/j.scitotenv.2023.168933).
- 131 L. Kim, T.-Y. Lee, H. Kim and Y.-J. An, Toxicity Assessment of Tire Particles Released from Personal Mobilities (Bicycles, Cars, and Electric Scooters) on Soil Organisms, *J. Hazard. Mater.*, 2022, **437**, 129362, DOI: [10.1016/j.jhazmat.2022.129362](https://doi.org/10.1016/j.jhazmat.2022.129362).
- 132 J. Ding, D. Zhu, H.-T. Wang, S. B. Lassen, Q.-L. Chen, G. Li, M. Lv and Y.-G. Zhu, Dysbiosis in the Gut Microbiota of Soil Fauna Explains the Toxicity of Tire Tread Particles, *Environ. Sci. Technol.*, 2020, **54**(12), 7450–7460, DOI: [10.1021/acs.est.0c00917](https://doi.org/10.1021/acs.est.0c00917).
- 133 K. Yang, S. Jing, Y. Liu, H. Zhou, Y. Liu, M. Yan, X. Yi and R. Liu, Acute Toxicity of Tire Wear Particles, Leachates and Toxicity Identification Evaluation of Leachates to the Marine Copepod, Tigriopus Japonicus, *Chemosphere*, 2022, **297**, 134099, DOI: [10.1016/j.chemosphere.2022.134099](https://doi.org/10.1016/j.chemosphere.2022.134099).
- 134 S. B. LaPlaca, C. D. Rice and P. van den Hurk, Chronic Toxicity of Tire Crumb Rubber Particles to Mummichog (Fundulus Heteroclitus) in Episodic Exposures, *Sci. Total Environ.*, 2022, **846**, 157447, DOI: [10.1016/j.scitotenv.2022.157447](https://doi.org/10.1016/j.scitotenv.2022.157447).
- 135 S. Garrard, J. Spicer and R. Thompson, Tyre Particle Exposure Affects the Health of Two Key Estuarine Invertebrates, *Environ. Pollut.*, 2022, **314**, 120244, DOI: [10.1016/j.envpol.2022.120244](https://doi.org/10.1016/j.envpol.2022.120244).
- 136 A. J. Bergmann, T. Masset, F. Breider, W. Dufefoi, K. Schirmer, B. J. D. Ferrari and E. L. M. Vermeirssen, Estrogenic, Genotoxic, and Antibacterial Effects of Chemicals from Cryogenically Milled Tire Tread, *Environ. Toxicol. Chem.*, 2024, **43**(9), 1962–1972, DOI: [10.1002/etc.5934](https://doi.org/10.1002/etc.5934).
- 137 M. Capolupo, L. Sørensen, K. D. R. Jayasena, A. M. Booth and E. Fabbri, Chemical Composition and Ecotoxicity of Plastic and Car Tire Rubber Leachates to Aquatic Organisms, *Water Res.*, 2020, **169**, 115270, DOI: [10.1016/j.watres.2019.115270](https://doi.org/10.1016/j.watres.2019.115270).
- 138 M. Capolupo, A. Rafiq, I. Coralli, T. Alessandro, P. Valbonesi, D. Fabbri and E. Fabbri, Bioplastic Leachates Characterization and Impacts on Early Larval Stages and Adult Mussel Cellular, Biochemical and Physiological Responses, *Environ. Pollut.*, 2023, **319**, 120951, DOI: [10.1016/j.envpol.2022.120951](https://doi.org/10.1016/j.envpol.2022.120951).
- 139 ECHA, *Guidance on the Application of the CLP Criteria, Parts 4 and 5*, Helsinki, 2024, DOI: [10.2823/4895485](https://doi.org/10.2823/4895485).
- 140 P. E. Redondo-Hasselerharm, V. N. d. Ruijter, S. M. Mintenig, A. Verschoor and A. A. Koelmans, Ingestion and Chronic Effects of Car Tire Tread Particles on Freshwater Benthic Macroinvertebrates, *Environ. Sci. Technol.*, 2018, **52**, 13986–13994, DOI: [10.1021/acs.est.8b05035](https://doi.org/10.1021/acs.est.8b05035).
- 141 F. R. Khan, L. L. Halle and A. Palmqvist, Acute and Long-Term Toxicity of Micronized Car Tire Wear Particles to Hyalella Azteca, *Aquat. Toxicol.*, 2019, **213**, 105216, DOI: [10.1016/j.aquatox.2019.05.018](https://doi.org/10.1016/j.aquatox.2019.05.018).
- 142 J. E. Weinstein, B. M. Ertel and A. D. Gray, Accumulation and Depuration of Microplastic Fibers, Fragments, and Tire Particles in the Eastern Oyster, Crassostrea Virginica: A Toxicokinetic Approach, *Environ. Pollut.*, 2022, **308**, 119681, DOI: [10.1016/j.envpol.2022.119681](https://doi.org/10.1016/j.envpol.2022.119681).
- 143 F. Hägg, D. Herzke, V. A. Nikiforov, A. M. Booth, K. H. Sperre, L. Sørensen, M. E. Creese and C. Halsband, Ingestion of Car Tire Crumb Rubber and Uptake of Associated Chemicals by Lumpfish (Cyclopterus Lumpus),



- Front. Environ. Sci.*, 2023, **11**, 1219248, DOI: [10.3389/fenvs.2023.1219248](https://doi.org/10.3389/fenvs.2023.1219248).
- 144 T. Masset, F. Breider, M. Renaud, J. Müller, A. Bergmann, E. Vermeirssen, W. Dufefoi, K. Schirmer and B. J. D. Ferrari, Effects of Tire Particles on Earthworm (*Eisenia Andrei*) Fitness and Bioaccumulation of Tire-Related Chemicals, *Environ. Pollut.*, 2025, 125780, DOI: [10.1016/j.envpol.2025.125780](https://doi.org/10.1016/j.envpol.2025.125780).
- 145 T. Masset, B. J. D. Ferrari, W. Dufefoi, K. Schirmer, A. Bergmann, E. Vermeirssen, D. Grandjean, L. C. Harris and F. Breider, Bioaccessibility of Organic Compounds Associated with Tire Particles Using a Fish *In Vitro* Digestive Model: Solubilization Kinetics and Effects of Food Coingestion, *Environ. Sci. Technol.*, 2022, **56**(22), 15607–15616, DOI: [10.1021/acs.est.2c04291](https://doi.org/10.1021/acs.est.2c04291).
- 146 K. Hiki, K. Asahina, K. Kato, T. Yamagishi, R. Omagari, Y. Iwasaki, H. Watanabe and H. Yamamoto, Acute Toxicity of a Tire Rubber-Derived Chemical, 6PPD Quinone, to Freshwater Fish and Crustacean Species, *Environ. Sci. Technol. Lett.*, 2021, **8**(9), 779–784, DOI: [10.1021/acs.estlett.1c00453](https://doi.org/10.1021/acs.estlett.1c00453).
- 147 M. Brinkmann, D. Montgomery, S. Selinger, J. G. P. Miller, E. Stock, A. J. Alcaraz, J. K. Challis, L. Weber, D. Janz, M. Hecker and S. Wiseman, Acute Toxicity of the Tire Rubber-Derived Chemical 6PPD-Quinone to Four Fishes of Commercial, Cultural, and Ecological Importance, *Environ. Sci. Technol. Lett.*, 2022, **9**(4), 333–338, DOI: [10.1021/acs.estlett.2c00050](https://doi.org/10.1021/acs.estlett.2c00050).
- 148 K. M. Unice, M. L. Kreider and J. M. Panko, Use of a Deuterated Internal Standard with Pyrolysis-GC/MS Dimeric Marker Analysis to Quantify Tire Tread Particles in the Environment, *Int. J. Environ. Res. Public Health*, 2012, **9**(11), 4033–4055, DOI: [10.3390/ijerph9114033](https://doi.org/10.3390/ijerph9114033).
- 149 S. Wagner, C. W. Funk, K. Müller and D. J. Raithel, The Chemical Composition and Sources of Road Dust, and of Tire and Road Wear Particles—A Review, *Sci. Total Environ.*, 2024, **926**, 171694, DOI: [10.1016/j.scitotenv.2024.171694](https://doi.org/10.1016/j.scitotenv.2024.171694).
- 150 E. S. Rødland, M. Gustafsson, D. Jaramillo-Vogel, I. Järllskog, K. Müller, C. Rauert, J. Rausch and S. Wagner, Analytical Challenges and Possibilities for the Quantification of Tire-Road Wear Particles, *TrAC, Trends Anal. Chem.*, 2023, **165**, 117121, DOI: [10.1016/j.trac.2023.117121](https://doi.org/10.1016/j.trac.2023.117121).
- 151 L. L. Trudsø, M. B. Nielsen, S. F. Hansen, K. Syberg, K. Kampmann, F. R. Khan and A. Palmqvist, The Need for Environmental Regulation of Tires: Challenges and Recommendations, *Environ. Pollut.*, 2022, **311**, 119974, DOI: [10.1016/j.envpol.2022.119974](https://doi.org/10.1016/j.envpol.2022.119974).
- 152 E. Abbate, A. I. Garmendia, G. Bracalente, L. Mancini, D. Tosches, K. Rasmussen, M. J. Bennett, H. Rauscher and S. Sala, *Safe and Sustainable by Design Chemicals and Materials - Methodological Guidance*, ISPRA, 2024, DOI: [10.2760/28450](https://doi.org/10.2760/28450).
- 153 M. Kovochich, M. Liong, J. A. Parker, S. C. Oh, J. P. Lee, L. Xi, M. L. Kreider and K. M. Unice, Chemical Mapping of Tire and Road Wear Particles for Single Particle Analysis, *Sci. Total Environ.*, 2021, **757**, 144085, DOI: [10.1016/j.scitotenv.2020.144085](https://doi.org/10.1016/j.scitotenv.2020.144085).
- 154 M. L. Kreider, J. M. Panko, B. L. McAtee, L. I. Sweet and B. L. Finley, Physical and Chemical Characterization of Tire-Related Particles: Comparison of Particles Generated Using Different Methodologies, *Sci. Total Environ.*, 2010, **408**(3), 652–659, DOI: [10.1016/j.scitotenv.2009.10.016](https://doi.org/10.1016/j.scitotenv.2009.10.016).
- 155 ISO, *ISO 472:2013(En) Plastics — Vocabulary*, 2013.

