

Cite this: *Chem. Sci.*, 2025, 16, 5058

All publication charges for this article have been paid for by the Royal Society of Chemistry

Received 10th January 2025

Accepted 8th February 2025

DOI: 10.1039/d5sc00237k

rsc.li/chemical-science

## Unusually air-stable copper(i) complexes showing high selectivity for carbon monoxide†

Borna Saeednia, ‡ Aria M. Sragow, ‡ Yannan Lin, ‡ Colton J. Sheehan, Amy S. Metlay, Michael R. Gau, Samantha A. Dye, ‡ Sarah P. O'Konski, ‡ Thomas E. Mallouk and Ivan J. Dmochowski \* ‡

We report two Cu(I)-tren host molecules with unusual air-stability, as revealed by strong preference for axial CO binding over bent O<sub>2</sub>. Spectroscopy, electrochemical, and X-ray crystal structure analyses indicate that the phenyl rotators of the capsule select for small axial ligands.

Structural features that regulate binding of diatomic molecules to metal centres are of central importance in many biochemical and chemical-industrial processes.<sup>1,2</sup> As dioxygen and carbon monoxide can compete for binding at protein metal centres, probing the structural origins of selectivity between O<sub>2</sub> and CO at metalloproteins is of interest.<sup>3–6</sup> Selective CO binding holds importance for sensing applications<sup>7,8</sup> and also for delivery, due to its pathophysiological and therapeutic roles.<sup>9</sup> Here we highlight copper proteins that bind dioxygen and perform vital functions such as production of norepinephrine (dopamine-β-hydroxylase, DBH), mitochondrial function (cytochrome c oxidase, COX), and O<sub>2</sub> transport in molluscs and arthropods (hemocyanin).<sup>10</sup> These and many other copper proteins carefully regulate the Cu(I)/Cu(II) redox couple, for purposes such as electron transfer, oxidation of substrates, and O<sub>2</sub> transport.<sup>11,12</sup> Synthetic models for the study of cuproproteins are well-precedented<sup>13,14</sup> and provide an opportunity to examine stereo-electronic features that control function.<sup>15</sup> However, Cu(I) systems demonstrating CO/O<sub>2</sub> selectivity are rare,<sup>16–18</sup> and control of selectivity by steric factors alone has, to the best of our knowledge, never been demonstrated.

Metalloproteins can achieve ligand selectivity by varying the size and accessibility of the binding pocket.<sup>15</sup> Many metalloproteins, including the hemocyanins (where a single dioxygen ligand coordinates two copper centres), possess deeply-buried active sites in which few guests bind.<sup>19</sup> Inspired by this selectivity, we were interested in creating a model system with a defined metallo-cavity for ligands to enter.<sup>20</sup> However, many classes of macromolecules possessing hydrophobic cavities such as cucurbiturils,<sup>21–23</sup> calixarenes,<sup>24–27</sup> resorcinarenes,<sup>28</sup> and

cyclodextrins<sup>29,30</sup> generally lack defined metal-binding sites. Hemicyptophanes are a class of small-molecule cages<sup>31,32</sup> that are excellent candidates for metalloprotein-inspired binding pockets, owing to their well-defined hydrophobic cavity created by a cyclotrimeratrylene (CTV) north pole<sup>33–36</sup> and the ease of attaching a south pole with a metal-binding site.<sup>31,37,38</sup> Furthermore, hemicyptophanes and other CTV-based supra-molecular structures have been used in copper-binding systems by the groups of Hardie and Dutasta.<sup>39–41</sup> In designing the south pole, we were inspired by myoglobin and haemoglobin. While CO/O<sub>2</sub> discrimination is an important feature of these iron-haem proteins, which has been mimicked in various small-molecule and protein models,<sup>42–45</sup> such discrimination has not yet been observed in an analogous monocopper protein.

First synthesised by Ogawa,<sup>46</sup> **L1** (Fig. 1) is a particularly versatile hemicyptophane. In addition to selective guest<sup>47</sup> and anion<sup>48</sup> binding with protonated **L1**, Zn(II)-coordinated **L1** catalyses the hydrolysis of activated alkyl carbonates<sup>46,49</sup> and Ru(III)-**L1** can catalyse the oxidation of primary alcohols.<sup>50</sup> Here, we sought to prepare Cu(I)-bound **L1** and investigate its ligand-binding properties (Fig. 1). **L1** provides an unusual example of a tripodal amine-substituted copper ligand. Many other tripodal amine- and amide-based Cu(I) systems are known, which use ligands such as tren, TMPA, and their derivatives<sup>51–55</sup> to explore the effects of alkyl groups, chelate ring size, and steric bulk on substrate binding.<sup>56</sup> However, with rare exceptions,<sup>57–60</sup> most



Fig. 1 Structures of **L1**, **L2**, [Cu**L1**]PF<sub>6</sub>, and [Cu**L2**]PF<sub>6</sub>.

Department of Chemistry, University of Pennsylvania, 231 S. 34th St., Philadelphia, PA, 19104, USA. E-mail: ivandmo@sas.upenn.edu

† Electronic supplementary information (ESI) available. CCDC 2359098–2359100. For ESI and crystallographic data in CIF or other electronic format see DOI: <https://doi.org/10.1039/d5sc00237k>

‡ These authors contributed equally.

notably from the Reinaud group,<sup>61–65</sup> these complexes lack a well-defined cavity at the metal site.

In this work, we synthesised the copper(I) complex  $[\text{CuL1}]\text{PF}_6$  from the previously-reported **L1**.<sup>46,47</sup> We observed air-stability and sought to elucidate the origin of this effect, as few examples of air-stable, 4-coordinate Cu(I) complexes with an open coordination site exist due to their typically high oxidation potential.<sup>66–70</sup> In similar complexes, portal size and/or steric blockage have been observed to control the size of the guests that can access the cavity.<sup>47</sup> Seeking to increase the portal size, we synthesised **L2** (Fig. 1), a tren-hemicryptophane identical to the **L1** ligand, but lacking the methoxy groups on the north pole (see ESI† for synthetic details). The resulting copper complex  $[\text{CuL2}]\text{PF}_6$  maintained an unusual degree of air-stability, despite its more accessible cavity.

Interestingly, no signs of oxidation in the solid state were observed for either  $[\text{CuL1}]\text{PF}_6$  or  $[\text{CuL2}]\text{PF}_6$  upon storage as the solid powder under air for several months (Fig. S2†). To assess stability in the solution phase, a solution of each Cu(I) compound was prepared in DMSO- $d_6$  under air. Both solutions were monitored by  $^1\text{H}$  NMR spectroscopy for two weeks, at the end of which the solutions were bubbled with air for 1 h (Fig. S3 and S4†). In the case of  $[\text{CuL1}]\text{PF}_6$ , no meaningful changes to the spectrum were observed after two weeks of air exposure (Fig. S3b†), nor was bubbling air observed to induce decomposition (Fig. S3c†). For  $[\text{CuL2}]\text{PF}_6$ , slight decomposition was observable after air exposure for two weeks (Fig. S4b†). Subsequent bubbling with air did not appear to cause significant additional decomposition (Fig. S4c†). However, repetition of these experiments with the internal standard revealed a 26% loss of compound over 14 days (Fig. S5 and S6†). This is presumably due to the formation of a paramagnetically broadened Cu(II) species. While susceptible to oxidation in solution over several weeks, the compound demonstrates an unusual lack of air-sensitivity compared to other  $\text{N}_3\text{Cu(I)}$  complexes, which generally undergo oxidation on the time scale of seconds to minutes.<sup>71,72</sup> Both Cu(I) complexes are unusually air-stable in the solution phase, but  $[\text{CuL1}]\text{PF}_6$  appears to be marginally more stable than  $[\text{CuL2}]\text{PF}_6$ . From this, it appears that the size of the portals is important, but cannot fully explain the diminished reactivity towards dioxygen in  $[\text{CuL1}]\text{PF}_6$ .

We hypothesised that increased air stability results from a cavity that is too constrained to easily accommodate Cu(I)- $\text{O}_2$  coordination. Binding to carbon monoxide—a ligand of similar size to dioxygen—was investigated for  $[\text{CuL1}]\text{PF}_6$  and  $[\text{CuL2}]\text{PF}_6$

using infrared (IR) spectroscopy in solution (Fig. 2). When purged with CO, in addition to the free CO peak at  $2137\text{ cm}^{-1}$ , a peak was observed at  $2079\text{ cm}^{-1}$  for both  $[\text{CuL1}]\text{PF}_6$  and  $[\text{CuL2}]\text{PF}_6$ , which we attribute to the formation of the Cu(I)-CO complex. Control experiments using fresh samples under dinitrogen showed neither IR peak. The  $2079\text{ cm}^{-1}$  stretching frequency for the bound peak is not differentiable between  $[\text{CuL1}]\text{PF}_6$  and  $[\text{CuL2}]\text{PF}_6$ , which is expected given the similar Cu(I)-tren ligand electronic environment. These vibrational frequencies are in line with other  $\text{N}_3\text{CuCO}$  complexes reported in the literature, which possess a  $\nu_{\text{CO}}$  of  $2040\text{--}2100\text{ cm}^{-1}$ .<sup>73,74</sup> The experiment was repeated using  $^{13}\text{C}$ -labeled CO, demonstrating the expected isotope shift (Fig. S7†). Binding studies were continued by comparing  $^1\text{H}$  NMR spectra for both compounds under air, after degassing through freeze-pump-thaw cycles, and upon pressurization with CO to 20 psi (Fig. 3). The proton NMR spectra of both compounds showed changes indicative of binding behaviour in the presence of CO, eliciting peak broadening and changes in chemical shift. The degassed and under-air spectra show no differences. As expected, the most-affected NMR peaks belong to the south pole and linker regions of the molecule, nearest the copper centre. In order to remove CO and restore the ligand-free NMR spectrum for both compounds, it was necessary to degas rigorously using the



Fig. 2 IR spectra of (a)  $[\text{CuL1}]\text{PF}_6$  under  $\text{N}_2$  (grey) and CO (red) and (b)  $[\text{CuL2}]\text{PF}_6$  under  $\text{N}_2$  (grey) and CO (red).



Fig. 3 (a)  $^1\text{H}$  NMR spectra (500 MHz,  $\text{CD}_3\text{CN}$ ) of  $[\text{CuL1}]\text{PF}_6$  under air (top), after degassing (middle) and under a 20 psi pressure of carbon monoxide (bottom). (b)  $^1\text{H}$  NMR spectra (500 MHz,  $\text{CD}_3\text{CN}$ ) of  $[\text{CuL2}]\text{PF}_6$  under air (top), after degassing (middle) and under a 20 psi pressure of carbon monoxide (bottom).



freeze–pump–thaw method. The changes to both the  $^1\text{H}$  and  $^{13}\text{C}$  NMR spectra (Fig. S8–S14<sup>†</sup>) of  $[\text{CuL1}]\text{PF}_6$  and  $[\text{CuL2}]\text{PF}_6$  can be attributed to the presence of the CO as the guest.<sup>75,76</sup> Notably, however, no signal corresponding to free or bound carbon monoxide was observed in the  $^{13}\text{C}$  NMR spectrum for either compound. Based on the evidence for a significant Cu(i)–CO bound population, we inferred that the CO off-rate is in an intermediate exchange regime resulting in extreme broadening of the CO-bound peak (Fig. S10 and S11<sup>†</sup>).<sup>77</sup> This hypothesis was confirmed by applying  $^{13}\text{C}$ -labeled CO to the sample, upon which one broad  $^{13}\text{C}$  NMR peak corresponding to bound and free CO was observed (Fig. S12 and S13<sup>†</sup>).

To examine carbon monoxide binding further, cyclic voltammetry (CV) experiments were performed in the absence and presence of carbon monoxide for  $[\text{CuL1}]\text{PF}_6$  and  $[\text{CuL2}]\text{PF}_6$  (Fig. 4). Both compounds displayed highly irreversible electron-transfer kinetics in ligand-free and ligand-bound states. We propose that the irreversible nature of the CVs is attributable to several factors, including the rigidity of the tren ligand within the capsule. In similar copper complexes, ligand rearrangement is necessary to facilitate redox changes at the metal.<sup>78,79</sup> Considerable evidence supports the role of a “rack” or entatic state in copper electron-transfer proteins, in which small movements in one or more axial ligands assist the complex in shifting between preferred coordination geometries, *i.e.*, tetrahedral Cu(i) and square planar/higher-coordinate Cu(II).<sup>11,80</sup> Our system may demonstrate the opposite: the movement of ligands is constrained by the macrocycle, hindering electron transfer. This phenomenon has been seen in similar TREN-based systems displaying irreversible electron-transfer kinetics, in which the constrained ligand enforces a particular coordination geometry upon the metal centre, thus favouring the corresponding oxidation state.<sup>58</sup> The very sluggish ET kinetics observed for these Cu model systems helps to explain their notable Cu(i) air-stability. We note that the extremely slow ET kinetics observed for both Cu(i)–CO species (Fig. 4) may be attributable to the high inner-sphere reorganisation energy associated with rapid CO ligand exchange.<sup>81</sup>

Titration of CO revealed a correlation between the CO concentration and reduction potential of both compounds. As



Fig. 4 Cyclic voltammograms for (a)  $[\text{CuL1}]\text{PF}_6$  and (b)  $[\text{CuL2}]\text{PF}_6$  in the presence of varying concentrations of carbon monoxide. All experiments were performed in anhydrous DMSO with 250 mM TBAPF<sub>6</sub> supporting electrolyte, glassy carbon working electrode, Pt mesh counter electrode, and Ag/AgNO<sub>3</sub> reference electrode calibrated to Fc<sup>+</sup>/Fc at a scan rate of 100 mV s<sup>-1</sup>. Arrows show the start point of the scan (−0.457 V vs. Fc<sup>+</sup>/Fc) and the scan direction.

more CO was titrated during the CV measurement, both apparent  $E_{1/2}$  values shifted anodically, providing additional evidence that CO is coordinating to the metal sites in  $[\text{CuL1}]\text{PF}_6$  and  $[\text{CuL2}]\text{PF}_6$ . Consistent with observations from  $^1\text{H}$  NMR, purging the sample with nitrogen did not revert the CV to its original state in either case. Finally, the Cu(i) oxidation potentials measured in both  $[\text{CuL1}]\text{PF}_6$  and  $[\text{CuL2}]\text{PF}_6$  indicate that oxidation by dioxygen should be thermodynamically favourable. This makes the relative Cu(i) air-stability all the more striking, particularly with such prominent CO binding. The larger portals in  $[\text{CuL2}]\text{PF}_6$  barely affect CO binding or air stability.

To shed light on the preference for CO binding and against O<sub>2</sub> binding, we sought to obtain various crystal structures to visualize the cavity. Crystals suitable for single crystal X-ray diffraction were obtained for  $[\text{CuL1}]\text{PF}_6$  by vapor diffusion of ether into acetonitrile (Fig. 5a). Aware of the “induced fit” model




Fig. 5 Crystal structures of (a)  $[\text{CuL1}]\text{PF}_6$ , (b)  $[\text{CuL1}(\text{MeCN})](\text{OTf})_2$  and (c)  $[\text{CuL1}(\text{N}_3)](\text{OTf})_2$  (counter ions have been removed for clarity). Mapped out cavities for (d)  $[\text{CuL1}]\text{PF}_6$ , (e)  $[\text{CuL1}(\text{MeCN})](\text{OTf})_2$  and (f)  $[\text{CuL1}(\text{N}_3)](\text{OTf})_2$  overlaid with the corresponding crystal structure. Guests and counter ions have been removed for clarity.



observed for related supramolecular complexes,<sup>82</sup> we suspected that a structure of the empty capsule **L1** would not accurately predict the size and shape of the cavity when a ligand is present. Therefore, we also sought to crystallise Cu(I)–ligand and Cu(II)–ligand complexes that should more closely resemble the putative Cu(II)–O<sub>2</sub> complex. Thus, **L1** was crystallised in the presence of Cu(OTf)<sub>2</sub>.<sup>†</sup> The resulting X-ray structure contained a molecule of acetonitrile residing in the cavity and bound axially to the copper (Fig. 5b), similar to the expected binding for CO. Crystallisation of **L1** was also attempted in the presence of Cu(OTf)<sub>2</sub> and *n*-Bu<sub>4</sub>NN<sub>3</sub> in hopes of capturing a ligand coordinated to the Cu centre in a bent fashion, similar to the expected dioxygen coordination geometry.<sup>53,83,84</sup> It was revealed that under these conditions an azide anion was bound to the copper(II) centre (Fig. 5c).

We mapped the surface of the ligand-accessible cavity for each of these crystal structures using Molovol.<sup>85</sup> Cavity surfaces overlaid with the crystal structure (not including the ligands) are presented in Fig. 5d–f. For the empty structure, we observed the cavity to be elongated and narrow near the copper centre, perfectly suited to accommodate acetonitrile (Fig. 5b) or carbon monoxide when bound axially. In fact, the cavity's oblong shape does not permit a bent ligand to bind entirely inside the host molecule. Instead, upon binding to azide at a bent angle, the aryl linkers rotate, allowing the azide ligand to thread between them such that it is only partially contained within the host cavity (Fig. 5c and f). While making space for the ligand, the linker conformational change comes at an entropic cost as it restricts the rotation of the linkers. Azide binding suggests that such binding angles are attainable, but the loss of entropy needs to be compensated for by enthalpic gains. As such, we propose that O<sub>2</sub> binding in [Cu**L1**]PF<sub>6</sub> and [Cu**L2**]PF<sub>6</sub> lacks sufficient enthalpy to fully compensate for the loss of entropy in these host molecules at rt.

A related phenomenon has been observed previously: in the Karlin group's seminal studies of Cu(I)–CO and –O<sub>2</sub> binding, the CO adduct formation rate constant is higher than that of the O<sub>2</sub> adduct,<sup>86</sup> which was attributed to a slower *k*<sub>off</sub> rate of CO for entropic reasons pertaining to ligand re-organisation. In comparison, the intermediate exchange that we observe in the <sup>13</sup>CO NMR may be due to the more constrained nature of the ligand, in which a smaller entropic penalty would be paid for the re-ligation of the TREN ligand than for a pendant pyridyl ligand.<sup>73</sup>

These observations carry implications for the study of cuproproteins and Cu-enzyme mimetics as a whole. Several model systems in the literature use CO as a redox-inactive ligand with similar binding behaviour to O<sub>2</sub>.<sup>15,87,88</sup> Our model system, while unusual, highlights the possibility that some monocopper proteins may bind CO and discriminate against O<sub>2</sub>, as is well known to occur in haemoglobin. Arthropod and molluscan hemocyanins display differing selectivities for CO and O<sub>2</sub>,<sup>89</sup> but this effect is presumably mediated by the cooperativity seen in anywhere from 6 to 160 dicopper sites.<sup>90</sup> The current study suggests the possibility of achieving discrimination in a monocopper protein based upon binding angle and cavity shape.

In conclusion, we have synthesised two unusually air-stable Cu(I) tren-hemicryptophane host molecules. We propose that our system is the first CO-binding Cu(I) complex for which O<sub>2</sub> binding/reaction is demonstrated to be electronically favourable, but is disfavoured due to steric factors alone. IR, NMR, and CV investigations yield strong evidence of discrimination for CO over O<sub>2</sub>. These findings shed insight on an alternate approach for stabilizing Cu(I) species, in which the coordination of a fifth ligand, as preferred by Cu(II), is made entropically unfavourable. This concept may be possible to adapt for small-molecule catalyst development and directed evolution of copper enzymes in the future, granting unusual ligand selectivity and reactivity.

## Data availability

Our laboratory maintains all laboratory notebooks in a secure site, and maintains electronic copies of all spectra on secure servers. We have provided the most critical data in the MS and ESI,<sup>†</sup> and we will be happy to provide any additional information relevant to this paper upon request from reviewers or the broader public, after publication.

## Author contributions


Conceptualization, Y. L., B. S., A. M. S., C. J. S., A. S. M., and I. J. D.; methodology, Y. L., B. S., A. M. S., S. A. D., and S. P. O.; investigation, Y. L., B. S., A. M. S., C. J. S., A. S. M., M. R. G., P. J. C.; formal analysis, B. S.; writing – original draft, A. M. S. and B. S.; writing – review and editing, A. M. S., B. S., C. J. S., A. S. M., and I. J. D.; supervision, I. J. D. and T. E. M.; project administration, I. J. D. and T. E. M.; funding acquisition, I. J. D. and T. E. M.

## Conflicts of interest

There are no conflicts to declare.


## Acknowledgements

This research was supported by NIH Grant R35-GM-131907 to I. J. D. NMR spectrometer use was supported by the NSF Major Research Instrumentation Program (NSF CHE-1827457), NIH supplement awards 3R01GM118510-03S1 and 3R01GM087605-06S1, and Vagelos Institute for Energy Science and Technology. We thank Dr Jun Gu and Dr Chad W. Lawrence for assistance with NMR spectroscopy experiments and Dr Charles W. Ross III for mass spectrometry expertise. The authors thank current members of the Dmochowski lab, and also Prof. Kang Du and Dr Walter Johnsen for fruitful discussions. I. J. D. fondly remembers being an undergraduate attending a lecture by George Whitesides in 1992 where GW described his recent publication in *Science*, “How To Make Water Run Uphill”.<sup>91</sup> The trap was set, and I. J. D. was caught! I. J. D. received wonderful training in the GW laboratory, and is still fascinated by studying complexity in molecular systems more than 30 years later.



## Notes and references

- 1 E. A. Lewis and W. B. Tolman, *Chem. Rev.*, 2004, **104**, 1047–1076.
- 2 L. Gonzaga de França Lopes, F. S. Gouveia Júnior, A. Karine Medeiros Holanda, I. Maria Moreira de Carvalho, E. Longhinotti, T. F. Paulo, D. S. Abreu, P. V. Bernhardt, M.-A. Gilles-Gonzalez, I. Cirino Nogueira Diógenes and E. Henrique Silva Sousa, *Coord. Chem. Rev.*, 2021, **445**, 214096.
- 3 A. C. Rosenzweig and M. H. Sazinsky, *Curr. Opin. Struct. Biol.*, 2006, **16**, 729–735.
- 4 L. M. Blomberg, M. R. A. Blomberg and P. E. M. Siegbahn, *J. Inorg. Biochem.*, 2005, **99**, 949–958.
- 5 T. Shimizu, D. Huang, F. Yan, M. Stranava, M. Bartosova, V. Fojtiková and M. Martinková, *Chem. Rev.*, 2015, **115**, 6491–6533.
- 6 T. G. Spiro, M. Z. Zgierski and P. M. Kozlowski, *Coord. Chem. Rev.*, 2001, **219–221**, 923–936.
- 7 J. Ohata, K. J. Bruemmer and C. J. Chang, *Acc. Chem. Res.*, 2019, **52**, 2841–2848.
- 8 X. Liu, N. Li, M. Li, H. Chen, N. Zhang, Y. Wang and K. Zheng, *Coord. Chem. Rev.*, 2020, **404**, 213109.
- 9 N. Bauer, X. Yang, Z. Yuan and B. Wang, *Chem. Sci.*, 2023, **14**, 3215–3228.
- 10 K. I. Tishchenko, E. K. Beloglazkina, A. G. Mazhuga and N. V. Zyk, *Rev. J. Chem.*, 2016, **6**, 49–82.
- 11 E. I. Solomon, D. E. Heppner, E. M. Johnston, J. W. Ginsbach, J. Cirera, M. Qayyum, M. T. Kieber-Emmons, C. H. Kjaergaard, R. G. Hadt and L. Tian, *Chem. Rev.*, 2014, **114**, 3659–3853.
- 12 P. K. Hota, A. Jose, S. Panda, E. M. Dunietz, A. E. Herzog, L. Wojcik, N. Le Poul, C. Belle, E. I. Solomon and K. D. Karlin, *J. Am. Chem. Soc.*, 2024, **146**, 13066–13082.
- 13 R. H. Holm and E. I. Solomon, *Chem. Rev.*, 2004, **104**, 347–348.
- 14 V. Mahadevan, R. K. Gebbink and T. D. P. Stack, *Curr. Opin. Chem. Biol.*, 2000, **4**, 228–234.
- 15 H. R. Lucas and K. D. Karlin, in *9 Copper-Carbon Bonds in Mechanistic and Structural Probing of Proteins as well as in Situations where Copper is a Catalytic or Receptor Site*, De Gruyter, 2015, pp. 295–362.
- 16 O. Green, B. A. Gandhi and J. N. Burstyn, *Inorg. Chem.*, 2009, **48**, 5704–5714.
- 17 G. Thiabaud, G. Guillemot, I. Schmitz-Afonso, B. Colasson and O. Reinaud, *Angew. Chem., Int. Ed.*, 2009, **48**, 7383–7386.
- 18 L. Le Clainche, Y. Rondelez, O. Sénèque, S. Blanchard, M. Campion, M. Giorgi, A. F. Duprat, Y. Le Mest and O. Reinaud, *Comptes Rendus Acad. Sci. – Ser. IIC Chem.*, 2000, **3**, 811–819.
- 19 K. A. Magnus, H. Ton-That and J. E. Carpenter, *Chem. Rev.*, 1994, **94**, 727–735.
- 20 M. J. Wiester, P. A. Ulmann and C. A. Mirkin, *Angew. Chem., Int. Ed.*, 2011, **50**, 114–137.
- 21 J. W. Lee, S. Samal, N. Selvapalam, H.-J. Kim and K. Kim, *Acc. Chem. Res.*, 2003, **36**, 621–630.
- 22 K. I. Assaf and W. M. Nau, *Chem. Soc. Rev.*, 2014, **44**, 394–418.
- 23 B. Tang, J. Zhao, J.-F. Xu and X. Zhang, *Chem.–Eur. J.*, 2020, **26**, 15446–15460.
- 24 G. Sachdeva, D. Vaya, C. M. Srivastava, A. Kumar, V. Rawat, M. Singh, M. Verma, P. Rawat and G. K. Rao, *Coord. Chem. Rev.*, 2022, **472**, 214791.
- 25 F. Sansone, L. Baldini, A. Casnati and R. Ungaro, *New J. Chem.*, 2010, **34**, 2715–2728.
- 26 O. Santoro and C. Redshaw, *Coord. Chem. Rev.*, 2021, **448**, 214173.
- 27 M. Durmaz, E. Halay and S. Bozkurt, *Beilstein J. Org. Chem.*, 2018, **14**, 1389–1412.
- 28 M. J. McIlldowie, M. Mocerino and M. I. Ogden, *Supramol. Chem.*, 2010, **22**, 13–39.
- 29 G. Crini, *Chem. Rev.*, 2014, **114**, 10940–10975.
- 30 E. Engeldinger, D. Armspach and D. Matt, *Chem. Rev.*, 2003, **103**, 4147–4174.
- 31 D. Zhang, A. Martinez and J.-P. Dutasta, *Chem. Rev.*, 2017, **117**, 4900–4942.
- 32 M. J. Hardie, in *Supramolecular Chemistry*, John Wiley & Sons, Ltd, 2012.
- 33 M. Yoshizawa, J. K. Klosterman and M. Fujita, *Angew. Chem., Int. Ed.*, 2009, **48**, 3418–3438.
- 34 A. J. Kirby, *Angew. Chem., Int. Ed. Engl.*, 1996, **35**, 706–724.
- 35 B. Ginovska, O. Y. Gutiérrez, A. Karkamkar, M.-S. Lee, J. A. Lercher, Y. Liu, S. Raugei, R. Rousseau and W. J. Shaw, *ACS Catal.*, 2023, **13**, 11883–11901.
- 36 P. Bhandari and P. S. Mukherjee, *ACS Catal.*, 2023, **13**, 6126–6143.
- 37 Y. Makita, T. Danno, K. Ikeda, H.-H. Lee, T. Abe, K. Sogawa, A. Nomoto, S. Fujiwara and A. Ogawa, *Tetrahedron Lett.*, 2017, **58**, 4507–4509.
- 38 S. A. Ikbāl, C. Colomban, D. Zhang, M. Delecluse, T. Brotin, V. Dufaud, J.-P. Dutasta, A. B. Sorokin and A. Martinez, *Inorg. Chem.*, 2019, **58**, 7220–7228.
- 39 T. K. Ronson, H. Nowell, A. Westcott and M. J. Hardie, *Chem. Commun.*, 2010, **47**, 176–178.
- 40 A. Schmitt, S. Collin, C. Bucher, V. Maurel, J.-P. Dutasta and A. Martinez, *Org. Biomol. Chem.*, 2015, **13**, 2157–2161.
- 41 O. Perraud, J.-B. Tommasino, V. Robert, B. Albela, L. Khrouz, L. Bonneviot, J.-P. Dutasta and A. Martinez, *Dalton Trans.*, 2013, **42**, 1530–1535.
- 42 C. K. Chang and T. G. Traylor, *Proc. Natl. Acad. Sci. U. S. A.*, 1975, **72**, 1166–1170.
- 43 D. H. Busch, L. L. Zimmer, J. J. Grzybowski, D. J. Olszanski, S. C. Jackels, R. C. Callahan and G. G. Christoph, *Proc. Natl. Acad. Sci. U. S. A.*, 1981, **78**, 5919–5923.
- 44 J. P. Collman, A. Dey, R. A. Decreau, Y. Yang, A. Hosseini, E. I. Solomon and T. A. Eberspacher, *Proc. Natl. Acad. Sci. U. S. A.*, 2008, **105**, 9892–9896.
- 45 J. P. Collman, A. Dey, C. J. Barile, S. Ghosh and R. A. Decréau, *Inorg. Chem.*, 2009, **48**, 10528–10534.
- 46 Y. Makita, K. Sugimoto, K. Furuyoshi, K. Ikeda, S. Fujiwara, T. Shin-ike and A. Ogawa, *Inorg. Chem.*, 2010, **49**, 7220–7222.
- 47 Y. Lin, M. R. Gau, P. J. Carroll and I. J. Dmochowski, *J. Org. Chem.*, 2022, **87**, 5158–5165.



- 48 Y. Lin, K. Du, M. R. Gau and I. J. Dmochowski, *Chem. Sci.*, 2023, **14**, 291–297.
- 49 Y. Makita, K. Ikeda, K. Sugimoto, T. Fujita, T. Danno, K. Bobuatong, M. Ehara, S. Fujiwara and A. Ogawa, *J. Organomet. Chem.*, 2012, **706–707**, 26–29.
- 50 Y. Makita, T. Fujita, T. Danno, M. Inoue, M. Ueshima, S. Fujiwara and A. Ogawa, *Supramol. Catal.*, 2012, **1**, 11–13.
- 51 M. P. Lanci, V. V. Smirnov, C. J. Cramer, E. V. Gauchenova, J. Sundermeyer and J. P. Roth, *J. Am. Chem. Soc.*, 2007, **129**, 14697–14709.
- 52 A. Poater and L. Cavallo, *Inorg. Chem.*, 2009, **48**, 4062–4066.
- 53 C. Würtele, E. Gaoutchenova, K. Harms, M. C. Holthausen, J. Sundermeyer and S. Schindler, *Angew. Chem., Int. Ed.*, 2006, **45**, 3867–3869.
- 54 A. K. Mishra and S. Verma, *Inorg. Chem.*, 2010, **49**, 3691–3693.
- 55 L. Zhou, D. Powell and K. M. Nicholas, *Inorg. Chem.*, 2007, **46**, 2316–2321.
- 56 T. Brückmann, J. Becker, C. Würtele, M. T. Seuffert, D. Heuler, K. Müller-Buschbaum, M. Weiß and S. Schindler, *J. Inorg. Biochem.*, 2021, **223**, 111544.
- 57 A. Mondal, K. P. Reddy, S. Som, D. Chopra and S. Kundu, *Inorg. Chem.*, 2022, **61**, 20337–20345.
- 58 D. K. Chand and P. K. Bharadwaj, *Inorg. Chem.*, 1998, **37**, 5050–5055.
- 59 L. Chaloner and X. Ottenwaelder, *Inorg. Chim. Acta*, 2018, **481**, 106–112.
- 60 L. Chaloner, A. Khomutovskaya, F. Thomas and X. Ottenwaelder, *Dalton Trans.*, 2016, **45**, 11109–11119.
- 61 S. Paria, Y. Morimoto, T. Ohta, S. Okabe, H. Sugimoto, T. Ogura and S. Itoh, *Eur. J. Inorg. Chem.*, 2018, **2018**, 1976–1983.
- 62 G. Izzet, B. Douziech, T. Prangé, A. Tomas, I. Jabin, Y. Le Mest and O. Reinaud, *Proc. Natl. Acad. Sci. U. S. A.*, 2005, **102**, 6831–6836.
- 63 G. De Leener, D. Over, C. Smet, D. Cornut, A. G. Porrás-Gutierrez, I. López, B. Douziech, N. Le Poul, F. Topić, K. Rissanen, Y. Le Mest, I. Jabin and O. Reinaud, *Inorg. Chem.*, 2017, **56**, 10971–10983.
- 64 G. Izzet, J. Zeitouny, H. Akdas-Killig, Y. Frapart, S. Ménage, B. Douziech, I. Jabin, Y. Le Mest and O. Reinaud, *J. Am. Chem. Soc.*, 2008, **130**, 9514–9523.
- 65 G. Izzet, M.-N. Rager and O. Reinaud, *Dalton Trans.*, 2007, 771–780.
- 66 D. Sengupta, P. Melix, S. Bose, J. Duncan, X. Wang, M. R. Mian, K. O. Kirlikovali, F. Joodaki, T. Islamoglu, T. Yildirim, R. Q. Snurr and O. K. Farha, *J. Am. Chem. Soc.*, 2023, **145**, 20492–20502.
- 67 M. P. Tang, L. Zhu, Y. Deng, Y. Shi, S. K.-M. Lai, X. Mo, X.-Y. Pang, C. Liu, W. Jiang, E. C. M. Tse and H. Y. Au-Yeung, *Angew. Chem., Int. Ed.*, 2024, **63**, e202405971.
- 68 S. Ostrowska, P. Arnaut, D. J. Liptrot, C. S. J. Cazin and S. P. Nolan, *Chem. Commun.*, 2023, **59**, 9126–9129.
- 69 L. Le Clainche, M. Giorgi and O. Reinaud, *Eur. J. Inorg. Chem.*, 2000, **2000**, 1931–1933.
- 70 C. Chuang, K. Lim, Q. Chen, J. Zubieta and J. W. Canary, *Inorg. Chem.*, 1995, **34**, 2562–2568.
- 71 S. Schindler, *Eur. J. Inorg. Chem.*, 2000, **2000**, 2311–2326.
- 72 K. Komiyama, H. Furutachi, S. Nagatomo, A. Hashimoto, H. Hayashi, S. Fujinami, M. Suzuki and T. Kitagawa, *Bull. Chem. Soc. Jpn.*, 2004, **77**, 59–72.
- 73 H. C. Fry, H. R. Lucas, A. A. Narducci Sarjeant, K. D. Karlin and G. J. Meyer, *Inorg. Chem.*, 2008, **47**, 241–256.
- 74 Y. Rondelez, O. Sénèque, M.-N. Rager, A. F. Duprat and O. Reinaud, *Chem.–Eur. J.*, 2000, **6**, 4218–4226.
- 75 K. E. Chaffee, H. A. Fogarty, T. Brotin, B. M. Goodson and J.-P. Dutasta, *J. Phys. Chem. A*, 2009, **113**, 13675–13684.
- 76 L. Garel, J.-P. Dutasta and A. Collet, *Angew. Chem., Int. Ed. Engl.*, 1993, **32**, 1169–1171.
- 77 L. Mureddu and G. W. Vuister, *FEBS J.*, 2019, **286**, 2035–2042.
- 78 P. J. Griffin, B. J. Charette, J. H. Burke, J. Vura-Weis, R. D. Schaller, D. J. Gosztola and L. Olshansky, *J. Am. Chem. Soc.*, 2022, **144**, 12116–12126.
- 79 G. Coullerez, E. Malmström and M. Jonsson, *J. Phys. Chem. A*, 2006, **110**, 10355–10360.
- 80 B. Dicke, A. Hoffmann, J. Stanek, M. S. Rampp, B. Grimm-Lebsanft, F. Biebl, D. Rukser, B. Maerz, D. Görries, M. Naumova, M. Biednov, G. Neuber, A. Wetzel, S. M. Hofmann, P. Roedig, A. Meents, J. Bielecki, J. Andreasson, K. R. Beyerlein, H. N. Chapman, C. Bressler, W. Zinth, M. Rübhausen and S. Herres-Pawlis, *Nat. Chem.*, 2018, **10**, 355–362.
- 81 C. Zong, C. J. Wilson, T. Shen, P. Wittung-Stafshede, S. L. Mayo and P. G. Wolynes, *Proc. Natl. Acad. Sci. U. S. A.*, 2007, **104**, 3159–3164.
- 82 O. Taratula, P. A. Hill, N. S. Khan, P. J. Carroll and I. J. Dmochowski, *Nat. Commun.*, 2010, **1**, 148.
- 83 Y. Kobayashi, K. Ohkubo, T. Nomura, M. Kubo, N. Fujieda, H. Sugimoto, S. Fukuzumi, K. Goto, T. Ogura and S. Itoh, *Eur. J. Inorg. Chem.*, 2012, **2012**, 4574–4578.
- 84 S. Kim, J. Y. Lee, R. E. Cowley, J. W. Ginsbach, M. A. Siegler, E. I. Solomon and K. D. Karlin, *J. Am. Chem. Soc.*, 2015, **137**, 2796–2799.
- 85 J. B. Maglic and R. Lavendomme, *J. Appl. Crystallogr.*, 2022, **55**, 1033–1044.
- 86 H. C. Fry, D. V. Scaltrito, K. D. Karlin and G. J. Meyer, *J. Am. Chem. Soc.*, 2003, **125**, 11866–11871.
- 87 C. D. Kline and N. J. Blackburn, *Biochemistry*, 2016, **55**, 6652–6661.
- 88 E. E. Chufán, S. T. Prigge, X. Siebert, B. A. Eipper, R. E. Mains and L. M. Amzel, *J. Am. Chem. Soc.*, 2010, **132**, 15565–15572.
- 89 H. Decker, P. R. Connelly, C. H. Robert and S. J. Gill, *Biochemistry*, 1988, **27**, 6901–6908.
- 90 H. Decker, N. Hellmann, E. Jaenicke, B. Lieb, U. Meissner and J. Markl, *Integr. Comp. Biol.*, 2007, **47**, 631–644.
- 91 M. K. Chaudhury and G. M. Whitesides, *Science*, 1992, **256**, 1539–1541.

