

COMMENTARY

View Article Online
View Journal | View IssueCite this: *Chem. Sci.*, 2025, **16**, 18478

All publication charges for this article have been paid for by the Royal Society of Chemistry

DOI: 10.1039/d5sc90203g

rsc.li/chemical-science

Open Access Article. Published on 02 October 2025. Downloaded on 04.02.26 16:38:45.
 This article is licensed under a Creative Commons Attribution 3.0 Unported Licence.

9-Azabicyclo[3.3.1]nonane-3-one *N*-oxyl (ketoABNO) was originally developed by Dupeyre and Rassa as a stable radical that persists at ambient temperature.¹ Later, Iwabuchi and co-workers demonstrated that sterically less-hindered *N*-oxyl radicals—such as AZADO and ABNO, stabilized by Bredt's rule—could broaden the substrate scope of TEMPO-catalyzed alcohol oxidation.² Building on these foundational studies, we reported in *Chemical Science* the implementation of ketoABNO as a catalyst for the aerobic oxidation of amines in conjunction with a CuBr-ligand complex (<https://doi.org/10.1039/C2SC20699D>).³ This study established a highly efficient method for the aerobic oxidation of amines to imines under mild conditions (Fig. 1a). Furthermore, by employing a chiral copper-bisoxazoline complex, catalytic asymmetric amino acid synthesis was achieved from a glycine derivative and a nitroalkane (Fig. 1b). At the time, although *N*-oxyl radicals such as TEMPO were gaining attraction as oxidation catalysts,⁴ ketoABNO itself had received

relatively little attention. While Kerton and co-workers independently reported CuBr₂/TEMPO-catalyzed aerobic oxidation of amines around the same time (Fig. 1c),⁵ the ketoABNO system featured broader substrate scope and practical scalability. To support further studies, we developed a concise, three-step synthesis of ketoABNO, enabling gram-scale preparation (Fig. 1d). These features contributed to the growing recognition of ketoABNO, which, over the past decade, has emerged as a powerful catalyst/reagent for both alcohol oxidation and selective modification of amino acid residues in proteins. In this Commentary, we reflect on these developments and consider future directions.


KetoABNO offers several advantages for oxidative transformations, including a high redox potential (+521 mV vs. Ag/Ag⁺) relative to TEMPO (+320 mV) and ABNO (+419 mV), as well as reduced steric hindrance (Fig. 1e). These properties enable selective and efficient aerobic oxidation of amines and alcohols in conjunction with copper complexes. For example, Zhang and co-workers employed a Cu salt/ketoABNO catalytic system in a micro-packed bed reactor for

the aerobic oxidation of ethyl lactate to ethyl pyruvate, a valuable intermediate for pharmaceuticals and nutritional supplements.⁶ Notably, this system outperformed conventional Cu salt/TEMPO systems in both activity and selectivity.


Organocatalytic aerobic alcohol oxidation conditions without the use of metal-complex catalysts were developed by Liang and Hu,⁷ Iwabuchi,^{2b} and Stahl,⁸ who introduced NaNO₂/O₂-based cocatalytic systems for the reoxidation of *N*-oxyl radicals. These advances were subsequently adapted to ketoABNO systems (Fig. 1f), enabling metal-free aerobic oxidations. A key benefit of this advancement was the expanded substrate scope to include aldehydes, whose oxidation had previously been hindered by the acidity of the resulting carboxylic acids (Fig. 1g).⁹ Taking advantage of the high oxidation potential of ketoABNO, we later demonstrated the aerobic organocatalytic oxidation of α -fluoroalkyl alcohols to the corresponding fluoroalkyl ketones at room temperature (Fig. 1h).¹⁰ Additional modifications, such as the use of bismuth nitrate as an NO_x mediator, were introduced by Li and co-workers.¹¹ Collectively, these contributions have

Graduate School of Pharmaceutical Sciences, The University of Tokyo, Japan. E-mail: kanai@mol.f.u-tokyo.ac.jp


a. Our work, 2012




b.




c. Kerton's work, 2012




d. Synthetic scheme of ketoABNO




e. Oxidation potential of N-oxyl radicals



f. Stahl's work, 2013



g. Stahl's work, 2016



h. Our work, 2015




Fig. 1 Utilities and properties of ketoABNO. (a) Catalytic aerobic oxidation of amines developed by our group. (b) Catalytic asymmetric oxidative coupling between amines and nitroalkanes. (c) Catalytic aerobic oxidation of amines by $\text{CuBr}_2/\text{TEMPO}$ catalysis developed by Kerton's group. (d) Synthetic route of ketoABNO. (e) Oxidation potentials of *N*-oxyl radicals (vs. Ag/Ag^+). (f) Catalytic aerobic oxidation of alcohols developed by Stahl's group. (g) Catalytic aerobic oxidation of aldehydes developed by Stahl's group. (h) Catalytic aerobic oxidation of fluoroalkyl alcohols developed by our group.

firmly established ketoABNO as a versatile reagent for mild alcohol oxidation. Nevertheless, for many alcohol oxidations, ABNO remains a practical alternative due to its comparable reactivity and broader synthetic accessibility, including its frequent use as a precursor to functionalized ABNO derivatives.¹²

In parallel with advances in the small-molecule-directed methodology, ketoABNO has also found unique applications in biomolecular chemistry. Owing to its mildness and aqueous compatibility—critical features for protein chemistry—ketoABNO has enabled amino acid residue-selective chemical modifications of peptides and proteins. We demonstrated an artificial peptidase-type reaction—the serine-selective aerobic cleavage of peptide bonds—using a $\text{CuI}/\text{ketoABNO}$ system, initiated by oxidation of the primary alcohol in the serine side chain (Fig. 2a).¹³ Oxidation of the hydroxymethyl group at

serine to an aldehyde triggers hydrolysis of the peptide backbone. This reaction was applicable to proteins, with no detectable cleavage observed at threonine residues.


During these studies, we serendipitously discovered that ketoABNO selectively forms an adduct with tryptophan residues, marking the first example of heavy-metal-free, tryptophan-selective protein modification.¹⁴ Under ketoABNO/ NaNO_2 conditions—without the use of a copper salt—in an acetonitrile/water/acetic acid mixture, the adduct was formed at the 3-position of tryptophan's indole ring in peptides and proteins (Fig. 2b(A)). The reaction proceeded *via* a Friedel-Crafts mechanism with a ketoABNO-derived oxoammonium intermediate. Building on this mechanistic insight, we developed milder methods for generating the reactive oxoammonium species, such as electrochemical oxidation (Fig. 2b(B))¹⁵ and

a ketoABNOH/TEMPO⁺ system (Fig. 2b(C)).¹⁶ The precise reason for the high chemoselectivity of ketoABNO-derived oxoammonium toward indoles remains to be elucidated.


Because tryptophan is the rarest of the canonical amino acids and is typically buried within proteins due to its hydrophobic characteristics, this method offers potential advantages for producing bioprobes with high homogeneity.¹⁷ The ketone moiety of ketoABNO serves as a versatile handle for straightforward derivatization, enabling attachment to cytotoxic agents for constructing antibody-drug conjugates (ADCs),¹⁸ gold nanoparticles for the construction of antibody-derived cryo-electron microscopy probes,¹⁶ and nucleic acids for generating DNA-protein conjugates¹⁹ (Fig. 2c). This synthetic methodology holds promise for broader application in the development of novel biological tools, diagnostic agents, and therapeutics.



a. Serine-selective peptide/protein cleavage



b. Tryptophan-selective modification



c. Application of tryptophan-selective modification




Fig. 2 Applications of ketoABNO to peptide and protein modifications. (a) Serine-selective peptide/protein cleavage. (b) Tryptophan-selective biconjugation. (c) Applications to life sciences.

In summary, we have outlined the evolution of ketoABNO chemistry since our initial report in 2012. The reagent's ease of use—requiring only simple mixing with a cocatalyst and/or oxidant, including molecular oxygen—combined with its water tolerance, mild and tunable reactivity, and high chemoselectivity, makes it a highly attractive tool in modern synthetic chemistry and pharmaceutical sciences. We anticipate continued growth of this platform at the interface of synthetic methodology, biology, and medicine.

Author contributions

M.T. and M.K. wrote the manuscript.

Conflicts of interest

There are no conflicts to declare.

Data availability

There is no additional data associated with this article.

References

- 1 R. Dupeyre and A. Rassa, *J. Am. Chem. Soc.*, 1966, **88**, 3180.
- 2 (a) M. Shibuya, M. Tomizawa, I. Suzuki and Y. Iwabuchi, *J. Am. Chem. Soc.*, 2006, **128**, 8412; (b) M. Shibuya, Y. Osada, Y. Sasano, M. Tomizawa and Y. Iwabuchi, *J. Am. Chem. Soc.*, 2011, **133**, 6497.
- 3 T. Sonobe, K. Oisaki and M. Kanai, *Chem. Sci.*, 2012, **3**, 3249.
- 4 Q. Cao, L. Dornan, L. Rogan, N. Hughes and M. Muldoon, *Chem. Commun.*, 2014, **50**, 4524.
- 5 Z. Hu and F. M. Kerton, *Org. Biomol. Chem.*, 2012, **10**, 1618.
- 6 C. Zhang, J. Luo, B. Xie, W. Liu and J. Zhang, *Chem. Eng. Sci.*, 2024, **292**, 11985.
- 7 R. Liu, X. Liang, C. Dong and X. Hu, *J. Am. Chem. Soc.*, 2004, **126**, 4113.
- 8 M. Lauber and S. Stahl, *ACS Catal.*, 2013, **3**, 2612.
- 9 K. Miles, M. Abrams, C. Landis and S. Stahl, *Org. Lett.*, 2016, **18**, 3590.



- 10 Y. Kadoh, M. Tashiro, K. Oisaki and M. Kanai, *Adv. Synth. Catal.*, 2015, **357**, 2193.
- 11 Y. Hu, L. Chen, G. Shen, J. Li, S. Li, H. Li and Y. Li, *Molecules*, 2022, **27**, 3727.
- 12 (a) L. Rogan, N. Hughes, Q. Cao, L. Dornan and M. Muldoon, *Catal. Sci. Technol.*, 2014, **4**, 1720; (b) B. Karimi, M. Ghahremani, H. Vali, R. Ciriminna and M. Pagliaro, *Chem. Commun.*, 2021, **57**, 8897.
- 13 Y. Seki, K. Tanabe, D. Sasaki, Y. Sohma, K. Oisaki and M. Kanai, *Angew. Chem., Int. Ed.*, 2014, **126**, 6619.
- 14 Y. Seki, T. Ishiyama, D. Sasaki, J. Abe, Y. Sohma, K. Oisaki and M. Kanai, *J. Am. Chem. Soc.*, 2016, **138**, 10798.
- 15 R. Kuroda, E. Toyama, T. Sugai, M. Imai, K. Watanabe, N. Konoue, K. Maruyama, M. Kondo, S. Masaoka, T. Saitoh, K. Oisaki and M. Kanai, *ChemRxiv*, 2025, preprint, DOI: [10.26434/chemrxiv-2025-s1wv9-v3](https://doi.org/10.26434/chemrxiv-2025-s1wv9-v3).
- 16 K. Malawska, S. Takano, K. Oisaki, H. Yanagisawa, M. Kikkawa, T. Tsukuda and M. Kanai, *Bioconjugate Chem.*, 2023, **34**, 781.
- 17 (a) J. deGruyter, L. Malins and P. Baran, *Biochemistry*, 2017, **56**, 3863; (b) E. Hoyt, P. Cal, B. Oliveira and G. Bernardes, *Nat. Rev. Chem.*, 2019, **3**, 147; (c) N. Kjærsgaard, T. Nielsen and K. Gothelf, *ChemBioChem*, 2022, **23**, e202200245.
- 18 (a) K. Maruyama, K. Malawska, N. Konoue, K. Oisaki and M. Kanai, *Synlett*, 2020, **31**, 784; (b) H. Tagawa, K. Maruyama, K. Sasaki, N. Konoue, A. Kishimura, M. Kanai, T. Mori, K. Oisaki and Y. Katayama, *RSC Adv.*, 2020, **10**, 16727.
- 19 A. Spampinato, D. Leone, R. Pohl, J. Tarábek, M. Soltysová, M. Polák, A. Kádek, V. Sykorová, P. Rezáková and M. Hocek, *Chem.-Eur. J.*, 2024, **30**, e202402151.

