



**Ferritin Exhibits Michaelis-Menten Behavior with Oxygen
but not with Iron During Iron Oxidation and Core
Mineralization**

Journal:	<i>Metallomics</i>
Manuscript ID	MT-ART-01-2019-000001.R1
Article Type:	Paper
Date Submitted by the Author:	24-Jan-2019
Complete List of Authors:	Bou-Abdallah, Fadi; SUNY at Potsdam, Chemistry Flint, Nicholas; SUNY Potsdam Wilkinson, Tyler; Clarkson University Salim, Samantha; Clarkson University Srivastava, Ayush; SUNY Potsdam Poli, Maura; Universita degli Studi di Brescia Arosio, Paolo; Universita degli Studi di Brescia Melman, Artem; Clarkson University, Chemistry + Biomolecular Science

Significance to metalloomics

Ferritins are highly conserved supramolecular protein nanostructures that play an important role in the biological management of iron. Thousands of iron atoms can be safely and reversibly sequestered inside its hollow cavity. Our results show that iron oxidation kinetics in mammalian ferritins do not obey a Michaelis-Menten saturation kinetic behavior, as previously proposed. Physiologically relevant values for K_m , O_2 ($\sim 1-2 \mu\text{M}$) are observed, suggesting that iron deposition inside ferritin occurs even under conditions of severe hypoxia. Our study provides fundamental insights into iron oxidation kinetics in ferritin and the general mechanism of iron metabolism.

Ferritin Exhibits Michaelis-Menten Behavior with Oxygen but not with Iron During Iron Oxidation and Core Mineralization

Fadi Bou-Abdallah*¹, Nicholas Flint¹, Tyler Wilkinson², Samantha Smith², Ayush Kumar Srivastava^{1,3}, Maura Poli³, Paolo Arosio³, Artem Melman*²

¹ Department of Chemistry, State University of New York, Potsdam, NY 13676, USA

² Department of Chemistry & Biomolecular Science, Clarkson University, Potsdam, NY 13699, USA

³ Department of Molecular and Translational Medicine, University of Brescia, 25121 Brescia, Italy

* where correspondence should be addressed: bouabdf@potdam.edu ; amelman@clarkson.edu

Abstract

The excessively high and inconsistent literature values for $K_{m,Fe}$ and K_{m,O_2} prompted us to examine the iron oxidation kinetics in ferritin, the major iron storage protein in mammals, and to determine whether a traditional Michaelis-Menten enzymatic behavior is obeyed. The kinetics of Fe(II) oxidation and mineralization catalyzed by three different types of ferritins (recombinant human homopolymer 24H, HuHF, human heteropolymer ~ 21H:3L, HL, and horse spleen heteropolymer ~ 3.3H:20.7L, HosF) were therefore studied under physiologically relevant O₂ concentrations, but also in the presence of excess Fe(II) and O₂ concentrations. The observed iron oxidation kinetics exhibited two distinct phases (Phase I and Phase II), neither of which obeyed Michaelis-Menten kinetics. While Phase I was very rapid and corresponded to the oxidation of approximately 2 Fe(II) ions per H-subunit, Phase II was much slower and varied linearly with the concentration of iron(II) cations in solution, independent of the size of the iron core. Under low oxygen concentration close to physiological, the iron uptake kinetics revealed a Michaelis-Menten behavior with K_{m,O_2} values in the low μ M range (i.e. ~ 1 – 2 μ M range). Our experimental K_{m,O_2} values are significantly lower than typical cellular oxygen, indicating that iron oxidation and mineralization in ferritin should not be affected by the oxygenation levels of cells, and should proceed even under hypoxic events. A kinetic model is proposed in which the inhibition of the protein's activity is caused by bound iron(III) cations at the ferroxidase center, with the rate limiting step corresponding to an exchange or a displacement reaction between incoming iron(II) cations and bound Fe(III) cations.

Introduction

Iron is an essential nutrient for virtually most forms of life and an important component of many cellular processes including oxygen transport, electron transfer reactions, energy metabolism, DNA synthesis and gene regulation¹. To mitigate problems associated with iron-initiated radical chemistry, and the insolubility of Fe(III) ions at neutral pHs, cells synthesize ferritin, a ubiquitous and well-characterized iron storage and detoxification protein, capable of sequestering thousands of iron atoms in the form of a biologically available ferric oxy/hydroxide mineral core²⁻⁷. For most organisms, ferritin is a polypeptide of 24-subunits that can vary, selectively, in amino acid sequence and subunit type, and can be either enzymatically active (due to di-iron ferroxidase centers present on H-subunits or H-like subunits having a MW of ~ 21,000 kDa) or enzymatically inactive (no ferroxidase centers on L-subunits having a MW of ~ 19,000 kDa). In mammals, the H- and L-subunits co-assemble in various H:L ratios with a tissue specific distribution to form a shell-like structure with 4/3/2 octahedral symmetry^{2, 5-8}. Examples of isoferritins from different human tissues include placenta (~ 20% H, 80% L), spleen (~ 10% H, 90% L), liver (~ 50% H, 50% L), and heart (~ 90% H, 10% L). The active H-subunit has a ferroxidase center that rapidly catalyzes the oxidation of Fe(II) to Fe(III), and ultimately its hydrolysis to form the iron inorganic core inside the protein shell, whereas the L-subunit does not. A recent study from our laboratory demonstrated a dual role for the L-subunit in facilitating iron turnover at the ferroxidase centers of H-subunits, and in mineralization of the iron core⁸.

While the majority of iron present in living systems is predominantly associated with enzymes and proteins, a minor fraction is chelatable and loosely bound to a heterogeneous population of ligands, including polypeptides and metal complexing groups such as carboxylates, phosphates, amides, thiolates, and hydroxylates^{9, 10}. This exchangeable pool of iron is referred to as cellular labile iron pool (LIP), is redox-active with an estimated concentration within mammalian cells in the low μM range (i.e. less than 5 % of the estimated total cell iron pool of 20 to 100 μM). In extracellular fluids, iron is tightly chelated by transferrin (i.e. affinity constant $\sim 10^{22} \text{ M}^{-1}$), and is safely carried to the interior of cells, via receptor-mediated endocytosis. This process involves the transferrin receptor which has nM affinity to the iron-loaded protein¹¹. The interaction of the iron-loaded transferrin with the transferrin receptor is a key cellular process that occurs during the normal course of iron metabolism. However, although iron bound to transferrin is redox-inactive, and virtually non-exchangeable, transferrin is only about 30% saturated with iron in normal human

1
2
3 serum. The approximately 25-50 μM transferrin present in the serum is unequally distributed
4 between fully saturated diferric-protein complex ($\sim 27\%$), half saturated monoferric-protein
5 complexes ($\sim 11\text{-}23\%$) and iron-free protein ($\sim 40\%$)¹².
6
7

8 Over the past several decades, numerous *in-vitro* studies have aimed at understanding the
9 mechanism of iron uptake, oxidation, and core formation in ferritins. In most ferritins, two widely
10 accepted models have been proposed, the protein catalysis model, and the crystal growth model,
11 or a combination thereof²⁻⁸, although key structural differences and disagreements on suggested
12 steps of the proposed models remain unresolved^{13, 14, 15}. Nonetheless, the overall and generally
13 accepted multistep mechanism involves the binding of ferrous ions at the protein catalytic sites,
14 followed by oxidation and movement of the resulting protein-ferric intermediate species ($\mu\text{-}1,2\text{-}$
15 peroxo-diFe(III) and $\mu\text{-oxo(hydroxo)-bridged diiron(III)}$) into the protein cavity, where a stable
16 nucleus of crystalline mineral (ferrihydrite) starts to form. The growing mineral core then provides
17 additional nucleation sites onto which incoming Fe(II) can be further deposited and oxidized.
18 While a large majority of these studies has been performed with recombinant homopolymeric
19 ferritin of H- or L-type subunits, the widespread occurrence of heteropolymeric ferritins in
20 mammals, necessitated a detailed mechanistic understanding of the complementary roles of H- and
21 L-subunits in the biological processing and management of iron by these proteins. A recent study
22 from our lab examined the kinetics of iron oxidation and mineralization reactions in recombinant
23 heteropolymeric human ferritin, using ferritin samples with different H to L compositions (i.e. H-
24 rich or $\sim 20\text{H:}4\text{L}$ and L-rich or $\sim 22\text{L:}2\text{H}$). The data demonstrated a critical role of L-subunits in
25 enhancing the activity of H-subunits, and increasing the capacity of the protein to take up iron, in
26 support of the different distributions of isoferritins of different subunits composition in tissues of
27 different organs⁸. However, while heteropolymeric H-rich ferritin sample outperformed the
28 homopolymeric H-subunits ferritin in terms of iron oxidation and core formation kinetics,
29 important enzymatic parameters including K_M , V_{max} , k_{cat} , and catalytic efficiency k_{cat}/K_M were
30 missing. Previously published values were either compromised by the experimental conditions
31 employed, including uncontrolled and high oxygen concentration, the chelating ability of the
32 buffer used^{16, 17}, or were excessively high (i.e. in the mM range)^{18, 19}, and physiologically
33 irrelevant, considering the low μM $[\text{O}_2]$ and cellular labile iron pool concentrations. Here, we
34 explore and compare the enzymatic activity, and kinetic parameters of three different ferritin
35 samples, including an H-rich heteropolymer ferritin ($\sim 20\text{-}21\text{H:}4\text{-}3\text{L}$), an exclusively
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 homopolymer H-chain ferritin (100% H), and a naturally occurring heteropolymer ferritin from
4 horse spleen (~21-22L:3-2H).
5
6
7

8 **Materials and Methods**

9

10 **Ferritin samples and chemicals.** The H-rich human heteropolymer (HL) and homopolymer H-
11 chain (HuHF) ferritin samples were produced and purified as described elsewhere^{8,25}. All of the
12 proteins employed in this study are iron-free (apo-protein), although we do not discount the
13 presence of minute amounts of iron inside the protein's cavity. Protein concentrations were
14 determined spectrophotometrically at 280 nm using a molar absorptivity of 24,000 M⁻¹cm⁻¹ per
15 subunit for HL, 21,000 M⁻¹cm⁻¹ per subunit for HuHF⁸, and 19,500 M⁻¹cm⁻¹ per subunit for HosF
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

The subunit composition of recombinant heteropolymer ferritin was quantified by SDS-CGE and SDS-PAGE⁸ and was found to contain ~ 15% L-subunits and ~ 85% H-subunits (i.e. ~ 20-21 H and 3-4 L). All chemicals employed in this study were reagent grade and used without further purification. MOPS (3-(*N*-morpholino)propanesulfonic acid) buffer was purchased from Research Organics (Cleveland, OH), ferrous sulfate heptahydrate from J. T. Baker (Phillipsburg, NJ), sodium dithionite, 2,2'-bipyridyl and iron-free horse spleen (HosF) ferritin from Sigma-Aldrich (St. Louis, MO). Fe(II) stock solutions were freshly prepared immediately before each experiment in a dilute HCl solution at pH 2.0.

Kinetic measurements. Conventional UV-vis spectroscopy was performed on a Varian Cary 50 Bio spectrophotometer from Agilent Technologies. All experiments were conducted at 25 °C, in 100 mM MOPS buffer and 100 mM NaCl, pH 7.4, and reagents concentrations for each experiments are given in the figure captions. All kinetic experiments were repeated two to four times using independent protein preparations to ensure reproducibility. The kinetic traces shown in the figures represent one of multiple individual runs.

Iron oxidation kinetics in the presence of excess O₂. The kinetics of iron oxidation in ferritin were followed at 305 nm where the Fe(III) oxo(hydroxo) species absorbs. The concentrations of HuHF, HL and HosF ferritin samples employed in this study were 0.5, 0.5, and 5 μM, respectively, and the Fe(II) concentrations added to these samples varied between 20 and 250 μM for HuHF and HL, and 30 to 215 μM for HosF. The use of 10-fold higher concentration of HosF was necessary to compensate for the low H-subunit content of the protein, and the overall lower oxidation rates. Under these conditions, the Fe(II)/ferritin ratios varied between a minimum of ~

1
2
3 2 Fe/H-subunit to a maximum of ~ 500 Fe/protein. The instrument was zeroed using the ferritin
4 solution, prepared in buffer as the blank. Typically two or three μL of a ferrous sulfate solution
5 prepared in deionized H_2O (pH 2) were injected into a 1.0 mL protein solution, with rapid spin bar
6 stirring under the conditions stated in the figure captions. Time-dependent absorbance kinetic
7 traces were collected at 25 °C and the data analyzed with Excel or OriginLab version 8.0
8 (OriginLab Corp.).
9

10
11
12 **Iron oxidation kinetics in the presence of excess Fe(II) and low O_2 concentrations.** Additional
13 kinetic experiments were performed in the presence of reduced O_2 concentrations (11 to 30 μM)
14 and a large excess of Fe(II) (at least ten-fold excess over O_2). Solutions of ferritin in 100 MOPS,
15 and 100 mM NaCl, pH 7.4, iron(II) sulfate in pH 5.0 water, and distilled water were deoxygenated
16 by repeated vacuuming under constant stirring and purging with pure nitrogen or argon gas [21].
17 The degassed solutions were transferred via gas-tight Hamilton syringes into a tightly septum-
18 sealed and nitrogen-filled (or argon-filled) quartz UV-vis cuvette at 25 °C, while avoiding the
19 formation of pockets of gas above the solution. The final experimental conditions after mixing all
20 reagents were as follow: 0.2-1 μM of ferritin in 50 mM MOPS, 50 mM NaCl, pH 7.4, 11-30 μM
21 dissolved oxygen, and 140 to 500 μM of Fe^{2+} . The method used to calculate the iron oxidation
22 rates and the concentrations of remaining Fe(II) and O_2 in solution are described below in detail.
23 In brief, the initial concentration of dissolved oxygen is calculated based on the final $[\text{Fe}^{3+}]$ cations
24 produced during the oxidation reaction, assuming either 4:1 or 2:1 $\text{Fe}^{\text{II}}:\text{O}_2$ stoichiometry. The
25 concentration of remaining oxygen in solution at any point during the kinetic is calculated based
26 on the difference between the initial calculated concentration of oxygen and one of the two $\text{Fe}^{\text{II}}:\text{O}_2$
27 presumed oxidation ratios (i.e. 4 $[\text{Fe}^{3+}]$ for the 4:1 $\text{Fe}^{\text{II}}:\text{O}_2$ ratio and 2 $[\text{Fe}^{3+}]$ for the 2:1 $\text{Fe}^{\text{II}}:\text{O}_2$
28 ratio)". We note that a direct measurement of $[\text{O}_2]$ in solution is not a practical approach given the
29 low sensitivity, inaccuracy, and responsiveness of commercially available oxygen electrodes,
30 particularly in the most interesting area of our kinetic studies (i.e. under low O_2 concentration). To
31 overcome this limitation, we adopted a new approach based on the known $\text{Fe}^{\text{II}}:\text{O}_2$ oxidation
32 stoichiometry, as discussed above.
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50

51 **Results**

52 We believe that maintaining about 1 μM concentration of labile iron pool must involve a dynamic
53 equilibrium process, and a constant overall rate of exchange between iron uptake and iron release.
54
55
56
57

While iron release is a complex process that involves the proteolytic degradation of ferritin in lysosome,^{20, 21} and possibly alternative processes of iron mobilization, either by reduction^{22, 23} or direct chelation,²⁴ the mechanism of iron uptake, oxidation, and mineralization inside ferritin is relatively well understood. In cell-free conditions, iron oxidation kinetics can be easily monitored by light absorption of the oxidized iron(III)-protein complex at 305 nm⁸. However, due to the low molar absorptivity of the iron-protein complex, this assay becomes insensitive to the typical 1 μM cytosolic concentration of labile Fe^{2+} cations. One solution is to extrapolate and deduce the iron oxidation rates from kinetics conducted at much higher iron concentrations. Assuming the iron oxidation reactions obey Michaelis-Menten kinetics, the values of K_m and V_{max} for Fe^{2+} and O_2 can be easily determined.

K_m for iron

In our measurements of the K_m value for iron in ferritin, the working protein solutions were exposed to atmospheric oxygen for a prolonged period. Under these conditions, the concentrations of dissolved oxygen is around 250 μM at 25 °C, a value much higher than the reported K_m for O_2 ^{16, 17, 19, 25}. Furthermore, considering either 2:1 or 4:1 $\text{Fe}^{\text{II}}:\text{O}_2$ ratios, the $[\text{O}_2]$ is in large excess even at the highest $\text{Fe}(\text{II})$ concentration (i.e. 250 μM) used in our experiments. Figure 1 shows the kinetic traces of the optical densities at 305 nm, where the Fe^{III} -oxo(hydroxo)-protein complex absorbs, as a function of reaction time.




Figure 1. Kinetic curves for the catalytic oxidation of $\text{Fe}(\text{II})$ by recombinant human homopolymer apo-HuHF (left), recombinant human heteropolymer apo-HL (middle), and apo-HosF (right). The experimental conditions are: HuHF (0.5 μM), HL (0.5 μM), HosF (5 μM), in 100 mM MOPS buffer and 100 mM NaCl, pH 7.4 and 25 °C. The ratio of $\text{Fe}(\text{II})/\text{protein}$ is shown on each panel and is color coded.

In these experiments, and at all levels of iron additions, the reaction proceeded quickly in all three ferritin samples with > 90% of $\text{Fe}(\text{II})$ mineralized within less than one minute, even at the largest concentration of $\text{Fe}(\text{II})$ employed (i.e. 250 μM). The 10-fold higher concentration of HosF translates into approximately one order of magnitude slower rates of iron mineralization, compared

1
2
3 to HuHF and HL, and is consistent with the approximately ten times higher number of H-subunits
4 present in these proteins.
5

6
7 As previously observed⁸, iron oxidation in ferritin proceeds with an initial maximal rate, and slows
8 down as the concentration of iron(II) cations is depleted from the medium, in favor of the formation
9 of the mineralized iron(III) hydroxide core. A common approach for calculating the rate constants
10 of these enzymatic reactions, and ultimately K_m and V_{max} parameters, involves fitting the kinetic
11 traces using polynomial or exponential fits provided in various software packages, or simply by
12 choosing the Michaelis–Menten kinetic equation in data analysis software like OriginLab. The
13 problem with such approach is that the reaction kinetic is implicitly assumed to follow Michaelis–
14 Menten behavior. To properly analyze the kinetics of iron(II) oxidation in ferritin, and to
15 determine the applicability of Michaelis–Menten equation, we calculated the reaction rates by
16 numerical differentiation of the kinetic curves. This calculation is accomplished by analyzing the
17 changes in the concentration of oxidized iron (calculated by dividing the absorbance value at 305
18 nm by $3000 \text{ M}^{-1}\text{cm}^{-1}$, which is the average molar absorptivity constant of the Fe(III)-ferritin
19 complex) observed over a period of 1.00 sec. Initially, the concentration of iron(II) cations added
20 to each ferritin sample is known; however, we determined the exact concentration of oxidized iron
21 inside ferritin, at the end of the kinetic run, using Beer Lambert's law, and a molar absorptivity of
22 $3000 \text{ M}^{-1}\text{cm}^{-1}$ ⁸. This allowed us to calculate the concentration of remaining iron(II) cations in
23 solution during the progress of the oxidation reaction. Clearly, this numerical differentiation
24 approach leads to a strong amplification of the noise in the original data. To minimize this effect,
25 we averaged the initial and the final concentrations of oxidized iron, by subsequently analyzing 10
26 successive absorption measurements, separated by 0.25 sec time interval. This method of
27 calculation of reaction rates, allows their measurement at any point during the kinetic run as a
28 function of the remaining Fe(II) concentrations in solution. The results of the three ferritin samples
29 (HuHF, HL and HosF) tested in this study are shown in Figure 2, and reveal a completely different
30 behavior than the presumed Michaelis-Menten kinetics. An initial short and very rapid reaction
31 phase is observed immediately after the addition of Fe(II) to an oxygenated ferritin solution in
32 buffer, followed by a second much slower phase that continues until all Fe(II) ions present in
33 solution are fully depleted.
34
35
36
37
38
39
40
41
42

43 Table 1 shows that at the transition point between Phase I and Phase II, the iron loading of the
44 three proteins is estimated at 41-71 Fe^{III} /shell for HuHF, 41-56 Fe^{III} /shell for HL, and 8-11
45 Fe^{III} /shell for HosF, corresponding to 2.6 ± 0.4 , 2.2 ± 0.3 , and $3.2 \pm 0.6 \text{ Fe}^{\text{III}}$ /H-subunit for the
46 three proteins, respectively.
47
48

49 The gradual decrease in reaction rates observed in Phase II of Figure 2 is proportional to the
50 concentration of remaining Fe(II) cations in solution. As more Fe(II) ions are oxidized by the
51 protein, and the inorganic mineral core continues to grow, the second phase of the kinetic traces
52 had little effect on the reaction rates, irrespective of the amount of added iron (Fig. 2), or the size
53 of the inorganic iron core (Fig. 3). For instance, in HL ferritin (Fig. 2), the reaction rates at $30 \mu\text{M}$
54 of remaining Fe(II) are in the range $1.6 \pm 0.3 \mu\text{M/s}$ at iron loadings of 250, 200, 150, 125, and 75
55
56
57
58
59
60

1
2
3 μM . In HuHF and HosF, at 30 μM remaining Fe(II), these rates were $3.5 \pm 0.3 \mu\text{M/s}$ and $2.9 \pm$
4 $0.5 \mu\text{M/s}$, respectively. The lack of a second phase (Phase II) under low Fe(II) concentrations (i.e.
5 stoichiometric amounts of ~ 2 Fe(II)/di-iron center or 42 Fe(II)/shell for HuHF and HL, and 4
6 Fe(II)/shell for HosF, Fig. 2), suggest the absence of a displacement reaction of bound Fe(III)
7 cations by incoming Fe(II) cations.
8
9

10
11 To obtain more insights into the origin of the transition between Phases I and II, and to examine
12 the effect of an existing iron core within the ferritin cavity, two back-to-back injections of Fe(II)
13 solutions were made to the same HuHF and HL ferritin samples (Fig.3). The results showed a
14 dramatically shortened (in the case of HuHF) or almost inexistent first phase (in the case of HL),
15 suggesting that a different iron oxidation mechanism had taken, in the presence of an existing iron
16 core. Here again, the reaction rates exhibit a linear dependency on the amount of Fe(II) present in
17 solution.
18
19

20
21 **Table 1.** Analysis of the iron oxidation kinetics in HuHF, HL, and HosF shown in Figures 1 and 2. The experimental
22 conditions are provided in the caption of Figure 1.
23

HuHF					(Assuming 24 H)
Fe ^{II} /shell added	Experimentally calculated [Fe ^{II}]	Remaining [Fe ^{II}] in solution	[Fe ^{III}] at the transition point (between Phase I and Phase II)	Fe ^{III} /shell at transition	Fe ^{III} /H-chain at transition
500	249 ± 13	214 ± 11	36 ± 2	71 ± 5	2.9 ± 0.2
400	211 ± 2	172 ± 3	39 ± 1	79 ± 2	3.3 ± 0.1
300	155 ± 2	122 ± 4	33 ± 2	66 ± 4	2.7 ± 0.1
200	101 ± 3	77 ± 1	24 ± 3	48 ± 5	2.0 ± 0.1
100	50 ± 2	30 ± 1	20 ± 1	41 ± 1	2.5 ± 0.1
	--			Average	2.6 ± 0.4
HL	--				(Assuming 21 H)
500	284 ± 37	256 ± 35	28 ± 2	56 ± 3	2.6 ± 0.1
400	195 ± 3	171 ± 2	24 ± 2	48 ± 3	2.2 ± 0.1
300	148 ± 5	120 ± 1	27 ± 5	55 ± 10	2.5 ± 0.4
250	122 ± 10	99 ± 3	23 ± 6	47 ± 13	2.1 ± 0.5
200	105 ± 1	83 ± 1	22 ± 1	43 ± 3	2.0 ± 0.2
150	73 ± 1	52 ± 3	21 ± 3	41 ± 6	1.9 ± 0.3
	--			Average	2.2 ± 0.3
HosF	--				(Assuming 3.3 H)
40	212 ± 6	157 ± 6	55 ± 1	11 ± 1	3.3 ± 0.03
32	172 ± 4	115 ± 1	57 ± 3	11 ± 1	3.3 ± 0.18
24	126 ± 2	77 ± 1	49 ± 1	10 ± 1	3.0 ± 0.003
16	94 ± 1	55 ± 4	38 ± 3	8 ± 1	2.4 ± 0.21
				Average	3.0 ± 0.5




Figure 2. Plots of the reaction rates vs. the concentration of remaining Fe(II) cations in solution. Top left: typical plot for HL ferritin with 400 Fe^{II}/shell; Top right and Bottom: overlay of several reaction rate plots at different ratios of Fe^{II}/shell for HuHF, Hl, and HosF, as indicated next to each curve. The experimental conditions are those of Figure 1.




Figure 3. Top: Representative kinetic curves for two successive injections of the same amount of Fe^{II}/shell, as indicated on each plot, in the presence of HuHF (left) or HL ferritin (right). Bottom: Plots of the reaction rates vs. the concentration of remaining iron(II) cations in solution following the second addition of Fe(II) to the same protein sample (as shown on each panel), HuHF (left), and HL (right). The experimental conditions are the same as in Figure 1.

Fe(II) oxidation under low concentration of O₂ - K_m for oxygen

The iron oxidation kinetics described above are based on the assumption that the reaction rate does not depend on the concentration of oxygen in solution. However, because of the large discrepancy in the literature values of K_m for O₂ (i.e. 6 - 140 μM)^{18,25}, we sought to investigate the effect of O₂ on the reaction rates. One of the reasons for this wide range of K_m values stems from the difficulty in maintaining and accurately monitoring the concentration of soluble oxygen in solution. In an earlier study, we used a sensitive oxygen microelectrode to measure and monitor in real time the concentration of oxygen²³ in solution, and its effect on the rates of the reductive release of iron from ferritin. Unfortunately, this method becomes unreliable when monitoring very low concentrations of oxygen, particularly if these concentrations are rapidly changing.

Nonetheless, to overcome this problem, we conducted kinetic experiments in a tightly sealed UV-vis cuvette, using partially deoxygenated ferritin solutions containing $[O_2]$ ranging between 20 and 40 μM . With this set up, the possibility of oxygen exchange with air is minimized²³, particularly considering the short reaction time (< 2 min), and the large excess of $[Fe(II)]$ employed. The reaction kinetics in the homogeneous solution progressed until all dissolved oxygen is depleted, and large excess of $Fe(II)$ remained in solution. Under these conditions, the ratio of $Fe(II)$ to O_2 is shown to occur at 4 $Fe(II)$ per 1 O_2 , and therefore it is possible to calculate both the concentration of oxygen at any time of the reaction as a one-fourth of the total concentration of $Fe(II)$ in solution. A typical absorbance vs. time kinetic trace for these types of reactions are shown in Figure 4a-c, top.




Figure 4. (Top row) Iron oxidation kinetics at 305 nm under low oxygen concentration using (a) HosF (1.0 μM), $Fe(II)$ (200 μM), and approximately 19 μM O_2 ; (b) HL (0.2 μM), $Fe(II)$ (140 μM), and approximately 11 μM of O_2 ; (c) HuHF (0.4 μM), $Fe(II)$ (500 μM), and approximately 30 μM O_2 . (Middle row): Dependence of the reaction rates on the concentration of remaining O_2 , calculated assuming 4:1 $Fe^{II}:O_2$ stoichiometry. Red and green curves are kinetics before and after correction (more details in the text). (Lower row) Michaelis-Menten fit to the corrected kinetic curves for the calculation of K_m and V_{max} parameters.

Calculation of K_{m,O_2} and V_{max} parameters for HuHF, HL, and HosF

Following the same methodology described above, the kinetic curves under low oxygen concentration (Fig. 4) were numerically differentiated for our three proteins (HuHF, HL, and HosF), to determine how the reaction rates change as a function of remaining $[O_2]$ in solution. To minimize the reaction noise, the reaction rates were calculated using changes in Fe(III) concentrations (calculated from optical densities as an average of four measurements within 1 s time interval) over 2.5 s time period, and then converted into $\mu\text{M/s}$. Once oxygen is fully depleted from the solution, the concentration of oxidized iron (i.e. $75.5 \mu\text{M}$ in the case of HosF) was calculated from the absorbance value of 0.226 at 60 seconds (Fig. 4 a, top). Assuming a 4:1 $\text{Fe}^{\text{II}}:\text{O}_2$ ratio, the initial solution concentration of oxygen (i.e. $75.5/4=18.9 \mu\text{M}$) is used to experimentally determine the concentration of dissolved oxygen at any time during the oxidation reaction. The reaction rates are then plotted as a function of remaining $[O_2]$ in solution, and are shown in Figure 4 (middle and bottom panels). However, because of the presence of 2 phases (Fig. 2), only the second slow phase of the reaction (dotted box of Fig. 4, top) is further analyzed. To make sure that only Phase II of the reaction is selected, the kinetic data were analyzed at a concentration of Fe(III) that is twice or more than that observed at the end of Phase I (i.e. $\geq 2 \times 11 \text{ Fe}^{\text{III}}/\text{shell}$ for HosF, Table 1). This corresponds to an $[\text{Fe}^{3+}]$ spanning between 22 and $75.5 \mu\text{M}$, and an $[O_2]$ concentration of 13 - 0 μM , respectively (assuming 4:1 $\text{Fe}^{\text{II}}:\text{O}_2$ stoichiometry). During that time frame (i.e. 13 – 60 s, Fig. 4a), the concentration of $[\text{Fe}(\text{II})]_{\text{remaining}}$ in solution ranges between 177 μM and 124.7 μM , and can be accounted for by multiplying the calculated reaction rates (Fig. 4, middle, red uncorrected curve) by the ratio of $[\text{Fe}(\text{II})]_{\text{initial}} (177 \mu\text{M}) / [\text{Fe}(\text{II})]_{\text{current}}$. This adjustment is possible because during Phase II, the reaction rates depend linearly on $[\text{Fe}(\text{II})]$. The adjusted kinetic traces (Fig. 4, middle panels, green corrected curves) are remarkably similar to classical Michaelis-Menten kinetics, the fit of which yielded the following parameters: $V_{\text{max}} = 1.94 \pm 0.01 \mu\text{M/s}$ and $K_{m,O_2} = 0.77 \pm 0.03 \mu\text{M}$.

Although the total concentration of available $[\text{Fe}(\text{II})]$ in solution is in large excess compared to the concentration of HosF (i.e. 22-75 $\text{Fe}^{\text{III}}/\text{shell}$), and although the stoichiometry of $\text{Fe}^{\text{II}}:\text{O}_2$ reported in the literature is close to 4:1, it behooved us to repeat the above analysis assuming an oxidation ratio of $2\text{Fe}^{\text{II}}:1\text{O}_2$. Indeed, the $\text{Fe}^{\text{II}}:\text{O}_2$ ratio may depend on the overall experimental conditions including total iron concentration present in solution, pH, iron chelators, buffer and protein type²⁷⁻³⁰. Because the concentration of oxygen in solution is calculated based on the $\text{Fe}^{\text{II}}:\text{O}_2$ stoichiometry, the value of K_{mO_2} will also depend on that stoichiometry. The most reliable and straightforward way to address the uncertainty in the value of K_{mO_2} is to assume that the actual stoichiometry of the reaction is anywhere between 4:1 and 2:1. A re-analysis of the kinetics assuming a $2\text{Fe}^{\text{II}}:1\text{O}_2$ oxidation stoichiometry showed an analogous Michaelis-Menten behavior; however, the value of K_{mO_2} was twice as large (i.e. $K_{mO_2} = 1.54 \pm 0.06 \mu\text{M}$). Based on these calculations, it can be concluded that the average K_{mO_2} for the iron oxidation reaction in HosF is $1.15 \pm 0.54 \mu\text{M}$ for $2 < \text{Fe}^{\text{II}}:\text{O}_2 < 4$.

1
2
3 A similar analysis was performed with recombinant heteropolymer HL ferritin (Fig. 4) under
4 deficient $[O_2]$, $[HL] = 0.2 \mu M$, and an initial total $[Fe(II)] = 140 \mu M$. The concentration of oxidized
5 iron at the end of the experiment in Fig. 4 was calculated to be $45.5 \mu M$, corresponding to an initial
6 O_2 concentration of $11.3 \mu M$, assuming a 4:1 $Fe^{II}:O_2$ ratio. In the case of HL ferritin (Fig. 2),
7 Phase II of the reaction occurs around 48 $Fe^{III}/shell$, which when adjusted for the $0.2 \mu M$
8 concentration of HL and the $2Fe^{II}:1O_2$, corresponds to a mineralized iron(III) concentration of
9 about $20 \mu M$. The remaining $[Fe(II)]$ (i.e. $120 \mu M$) correlates with a reaction time of about 35
10 seconds (Fig. 4b, top). The plot of the reaction rates as a function of $[O_2]$ in the solution was
11 calculated as previously described, and is shown in Fig. 4b, middle. Here again, a fit using the
12 Michaelis-Menten equation yielded a V_{max} value of $0.52 \pm 0.03 \mu M/s$ and a K_{mO_2} of 0.46 ± 0.02
13 μM , with an average K_{mO_2} value for HL of $0.69 \pm 0.32 \mu M$ for $2 < Fe^{II}:O_2 < 4$. Analogous fitting
14 of the HuHF experiments yielded a V_{max} value of $1.61 \pm 0.03 \mu M/s$ and a K_{mO_2} of 1.30 ± 0.08
15 μM assuming a $4Fe^{II}:1O_2$, and an average K_{mO_2} value for HuHF of $1.95 \pm 0.92 \mu M$ for $2 < Fe^{II}:O_2$
16 < 4 .
17
18
19
20
21
22

23 Discussion

24
25 Under our experimental conditions, our data show that the rates of iron oxidation as a function of
26 $Fe(II)$ concentration, in the three types of ferritin, HuHF, HL, and HosF examined here, do not
27 follow Michaelis-Menten kinetics. The reaction rates were substantially higher at high $Fe(II)$
28 concentrations, suggesting pseudo-first order kinetics. However, the existence of two distinct iron
29 oxidation phases indicates that the reaction mechanism is much more complex. The rapid decrease
30 of reaction rates during Phase I could not be attributed to a decrease in the concentration of $Fe(II)$,
31 since it does not dramatically change during that phase, particularly under high $Fe(II)$
32 concentrations (i.e. 400 or 500 $Fe^{II}/shell$ additions in HuHF or HL ferritin, and 40 $Fe^{II}/shell$ in
33 HosF). Instead, we believe that the rapid decrease of reaction rates during Phase I is caused by the
34 inhibition of the catalytic activity of the ferroxidase centers when the oxidized $Fe(III)$ cations
35 remain bound to these centers. Table 1 shows that the Fe^{III}/H -subunit inhibition stoichiometry is
36 between 2 and 2.5 for HuHF and HL, and around 3.5 for HosF. The oxidized $Fe(III)$ cations at the
37 ferroxidase centers of the H-subunit must either dissociate (i.e. detach from the center), or the
38 ferroxidase centers act as a cofactor for the oxidation of incoming $Fe(II)$ ions^{4, 13-15}. In either
39 case, more $Fe(II)$ ions would get oxidized and ultimately join the growing inorganic ferrihydrite
40 core inside the ferritin cavity, in analogy to what has been previously proposed for bacterioferritins
41⁴. The dissociation of $Fe(III)$ cations represents the rate-limiting step of the reaction that can be
42 accelerated by other metal cations such as $Fe(II)$, or by chelate ligands capable of binding $Fe(III)$
43 ions. This suggested reaction mechanism is compatible with several observations including, (1)
44 the pseudo-first order kinetics during Phase II, (2) the absence (or significantly depressed) Phase
45 I following the second addition of $Fe(II)$ cations, (3) the previously reported enhanced effect of
46 phosphate, a known $Fe(III)$ chelate ligand³¹ on the rate of $Fe(II)$ oxidation³².
47
48
49
50
51
52
53

54 An earlier study using an *E.coli* ferritin (i.e. bacterioferritin) found that the oxidized di- $Fe(III)$ -
55 protein complex that forms at the ferroxidase center of the protein is stable, and acts as a catalytic
56
57

1
2
3 site for O₂ reduction, but that in *P. aeruginosa* bacterioferritin, the di-iron complex is a transit site
4 for the transfer of Fe(III) ions into the central cavity. In *E. coli*, and with the assistance of a tyrosyl
5 radical, two additional Fe(II) sites provide the necessary electrons to reduce the di-Fe(III)-protein
6 complex to a di-Fe(II)-complex. The hydrolysis of the oxidized iron at the two additional iron
7 sites contributes to the formation of the ferritin mineral core. The reduced di-Fe(II)-protein
8 complex further reacts with O₂ or H₂O₂, and the cycle is repeated until all of the Fe(II) ions are
9 oxidized⁴.
10
11
12

13 The notion that the rate limiting step during iron oxidation in ferritin might involve an exchange
14 reaction between Fe(III) ions at the ferroxidase centers (or at nearby oxidation sites) by incoming
15 Fe(II), is at variance with the previously proposed surface catalytic activity of the inorganic iron
16 core¹⁷. To the best of our knowledge, ferroxidase activity has never been observed on isolated
17 iron(III) cores. Furthermore, comparison of Fe(II) oxidation rates catalyzed by HL ferritin, using
18 initial concentrations of 150 to 500 Fe^{II}/shell, shows very similar reaction rates (i.e. ~ 2.5 μM/s)
19 at 50 μM [Fe(II)] remaining in solution (Fig. 2 left, Fig. 3, right), suggesting that the protein
20 ferroxidase activity is independent on the size of the iron(III) hydroxide core. Similar results were
21 observed with HuHF and HosF (Figs. 2, 3).
22
23
24
25

26 The results of Figures 1 and 2 show that the Fe(III) turnover numbers for HuHF, HL, and HosF
27 are 40 s⁻¹, 24 s⁻¹, and 7 s⁻¹, corresponding to 1.7 s⁻¹, 1.2 s⁻¹ and 2.1 s⁻¹ per H-subunit, assuming 24,
28 21, and 3.3 H subunits for HuHF, HL, and HosF, respectively. More specifically, the concentration
29 of Fe(II) at the beginning of Phase I is approximately 200 μM for the three experiments involving
30 0.5 μM HuHF or HL (400 Fe^{II}/shell), and 5 μM HosF (40 Fe^{II}/shell). These conditions correspond
31 to about 12 Fe^{II}/H-subunit with initial rates of the iron oxidation reactions of approximately 20
32 μM/s, 12 μM/s, and 35 μM/s, for HuHF, HL, and HosF, respectively (Fig. 2). When a second
33 injection of Fe(II) is made to the same protein sample (i.e. another 400 Fe^{II}/shell, for HuHF or HL)
34 immediately after the 1st Fe(II) oxidation reaction is completed, turnover numbers of 1.0 s⁻¹ for
35 HuHF, and 0.8 s⁻¹ for HL are observed (Fig. 3).
36
37
38
39

40 In contrast to Phase I, the reaction rates during Phase II slow down dramatically, presumably
41 because most of the protein's ferroxidase centers are already occupied by Fe(III) cations, and are
42 thus "inactive", until Fe(III) ions are displaced (or exchanged) by new incoming Fe(II) ions. Under
43 oxygen deficient conditions (i.e. low μM range), direct measurements of the concentration of
44 dissolved oxygen in solution using commercial oxygen microelectrodes is challenging, due to their
45 relatively slow response and poor precision. However, the reaction rates as a function of [O₂] can
46 be easily calculated, although indirectly, by monitoring the concentration of oxidized iron inside
47 ferritin at 305 nm. Figure 4 shows that the reaction rates show a Michaelis-Menten type kinetics
48 at [O₂] less than 10 μM, with the sharpest change occurring at less than 2 μM [O₂]. Because the
49 Fe^{II}:O₂ stoichiometry can vary between 2:1 and 4:1 (depending on the experimental conditions and
50 the ratio of the Fe^{II}/shell added), the calculated [O₂] concentration will consequently vary, resulting
51 in as much as doubling of the experimentally calculated K_{m,O2} values. Assuming 4Fe^{II}:1O₂
52
53
54
55
56
57
58
59
60

1
2
3 stoichiometry, the experimentally calculated K_{m,O_2} values for HuHF, HL, and HosF are 1.95 ± 0.92
4 μM , 0.69 ± 0.32 μM , and 1.15 ± 0.54 μM , respectively. These values are m

6
7 uch lower than the lowest reported K_{m,O_2} values in the literature (i.e. 6 ± 2 μM for HuHF, 60 ± 12
8 μM for HL, and 140 ± 30 μM for HosF)¹⁸. Considering the similarity in the structure-function
9 relationships of the three types of ferritin tested here, we believe that these relatively similar K_{m,O_2}
10 values are much more reasonable than the reported literature values.
11

12
13 Clearly, our experimental K_{m,O_2} values have important physiological implications. For instance,
14 the concentration of oxygen in living cells is in the range of 1.3-2.5 kPa³³ (i.e. ~ 18 -34 μM), but
15 much lower (i.e. ~ 7 μM) inside mitochondria. Our measured K_{m,O_2} values are much lower than
16 these cellular O_2 concentrations, suggesting that neither iron mineralization inside ferritin, nor the
17 iron oxidation rates, should be affected by variations in cellular oxygenation level, including under
18 conditions of severe hypoxia.
19
20

21 **Conclusions**

22
23 The kinetics of iron(II) oxidation catalyzed by three different types of ferritins (HuHF, HL, and
24 HosF) were studied in detail, using numerical differentiation of the change in optical density over
25 time. In the presence of excess $[O_2]$, the iron oxidation rates were continuously monitored until
26 all iron(II) cations are completely depleted from solution. The observed iron oxidation reactions
27 proceeded through two distinct phases (Phase I and Phase II), neither of which obeyed Michaelis-
28 Menten kinetics. The initial and short Phase I reaction showed a rapid drop in reaction rates until
29 approximately 2 Fe(II) ions have been oxidized per H-subunit. The subsequent and much slower
30 Phase II reaction exhibited a significant decrease in reaction rates, proportionally to the
31 concentration of remaining iron(II) cations in solution, but independently of the size of the iron
32 core. A kinetic model is proposed in which the inhibition of the protein's activity is caused by
33 bound iron(III) cations at the ferroxidase center, with the rate limiting step corresponding to an
34 exchange or a displacement reaction between incoming iron(II) cations and bound Fe(III) cations.
35 On the other hand, and in the presence of limited $[O_2]$, the kinetics of iron oxidation proceeded
36 until all dissolved oxygen is fully depleted from solution, and exhibited Michaelis-Menten
37 behavior, with K_{m,O_2} values in the low μM range (i.e. $\sim 1 - 2$ μM). The experimentally measured
38 values of K_{m,O_2} are considerably lower than typical cellular oxygen concentrations in normoxic
39 and hypoxic cells, indicating that the process of iron mineralization in ferritin should not be
40 affected by hypoxic events.
41
42
43
44
45
46
47

48 **Acknowledgements**

49
50 This work is supported by the National Institute of Health, Award Numbers R15GM104879
51 (F.B.A.), and NSF award CHE 1150768 (A.M.).
52
53
54
55
56
57
58
59
60

References

1. B. Silva and P. Faustino, An overview of molecular basis of iron metabolism regulation and the associated pathologies, *Biochimica et Biophysica Acta (BBA)-Molecular Basis of Disease*, 2015, **1852**, 1347-1359.
2. F. Bou-Abdallah, The iron redox and hydrolysis chemistry of the ferritins, *Biochimica et Biophysica Acta (BBA)-General Subjects*, 2010, **1800**, 719-731.
3. I. Bertini, D. Lalli, S. Mangani, C. Pozzi, C. Rosa, E. C. Theil and P. Turano, Structural insights into the ferroxidase site of ferritins from higher eukaryotes, *J. Am. Chem. Soc.*, 2012, **134**, 6169-6176.
4. J. M. Bradley, N. E. Le Brun and G. R. Moore, Ferritins: furnishing proteins with iron, *Journal of Biological Inorganic Chemistry*, 2016, **21**, 13-28.
5. P. M. Harrison and P. Arosio, The ferritins: molecular properties, iron storage function and cellular regulation, *Biochimica et Biophysica Acta (BBA)-Bioenergetics*, 1996, **1275**, 161-203.
6. N. D. Chasteen, Ferritin. Uptake, storage, and release of iron, *Metal ions in biological systems*, 1998, **35**, 479.
7. P. Arosio, R. Ingrassia and P. Cavadini, Ferritins: a family of molecules for iron storage, antioxidation and more, *Biochimica et Biophysica Acta (BBA)-General Subjects*, 2009, **1790**, 589-599.
8. M. Mehlenbacher, M. Poli, P. Arosio, P. Santambrogio, S. Levi, N. D. Chasteen and F. Bou-Abdallah, Iron oxidation and core formation in recombinant heteropolymeric human ferritins, *Biochemistry*, 2017, **56**, 3900-3912.
9. B. P. Esposito, S. Epsztejn, W. Breuer and Z. I. Cabantchik, A review of fluorescence methods for assessing labile iron in cells and biological fluids, *Analytical Biochemistry*, 2002, **304**, 1-18.
10. Z. I. Cabantchik, Labile iron in cells and body fluids: physiology, pathology, and pharmacology, *Frontiers in pharmacology*, 2014, **5**, 45.
11. F. Bou-Abdallah and T. R. Terpstra, The thermodynamic and binding properties of the transferrins as studied by isothermal titration calorimetry, *Biochimica et Biophysica Acta (BBA)-General Subjects*, 2012, **1820**, 318-325.
12. J. Williams and K. Moreton, The distribution of iron between the metal-binding sites of transferrin human serum, *Biochemical Journal*, 1980, **185**, 483-488.
13. W. R. Hagen, P.-L. Hagedoorn and K. H. Ebrahimi, The workings of ferritin: a crossroad of opinions, *Metallomics*, 2017, **9**, 595-605.
14. J. M. Bradley, G. R. Moore and N. E. Le Brun, Mechanisms of iron mineralization in ferritins: one size does not fit all, *Journal of Biological Inorganic Chemistry*, 2014, **19**, 775-785.
15. K. H. Ebrahimi, P.-L. Hagedoorn and W. R. Hagen, Phosphate accelerates displacement of Fe (III) by Fe (II) in the ferroxidase center of *Pyrococcus furiosus* ferritin, *FEBS letters*, 2013, **587**, 220-225.
16. K. Orino, S. Harada, M. Natsuhori, K. Takehara and K. Watanabe, Kinetic analysis of bovine spleen apoferritin and recombinant H and L chain homopolymers: Iron uptake suggests early stage H chain ferroxidase activity and second stage L chain cooperation, *Biometals*, 2004, **17**, 129-134.
17. K. Orino, S. Kamura, M. Natsuhori, S. Yamamoto and K. Watanabe, Two pathways of iron uptake in bovine spleen apoferritin dependent on iron concentration, *Biometals*, 2002, **15**, 59-63.
18. S. Sun, P. Arosio, S. Levi and N. D. Chasteen, Ferroxidase kinetics of human liver apoferritin, recombinant H-chain apoferritin, and site-directed mutants, *Biochemistry*, 1993, **32**, 9362-9369.
19. I. G. Macara, T. G. Hoy and P. M. Harrison, The formation of ferritin from apoferritin. Catalytic action of apoferritin, *Biochem. J.*, 1973, **135**, 343-348.
20. M. C. Linder, Mobilization of stored iron in mammals: a review, *Nutrients*, 2013, **5**, 4022-4050.

- 1
 - 2
 - 3
 - 4
 - 5
 - 6
 - 7
 - 8
 - 9
 - 10
 - 11
 - 12
 - 13
 - 14
 - 15
 - 16
 - 17
 - 18
 - 19
 - 20
 - 21
 - 22
 - 23
 - 24
 - 25
 - 26
 - 27
 - 28
 - 29
 - 30
 - 31
 - 32
 - 33
 - 34
 - 35
 - 36
 - 37
 - 38
 - 39
 - 40
 - 41
 - 42
 - 43
 - 44
 - 45
 - 46
 - 47
 - 48
 - 49
 - 50
 - 51
 - 52
 - 53
 - 54
 - 55
 - 56
 - 57
 - 58
 - 59
 - 60
21. T. Z. Kidane, E. Sauble and M. C. Linder, Release of iron from ferritin requires lysosomal activity, *American Journal of Physiology-Cell Physiology*, 2006, **291**, C445-C455.
22. L. E. Johnson, T. Wilkinson, P. Arosio, A. Melman and F. Bou-Abdallah, Effect of chaotropes on the kinetics of iron release from ferritin by flavin nucleotides, *Biochimica et Biophysica Acta (BBA) - General Subjects*, 2017.
23. G. Melman, F. Bou-Abdallah, E. Vane, P. Maura, P. Arosio and A. Melman, Iron release from ferritin by flavin nucleotides, *Biochimica et Biophysica Acta (BBA) - General Subjects* 2013, **1830**, 4669-4674.
24. P. Sanchez, N. Galvez, E. Colacio, E. Minones and J. Dominguez-Vera, Catechol releases iron (III) from ferritin by direct chelation without iron (II) production, *Dalton Transactions*, 2005, 811-813.
25. S. Sun and N. D. Chasteen, Ferroxidase kinetics of horse spleen apoferritin, *Journal of Biological Chemistry*, 1992, **267**, 25160-25166.
26. G. Zhao, M. Su and N. D. Chasteen, Mu-1,2-Peroxo Diferric Complex Formation in Horse Spleen Ferritin. A Mixed H/L-Subunit Heteropolymer, *Journal of molecular biology*, 2005, **352**, 467-477.
27. G. Zhao, F. Bou-Abdallah, X. Yang, P. Arosio and N. D. Chasteen, Is hydrogen peroxide produced during iron (II) oxidation in mammalian apoferritins?, *Biochemistry*, 2001, **40**, 10832-10838.
28. M. E. Van Eden and S. D. Aust, The consequences of hydroxyl radical formation on the stoichiometry and kinetics of ferrous iron oxidation by human apoferritin, *Free Radical Biology and Medicine*, 2001, **31**, 1007-1017.
29. G. Zhao, F. Bou-Abdallah, P. Arosio, S. Levi, C. Janus-Chandler and N. D. Chasteen, Multiple pathways for mineral core formation in mammalian apoferritin. The role of hydrogen peroxide, *Biochemistry*, 2003, **42**, 3142-3150.
30. K. D. Welch, T. Z. Davis and S. D. Aust, Iron autoxidation and free radical generation: effects of buffers, ligands, and chelators, *Archives of Biochemistry and Biophysics*, 2002, **397**, 360-369.
31. J. Hegenauer, P. Saltman and G. Nace, Iron (III)-phosphoprotein chelates: stoichiometric equilibrium constant for interaction of iron (III) and phosphorylserine residues of phosvitin and casein, *Biochemistry*, 1979, **18**, 3865-3879.
32. C. Cutler, A. Bravo, A. D. Ray and R. K. Watt, Iron loading into ferritin can be stimulated or inhibited by the presence of cations and anions: A specific role for phosphate, *Journal of inorganic biochemistry*, 2005, **99**, 2270-2275.
33. J. Gleadle and P. J. Ratcliffe, in *Encyclopedia of life sciences*, John Wiley & Sons, Ltd, Chichester, 2001.