


 Cite this: *RSC Adv.*, 2020, 10, 43103

In silico identification of SARS-CoV-2 spike (S) protein–ACE2 complex inhibitors from eight *Tecoma* species and cultivars analyzed by LC-MS†

 Seham S. El Hawary,^a Amira R. Khattab,^b Hanan S. Marzouk,^c Amira S. El Senousy,^a Mariam G. A. Alex,^c Omar M. Aly,^d Mohamed Teleb^e and Usama Ramadan Abdelmohsen^{b,fg}

Coronavirus (CoV) is a positive RNA genome virus causing a global panic nowadays. *Tecoma* is a medicinally-valuable genus in the Bignoniaceae family, with some of its species exhibiting anti-HIV activity. This encouraged us to conduct an *in silico* exploration of some phytochemicals in *Tecoma* species cultivated in Egypt, namely *Tecoma capensis* and its four varieties *i.e.* yellow, harmony, pink and red, *T. grandiflora* Loisel., *T. radicans* L., and one hybrid *i.e.* *Tecoma* × *smithii* W. Watson. LC/MS-based metabolite profiling of the studied *Tecoma* plants resulted in the dereplication of 12 compounds (1–12) belonging to different phytochemical classes *viz.* alkaloids, iridoids, flavonoids and fatty acid esters. The *in silico* inhibitory action of these compounds against SARS-CoV-2 spike protein C-terminal domain in complex with human ACE2 was assessed *via* molecular docking. Succinic acid decyl-3-oxobut-2-yl ester (10), a fatty acid ester, possessed the best binding affinity (−6.77 kcal mol^{−1}), as compared to hesperidin (13) (−7.10 kcal mol^{−1}).

 Received 22nd October 2020
 Accepted 23rd November 2020

DOI: 10.1039/d0ra08997d

rsc.li/rsc-advances

1. Introduction

Coronaviruses (CoVs) are positive RNA genome viruses, belonging to the Coronaviridae family of the Nidovirales order, which is divided into four genera (A, B, C and D). SARS-CoV-2 belongs to the B genus. CoV possess four structural proteins: spike protein, envelope protein, membrane protein, and nucleocapsid protein.¹ Spike protein promotes host attachment and viral cell membrane fusion during virus infection.² Potential anti-coronavirus treatments can be divided into two main categories, one operating on the human immune system or human cells, and the other on the coronavirus itself.³ Viruses

often bind to receptor proteins on the surface of cells in order to enter human cells, for example, the SARS virus links with human angiotensin-converting enzyme 2 (hACE2) receptor.⁴ Protein–protein docking showed that SARS-CoV-2 spike proteins have a strong affinity for hACE2.⁵ Through virtual screening many compounds could be identified as inhibitors of hACE2, however these potential hACE2 inhibitors might not be beneficial for the management of SARS-CoV-2 infection due to the protective role of hACE2 in lung injury. Hesperidin was the only compound reported till now that could target the binding interface between spike protein and hACE2. It was reported that hesperidin can lie on the middle shallow pit of the surface of the receptor-binding domain (RBD) of spike protein.⁶ Shang J. *et al.*, determined the crystal structure of RBD of the spike protein of SARS-CoV-2 (engineered to facilitate crystallization) in complex with hACE2. In comparison with the SARS-CoV RBD, an hACE2-binding ridge in SARS-CoV-2 RBD has a more compact conformation; moreover, several residues changes in the SARS-CoV-2 RBD stabilize two virus-binding hotspots at the RBD–hACE2 interface. These structural features of SARS-CoV-2 RBD increase its hACE2 binding affinity.⁷

Tecoma genus is one of the medicinally-valuable members in Bignoniaceae family, embracing fourteen species of shrubs and small trees, of which twelve are native to America and two to Africa. The leaves were traditionally used by people in Latin America for diabetes management.⁸ Later on, several *Tecoma* species were reported to possess a wide variety of pharmacological actions *viz.* anti-inflammatory,

^aPharmacognosy Department, Faculty of Pharmacy, Cairo University, Egypt

^bPharmacognosy Department, College of Pharmacy, Arab Academy for Science, Technology and Maritime Transport, Alexandria 1029, Egypt. E-mail: Dr_amira_khattab@aast.edu

^cPharmacognosy Department, Faculty of Pharmacy, Pharos University in Alexandria, Egypt

^dMedicinal Chemistry Department, Faculty of Pharmacy, Minia University, Minia, 61519, Egypt

^eDepartment of Pharmaceutical Chemistry, Faculty of Pharmacy, Alexandria University, Alexandria, 21521, Egypt

^fPharmacognosy Department, Faculty of Pharmacy, Deraya University, Minia 61111, Egypt. E-mail: usama.ramadan@mu.edu.eg

^gPharmacognosy Department, Faculty of Pharmacy, Minia University, Minia, 61519, Egypt

† Electronic supplementary information (ESI) available. See DOI: 10.1039/d0ra08997d



performed using a BEH C18 column (2.1 × 100 mm, 1.7 mm particle size) and a guard column (2.1 × 5 mm, 1.7 mm particle size) using the method previously described in ref. 19. Fig. S1† depicts the total ion chromatograms of the eight studied plants. MZmine 2.12 was used for processing the obtained raw data by the negative ionization mode. The processed data set was then subjected to molecular formula prediction and peak identification *via* dereplication using online METLIN²⁵ and Dictionary of Natural Products (DNP)²⁶ databases.

2.3. Docking studies

2.3.1. Docking of the target phytochemicals. Molecular docking study was carried out using Molecular Operating Environment (MOE) software package version 2016.10, Chemical Computing Group, Montreal, Canada. The crystal structure of SARS-CoV-2 spike (S) protein C-terminal domain (SARS-CoV-2-CTD) in complex with human ACE2 (hACE2) was obtained from the protein data bank (PDB ID 6LZG).²⁷ Unwanted residues, ligands and solvents were eliminated, then the complex was prepared employing the default “Structure preparation” module settings. ‘Site Finder’ feature of MOE 2016.10 was employed in search for the receptor site in the SARS-CoV-2-CTD binding interface. MDB database file of the target phytochemicals was subjected to default energy minimization and geometry optimization. Triangular matcher algorithm was applied to set the ligand placement. The default scoring function was alpha HB which generated the top 5 non-redundant poses of the lowest binding energy conformers of the tested phytoligands. Docking was conducted with induced fitting protocol to record the best possible molecular interactions. Results were listed based on the *S*-scores with RMSD value < 2 Å. Graphical representations of the phytoligands interactions were then generated and inspected.

3. Results and discussion

3.1. LC-HRESIMS based metabolic profiling

Twelve phytochemicals belonging to different phytochemical classes *viz.* alkaloids, iridoids, flavonoids, and fatty acid esters, have been identified by dereplication of the obtained LC-HRESIMS derived metabolite profiles of the eight *Tecoma* species and cultivars (Table 1). These metabolites have been annotated using databases (METLIN and DNP databases). From these databases, the molecular ion peaks appeared at *m/z* 325.1835, 349.1141, 327.2163 and 311.1672, with their predicted molecular formulas of C₂₁H₂₆O₃, C₁₉H₂₆O₆, C₁₈H₃₂O₅ and C₂₀H₄₀O₂ were dereplicated to be corresponding to four fatty acid esters *viz.* octanoic acid, 4-benzyloxyphenyl ester (**1**), fumaric acid, 3,4-dimethoxyphenyl heptyl ester (**2**), succinic acid, decyl 3-oxobut-2-yl ester (**10**) and valeric acid, pentadecyl ester (**12**), respectively.

Besides, four alkaloids were dereplicated; boschniakine (**3**) with its molecular ion peak detected at *m/z* 160.5217, and a predicted molecular formula of C₁₀H₁₁NO, which was formerly identified in *T. stans*.^{28,29} Besides, actinidine (**5**), skytanthine (**6**), and tecomanine (**7**) were identified based on their molecular ion peaks appearing at *m/z* 147.9783, 166.3757 and 178.0705 and possessing predicted molecular formulas of C₁₀H₁₃N, C₁₁H₂₁N and C₁₁H₁₇NO which were previously isolated from several *Tecoma* spp. *i.e.* *T. stans* and *T. arequipensis*.^{28,30} Three iridoid glucosides were tentatively identified as being 7-*O*-(*p*-OH)cinnamoyltecomoside (**8**), 7-*O*-(*p*-MeO)cinnamoyltecomoside (**9**) and 7-*O*-(*p*-OH)benzoyl tecomoside (**11**) based on their respective molecular ion peaks at *m/z* 521.1938, 535.2093 and 495.2212, and molecular formulas of C₂₅H₃₀O₁₂, C₂₆H₃₂O₁₂ and C₂₃H₂₈O₁₂. These compounds were previously isolated from *T. capensis*,³¹ however, they are reported here in most of the studied *Tecoma* species for the first time. One flavonoid was identified as being luteolin 7-*O*-β-D-glucopyranoside (**4**) by its molecular ion peak depicted at *m/z* 447.1537,

Table 2 Docking simulations results of the studied *Tecoma* phytochemicals

No.	Name of phytoligands	ΔG^a (kcal mol ⁻¹)	Interactions at the binding interface ^b	
			hACE2 residues	SARSCoV-2-CTD residues
1	Octanoic acid, 4-benzyloxyphenyl ester	-6.66	No interaction	Gln493
2	Fumaric acid, 3,4-dimethoxyphenyl heptyl ester	-5.28	No interaction	Glu406, Arg408, Lys417
3	Boschniakine	-3.76	Lys353	Gln493
4	Luteolin 7- <i>O</i> -β-D-glucopyranoside	-4.84	Glu37, Lys353	Glu406, Gln493
5	Actinidine	-4.80	No interaction	No interaction
6	Skytanthine	-4.47	No interaction	Arg403
7	Tecomanine (syn. tecomine)	-4.64	Lys353	Gly496
8	7- <i>O</i> -(<i>p</i> -OH)cinnamoyltecomoside	-6.73	No interaction	Glu406, Gln409
9	7- <i>O</i> -(<i>p</i> -MeO)cinnamoyltecomoside	-5.44	His34 , Ala389, Arg393	Arg403
10	Succinic acid, decyl 3-oxobut-2-yl ester	-6.77	His34 , Glu35	Gln493
11	7- <i>O</i> -(<i>p</i> -OH)benzoyl tecomoside	-6.48	Glu37, Lys353	Asp405, Arg408, Gly496
12	Valeric acid, pentadecyl ester	-6.73	Lys353	Gly496
13	Hesperidin	-7.10	His34 , Ala387	Gln409, Lys417 , Ser494

^a The ligand–receptor complex binding free energy at RMSD < 2 Å. ^b The key residues involved in the SARS-CoV-2-CTD–hACE2 complex formation are listed in bold.





Fig. 1 (A) 3D binding mode of **1** (cyan sticks), (B) 2D binding mode of **1**, (C) 3D binding mode of **2** (yellow sticks), (D) 2D binding mode of **2**, (E) 3D binding mode of **3** (magenta sticks), (F) 2D binding mode of **3**, (G) 3D binding mode of **4** (orange sticks), (H) 2D binding mode of **4**, (I) 3D binding mode of **6** (light pink sticks), (J) 2D binding mode of **6**, (K) 3D binding mode of **7** (green sticks), (L) 2D binding mode of **7**, (M) 3D binding mode of **8** (deep pink sticks), (N) 2D binding mode of **8**, (O) 3D

binding mode of **9** (white sticks), (P) 2D binding mode of **9**, (Q) 3D binding mode of **10** (red sticks), (R) 2D binding mode of **10**, (S) 3D binding mode of **11** (deep yellow sticks), (T) 2D binding mode of **11**, (U) 3D binding mode of **12** (pink sticks), (V) 2D binding mode of **12** (W) 3D binding mode of **13** (blue sticks), (X) 2D binding mode of **13** in the binding interface of SARS-CoV-2-CTD in complex with hACE2 (PDB ID 6LZG).

3.2. Molecular docking analysis of selected phytocompounds from *Tecoma* species and cultivars

Docking simulations results (Table 2) showed that most of the studied phytocompounds displayed moderate to promising binding affinities compared to hesperidin (**13**). Succinic acid, decyl-3-oxobut-2-yl ester (**10**) came at the top of the list due to its best binding affinity ($-6.77 \text{ kcal mol}^{-1}$) among the studied phytocompounds. 7-*O*-(*p*-OH)cinnamoyltecomoside (**8**), valeric acid, pentadecyl ester (**12**), octanoic acid, 4-benzyloxyphenyl ester (**1**), 7-*O*-(*p*-OH)benzoyl tecomoside (**11**) showed slightly less binding affinities ranging from -6.73 to $-6.48 \text{ kcal mol}^{-1}$. Moderate fitting was observed in case of the two phytocompounds *viz.* 7-*O*-(*p*-MeO)cinnamoyltecomoside (**9**), and fumaric acid, 3,4-dimethoxyphenyl heptyl ester (**2**) with binding affinities of -5.44 and $-5.28 \text{ kcal mol}^{-1}$, respectively. The remaining phytocompounds showed relatively low binding affinities.

Inspecting the binding modes of the promising phytocompounds revealed that most of them were able to accommodate into the interface and interact with the key amino acids, in most cases, through hydrogen bonds (Fig. 1 and 2). Thus, they may destabilize or even prevent the virus-receptor engagement that is generally dominated by polar contacts mediated by these key hydrophilic amino acid residues.²⁷ Succinic acid, decyl-3-oxobut-2-yl ester (**10**), showing the best binding affinity in the current study, exhibited H- π interactions with His34 of the hACE-2 involved in the complex formation (Fig. 1Q and R). Among the studied phytocompounds, the second most promising *in silico* activity was recorded by valeric acid, pentadecyl ester (**12**) which interacted with SARS-CoV-2-CTD Gly496 and hACE2 Lys353 (Fig. 1U and V). Similarly, 7-*O*-(*p*-OH)benzoyl tecomoside (**11**) exhibited these key interactions in addition to binding hACE-2 Glu37 (Fig. 1S and T). It is worth mentioning that a single Lys353 mutation was reported to be sufficient to abolish the interactions at the interface. Interestingly, boschniakine (**3**; Fig. 1E and F), luteolin 7-*O*-*D*-glucopyranoside (**4**; Fig. 1G and H), tecomanine (**7**; Fig. 1K and L) were able to interact with Lys353 despite their relatively low binding affinities to the complex. Notably, fumaric acid, 3,4-dimethoxyphenyl heptyl ester (**2**) showed interactions with SARS-CoV-2 CTD through its key residue Lys417 as well as Glu406 with no interactions with the hACE-2 side (Fig. 1C and D). This observation was also recorded in case of octanoic acid, 4-benzyloxyphenyl ester (**1**; Fig. 1A and B), skytanthine (**6**; Fig. 1I and J) and 7-*O*-(*p*-OH)cinnamoyltecomoside (**8**; Fig. 1M and N). On the

binding mode of **9** (white sticks), (P) 2D binding mode of **9**, (Q) 3D binding mode of **10** (red sticks), (R) 2D binding mode of **10**, (S) 3D binding mode of **11** (deep yellow sticks), (T) 2D binding mode of **11**, (U) 3D binding mode of **12** (pink sticks), (V) 2D binding mode of **12** (W) 3D binding mode of **13** (blue sticks), (X) 2D binding mode of **13** in the binding interface of SARS-CoV-2-CTD in complex with hACE2 (PDB ID 6LZG).





Fig. 2 Overlay of 1 (cyan sticks), 2 (yellow sticks), 3 (magenta sticks), 4 (orange sticks), 6 (light pink sticks), 7 (green sticks), 8 (deep pink sticks), 9 (white sticks), 10 (red sticks), 11 (deep yellow sticks), 12 (pink sticks) and 13 (blue sticks) in the binding interface of SARS-CoV-2-CTD in complex with hACE2 (PDB ID 6LZG).

other hand, actinidine (5) did not exhibit any considerable interactions with its residues.

4. Conclusions

In the current study, *in silico* exploration was conducted for some of *Tecoma* phytochemicals that could inhibit SARS-CoV entry to its host cells. Twelve compounds from eight *Tecoma* plants belonging to different phytochemical classes *viz.* alkaloids, iridoids, flavonoids, and fatty acid esters were derelicated using LC-HRESIMS, among which a fatty acid ester named succinic acid, decyl-3-oxobut-2-yl ester (10) was reported to possess the best binding affinity ($-6.77 \text{ kcal mol}^{-1}$), followed by 7-*O*-(*p*-OH)cinnamoyltecomoside (8), valeric acid, pentadecyl ester (12), octanoic acid, 4-benzyloxyphenyl ester (1), 7-*O*-(*p*-OH) benzoyl tecomoside (11) which showed slightly less binding affinities ranging from -6.73 to $-6.48 \text{ kcal mol}^{-1}$, as compared to hesperidin (13). These phytochemicals could serve as potential candidates for the discovery of anti-SARS drugs that possess a preventive potential, however the therapeutic potential is yet to be validated using *in vitro* and *in vivo* studies.

Conflicts of interest

The authors declare no conflict of interest.

Acknowledgements

We thank Prof. Tobias Gulder (University of Dresden) for LC-MS measurement.

References

- 1 B. J. Bosch, R. van der Zee, C. A. M. de Haan and P. J. M. Rottier, *J. Virol.*, 2003, **77**, 8801–8811.
- 2 A. M. Sayed, A. R. Khattab, A. M. AboulMagd, H. M. Hassan, M. E. Rateb, H. Zaid and U. R. Abdelmohsen, *RSC Adv.*, 2020, **10**, 19790–19802.

- 3 A. S. Omrani, M. M. Saad, K. Baig, A. Bahloul, M. Abdul-Matin, A. Y. Alaidaroos, G. A. Almakhlafi, M. M. Albarrak, Z. A. Memish and A. M. Albarrak, *Lancet Infect. Dis.*, 2014, **14**, 1090–1095.
- 4 X. Y. Ge, J. L. Li, X. Lou Yang, A. A. Chmura, G. Zhu, J. H. Epstein, J. K. Mazet, B. Hu, W. Zhang, C. Peng, Y. J. Zhang, C. M. Luo, B. Tan, N. Wang, Y. Zhu, G. Cramer, S. Y. Zhang, L. F. Wang, P. Daszak and Z. L. Shi, *Nature*, 2013, **503**, 535–538.
- 5 X. Xu, P. Chen, J. Wang, J. Feng, H. Zhou, X. Li, W. Zhong and P. Hao, *Sci. China: Life Sci.*, 2020, **63**, 457–460.
- 6 C. Wu, Y. Liu, Y. Yang, P. Zhang, W. Zhong, Y. Wang, Q. Wang, Y. Xu, M. Li, X. Li, M. Zheng, L. Chen and H. Li, *Acta Pharm. Sin. B*, 2020, **10**(5), 766–788.
- 7 J. Shang, G. Ye, K. Shi, Y. Wan, C. Luo, H. Aihara, Q. Geng, A. Auerbach and F. Li, *Nature*, 2020, **581**, 221–224.
- 8 L. Aguilar-Santamaría, G. Ramírez, P. Nicasio, C. Alegría-Reyes and A. Herrera-Arellano, *J. Ethnopharmacol.*, 2009, **124**, 284–288.
- 9 S. Verma, *Phytochemical and pharmacological review study on Tecoma stans Linn*, 2016, vol. 4.
- 10 M. Rahmatullah, W. Samarrai, R. Jahan, S. Rahman, Z. U. M. E. U. Miajee, M. H. Chowdhury, S. Bari, A. B. M. A. Bashar, A. K. Azad and S. Ahsan, *DNA Repair*, 2010, **4**, 236–253.
- 11 L. Zhang, D. Lin, X. Sun, U. Curth, C. Drosten, L. Sauerhering, S. Becker, K. Rox and R. Hilgenfeld, *Science*, 2020, **368**, 409–412.
- 12 A. K. Ghosh, M. Brindisi, D. Shahabi, M. E. Chapman and A. D. Mesecar, *ChemMedChem*, 2020, **15**, 907–932.
- 13 R. Alexpandi, J. F. De Mesquita, S. K. Pandian and A. V. Ravi, *Front. Microbiol.*, 2020, **11**, DOI: 10.3389/fmicb.2020.01796.
- 14 W. R. Ferraz, R. A. Gomes, A. L. S. Novaes and G. H. Goulart Trossini, *Future Med. Chem.*, 2020, **12**, 1815–1828.
- 15 D. Gentile, V. Patamia, A. Scala, M. T. Sciortino, A. Piperno and A. Rescifina, *Mar. Drugs*, 2020, **18**(4), DOI: 10.3390/md18040225.
- 16 S. Shahinshavali, K. A. Hossain, A. V. D. N. Kumar, A. G. Reddy, D. Kolli, A. Nakhi, M. V. B. Rao and M. Pal, *Tetrahedron Lett.*, 2020, **61**(40), DOI: 10.1016/j.tetlet.2020.152336.
- 17 M. Hagar, H. A. Ahmed, G. Aljohani and O. A. Alhaddad, *Int. J. Mol. Sci.*, 2020, **21**(11), DOI: 10.3390/ijms21113922.
- 18 S. T. Ngo, N. Quynh Anh Pham, L. Thi Le, D.-H. Pham and V. V. Vu, *J. Chem. Inf. Model.*, 2020, DOI: 10.1021/acs.jcim.0c00491.
- 19 A. I. Owis, M. S. El-Hawary, D. El Amir, O. M. Aly, U. R. Abdelmohsen and M. S. Kamel, *RSC Adv.*, 2020, **10**, 19570–19575.
- 20 F. M. Abd El-Mordy, M. M. El-Hamouly, M. T. Ibrahim, G. A. El-Rheem, O. M. Aly, A. M. Abd El-Kader, K. A. Youssif and U. R. Abdelmohsen, *RSC Adv.*, 2020, **10**, 32148–32155.
- 21 A. S. Achutha, V. L. Pushpa and S. Suchitra, *J. Proteome Res.*, 2020, **19**(11), 4706–4717, DOI: 10.1021/acs.jproteome.0c00683.
- 22 R. Kumawat, S. Sharma and S. Kumar, *Acta Pol. Pharm.*, 2012, **69**, 993–996.



- 23 W. M. Abdel-Mageed, E. Y. Backheet, A. A. Khalifa, Z. Z. Ibraheim and S. A. Ross, *Fitoterapia*, 2012, **83**, 500–507.
- 24 A. H. Elmaidomy, R. Mohammed, H. M. Hassan, A. I. Owis, M. E. Rateb, M. A. Khanfar, M. Krischke, M. J. Mueller and U. R. Abdelmohsen, *Metabolites*, 2019, **17**(8), DOI: 10.3390/md17080465.
- 25 C. Guijas, J. R. Montenegro-Burke, X. Domingo-Almenara, A. Palermo, B. Warth, G. Hermann, G. Koellensperger, T. Huan, W. Uritboonthai, A. E. Aisporna, D. W. Wolan, M. E. Spilker, H. P. Benton and G. Siuzdak, *Anal. Chem.*, 2018, **90**, 3156–3164.
- 26 M. Whittle, P. Willett, W. Klaffke and P. Van Noort, *J. Chem. Inf. Comput. Sci.*, 2003, **43**, 449–457.
- 27 Q. Wang, Y. Zhang, L. Wu, S. Niu, C. Song, Z. Zhang, G. Lu, C. Qiao, Y. Hu, K. Y. Yuen, Q. Wang, H. Zhou, J. Yan and J. Qi, *Cell*, 2020, **181**(4), 894–904.
- 28 L. Costantino, L. Raimondi, R. Pirisino, T. Brunetti, P. Pessotto, F. Giannessi, A. P. Lins, D. Barlocco, L. Antolini and S. A. El-Abady, *Farmaco*, 2003, **58**, 781–785.
- 29 A. Al-Azzawi, A. Al-Guboori, A. Abdul-Sada and M. Al-Azzawi, *Planta Med.*, 2010, **76**(12), DOI: 10.1055/s-0030-1264698.
- 30 G. H. Harris, E. C. Fixman, F. R. Stermitz and L. Castedo, *J. Nat. Prod.*, 1988, **51**, 543–548.
- 31 M. Guiso, C. Marra, F. Piccioni and M. Nicoletti, *Phytochemistry*, 1997, **45**, 193–194.
- 32 M. Marzouk, A. Gamal-Eldeen, M. Mohamed and M. El-Sayed, *Z. Naturforsch., C: J. Biosci.*, 2006, **61**, 783–791.

