


 Cite this: *RSC Adv.*, 2022, 12, 30278

Recovery of bacterioruberin and proteins using aqueous solutions of surface-active compounds†

 Bárbara M. C. Vaz,^{‡a} Mariam Kholany,^{‡a} Diana C. G. A. Pinto,^{Ⓜ^b} Inês P. E. Macário,^{ac} Telma Veloso,^{ac} Tânia Caetano,^{Ⓜ^c} Joana L. Pereira,^{Ⓜ^c} João A. P. Coutinho^{Ⓜ^a} and Sónia P. M. Ventura^{Ⓜ^{*a}}

Haloarchaea microorganisms are little explored marine resources that can be a promising source of valuable compounds with unique characteristics, due to their adaptation to extreme environments. In this work, the extraction of bacterioruberin and proteins from *Haloferax mediterranei* ATCC 33500 was investigated using aqueous solutions of ionic liquids and surfactants, which were further compared with ethanol. Despite the good performance of ethanol in the extraction of bacterioruberin, the use of aqueous solutions of surface-active compounds allowed the simultaneous release of bacterioruberin and proteins in a multi-product process, with the non-ionic surfactants being identified as the most promising. The optimum operational conditions allowed a maximum extraction yield of 0.37 ± 0.01 mg_{bacterioruberin} g_{wet biomass}⁻¹ and 352 ± 9 mg_{protein} g_{wet biomass}⁻¹ with an aqueous solution of Tween® 20 (at 182.4 mM) as the extraction solvent. In addition, high purities of bacterioruberin were obtained, after performing a simple induced precipitation using ethanol as an antisolvent to recover the proteins present in the initial extract. Finally, a step for polishing the bacterioruberin was performed, to enable solvent recycling, further closing the process to maximize its circularity.

 Received 22nd April 2022
 Accepted 2nd October 2022

DOI: 10.1039/d2ra02581g

rsc.li/rsc-advances

Introduction

Our continued reliance on a petroleum-based economy to satisfy humankind's needs cannot be sustained anymore.¹ The traditional model of take, make and dispose using fossil-based feedstocks has severe implications for human health, the depletion of fossil resources, and as a driver of climate change.² In this sense, the development of sustainable technologies and practices to ensure a gradual transition to a renewable (bio-based) economy should be encouraged.^{1,3} By considering the concepts of circular economy and integrated biorefineries, the complete valorization of all biomass fractions could be achieved. Besides maximizing the value of raw materials, waste production could be minimized or avoided, which leads to an increase in the economic viability of downstream processes for industrial implementation.⁴

Recently, marine resources like algae and cyanobacteria have gained significant interest as renewable feedstocks to produce several natural bioactive compounds. Their sustainable use can easily fit the priorities of the 2030 Agenda for Sustainable Development.⁵ They can generate biofuels and bioenergy and integrate bioactive medicinal products, food, and materials.⁶ In this context, halophilic microorganisms from the domain Archaea, with high commercial potential, have emerged as underexplored marine resources, easily found around the globe, mostly in hypersaline environments. Additionally, they are natural producers of several relevant high-demand products, such as proteins, poly(3-hydroxybutyrate), polyhydroxyalkanoates, and carotenoids.⁷

Proteins are complex molecules with a unique three-dimensional structure composed of numerous combinations of the 20 different amino acids attached to form long chains.⁸ There is a problem in searching for new sources of proteins for food purposes. However, due to their multiple biological roles, their biotechnological interest is vast,^{9,10} and thus, depending on the final application, their purification may be required. In halophilic archaea (or haloarchaea), halophilic proteins assist adaptation to extreme saline conditions by modifying their structure. The presence of acidic amino acids on the protein surface confers a more hydrophilic and flexible surface, capable of increasing the hydration of the protein. Hence, its aggregation is prevented, and the protein's functionality is retained.¹¹ On the other hand, carotenoids are natural colorants well

^aCICECO – Aveiro Institute of Materials, Department of Chemistry, University of Aveiro, Campus Universitário de Santiago, 3810-193 Aveiro, Portugal. E-mail: spventura@ua.pt

^bLAQV – REQUIMTE, Department of Chemistry, University of Aveiro, 3810-193 Aveiro, Portugal

^cCESAM – Centre for Environmental and Marine Studies, Department of Biology, University of Aveiro Campus Universitário de Santiago, 3810-193 Aveiro, Portugal

† Electronic supplementary information (ESI) available. See DOI: <https://doi.org/10.1039/d2ra02581g>

‡ These authors contributed equally to this manuscript.



known for their biotechnological applications and key properties for human health.^{12,13} The haloarchaea group can produce many carotenoids, including β -carotene, salinixanthin, bacterioruberin, and its precursors lycopene and phytoene.^{7,12} Among these, bacterioruberin represents the most abundant carotenoid produced, since all the species with sequenced genomes encode its biosynthetic pathway.¹⁴ Bacterioruberin can influence the membrane fluidity, acting as a water barrier and allowing the permeability of oxygen and other molecules needed for the cell's survival in hypersaline environments. Consequently, the membrane structure rigidity is increased.^{7,15,16} Of great relevance, this C₅₀ isoprenoid, composed of 13 conjugated double bonds and 4 hydroxyl groups, has a higher antioxidant activity than other molecules. Therefore, higher protection can be achieved against intense light, gamma irradiation, and DNA damaging agents such as radiography, UV-irradiation, and H₂O₂ exposure.^{7,15,16} These characteristics make bacterioruberin an appealing bioactive compound to be applied in the food, cosmetic and pharmaceutical industries.¹⁷ Moreover, their commercial potential goes beyond the applications previously mentioned. More recently, some of us have investigated its optical potential to develop optical sensors and retinal gene therapy formulations (ongoing work), with promising results. Despite the high potential of bacterioruberin and halophilic proteins, efficient methods of extraction and purification are still at an early stage. Most literature focuses on exploring protein structures and their adaptations to extreme environments,^{18–23} and optimizing bacterioruberin production and its possible applications.^{15,17,24–26} The studies found just quantify bacterioruberin extracted using organic solvents, without considering biomass valorization. Regarding the purification step, some works used thin-layer chromatography.^{17,27–32} However, this technique, despite being

simple and efficient at the lab scale, is not industrially adequate. In this sense, more efforts are needed to develop efficient and sustainable approaches with scale-up potential encompassing the valorization of multiple products obtained from this biomass.

In this work, aqueous solutions of surfactants and ionic liquids (ILs) were investigated with the prospect of simultaneously recovering different products from the same biomass in a more sustainable way. As surfactants and tensioactive ILs are amphiphilic molecules,³³ they become effective solvents in the extraction of both more hydrophilic compounds, *e.g.* proteins,^{10,34–37} and more hydrophobic molecules,^{37–41} such as bacterioruberin. Furthermore, their use has several advantages, namely milder process conditions while using water as the main solvent, which could also be attractive for cell disruption, as most haloarchaea lyse rapidly when suspended in water.⁷ After the extraction of multiple compounds, a purification step is usually needed to obtain each compound with the appropriate purity for the specific application. Induced protein precipitation was applied in this work. The addition of a precipitating agent lowers the solubility of proteins in water, causes their precipitation⁴² and allows the separation of the solvent. As well as its simplicity, it has industrial potential,⁴³ being recognized as having a low carbon footprint and economic impact.^{44,45} In the end, a polishing step was also applied to bacterioruberin, allowing the solvent to be recycled.

Results and discussion

Screening of surfactants and ILs as solvents

A large screening of aqueous solutions of surface-active compounds belonging to different classes (cationic, anionic, non-ionic) and other non-surfactant agents was performed at




Fig. 1 Bacterioruberin yield of extraction using aqueous solutions of cationic, anionic, and non-ionic compounds at 100 mM, as well as ethanol, used as a control solvent, at room temperature (20–25 °C), under a constant vertical rotation at 50 rpm, for 45 min, protected from light exposure, and at a fixed SLR of 0.1 g_{wet biomass} mL_{solvent}⁻¹.



100 mM to assess their ability to extract bacterioruberin (the most valuable compound present). The outcomes were benchmarked against ethanol. The surfactants tested were pre-selected considering previous works that identified them to be efficient in the extraction of biomolecules, namely carotenoids.^{37,38,46} Table S1 in the ESI† reports the ability of the surface-active compounds tested (30 in total) to extract bacterioruberin. Fig. 1 shows the yields of extraction ($\text{mg}_{\text{bacterioruberin}} \text{g}_{\text{wet biomass}}^{-1}$) of the systems with extraction capability, in terms of the ability of the surfactants/ILs to induce the release of the target compound. Photographs of all systems can be found in Fig. S1 in the ESI.†

Ethanol displayed a better bacterioruberin yield of extraction than aqueous solutions of surface-active compounds (Fig. 1). Ethanol is recognized for its excellent performance in pigment extraction and membrane solubilization.^{40,47} Thus, and also owing to the presence of bacterioruberin in the cell membrane, which endows it with a low water permeability,⁷ the performance of ethanol in bacterioruberin extraction was expected.

Regarding the performance of the aqueous solutions of surface-active compounds as depicted in Fig. 1 and Table S1 in the ESI,† almost none of the anionic or the non-tensioactive compounds were able to efficiently extract bacterioruberin. The non-tensioactive ILs have a low capability to disrupt the phospholipidic cell membranes due to their hydrophilic nature.³⁸ The low performance of the anionic surfactants can be attributed to the fact that both types of tensioactive compounds and the phospholipids (components of the cell membrane) are negatively charged, creating repulsive forces.³⁷ In contrast, almost all cationic and non-ionic compounds were able to release the compound of interest. This can be explained by their ability to establish attractive electrostatic interactions between the IL cation and the negatively charged head of the phospholipids.^{38,39} Bacterioruberin is almost exclusively present in its

non-ionic form. This allows for more stable interactions between the pigment and the micelle hydrophobic core. However, with such conditions (with a surfactant concentration of 100 mM), a high viscosity was observed in most solutions obtained after extraction (see Table S1 in the ESI†), which is normally correlated with the release of chromosomal DNA during cell disruption.^{48,49} This phenomenon, whose origin is yet to be proved, imposed some experimental constraints. To overcome this issue and better assess the extraction capacity of the most efficient solvents (here, the non-ionic surfactants), a second screening at 250 mM surfactant concentration followed, with the respective results shown in Fig. 2. The UV-Vis spectra of the resulting extracts can be found in Fig. S2 in the ESI.†

Since the non-ionic surfactants tested are viscous liquids at room temperature, an increase in their concentration results in a more viscous aqueous solution. However, the viscosity of the final solutions (after bacterioruberin extraction) at 250 mM was revealed to be lower and more manageable than that of the same solutions at 100 mM. On the other hand, a very high surfactant concentration in the initial aqueous solution can also lead to viscosity issues. Therefore, the choice of an adequate concentration of surfactant seems to be a key factor in this work's success. Fig. 2 shows that the highest performances at these higher concentrations were once again accomplished by Tween® 20 and ethanol. Nevertheless, ethanol lacks the desired capacity to simultaneously extract bacterioruberin and the proteins present in the biomass, thus failing to achieve the envisioned multi-product exploitation scenario. Aqueous solutions of surfactants provide a milder environment, thus allowing the extraction of different fractions of interest (proteins and pigments), both with applications related to the food industry. Ethanol leads to a protein extraction yield of approximately $24 \text{ mg}_{\text{protein}} \text{g}_{\text{wet biomass}}^{-1}$, while Tween® 20 leads to a protein




Fig. 2 Bacterioruberin yield of extraction using aqueous solutions of non-ionic compounds at 250 mM, as well as using ethanol as the control solvent, at room temperature (20–25 °C), under a constant vertical rotation at 50 rpm, for 45 min, protected from light exposure, and at a fixed SLR of $0.1 \text{ g}_{\text{wet biomass}} \text{ mL}_{\text{solvent}}^{-1}$.



extraction yield of approximately $352 \text{ mg}_{\text{protein}} \text{ g}_{\text{wet biomass}}^{-1}$, increasing by over 14 times the amount of protein obtained. As a result, the following work used Tween® 20 as the best solvent.

Optimization of the solid–liquid extraction

An optimization of the operational conditions was performed based on a central composite rotatable design (CCRD-2³) comprising three independent variables, namely SLR (in $\text{g}_{\text{wet biomass}} \text{ mL}_{\text{solvent}}^{-1}$, X1), time of extraction (t in min, X2) and concentration of surfactant in water (C_{surf} in mM, X3). Eighteen assays with four central (level 0) and axial points (-1.68 and $+1.68$ levels) were studied in terms of bacterioruberin yield of extraction (in $\text{mg}_{\text{bacterioruberin}} \text{ g}_{\text{wet biomass}}^{-1}$) ranging from $0.111 \text{ mg}_{\text{bacterioruberin}} \text{ g}_{\text{wet biomass}}^{-1}$ in assay 5 to $0.331 \text{ mg}_{\text{bacterioruberin}} \text{ g}_{\text{wet biomass}}^{-1}$ in assay 9 (Table S2 in the ESI†). The fitted model expressed in eqn (1), obtained using the SS residual from Analysis of Variance (ANOVA), revealed good predictability at a confidence level of 95% with $R^2 = 0.80912$ and $F_{\text{calculated}} > F_{\text{tabulated}}$.

$$\text{Yield of extraction (mg}_{\text{bacterioruberin}} \text{ g}_{\text{wet biomass}}^{-1}) = 0.147055 - 0.270072(X1) + 0.002002(X3)^2 \quad (1)$$

The response surfaces plotted in Fig. 3 show low effects on yield imposed by the extraction time. Therefore, the minimum time tested is the obvious choice to reduce the energy consumption of

the process. The concentration of Tween® 20 has a great influence in the extraction process, where an optimum value was reached at 182.4 mM, the minimum level tested for this variable. These results suggest that a higher extraction yield could be achieved at concentrations of Tween® 20 below 182.4 mM, although an increase in the tested range would not be suitable due to the viscosity issues noticed (see above). SLR has a positive influence on the yield of extraction as predicted by the model, with the best response being obtained at $0.06224 \text{ g}_{\text{wet biomass}} \text{ mL}_{\text{solvent}}^{-1}$. These results can be confirmed by the data present in Table S3† in the ESI†, the Pareto chart (Fig. S3 in the ESI†), and the predicted *vs.* observed values present in Fig. S4 in the ESI†. Here, it becomes evident that the solid–liquid ratio and the concentration of Tween® 20 in water were the variables that most constrained the model regarding the bacterioruberin yield of extraction. Therefore, after finding the optimal operational conditions (8.04 min, 182.4 mM, and $0.06224 \text{ g}_{\text{wet biomass}} \text{ mL}_{\text{solvent}}^{-1}$), a model validation was developed. A bacterioruberin yield of extraction of $0.37 \pm 0.01 \text{ mg}_{\text{bacterioruberin}} \text{ g}_{\text{wet biomass}}^{-1}$ was obtained experimentally (Table S4 in the ESI†), encompassing a mean relative deviation of 11.6%. Under these conditions, a protein yield of extraction of $352 \pm 9 \text{ mg}_{\text{protein}} \text{ g}_{\text{wet biomass}}^{-1}$ was achieved.

Protein induced precipitation

Ethanol is a common solvent used as a precipitating agent since it possesses a lower dielectric constant than water,⁵⁰ which leads



Fig. 3 Response surface plots obtained for the CCRD (2³) using an aqueous solution of Tween® 20 regarding SLR ($\text{g}_{\text{wet biomass}} \text{ mL}_{\text{solvent}}^{-1}$), concentration of surfactant in water (C_{surf} , mM), and time of extraction (t , min) in terms of bacterioruberin yield of extraction (yield of extraction, $\text{mg}_{\text{bacterioruberin}} \text{ g}_{\text{wet biomass}}^{-1}$).



to an increase in the attraction forces between proteins, specifically at low temperatures.^{51–53} To achieve the minimum amount of protein in the supernatant fraction (SF), an evaluation of the process conditions, regarding temperature, volume of ethanol added to the initial extract, and time of the precipitation, was performed between -20 and 30 °C, 0.5 and 4 mL and 2 and 30 min, respectively (Fig. 4).

The influence of temperature on the precipitation was assessed at a fixed time of 20 min and a volume of ethanol of 3 mL. Lower temperatures, particularly 4 °C, slightly reduce the protein losses of the process (Fig. 4A). However, as the difference in the protein content between 4 °C and 25 °C corresponds to only 3.9% , a temperature of 25 °C was seen as preferable, as it lowers the energetic burden of the overall process. Taking this into account, the precipitation time was evaluated at a fixed temperature of 25 °C and 3 mL of ethanol. The data obtained (Fig. 4B) show an almost constant protein content with a minimum at 2 min, and this time was chosen to proceed.




Fig. 4 Protein content in the supernatant fraction (SF) after protein precipitation considering different temperatures (A), times of precipitation (B), and volumes of ethanol added to the initial extract (C).

Finally, the volume of ethanol was considered (Fig. 4C) under fixed conditions of 25 °C and 2 min, and it was shown that the volume of ethanol is the variable that most influences the precipitation process: protein content decreases with an increase in the ethanol volume added to the initial extract. A volume of 3 mL of ethanol was selected as the difference between 3 mL and 4 mL corresponds to only 1.5% protein content. Thus, with a set of optimum conditions of 25 °C, 2 min, and 3 mL of ethanol, a yield of 234.5 ± 0.9 $\text{mg}_{\text{protein}} \text{g}_{\text{wet biomass}}^{-1}$ (quantified and calculated after protein redissolution in PBS), which corresponds to 91% of the proteins present in the initial extract, was obtained.

Moreover, further studies were conducted to understand the effect of consecutive precipitations (Fig. S5 in the ESI†). After the first precipitation, and consequent recovery of the pellet fraction, a second precipitation was carried out by adding another 3 mL to the supernatant fraction and repeating the procedure. Only a very small fraction of approximately 1.0% of the total proteins present in the initial extract was obtained in the second precipitation. This corresponds to a yield of 2.7 ± 0.3 $\text{mg}_{\text{protein}} \text{g}_{\text{wet biomass}}^{-1}$ (quantified and calculated after protein redissolution in PBS), which is markedly lower when compared to the first, and thus the application of a second step of precipitation was disregarded. This further confirms that over 90% of the protein fraction was recovered in a single step of precipitation, as also visible by the abundance of proteins separated through SDS-PAGE electrophoresis of the precipitated proteins after dissolution in PBS ($1\times$, pH 7.4), Fig. S6 in the ESI.† Finally, the bacterioruberin content was measured, after ethanol evaporation, to investigate bacterioruberin losses during induced precipitation, with a loss fixed at 12% .

Bacterioruberin polishing and conceptual process design

A bacterioruberin polishing step was accomplished to achieve the recycling of Tween® 20. Here, the objective was the separation of the bacterioruberin and the solvent, thus facilitating its further application. Briefly, to perform this step, the water content was first removed from the bacterioruberin extract (extract obtained after induced precipitation and ethanol evaporation) by freeze-drying. Then, the formation of two phases was achieved by adding ethanol to the sample [$1 : 20$ ($V_{\text{bacterioruberin extract}} : V_{\text{ethanol}}$)]. Afterwards, the solution obtained was stored in the freezer at -80 °C for three days. Thus, bacterioruberin migrated to the ethanol-rich (top) phase, while Tween® 20 remained in the bottom phase. At the end, the ethanol-rich phase was analyzed by ^1H NMR spectroscopy (Fig. S7 in the ESI†) to evaluate if Tween® 20 was fully separated, and no surfactant was detected. The presence of bacterioruberin in the top phase of the collected sample was confirmed by UHPLC-MS analysis. It was possible to identify a mixture of bacterioruberin ($m/z = 741$) and monoanhydrobacterioruberin ($m/z = 724$) at around 12.59 min, with bacterioruberin being the major compound in the mixture (85%). However, under the conditions used (positive mode), these two compounds' pseudomolecular ions appear as $[\text{M} + \text{H}]^+$ and easily produce more stable fragment ions with $m/z = 635$ and 591 , respectively (see Fig. S8 in



the ESI⁺). Finally, the bottom phase was analyzed by UV-Vis spectroscopy, with only 2.8% of bacterioruberin detected, indicating that around 97.2% of bacterioruberin was successfully concentrated in the top phase.

A final diagram of the process developed in this work was proposed (Fig. 5), in which all steps were considered. In the end, the colorant can be provided in ethanolic solution or in its dry form, depending on the desired end application. Drying should be performed using a vacuum dryer at low temperature and pressure. This not only prevents pigment degradation, thus allowing the recycling of the solvents, Tween® 20 and ethanol, but also allows easy scale-up.

Experimental

Materials and methods

Archaea cultivation. *Haloferax mediterranei* ATCC 33500 was produced under controlled conditions targeting maximum bacterioruberin yield, in YPC-Hv (yeast, peptone and casamino acids media) culture medium: peptone of meat 1.0 (g L⁻¹); casamino acids 1.0 (g L⁻¹); yeast extract 5.0 (g L⁻¹); NaCl 144 (g L⁻¹); MgSO₄·7H₂O 21.0 (g L⁻¹); MgCl₂·6H₂O 18.0 (g L⁻¹); KCl 4.2 (g L⁻¹); Tris-HCl 12 mM (pH 7.5); KOH 1 M, and 0.5 M CaCl₂.⁵⁴ The medium was sterilized at 121 °C prior to Archaea culturing. A single colony was selected from the agar plate (YPC-Hv broth with 1.5% agar) and inoculated into 25 mL of YPC-Hv broth in a 100 mL Erlenmeyer flask and cultured at 37 °C, 180 rpm, under continuous light (4000 lux) for 72 h. To increase the cell mass concentration, 20 mL of this pre-inoculum were resuspended in 400 mL of fresh broth in a 1 L Erlenmeyer flask

and incubated under the same conditions (*i.e.* 37 °C, 180 rpm, 4000 lux) for 96 h. Archaea growth was monitored by optical density determined at 600 nm. After the batch culturing period, the biomass was centrifuged in a Thermo Scientific Heraeus Megafuge 16R centrifuge at 18 894*g* for 15 min at room temperature (20–25 °C). The supernatant was discarded, and the pellet was stored in the dark at –20 °C until further use. Some natural variability was found in the data due to the use of different cultivation batches.

Chemicals. Ethanol absolute (analytical reagent grade, CAS 64-17-5) used in solid-liquid extraction and in induced protein precipitation was acquired from Fisher Scientific. Sodium chloride (NaCl, 99.5 wt%, CAS 7647-14-5), potassium chloride (KCl, 99.5 wt%, CAS 7447-40-7), anhydrous di-sodium hydrogen phosphate (Na₂HPO₄, 99 wt%, CAS 7558-79-4) and monobasic potassium phosphate (KH₂PO₄, 99.5–100.5 wt%, CAS 7778-77-0), used to prepare phosphate-buffered saline (PBS), were purchased from Fisher Scientific, Chem-Lab, Panreac, and Honeywell, respectively. To prepare the loading buffer for SDS-PAGE analysis, glycerol (99.88 wt%, CAS 56-81-5) acquired from Fisher Chemical, bromophenol blue sodium salt (CAS 34725-61-6) from Merck, dithiothreitol (DTT, 98 wt%, CAS 3483-12-3) from NZYtech, and tris(hydroxymethyl)-aminomethane ((HOCH₂)₃CNH₂, 99 wt%, CAS 77-86-1) from Alfa Aesar, were used. RunBlue Teo 20× Teo Tricine SDS and RunBlue SDS gel 4–12% 10 cm × 10 cm were supplied by Expedeon. BlueSafe used to stain the proteins and Amersham™ ECL™ Rainbow™ Marker – full range were acquired from NZYtech and Cytiva, respectively. Information regarding the ILs and surfactants screened in this work such as purity, CAS number and supplier




Fig. 5 Schematic diagram of the downstream process developed in this work, consisting of (i) cell disruption/solid-liquid extraction of bacterioruberin and proteins using Tween® 20 (aq) as the solvent, (ii) protein induced precipitation with ethanol, and redissolution of proteins in PBS, and (iii) bacterioruberin polishing and recycling of the solvents. Dashed lines are only a suggestion of how the process could be industrially implemented, and these steps were not experimentally performed in this work.



can be found in Table S5 in the ESI.† The molecular structures are shown in Fig. S9 in the ESI.†

Cell disruption and solid–liquid extraction. Cell disruption and solid–liquid extraction steps were performed simultaneously following an adapted protocol previously developed by us.³⁸ Briefly, this step was carried out in a shaker (IKA Trayster Digital) at room temperature (20–25 °C) under a constant vertical rotation at 50 rpm, for 45 min, protected from light exposure. Several solvents, including aqueous solutions of ILs and surfactants (cationic, anionic, non-ionic, and non-tensioactive), were screened at 100 mM to assess their capacity to release first bacterioruberin (the most valuable compound) from the biomass. The critical micelle concentration (CMC) of all tested surface-active compounds is reported in Table S5 in the ESI.† Finally, a control extraction using pure ethanol was also performed.

An additional screening of aqueous solutions of non-ionic surfactants at 250 mM was also carried out. All tests were conducted at a fixed solid–liquid ratio (SLR) of 0.1 g_{wet biomass} mL_{solvent}⁻¹, i.e. 0.1 g of wet biomass and 1 mL of the respective solvent. Finally, a centrifugation step was performed at 16 200g for 15 min in a VWR Microstar 17 centrifuge, at room temperature (20–25 °C). The resulting supernatant was recovered, and the biomass debris was discarded. All the assays were done in duplicate.

Optimization of the cell disruption/solid–liquid extraction step. A response surface methodology (RSM) was applied to simultaneously study different variables and to identify the most significant parameters and their interaction, aiming at defining the optimum conditions that maximize the yield of bacterioruberin extraction. A central composite rotatable design (CCRD-2³) was applied to the best cell disruption/solid–liquid extraction solvent in a total of 18 assays with 4 central points and axial points. The independent variables studied were the SLR (1 g_{wet biomass} mL_{solvent}⁻¹), the concentration of surfactant in water (*C*_{surf}, mM), and time of extraction (*t*, min), with the dependent variable being bacterioruberin yield of extraction (in mg_{bacterioruberin} g_{wet biomass}⁻¹). The results were statistically analyzed using Statistica® 7 software and with a 95% confidence level, following the theory proposed by Dean *et al.*⁵⁵ and Rodrigues and Lemma.⁵⁶ The real values are presented in Table S2 in the ESI.† Lastly, using the means of relative deviation (%) the optimum conditions determined were validated in triplicate.

Bacterioruberin quantification. The quantification of bacterioruberin was determined using a UV-Vis microplate reader (Synergy HT microplate reader – BioTek). The absorption spectra were measured between 350 and 700 nm and bacterioruberin content was determined by eqn (2), using calibration curves previously obtained at 494 nm for ethanolic solutions and 504 nm for aqueous solutions. The bacterioruberin standard, used to determine the calibration curves, was obtained by preparative thin-layer chromatography (TLC) in silica gel 60. A mobile phase of acetone : dichloromethane (1 : 1) was used, after the solid–liquid extraction of bacterioruberin with ethanol from *Haloflex mediterranei* ATCC 33500, and subsequent redissolution of the resulting extract in dichloromethane.

$$\text{Yield of extraction (mg}_{\text{bacterioruberin}} \text{ g}_{\text{wet biomass}}^{-1}) = \frac{[\text{bacterioruberin}] \times \text{volume}}{\text{weight}} \quad (2)$$

where “[bacterioruberin]” corresponds to the concentration of bacterioruberin in the extract (mg mL⁻¹), “volume” is the volume of solvent (mL) and “weight” is the weight of the wet cells tested (g).

Protein induced precipitation. Protein precipitation was performed using ethanol as the precipitating agent which was added to the initial extract (extract obtained from the solid–liquid extraction step). Different operational conditions, namely temperature (°C), volume of ethanol added (mL), and time for precipitation (min), were evaluated in a range of –20 to 30 °C, 0.5 to 4 mL, and 2 to 30 min, respectively. The samples were centrifuged, and two different fractions were obtained, a pellet and a supernatant, using a Thermo Scientific Heraeus Megafuge 16R centrifuge at 4700g, for 15 min. The protein recovery results were analyzed in terms of the remaining protein content present in the supernatant fraction relative to the protein content in the initial extract (in percent, %). The pellet fraction from the optimized precipitation was resuspended in 2 mL of PBS (1×, pH 7.4) for further analysis. The yield of precipitation (mg_{protein} g_{wet biomass}⁻¹) was quantified using a calibration curve previously obtained with bovine serum albumin (BSA) as a standard protein at 280 nm. Finally, the ethanol content of the supernatant fraction was evaporated to evaluate bacterioruberin content after precipitation and to further pursue its polishing.

Sodium dodecyl sulfate-polyacrylamide gel electrophoresis (SDS-PAGE). The resulting protein extract (pellet fraction resuspended in PBS) was analyzed by SDS-PAGE following the procedure of Laemmli (1970).⁵⁷ First, the sample was diluted (1 : 1) in loading buffer (4% (w/v) SDS, 20% (w/v) glycerol, 0.5 mM Tris-HCl pH 6.8, 0.02% (w/v) bromophenol blue, 3.1% (w/v) DTT) and incubated for 5 min at 95 °C for denaturation to occur. Then, it was loaded into a polyacrylamide gel along with a molecular weight marker (Amersham™ ECL™ Rainbow™ Marker – Full range) and left to run for 2 h at 110 V. Finally, to stain the proteins, incubation with BlueSafe was carried out under mild agitation for 2 h.

Bacterioruberin polishing. The bacterioruberin polishing procedure used was adapted from the literature.^{58,59} First, the water content present in the bacterioruberin extract was removed by freeze-drying the sample. Then, ethanol was added in the proportion of 1 : 20 (*V*_{bacterioruberin extract} : *V*_{ethanol}) and the solution was kept for three days at –80 °C, to induce bacterioruberin release from the micellar complex present in Tween® 20. After this period, two phases were formed, a bottom viscous surfactant-rich phase and a top ethanol-rich phase, to which the pigment migrated. The bacterioruberin presence in the top phase was further evaluated by collecting a sample, removing the ethanol, and redissolving it in a controlled volume of 0.5 mL of ethanol. In addition, a ¹H NMR analysis was also performed to assess Tween® 20 content in the top phase using a Bruker AC 30 spectrometer (250 MHz) at room temperature, where deuterium oxide was used as the solvent.



Ultra-high performance liquid chromatography coupled mass spectrometry (UHPLC-MS) analysis. After the polishing step, the bacterioruberin extract was analyzed by UHPLC-MS using a Thermo Scientific Ultimate 3000RSLC (Dionex) equipped with a Dionex UltiMate 3000 RS diode array detector and coupled to a mass spectrometer. The analysis was performed in positive mode. The separation of the compounds was carried out with a gradient elution program at a flow rate of 2 mL min⁻¹, at 30 °C, by using a Hypersil Gold C₁₈ column (100 × 2.1 mm i.d.; 1.9 μm particle diameter, Thermo Fisher). The injection volume in the UHPLC system was 5 μL and the mobile phase consisted of 0.1% formic acid in water (A) and acetonitrile (30) : methanol (70) (B).

Conclusions

In this work, the recovery of bacterioruberin and proteins from *Haloferax mediterranei* was successfully achieved using tensioactive solvents. Among the aqueous solutions of surfactants and ILs studied, Tween® 20 distinguished itself as the best. This non-ionic solvent is not only able to interact with the negatively charged head of the phospholipids allowing the release of the intracellular compounds, but also increases the solubility of bacterioruberin (hydrophobic molecule) in aqueous media, by forming micelles above the CMC, thus allowing for the design of a multi-product biorefinery process.

Next, through optimization of the process operational conditions, maximum extraction yields of bacterioruberin of 0.37 ± 0.01 mg_{bacterioruberin} g_{wet biomass}⁻¹ and proteins of 352 ± 9 mg_{protein} g_{wet biomass}⁻¹ were achieved. Furthermore, in a single protein precipitation step, over 90% of the proteins were successfully separated from bacterioruberin. Finally, it was possible to perform a bacterioruberin polishing step and propose a conceptual design of the final process.

Author contributions

Bárbara M. C. Vaz: investigation, formal analysis, validation, writing – original draft. Mariam Kholany: conceptualization, methodology, supervision, writing – review & editing. Diana C. G. A. Pinto: resources. Inês P. E. Macário: investigation, resources. Telma Veloso: investigation, resources. Tânia Caetano: resources, writing – review & editing. Joana L. Pereira: resources, writing – review & editing. João A. P. Coutinho: resources, supervision. Sónia P. M. Ventura: conceptualization, methodology, resources, writing – review & editing, supervision, funding acquisition.

Conflicts of interest

There are no conflicts to declare.

Acknowledgements

This work was developed within the scope of the project CICECO-Aveiro Institute of Materials, UIDB/50011/2020, UIDP/50011/2020 & LA/P/0006/2020, financed by national funds through the FCT/MEC (PIDDAC). The authors acknowledge

FTC/MCTES for the financial support to CESAM, UIDB/50017/2020 + UIDP/50017/2020 + LA/P/0094/2020, financed by national funds. The authors are also grateful to the FCT for the doctoral grants of M. Kholany (SFRH/BD/138413/2018), I. P. E. Macário (SFRH/BD/123850/2016), and T. Veloso (SFRH/BD/147346/2019). T. Caetano was funded by national funds (OE), through FCT, in the scope of the framework contract foreseen in the numbers 4, 5, and 6 of article 23, of the Decree-Law 57/2016, of August 29, changed by Law 57/2017, of July (CEECIND/01463/2017). The National NMR Network, funded within the framework of the National Program for Scientific Re-equipment, contract REDE/1517/RMN/2005 with funds from POCI 2010 (FEDER) and FCT, is also acknowledged.

References

- 1 R. Katakai, R. S. Chutia, M. Mishra, N. Bordoloi, R. Saikia and T. Bhaskar, in *Recent Advances in Thermo-Chemical Conversion of Biomass*, ed. A. Pandey, T. Bhaskar, M. Stöcker and R. K. Sukumaran, Elsevier, Amsterdam, 1st edn, 2015, pp. 31–74.
- 2 J. B. Zimmerman, P. T. Anastas, H. C. Erythropel and W. Leitner, *Science*, 2020, **367**, 397–400.
- 3 P. Stegmann, M. Londo and M. Junginger, *Resour., Conserv. Recycl.*, 2020, **6**, 100029.
- 4 G. P. 't Lam, M. H. Vermuë, M. H. M. Eppink, R. H. Wijffels and C. van den Berg, *Trends Biotechnol.*, 2018, **36**, 216–227.
- 5 N. K. Arora and I. Mishra, *Environ. Sustainability*, 2019, **2**, 339–342.
- 6 M. I. Khan, J. H. Shin and J. D. Kim, *Microb. Cell Fact.*, 2018, **17**, 36.
- 7 M. Giani, I. Garbayo, C. Vilchez and R. M. Martínez-Espinosa, *Mar. Drugs*, 2019, **17**, 524.
- 8 I. Hamed, F. Özogul, Y. Özogul and J. M. Regenstein, *Compr. Rev. Food Sci. Food Saf.*, 2015, **14**, 446–465.
- 9 M. N. Gupta, in *Natural-Based Polymers for Biomedical Applications*, ed. R. L. Reis, N. M. Neves, J. F. Mano, M. E. Gomes, A. P. Marques and H. S. Azevedo, Woodhead Publishing, Cambridge, 1st edn, 2008, pp. 54–84.
- 10 J. C. F. Nunes, M. R. Almeida, J. L. Faria, C. G. Silva, M. C. Neves, M. G. Freire and A. P. M. Tavares, *J. Solution Chem.*, 2022, **51**, 243–278.
- 11 N. Sharma, M. S. Farooqi, K. K. Chaturvedi, S. B. Lal, M. Grover, A. Rai and P. Pandey, *Database*, 2014, 1–9.
- 12 M. Rodrigo-Baños, I. Garbayo, C. Vilchez, M. J. Bonete and R. M. Martínez-Espinosa, *Mar. Drugs*, 2015, **13**, 5508–5532.
- 13 C. Vilchez, E. Forján, M. Cuaresma, F. Bédmar, I. Garbayo and J. M. Vega, *Mar. Drugs*, 2011, **9**, 319–333.
- 14 S. Serrano, S. Mendo and T. Caetano, *Res. Microbiol.*, 2022, **173**, 103919.
- 15 M. de la Vega, A. Sayago, J. Ariza, A. G. Barneto and R. León, *Biotechnol. Prog.*, 2016, **32**, 592–600.
- 16 R. Calegari-Santos, R. A. Diogo, J. D. Fontana and T. M. B. Bonfim, *Curr. Microbiol.*, 2016, **72**, 641–651.
- 17 M. Giani, Z. Montero-Lobato, I. Garbayo, C. Vilchez, J. M. Vega and R. M. Martínez-Espinosa, *Mar. Drugs*, 2021, **19**, 100.



- 18 K. L. Britton, P. J. Baker, M. Fisher, S. Ruzhenikov, D. J. Gilmour, M.-J. Bonete, J. Ferrer, C. Pire, J. Esclapez and D. W. Rice, *Proc. Natl. Acad. Sci. U. S. A.*, 2006, **103**, 4846–4851.
- 19 M. Mevarech, F. Frolow and L. M. Gloss, *Biophys. Chem.*, 2000, **86**, 155–164.
- 20 G. Payá, V. Bautista, M. Camacho, M.-J. Bonete and J. Esclapez, *Biochimie*, 2021, **187**, 33–47.
- 21 C. Ebel, D. Madern and G. Zaccai, *Extremophiles*, 2009, **2**, 278–297.
- 22 A. Kumar, A. Alam, D. Tripathi, M. Rani, H. Khatoon, S. Pandey, N. Z. Ehtesham and S. E. Hasnain, *Semin. Cell Dev. Biol.*, 2018, **84**, 147–157.
- 23 A. Siglioccolo, A. Paiardini, M. Piscitelli and S. Pascarella, *BMC Struct. Biol.*, 2011, **11**, 50.
- 24 M. Abbes, H. Baati, S. Guermazi, C. Messina, A. Santulli, N. Gharsallah and E. Ammar, *BMC Complementary Altern. Med.*, 2013, **13**, 255.
- 25 L. Zalazar, P. Pagola, M. V. Miró, M. S. Churio, M. Cerletti, C. Martínez, M. Iniesta-Cuerda, A. J. Soler, A. Cesari and R. De Castro, *J. Appl. Microbiol.*, 2019, **126**, 796–810.
- 26 W. J. Müller, M. S. Smit, E. van Heerden, M. D. Capes and S. DasSarma, *Front. Microbiol.*, 2018, **9**, 2563.
- 27 J. J. Alvares and I. J. Furtado, *J. Anal. Sci. Technol.*, 2021, **12**, 44.
- 28 M. Manikandan, N. Hasan and H.-F. Wu, *Talanta*, 2013, **107**, 167–175.
- 29 S. Mishra, C. S. Chanotiya, K. Shanker and A. K. Tripathi, *FEMS Microbiol. Lett.*, 2021, **368**, 5.
- 30 L. Higa Herminia, P. Schilrreff, A. Martín Briski, H. Emanuel Jerez, M. Alexandre de Farias, R. Villares Portugal, E. Lilia Romero and M. Jose Morilla, *Colloids Surf., B*, 2020, **191**, 110961.
- 31 F. I. Kesbiç and N. Gültepe, *Folia Microbiol.*, 2022, **67**, 71–79.
- 32 J. Biswas, F. N. Haque and A. K. Paul, *J. Adv. Microbiol.*, 2016, **1**, 1–11.
- 33 S. Ghosh, A. Ray and N. Pramanik, *Biophys. Chem.*, 2020, **265**, 106429.
- 34 Y. Li, F. Fang, M. Sun, Q. Zhao, Y. Hu, Z. Sui, Z. Liang, L. Zhang and Y. Zhang, *Talanta*, 2020, **213**, 120848.
- 35 D. Danwanichakul, O. Rattanaphan, J. Srisatjang and P. Danwanichakul, *J. Appl. Polym. Sci.*, 2014, **131**, 39900.
- 36 X. Lin, Y. Wang, Q. Zeng, X. Ding and J. Chen, *Analyst*, 2013, **138**, 6445–6453.
- 37 M. Martins, W. C. Ooi, M. C. Neves, J. F. Pereira and J. A. Coutinho, *J. Chem. Technol. Biotechnol.*, 2018, **93**, 1864–1870.
- 38 M. Kholany, P. Trébulle, M. Martins, S. P. M. Ventura, J.-M. Nicaud and J. A. P. Coutinho, *J. Chem. Technol. Biotechnol.*, 2020, **95**, 1126–1134.
- 39 M. Martins, L. M. de S. Mesquita, B. M. C. Vaz, A. C. R. V. Dias, M. A. Torres-Acosta, B. Quéguineur, J. A. P. Coutinho and S. P. M. Ventura, *ACS Sustainable Chem. Eng.*, 2021, **9**, 6599–6612.
- 40 M. Martins, A. P. M. Fernandes, M. A. Torres-Acosta, P. N. Collén, M. H. Abreu and S. P. M. Ventura, *Sep. Purif. Technol.*, 2021, **254**, 117589.
- 41 L. M. de S. Mesquita, M. Martins, É. Maricato, C. Nunes, P. S. G. N. Quinteiro, A. C. R. V. Dias, J. A. P. Coutinho, L. P. Pisani, V. V. de Rosso and S. P. M. Ventura, *ACS Sustainable Chem. Eng.*, 2020, **8**, 4085–4095.
- 42 R. R. Burgess, in *Methods in Enzymology*, ed. R. R. Burgess and M. P. Deutscher, Elsevier, Amsterdam, 2nd edn, 2009, vol. 463, pp. 331–342.
- 43 D. J. Bell, M. Hoare and P. Dunnill, in *Downstream Processing, Advances in Biochemical Engineering/Biotechnology*, ed. A. Fiechter, Springer, Berlin, Heidelberg, 1st edn, 1983, pp. 1–72.
- 44 M. Martins, B. P. Soares, J. H. P. M. Santos, P. Bharmoria, M. A. Torres Acosta, A. C. R. V. Dias, J. A. P. Coutinho and S. P. M. Ventura, *ACS Sustainable Chem. Eng.*, 2021, **9**, 3942–3954.
- 45 M. J. Castro-Alonso, L. E. Montañez-Hernandez, M. A. Sanchez-Muñoz, M. R. M. Franco, R. Narayanasamy and N. Balagurusamy, *Front. Mater.*, 2019, **6**, 126.
- 46 F. A. Vieira, R. J. R. Guilherme, M. C. Neves, H. Abreu, E. R. O. Rodrigues, M. Maraschin, J. A. P. Coutinho and S. P. M. Ventura, *Sep. Purif. Technol.*, 2017, **172**, 268–276.
- 47 M. Martins, F. A. Vieira, I. Correia, R. A. S. Ferreira, H. Abreu, J. A. P. Coutinho and S. P. M. Ventura, *Green Chem.*, 2016, **18**, 4287–4296.
- 48 J. M. Newton, D. Schofield, J. Vlahopoulou and Y. Zhou, *Biotechnol. Prog.*, 2016, **32**, 1069–1076.
- 49 B. Balasundaram, D. Nesbeth, J. M. Ward, E. Keshavarz-Moore and D. G. Bracewell, *Biotechnol. Bioeng.*, 2009, **104**, 134–142.
- 50 M. Mohsen-Nia, H. Amiri and B. Jazi, *J. Solution Chem.*, 2010, **39**, 701–708.
- 51 C. J. van Oss, *J. Protein Chem.*, 1989, **8**, 661–668.
- 52 S. Taskila, M. Ahokas, J. Järvinen, J. Toivanen and J. P. Tanskanen, *Scientifica*, 2017, 5120947.
- 53 G. Schilcher, A. Schlagenhauf, D. Schneditz, H. Scharnagl, W. Ribitsch, R. Krause, A. R. Rosenkranz, T. Stojakovic and J. H. Horina, *PLoS One*, 2013, **8**, e84869.
- 54 T. Allers, H.-P. Ngo, M. Mevarech and R. G. Lloyd, *Appl. Environ. Microbiol.*, 2004, **70**, 943–953.
- 55 A. Dean and D. Voss, *Design and Analysis of Experiments*, Springer, New York, 1st edn, 1999.
- 56 M. I. Rodrigues and A. F. Lemma, *Planejamento de Experimentos e Otimização de Processos*, Casa do Pão Editora, 3rd edn, 2014.
- 57 U. K. Laemmli, *Nature*, 1970, **227**, 680–685.
- 58 L. M. De Souza Mesquita, S. P. M. Ventura, A. R. C. Braga, L. P. Pisani, A. C. R. V. Dias and V. V. De Rosso, *Green Chem.*, 2019, **21**, 2380–2391.
- 59 B. M. C. Vaz, M. Martins, L. M. de Souza Mesquita, A. P. M. Fernandes, D. C. G. A. Pinto, M. G. P. M. S. Neves, M. C. Neves, J. A. P. Coutinho and S. P. M. Ventura, *Chem. Eng. J.*, 2022, **428**, 131073.

