


 Cite this: *Chem. Commun.*, 2023, 59, 4915

 Received 11th February 2023,  
 Accepted 27th March 2023

DOI: 10.1039/d3cc00637a

rsc.li/chemcomm

When the basic ligand **3** was added to the heteroleptic three-component slider-on-deck  $[Ag_3(1)(2)]^{3+}$  (sliding frequency  $k_{298} = 57$  kHz), it operated as a moderate brake pad ( $k_{298} = 45$  kHz). Due to motion in the resulting four-component slider-on-deck  $[Ag_3(1)(2)(3)]^{3+}$ , both ligand **3** and silver(I) were continuously exposed and became catalytically active in a concurrent tandem Michael addition/hydroalkoxylation.

Inside multicellular organisms, enzymes catalyze a gamut of reactions, ranging from simple to cascaded<sup>1</sup> and concurrent tandem reactions.<sup>2</sup> In many cases, enzymatic activity critically depends on protein mobility, *e.g.*, for substrate binding or product release.<sup>3</sup> If enzymes with multiple components are concerned, then mobility may even involve cooperative motions of distinct parts for achieving rate and efficiency increase,<sup>4,5</sup> as convincingly demonstrated by the ATP synthase.<sup>6</sup>

Only recently, chemists have been able to mimic such spectacular enzyme capabilities by developing multicomponent catalysts,<sup>7–9</sup> but few have succeeded to connect catalytic function with intrinsic motion.<sup>10</sup> Herein, we illustrate the suitability of four-component machinery to act as a dual catalytic effector for concurrent tandem catalysis with both processes occurring in one solution. Dual catalytic activity of a single slider-on-deck – instigated only by its dynamics – has not yet been realized.<sup>11</sup>

In earlier work, we demonstrated that the superior action of nanorotors and slider-on-deck systems as catalysts in both base-<sup>12</sup> or metal-catalyzed<sup>13</sup> reactions depended crucially on their motional speed. Notably, the speed correlated with the ability of the machinery to free the catalyst<sup>12</sup> or product<sup>13</sup> (reducing product inhibition). Such effect is conceptually

# Concurrent tandem catalysis enabled by nanomechanical motion in heteroleptic four-component dual-catalyst machinery†

 Emad Elramadi,<sup>‡</sup> Sohom Kundu,<sup>‡</sup> Debabrata Mondal and Michael Schmittl<sup>\*,†</sup>

different from static liberation of a catalyst as used in switchable catalysis.<sup>14,15</sup>

In detail, the four-component slider-on-deck was constructed from deck **1**,<sup>16</sup> biped **2**, silver(I) ions and the chelating base **3** as shown in Scheme 1 (for synthesis, see ESI,<sup>†</sup> Chapter 1). The resulting catalytic machinery  $[Ag_3(1)(2)(3)]^{3+}$  exhibited a stochastic sliding motion that in parallel liberated the silver(I) sites and base **3** for catalysis. The resulting dual-catalyst machinery proved to be



**Scheme 1** (a) Chemical structure of ligands **1–3**. (b) Transformation of the three- into the four-component dual-catalyst machinery. Both organobase **3** and the silver(I) centers are liberated due to the sliding motion in  $[Ag_3(1)(2)(3)]^{3+}$ , which enables concurrent tandem catalysis. A & B represent the substrates, C denotes the intermediate product and D the final product of the concurrent tandem reaction.

Center of Micro and Nanochemistry and (Bio)Technology, Organische Chemie I, School of Science and Technology, University of Siegen, Adolf-Reichwein-Str. 2, Siegen D-57068, Germany. E-mail: schmittl@chemie.uni-siegen.de;

Tel: +49(0) 2717404356

† Electronic supplementary information (ESI) available: Experimental procedures, characterization, VT-NMR, spectral data. See DOI: <https://doi.org/10.1039/d3cc00637a>

‡ Emad Elramadi and Sohom Kundu contributed equally.







Fig. 2 Comparison of partial  $^1\text{H}$ -NMR spectra (600 MHz,  $\text{CD}_2\text{Cl}_2$ , 298 K) of **3**, silver(I) phenanthroline complex  $[\text{Ag}(\mathbf{3})(\mathbf{11})]^+$ , three-component slider-on-deck  $[\text{Ag}_3(\mathbf{1})(\mathbf{2})(\mathbf{3})]^{3+}$ , and four-component slider-on-deck  $[\text{Ag}_3(\mathbf{1})(\mathbf{2})(\mathbf{3})]^{3+}$ .

both slider-on-deck systems, one can deduce that  $[\text{Ag}_3(\mathbf{1})(\mathbf{2})(\mathbf{3})]^{3+}$  is also in motion despite the addition of ligand **3**. By comparison of the shifts of proton signals  $i'$ ,  $j'$ -H in **3** ( $\delta_{i',j'} = 3.75, 3.83$  ppm),  $[\text{Ag}_3(\mathbf{1})(\mathbf{2})(\mathbf{3})]^{3+}$  ( $\delta_{i',j'} = 3.33$  ppm) and reference  $[\text{Ag}(\mathbf{3})(\mathbf{11})]^+$  ( $\delta_{i',j'} = 3.21$  ppm), similar upfield shifts were observed for both complexes attesting that **3** is situated in the shielding pocket of the silver(I) phenanthroline sites (Fig. 2). As expected from product liberation in moving rotors,<sup>13</sup> ligand **3** in  $[\text{Ag}_3(\mathbf{1})(\mathbf{2})(\mathbf{3})]^{3+}$  shows a shift that lies in between those of free **3** and  $[\text{Ag}(\mathbf{3})(\mathbf{11})]^+$ , indicating that some is liberated by the motion in the four-component slider-on-deck.

By means of variable temperature (VT)  $^1\text{H}$  NMR spectroscopy, we determined the exchange frequency of both slider-on-deck assemblies. In case of the three-component  $[\text{Ag}_3(\mathbf{1})(\mathbf{2})]^{3+}$ , the 4-H proton signal split at  $-60$  °C into two peaks (ratio 2 : 1) due to freezing the sliding of biped **2** across deck **1** on the NMR time scale (Fig. 3). Using WinDNMR,<sup>18</sup> the  $^1\text{H}$  NMR traces were simulated over a large temperature range, providing the activation data as  $\Delta H^\ddagger = 45.8$  kJ mol $^{-1}$ ,  $\Delta S^\ddagger = 0.7$  J mol $^{-1}$  K $^{-1}$  and  $\Delta G_{298}^\ddagger = 45.6$  kJ mol $^{-1}$  as well as the exchange rate  $k_{298} = 57$  kHz. Contrastingly, in the four-component slider-on-deck  $[\text{Ag}_3(\mathbf{1})(\mathbf{2})(\mathbf{3})]^{3+}$ , the proton 4-H signal split already at  $-40$  °C. The kinetic analysis afforded  $k_{298} = 45$  kHz along with the activation data  $\Delta H^\ddagger = 47.8$  kJ mol $^{-1}$ ,  $\Delta S^\ddagger = 5.4$  J mol $^{-1}$  K $^{-1}$  and  $\Delta G_{298}^\ddagger = 46.3$  kJ mol $^{-1}$ . As a result of the addition of **3** to  $[\text{Ag}_3(\mathbf{1})(\mathbf{2})]^{3+}$ , the motion thus slowed down by 21% at room temperature and even more at lower temperature. This trend is



Fig. 3 Experimental (left) and simulated (right) partial VT  $^1\text{H}$  NMR spectra ( $\text{CD}_2\text{Cl}_2$ , 600 MHz) of (a)  $[\text{Ag}_3(\mathbf{1})(\mathbf{2})]^{3+}$  and (b) of  $[\text{Ag}_3(\mathbf{1})(\mathbf{2})(\mathbf{3})]^{3+}$ .

Table 2 Product yield in both concurrent and sequential reactions catalyzed by  $[\text{Ag}_3(\mathbf{1})(\mathbf{2})(\mathbf{3})]^{3+}$

Time (h)	Yield of <b>6</b> (base cat.) (%)	Yield of <b>8</b> ( $\text{Ag}^+$ cat.) (%)	Yield of <b>10</b> (dual seq. cat.) (%)
0	0	0	0
2	7	17	2
4	18	35	7
6	25	47	12
8	29	53	15
10	33	61	21
12	35	64	27
14	38	67	34

a result of the higher positive activation entropy of  $[\text{Ag}_3(\mathbf{1})(\mathbf{2})(\mathbf{3})]^{3+}$  that is increased due to the liberation of **3** into solution in the rate-determining step. As in the rate-limiting step the biped must depart from one silver(I) binding site to kick out the base **3**, both the silver(I) and **3** are temporarily available for catalysis.

To investigate whether both catalytic units in  $[\text{Ag}_3(\mathbf{1})(\mathbf{2})(\mathbf{3})]^{3+}$  are simultaneously active we tested for concurrent base and silver(I) catalysis. Thus, the slider-on-deck  $[\text{Ag}_3(\mathbf{1})(\mathbf{2})(\mathbf{3})]^{3+}$  and reactants **4**, **5**, and **7** were combined (1 : 30 : 30 : 30) in  $\text{CD}_2\text{Cl}_2$  for running both transformations parallel (Scheme 2a). The reactions were monitored by  $^1\text{H}$  NMR over 14 h at room temperature, showing formation of the silver(I)-catalyzed product **8** (67%) and base-catalyzed product **6** (38%) as displayed in Table 2. The data demonstrate that the base-catalyzed formation of **6** is slower than the silver-catalyzed reaction affording **8**.

Since dual activity of this slider-on-deck is increased in comparison with the static model compounds, we probed a concurrent tandem catalysis consisting of a Michael addition followed by hydroalkoxylation (see Scheme 2b). Hence, the slider-on-deck  $[\text{Ag}_3(\mathbf{1})(\mathbf{2})(\mathbf{3})]^{3+}$  and reactants **4** & **9** (1 : 30 : 30) were mixed in  $\text{CD}_2\text{Cl}_2$ . The kinetic profile of this reaction was monitored at room temperature by  $^1\text{H}$  NMR over 14 h at 2 h intervals.  $^1\text{H}$  NMR spectra showed 34% of **10** after 14 h (black curve in Fig. 4).

Closer inspection revealed several surprising features. Initially, the base catalyzed reaction is the rate-determining step in the tandem reaction to **10**, because intermediate **10'** created by




Fig. 4 Yield of **10** with time in the tandem reaction catalysed by  $[\text{Ag}_3(\mathbf{1})(\mathbf{2})(\mathbf{3})]^{3+}$ .



base catalysis does not show up in the  $^1\text{H}$  NMR. Only, when product **10** of the tandem base- and silver(i) catalysis had formed to more than 15%, then intermediate **10'** (Scheme 2b) became visible. Furthermore, the kinetic profile of the tandem reaction showed an increasing rate the more product **10** had formed (Fig. 4), indicative of some autocatalytic effect. To check whether it is the final product **10** that impacts on the rate, the slider-on-deck  $[\text{Ag}_3(\mathbf{1})(\mathbf{2})(\mathbf{3})]^{3+}$ , reactants **4**, **9** and product **10** were mixed in a 1:30:30:3 ratio in  $\text{CD}_2\text{Cl}_2$ . Now 59% of **10** (49% after subtracting 10% of the initially added **10**) was afforded after 14 h with higher rate (initial rate at  $t = 0$ :  $\nu_0 = 2.8 \mu\text{M s}^{-1}$ ) than in absence of product **10** ( $\nu_0 = 2.1 \mu\text{M s}^{-1}$ ) (Fig. 4). This effect is attributed to the competitive displacement of base **3** at the silver(i)-loaded phenanthroline by product **10** as it may act as ligand as well. Thus, in presence of sliding motion and 1 equiv. of **10**, more base **3** is released.

In conclusion, we report a three-component slider-on-deck assembly  $[\text{Ag}_3(\mathbf{1})(\mathbf{2})]^{3+}$  with a sliding speed of  $k_{298} = 57 \text{ kHz}$ . Addition of amine **3** to the slider-on-deck converted it into a four-component dual-catalyst machinery  $[\text{Ag}_3(\mathbf{1})(\mathbf{2})(\mathbf{3})]^{3+}$  with a reduced sliding speed ( $k_{298} = 45 \text{ kHz}$ ). The sliding motion in the machinery instigated two simultaneous events that enabled concurrent tandem catalysis: (a) liberation of base **3**, and (b) exposure of the silver(i) sites, so that they could act synergistically<sup>19</sup> as dual catalyst in the two-step synthesis of **10**. Thus, the present system portrays a lucid example of unleashing multiple catalytic functions instigated by a single nanomechanical motion within artificial catalytic machinery.

We thank the DFG (Schm 647/22-1, no. 491092614) and the Univ. of Siegen for continued support. Moreover, we are indebted to Dr Paululat (Siegen) for the VT- $^1\text{H}$  NMR measurements.

## Conflicts of interest

There are no conflicts to declare.

## Notes and references

- 1 Y. Huang, A. M. Walji, C. H. Larsen and D. W. C. MacMillan, *J. Am. Chem. Soc.*, 2005, **127**, 15051–15053.

- 2 J.-C. Wasilke, S. J. Obrey, R. T. Baker and G. C. Bazan, *Chem. Rev.*, 2005, **105**, 1001–1020.
- 3 P. Ojeda-May, A. U. Mushtaq, P. Rogne, A. Verma, V. Ovchinnikov, C. Grundström, B. Dulko-Smith, U. H. Sauer, M. Wolf-Watz and K. Nam, *Biochemistry*, 2021, **60**, 2246–2258.
- 4 M. J. Wiester, P. A. Ulmann and C. A. Mirkin, *Angew. Chem., Int. Ed.*, 2011, **50**, 114–137.
- 5 K. A. Henzler-Wildman, V. Thai, M. Lei, M. Ott, M. Wolf-Watz, T. Fenn, E. Pozharski, M. A. Wilson, G. A. Petsko, M. Karplus, C. G. Hübner and D. Kern, *Nature*, 2007, **450**, 838–844.
- 6 P. Neupane, S. Bhuju, N. Thapa and H. K. Bhattarai, *Biomol. Concepts*, 2019, **10**, 1–10.
- 7 (a) Z. Dong, Q. Luo and J. Liu, *Chem. Soc. Rev.*, 2012, **41**, 7890–7908; (b) M. Raynal, P. Ballester, A. Vidal-Ferran and P. W. N. M. van Leeuwen, *Chem. Soc. Rev.*, 2014, **43**, 1734–1787; (c) D. Zhang, T. K. Ronson and J. R. Nitschke, *Acc. Chem. Res.*, 2018, **51**, 2423–2436; (d) N. Mittal, M. S. Özer and M. Schmittel, *Inorg. Chem.*, 2018, **57**, 3579–3586; (e) F. J. Rizzuto, L. K. S. von Krbeke and J. R. Nitschke, *Nat. Rev. Chem.*, 2019, **3**, 204–222; (f) G. Olivo, G. Capocasa, D. Del Giudice, O. Lanzalunga and S. Di Stefano, *Chem. Soc. Rev.*, 2021, **50**, 7681–7724; (g) P. Howlader and M. Schmittel, *Beilstein J. Org. Chem.*, 2022, **18**, 597–630; (h) A. W. Heard, J. M. Suárez and S. M. Goldup, *Nat. Rev. Chem.*, 2022, **6**, 182–196; (i) R. Saha, B. Mondal and P. S. Mukherjee, *Chem. Rev.*, 2022, **122**, 12244–12307.
- 8 *Supramolecular Catalysis: New Directions and Developments*, ed. P. W. N. M. van Leeuwen and M. Raynal, Wiley-VCH GmbH, Weinheim, 1st edn, 2022, ISBN: 9783527349029.
- 9 (a) N. Mittal, S. Pramanik, I. Paul, S. De and M. Schmittel, *J. Am. Chem. Soc.*, 2017, **139**, 4270–4273; (b) Y. Li, X. Caumes, M. Raynal and L. Bouteiller, *Chem. Commun.*, 2019, **55**, 2162–2165; (c) E. Iniesta and A. Vidal-Ferran, *Chem. Commun.*, 2020, **56**, 6364–6367.
- 10 L. van Dijk, M. J. Tilby, R. Szpera, O. A. Smith, H. A. P. Bunce and S. P. Fletcher, *Nat. Rev. Chem.*, 2018, **2**, 1–18.
- 11 V. V. Rajasekaran, A. Ghosh, S. Kundu, D. Mondal, T. Paululat and M. Schmittel, *Angew. Chem., Int. Ed.*, 2022, **61**, e2022124.
- 12 I. Paul, A. Goswami, N. Mittal and M. Schmittel, *Angew. Chem., Int. Ed.*, 2018, **57**, 354–358.
- 13 P. K. Biswas, S. Saha, T. Paululat and M. Schmittel, *J. Am. Chem. Soc.*, 2018, **140**(29), 9038–9041.
- 14 S. Gaikwad, A. Goswami, S. De and M. Schmittel, *Angew. Chem., Int. Ed.*, 2016, **55**, 10512–10517.
- 15 S. De, S. Pramanik and M. Schmittel, *Angew. Chem., Int. Ed.*, 2014, **53**, 14255–14259.
- 16 E. Elramadi, A. Ghosh, I. Valiyev, P. K. Biswas, T. Paululat and M. Schmittel, *Chem. Commun.*, 2022, **58**, 8073–8076.
- 17 U. Kaya, P. Chauhan, K. Deckers, R. Puttreddy, K. Rissanen, G. Raabe and D. Enders, *Synthesis*, 2016, 3207–3216.
- 18 H. J. Reich, *NMR Spectrum Calculations: WinDNMR, Version 7.1.13*, Department of Chemistry, University of Wisconsin.
- 19 L. Zhou and L. Liu, *Chem. Commun.*, 2021, **57**, 5690–5693.

