

Chemical Science

Volume 15
Number 15
21 April 2024
Pages 5405–5788

rsc.li/chemical-science



ISSN 2041-6539

PERSPECTIVE

Christophe Léger *et al.*
Outer-sphere effects on the O₂ sensitivity, catalytic bias and
catalytic reversibility of hydrogenases

Cite this: *Chem. Sci.*, 2024, 15, 5418

All publication charges for this article have been paid for by the Royal Society of Chemistry

Received 29th January 2024
Accepted 15th March 2024

DOI: 10.1039/d4sc00691g

rsc.li/chemical-science

Outer-sphere effects on the O₂ sensitivity, catalytic bias and catalytic reversibility of hydrogenases†

Andrea Fasano,^{ID} Vincent Fourmond^{ID} and Christophe Léger^{ID}*

The comparison of homologous metalloenzymes, in which the same inorganic active site is surrounded by a variable protein matrix, has demonstrated that residues that are remote from the active site may have a great influence on catalytic properties. In this review, we summarise recent findings on the diverse molecular mechanisms by which the protein matrix may define the oxygen tolerance, catalytic directionality and catalytic reversibility of hydrogenases, enzymes that catalyse the oxidation and evolution of H₂. These mechanisms involve residues in the second coordination sphere of the active site metal ion, more distant residues affecting protein flexibility through their side chains, residues lining the gas channel and even accessory subunits. Such long-distance effects, which contribute to making enzymes efficient, robust and different from one another, are a source of wonder for biochemists and a challenge for synthetic bioinorganic chemists.

Introduction


Metalloenzymes that use active sites based on transition metals to catalyse the production or consumption of small molecules such as H₂, O₂, CO₂ and N₂ have attracted much interest over the last few decades because of the need for cheap and efficient synthetic catalysts for the production and use of solar fuels. There is much hope that the knowledge acquired by characterising these enzymes will be useful to design efficient catalysts. Independently of these technological challenges, metalloenzymes are also useful as model systems, to observe

the limits of what chemistry can achieve, and learn about how Nature does it.

Hydrogenases are enzymes that use sophisticated inorganic active sites made of transition metals, such as iron and nickel, to catalyse the conversion between molecular hydrogen and protons. Two main classes of hydrogenases exist, named after the metal content of their inorganic active sites: the so-called FeFe hydrogenases have an active site, called the H-cluster, that consists of a dinuclear cluster of Fe covalently attached by a cysteine sulphur to a [4Fe4S] cluster (Fig. 1A); NiFe hydrogenases use a dinuclear cluster of Ni and Fe that is attached to the protein by four cysteine residues (Fig. 1B). These

Laboratoire de Bioénergétique et Ingénierie des Protéines, CNRS, Aix Marseille Université, UMR 7281, Marseille, France. E-mail: leger@imm.cnrs.fr

† Electronic supplementary information (ESI) available. See DOI: <https://doi.org/10.1039/d4sc00691g>



Christophe Léger (left),
Vincent Fourmond
(middle) and Andrea Fasano (right)

Dr Christophe Léger (left) obtained his PhD in physical chemistry from the University of Bordeaux and was a postdoc in the group of Fraser Armstrong. Dr Vincent Fourmond (middle) obtained his PhD in the interface of physics and biology from Université Paris Diderot in 2007. He held postdoctoral positions first in the group of Christophe Léger in Marseille and then Vincent Artero in Grenoble, before coming back to Marseille as a permanent CNRS researcher. CL and VF are both Directeurs de Recherche at CNRS. Dr Andrea Fasano (right) obtained a BSc in 2017 in Biotechnology at the Alma Mater Studiorum university of Bologna, an MSc in Industrial Biotechnology at the University of Padua in 2019, and a PhD at the University of Aix Marseille in 2023. In 2024, he was awarded the PhD prizes of the regional section of the French Chemical Society and of the national CNRS research group on bioinorganic chemistry (FrenchBIC).




Fig. 1 The active sites of FeFe and NiFe hydrogenases, in panels (A) and (B) respectively. In panel (A), the proximal and distal Fe ions are labelled. Colour code: S (yellow), Fe (orange), C (white), O (red), N (blue) and Ni (green). PDB: 4XDC for the FeFe active site and 3UQY for the NiFe active site.

active sites are produced and inserted into the apo-enzyme by dedicated enzymes that are part of a complex maturation machinery. The active site of most hydrogenases is buried in the protein matrix and connected to the solvent by a chain of accessory FeS clusters and a network of acidic residues and water molecules, which mediate long range electron and proton transfers. Preferred pathways (sometimes called “gas channels”) for the diffusion of the substrate H_2 and inhibitors (CO , O_2 , H_2S , and Cl^-) to and from the active site have also been identified with varying degrees of confidence.

Dihydrogen is a very important energy vector in the microbial world: it is produced as a by-product of fermentation and

nitrogen fixation,^{1,2} and reoxidized in various metabolic pathways, related to *e.g.* respiration (ATP production)³ or the non-photosynthetic reduction of CO_2 . As a consequence, most microorganisms produce at least one hydrogenase, but many produce several: *Escherichia coli* has 4 NiFe hydrogenases,⁴ and *Solidesulfovibrio fructosivorans* (Sf, previously known as *Desulfovibrio fructosovorans*, a model sulfate reducing bacterium) expresses 2 NiFe and 4 FeFe hydrogenases.⁵ The enzymes in each of the two families are called “homologous”: they embed the same active site (either FeFe or NiFe), but they have similar but distinct amino acid sequences, and therefore distinct structures and a variable number of accessory FeS clusters. They have been classified into groups based on phylogenetic and functional analyses.^{6,7} Some are monomeric, others may be part of membrane bound or multifunctional protein complexes from which they can be dissociated.

Hydrogenases were discovered nearly a century ago⁸ and extensively studied using biophysical methods over the last fifty years,^{9–11} but their biodiversity has been explored only recently.^{12,13} At the turn of the XXIst century, only about half a dozen of very similar hydrogenases had been used as model enzymes for mechanistic studies. Over the last twenty years, their number has rapidly increased. This work was probably mostly driven by the desire to use hydrogenases for H_2 oxidation (in fuel cells) or H_2 production (*e.g.* in photoelectrochemical

Table 1 The FeFe hydrogenases discussed in this work

Group ⁶	Source	Short name	Structure	Properties
A	<i>Chlamydomonas reinhardtii</i>	Cr HydA1	Group M1 (ref. 7) (no accessory FeS cluster), Fig. 3A, pdb 3LX4 (apo form)	Bidirectional (Fig. 2A), O_2 -sensitive, damaged by UV B ¹⁹ , readily inhibited by HS^- , ^{16,82} Cl^- ¹⁷
	<i>Desulfovibrio desulfuricans</i>	Dd	Group M2 (ref. 7) (two accessory clusters), dimeric, Fig. 3A, pdb 1HFE	Bidirectional, damaged by visible light, ^{18,125} readily inhibited by HS^- ⁸²
	<i>C. pasteurianum</i> I	CpI	Group M3 (ref. 7) (four accessory clusters forming a Y-shaped electron transfer conduit ¹²⁶), Fig. 3A, pdb 4XDC	Bidirectional
	<i>Clostridium acetobutylicum</i>	CaI	Group M3	Bidirectional (Fig. 2B), damaged by UV B ¹⁹ , reacts slowly with inhibitors (HS^- , Cl^- , CO and O_2)
	<i>Megasphaera elsdenii</i> I	MeI	Group M2, monomeric	Bidirectional, damaged by UV B ¹⁹ , reacts slowly with inhibitors (CO and O_2) ⁶²
	<i>Acetobacterium woodii</i>		M2-type hydrogenase associated with a formate dehydrogenase	Bidirectional, reacts slowly with O_2 (ref. 63)
	<i>C. pasteurianum</i> II	CpII	Group M2	Catalyses H_2 oxidation and evolution irreversibly (Fig. 2D) ²⁴
B2	<i>C. beijerinckii</i>	Cb	3 accessory clusters (incl. one in a SLBB domain), Fig. 3A, pdb 6TTL	Bidirectional (Fig. 2E), oxygen-stable ^{25,26,80}
	<i>C. pasteurianum</i> III	CpIII	M2, subgroup B2 because they bear a TSCCCP motif ²⁷	Bidirectional ²⁴ (Fig. 2G), O_2 -sensitive ²⁷
C	<i>Megasphaera elsdenii</i> II	MeII		Bidirectional (Fig. 2G) ²⁷
D	<i>Thermotoga maritima</i>	Tm HydS	Proton transfer pathway different from that in group A ^{111,127}	Bidirectional, very low activity, catalyses H_2 oxidation and evolution irreversibly (Fig. 2C) ¹²⁷
	<i>Thermoanaerobacter mathranii</i>	Tam HydS	Proton transfer pathway different from that in group A (Fig. 3D) ^{36,128}	Bidirectional, very low activity, catalyses H_2 oxidation and evolution irreversibly ^{23,33}



Table 2 The NiFe hydrogenases discussed in this work

Group		Structure	pdb	
1b	<i>Desulfovibrio gigas</i> <i>D. vulgaris</i> Miyazaki F <i>Solidesulfovibrio fructosivorans</i> , (formerly <i>Desulfovibrio fructosovorans</i>)	Dg Sf Hyn	Dimeric form (Fig. 4A), large subunit with the NiFe active site (Fig. 1B), small subunit with three FeS clusters: a 4-cysteiny [4Fe4S] cluster proximal to the active site (Fig. 4C); medial, high redox potential [3Fe4S] cluster; distal [4Fe4S] cluster coordinated by 3 cysteines and one histidine (Fig. 4F)	1FRV 4U9H 1FRF
1c	<i>Escherichia coli</i> hydrogenase 2	Hyd 2		6EN9
1e	<i>Allochrochromatium vinosum</i>	Av		3MYR
1a	<i>Desulfovibrio vulgaris</i> Hildenborough <i>D. baculatum</i>	DvH	Dimer. three cysteine and one selenocysteine ligands ¹³⁰ in the 1st coordination sphere of the Ni ion, instead of four cysteines. A medial [4Fe4S] cluster in the small subunit	5JSH 1CC1
1d	<i>Cupriavidus necator</i> (formerly <i>R. eutropha</i>) membrane bound hydrogenase <i>E. coli</i> ^{14,70} <i>Aquifex aeolicus</i> hydrogenase I ¹³⁵ <i>Cupriavidus metallidurans</i> (formerly <i>Ralstonia metallidurans</i>) <i>Salmonella enterica</i> hydrogenase 5	Cn MBH Ec Hyd 1 Aa h2ase I Cm Se Hyd 5 (ref. 42)	Small subunit with three FeS clusters, incl. a unique [3Fe4S] proximal cluster, coordinated by 6 cysteine residues ^{43–45} (Fig. 4D)	3RGW 3UQY 4C3O
1h	<i>C. necator</i> actinobacterial-type hydrogenase	Cn AH	Small subunit with three FeS clusters, incl. the proximal [4Fe4S] cluster coordinated by three cysteines and one aspartate (Fig. 4E) ⁴⁶	5AA5
2a	<i>Mycobacterium smegmatis</i>	HucSL	Embeds three [3Fe4S] clusters in each HucSL heterodimer	8DQV
2c	<i>C. necator</i> regulatory hydrogenase RH ⁴⁰ <i>Rhodobacter capsulatus</i> regulatory hydrogenase ⁴¹	Cn RH		
3d	<i>C. necator</i> soluble hydrogenases	Cn SH	The initial hypothesis that the active site of Cn SH is coordinated to four (rather than 2) cyanides ¹³⁶ was ruled out ¹³⁷	

Bidirectional (see *e.g.* green CV in Fig. 6A,⁷⁰ red in Fig. 6B,^{21,47} although Av is strongly biased towards H₂ oxidation¹²⁹) and O₂-sensitive (grey in Fig. 5B): inhibition by O₂ produces a mixture of inactive states, two of which are EPR active (the signatures are called NiA and NiB, the former reactivates more slowly than the latter^{51,52})

Bidirectional. React with O₂ to produce EPR-silent inactive states^{131,132}

Unidirectional (blue in Fig. 6A), O₂-tolerant: they sustain H₂ oxidation in the presence of O₂ (dark blue in Fig. 5A and B). Inhibition by O₂ is reversible and produces mainly the NiB state Cm has high affinity for H₂ (ref. 133 and 134) Aa H2ase I is thermostable, and reactivated upon irradiation by violet light²⁰

O₂-insensitive, low activity (<0.5 μmol of H₂ per min per mg), low affinity for H₂ (K_M ≈ 4 μM)⁷⁴

O₂-insensitive, low activity (≈ 4 μmol of H₂ per mg) and high affinity for H₂ (K_m in the 0.1 μM range, compared to ≈ 10 μM for Sf Hyn^{55,56})

Low activity (≈ 2 μmol of H₂ per mg (ref. 40)) and do not oxidatively inactivate upon exposure to O₂ (ref. 40 and 41) Bidirectional. Couples H₂ oxidation to NAD⁺ reduction.¹³⁸ O₂-tolerant (is inactivated by O₂ (ref. 137)). In a series of homologous SH hydrogenases, Cn SH is the enzyme that has the largest activity (200 μmol of H₂ per min per mg (ref. 139)) and retains the greatest activity in the presence of O₂ (ref. 86)



Table 2 (Contd.)

Group		Structure	pdb	
	<i>Hydrogenophilus thermoluteolus</i> soluble hydrogenase	Ht SH	These soluble hydrogenases embed only low-potential accessory clusters ([2Fe2S] and [4Fe4S])	5XFA
				Thermophilic. Oxidatively inactivated to a peculiar inactive state, initially called Ni-B-like ⁸⁶ and then demonstrated to involve an overoxidized Ni(IV) ion ^{39,85} (pdb 5XF9, orange in Fig. 4B)
3b	<i>Synechocystis</i> sp. PCC 6803 <i>Pyrococcus furiosus</i> hydrogenase I	HoxEFUYH PfSHI		Bidirectional ⁹² Bidirectional. Distinct aerobic inactivation kinetics ¹⁴⁰

cells), and it was made possible by a better understanding of their biosynthesis. Tables 1 and 2 list the hydrogenases that we discuss in this review and summarise their main structural features and catalytic properties. The hydrogenases that have been isolated most recently are still not representative of the entire biodiversity, because one tends to focus on small and soluble enzymes that are more easily produced, and yet their characterization studies have revealed an unexpected functional diversity.

Some hydrogenases oxidise a few hydrogen molecules per second, while others exhibit turnover frequencies in excess of thousands per second. Some are tolerant to O₂,¹⁴ whereas the active site of others is destroyed upon exposure to oxygen.¹⁵ Much variability has also been observed regarding the reactivity with other inhibitors such as sulfide,¹⁶ chloride¹⁷ or CO. Some

FeFe hydrogenases are damaged by light in the visible range¹⁸ whereas others are only affected by UV B,¹⁹ and one NiFe hydrogenase was reported to be activated by light.²⁰ All hydrogenases inactivate, more or less reversibly, under very oxidising or reducing conditions, but again this varies from one hydrogenase to another. Some hydrogenases are unidirectional (they are only active in one direction of the reaction, H₂ oxidation or evolution) whereas others are bidirectional.²¹ And some are active in response to a very small departure from equilibrium, whereas others catalyse H₂ oxidation or evolution only in response to a large overpotential (these behaviours have been termed reversible and irreversible catalysis, respectively²²).

In Marseille, we have been particularly active in using direct electrochemistry to study and compare hydrogenases from various sources.²⁸ In this technique the enzyme is adsorbed




Fig. 2 The electrochemical diversity of FeFe hydrogenases, illustrated by catalytic voltammograms recorded under conditions of direct electron transfer and using a rotating disc electrode, all under one atm. of H₂. (A) *C. reinhardtii* HydA1.¹⁷ (B) *C. acetobutylicum* hydrogenase I. (C) *Thermoanaerobacter mathranii* HydS.²³ (D) *C. pasteurianum* hydrogenase II.²⁴ (E) *C. beijerinckii* hydrogenase, and the C367D variant. (F) Variants of Cb, A561F (green); P386L (blue); L364F (dark red); M382E (orange).^{25,26} (G) *C. pasteurianum* hydrogenase III and *M. elsdenii* hydrogenase II. (H) The variant of *C. pasteurianum* hydrogenase III where the supernumerary cysteine (yellow in Fig. 3C) is deleted.²⁷ All the voltammograms were recorded in a mixed buffer (MES, CHES, HEPES, TAPS, and Na acetate all [5 mM], and Na₂SO₄ (0.1 M), or NaCl (0.1 M) if specified). Panels (A), (B), (C), (D), (G), and (H) conditions: pH 7; 30 °C; 20 mV s⁻¹; 3000 rpm. Panels (E) and (F) conditions: pH 7; 5 °C; 20 mV s⁻¹; 3000 rpm, currents normalised at -559 mV.




Fig. 3 The structures of various FeFe hydrogenases. (A) The overall structures of the enzymes from *C. reinhardtii* (pdb 3LX4), *D. desulfuricans* (the two subunits are shown in grey and yellow, pdb 1HFE), *C. pasteurianum* (Cpl, pdb 6N59), *C. beijerinckii* (Cb, AlphaFold model, since the structure pdb 6TTL is incomplete), CplII (AlphaFold model), and *T. mathranii* HydS (AlphaFold model). (B) The residues that define the conformational change in Cb^{25,26} (P386; L364; A561; M382, Cb numbering) and CO access to the active site in Cr HydA1 (ref. 35) (F417 and V423, Cpl numbering). (C) The supernumerary cysteine of CplII (C221, CplII numbering). (D) The proton transfer pathways in Cpl (gray) and Tam HydS (red).³⁶ Regarding the AlphaFold predictions shown in this figure, ESI Fig. 1† shows the confidence values along the peptide chain.

onto and undergoes direct electron transfer with an electrode, which is spun to avoid H₂ depletion or accumulation near the electrode surface.^{28–31} The current is proportional to the activity, which can be monitored either as a function of time at a constant electrode potential (*E*), or as a function of electrode potential in experiments called cyclic voltammetry where the electrode potential is repeatedly swept up and down at a certain scan rate (Fig. 2). The sign of the current indicates the direction of electron flow (we count positive and negative the H₂ oxidation and evolution currents, respectively), and the magnitude of the current is proportional to turnover frequency (TOF) times the amount of enzyme that contributes:

$$i \propto \text{TOF} \times (\text{surface coverage of the active enzyme}) \quad (1)$$

The turnover frequency depends on the electrode potential (and other experimental parameters such as *T*, pH, and [H₂]) and this dependence informs on the properties of the intermediates in the catalytic cycle.^{32,33} The second term in the right-hand side of eqn (1), the surface concentration of the active enzyme, is smaller than the total amount of enzyme adsorbed on the electrode, first because some enzyme molecules may be adsorbed in a configuration that does not allow electrical communication with the electrode (and hence do not contribute to the current), and second because some enzyme molecules may (in)activate in a certain range of electrode potentials. If such a redox-driven (in)activation process is slow on the time scale of the experiment, it results in a hysteresis in the voltammetric signature.³⁴

The cyclic voltammograms recorded with various homologous FeFe hydrogenases, shown in Fig. 2, display a striking diversity of shapes. All enzymes appear to catalyse H₂ oxidation and reduction, although to various extents. Panel (D) illustrates the case of an enzyme whose response is nearly unidirectional (it is mostly active for H₂ oxidation, see also Fig. 6 below). Panel (C) shows the irreversible response of a bidirectional enzyme. Very strong hystereses are seen in panels (A), (E)–(G); they illustrate the effects of slow, reversible, redox-driven transformations between active and inactive forms of the enzyme.³⁴

The immediate implication of the observation that distinct hydrogenases exhibit distinct catalytic properties is that in designing a device that would use a hydrogenase for a particular purpose, care must be taken to choose the right enzyme: not an O₂-sensitive enzyme if the system operates in air, or not a photosensitive enzyme if the goal is to photo-produce H₂. There is still much hope that by exploring the biodiversity of these enzymes further, one will identify “new” hydrogenases that are even more robust or better suited for a given application.

From a more fundamental perspective, this functional diversity clearly demonstrates that the properties of each enzyme are not dictated by the active site itself. Although inorganic chemists are used to the idea that ligands that do not directly coordinate the metal may have an influence on catalysis, these effects are usually not considered beyond the so-called second coordination sphere.⁴⁹ In contrast, research conducted over the past two decades describing the comparison of distinct hydrogenases and their modification by protein engineering has demonstrated that the molecular factors



influencing their reaction with inhibitors, catalytic bias and catalytic reversibility are very diverse. These factors include structural features that may be distant from the active site and may have an influence even across different subunits in the same enzyme complex.

O₂-tolerance

Most of the hydrogenases isolated so far – and all the highly active ones – are inhibited by O₂, which is recognized as a main obstacle to using these enzymes. The reactions of hydrogenases with O₂ are complex, and the actual effect of O₂ varies greatly between FeFe and NiFe hydrogenases, and from one enzyme to another in the same family.

Dioxygen irreversibly damages the active site of many FeFe hydrogenases (in a complex, multistep reaction^{15,35,50}) and it oxidises the active site of NiFe hydrogenases into one or a mixture of inactive species, two of which (called “ready” and “unready”) can be distinguished by the rates at which they reactivate under reductive conditions.^{51,52} These two NiFe hydrogenase inactive states can also be obtained upon anaerobic oxidation of the enzyme.⁵³ In addition to this, in some hydrogenases, dioxygen may damage the accessory FeS clusters involved in mediating long range ET. Comparing the O₂ sensitivity of hydrogenases is therefore not trivial because there is no such thing as an “overall” sensitivity that could be quantified by a single parameter, such as the value of just one rate constant or one inhibition constant. The expression “O₂-tolerance” refers to the observation that a hydrogenase can oxidise H₂ in the presence of O₂ for a significant amount of time. This implies that dioxygen acts as a reversible inhibitor (the ratio of reversible

inactivation over reactivation rate constants defines an apparent inhibition constant, the magnitude of which is related to the extent of inhibition) and that if any irreversible inactivation occurs, it is slow on the time scale of the particular experiment where the aerobic H₂ oxidation activity is monitored.

Blocking the gas channel of an O₂-sensitive NiFe hydrogenase may hinder O₂ access, but does not make the enzyme O₂-tolerant

Hypotheses regarding the reason why the NiFe hydrogenase sensors (RH) from *Rhodobacter capsulatus* and *Cupriavidus necator* (previously known as *Ralstonia eutropha*) resist O₂ came from the structural description of the gas channels that connect the active site to the solvent. Volbeda and coworkers observed that in many oxygen sensitive NiFe hydrogenases, a bottleneck at the end of this channel is shaped by the side chains of two conserved residues, a leucine and a valine (grey in Fig. 4B), whereas bulkier phenylalanine and isoleucine residues are present at the same positions in the O₂-resistant NiFe H₂ sensors (yellow in Fig. 4B). Volbeda's hypothesis, that a narrow pathway may hamper O₂ access to the active site and make the enzyme resistant to O₂,⁵⁴ was supported by two site-directed mutagenesis (SDM) studies of these H₂ sensors (group 2c): the double replacement of the phenylalanine and isoleucine with leucine and valine makes the enzymes susceptible to oxidative inhibition.^{40,41} However, the rates of aerobic inactivation of the double mutants remain orders of magnitude slower than that of prototypical hydrogenases (about 10^{−3} s^{−1} (atm O₂)^{−1} based on the data in Fig. 4 of ref. 40, 10^{−5} s^{−1} (atm O₂)^{−1} from Fig. 2B of




Fig. 4 The NiFe hydrogenase dimer. (A) Overview of the structure of prototypical hydrogenases, the large subunit (light grey) embeds the active site, and the small subunit (dark grey) a chain of FeS clusters. (B) The environment of the active site, showing the V74/L122/E25 (Sf Hyn numbering) residues in prototypical hydrogenases,^{37,38} the residue E32 binding the Ni in the oxidised form of Ht SH,³⁹ the I62/F110 residues in *Cupriavidus necator* (Cn, previously known as *Ralstonia eutropha*) MBH^{40,41} (based on an AlphaFold model), and the E73 and H229 residues in *S. enterica* (Se) Hyd 5.⁴² (C) The proximal [4Fe4S] cluster in prototypical NiFe hydrogenases. (D) The proximal [4Fe3S] cluster in group 1d hydrogenases, coordinated by two supernumerary cysteines, C19 and C120 (*E. coli* Hyd 1 numbering).^{43–45} (E) The proximal cluster coordinated by 3 cysteines and one aspartate in the AH enzyme from *C. necator*.⁴⁶ (F) The histidine ligated distal cluster in prototypical NiFe hydrogenases,⁴⁷ and the arginine residue substituted in *E. coli* Hyd 1.⁴⁸



ref. 41, compared to $\approx 10 \text{ s}^{-1} (\text{atm O}_2)^{-1}$ for Sf Hyn⁵⁵), which shows that these two residues are not the only key.

The gas channels of the recently characterised *Mycobacterium smegmatis* NiFe hydrogenases Huc (group 2a)⁵⁶ and of the actinobacterial-type hydrogenase from *C. necator* (AH, group 1h)⁴⁶ are actually reported to be particularly narrow; it has been suggested that this may play a role in ref. 46 (or result in ref. 56) O₂ tolerance, but these hypotheses have not been tested by SDM. A computational comparison of NiFe hydrogenases from group 1 led to the hypothesis that O₂ sensitivity correlates with a more complex tunnel matrix into the protein, with an increased number of openings toward the solvent compared to the O₂-tolerant hydrogenases.⁵⁷ The calculated O₂-diffusion pathways are also different in NiFe and NiFeSe hydrogenases,⁵⁸ and include a *hydrophilic* channel in the latter.⁵⁹

In attempts to increase the O₂ resistance of the O₂-sensitive NiFe hydrogenase from *S. fructosivorans* (Sf Hyn) by blocking the gas channel, we designed many site-directed variants where we replaced the valine (at position 74) or the leucine (122, grey in Fig. 4B) with bulkier residues, and we used two kinetic methods to experimentally evaluate the rates of diffusion along these modified channels. One method consists in using mass spectrometry to monitor the “isotope exchange reaction” ($\text{D}_2 + 2\text{H}^+ \rightarrow \text{H}_2 + 2\text{D}^+$), whose kinetics depends on the rate of diffusion in the channel; the other consists in using electrochemistry to measure the bimolecular rate constant of inhibition by CO, a competitive inhibitor that uses the same gas channel as O₂ to access the active site.⁶⁰ Single-point mutations (especially at position 74) alter the rate of intramolecular diffusion by more than three orders of magnitude, and both the charge and the size of the side chains at positions 74 and 122 matter.^{60,61} However, the effect of the mutations on the rate of inhibition by O₂ is rather small (only up to ten-fold), because in the WT enzyme and in most variants, the diffusion of O₂ along the gas channel is much faster than the rate of reaction of O₂ at the active site, and it is not the rate limiting step of the inhibition reaction.⁶¹ Mutations that have the most severe effect on the diffusion rate do slow down O₂ inhibition, but this effect alone only delays the loss of enzyme activity upon exposure to O₂,⁶¹ because the formation of the oxidised inactive states of the active site remains irreversible under the experimental conditions. We concluded that, at least in the particular case of this hydrogenase, slowing diffusion along the gas channel is not a strategy that makes the enzyme O₂-tolerant.

Inhibitor access to the active site of FeFe hydrogenases depends on unidentified details of the protein matrix

Regarding FeFe hydrogenases, the rate of inhibition by CO and O₂ is three orders of magnitude slower for MeI hydrogenase than for Dd⁶² (cf. ESI Table S3 in ref. 62). The trend is the same regarding the rate of inhibition by sulfide¹⁶ and chloride,¹⁷ with the bimolecular rates of inhibition increasing in the order Cpl < Cr HydA1 < Dd. The FeFe hydrogenase from *Acetobacterium woodii* also reacts slowly with O₂.⁶³ We believe that these differences are related to the rates of diffusion through the protein matrix, but which residues are responsible for this is

unknown. The mutation of a phenylalanine residue that is in the gas channel (F417 in Fig. 3B) to tyrosine in Cr HydA1 slows both the inhibition by O₂ (two-fold) and CO (ten-fold).³⁵ Other mutations of FeFe hydrogenases have been shown to moderately slow down O₂ inhibition,^{64–66} but the link between O₂ diffusion kinetics and protection has not been demonstrated. When random mutagenesis was used to identify O₂-resistant variants of Cpl, the hot-spots were found close to the accessory clusters, rather than close to the putative gas channel.^{65,66}

A non-natural O₂-protection mechanism of NiFe hydrogenases: the substitution of a conserved valine in the second coordination sphere of the Ni ion affects the rates of reactivation after O₂ inhibition

Modifying the V74 and L122 residues in the large subunit of Sf Hyn (grey in Fig. 4B), with the initial aim of blocking the gas channel, had two unexpected effects, one of which is an improvement of O₂ tolerance that is not related to the rate of O₂ access: many substitutions of the position 74 valine increase the rate at which the enzyme reactivates after inhibition by O₂. The V74C⁶⁷ and V74H³⁸ replacements are particularly effective (Fig. 5A). The most significant enhancements of the reactivation rates are observed when the side chain of the position-74 residue is hydrophilic³⁸ (although this rule is not strict⁶⁸), but




Fig. 5 Chronoamperometric responses of NiFe hydrogenases to transient exposures to O₂. (A) Various mutants of Sf Hyn, obtained by replacing the position 74 valine, compared to the O₂-tolerant enzyme from *Aquifex aeolicus* (Fig. 4B).³⁸ (B) *E. coli* Hyd 1 and Hyd 2, the C19G/C120G (called GG) variant of *E. coli* Hyd 1 (Fig. 4D) where the proximal cluster is a standard cubane cluster, the chimeric “L1S2mod” dimer made of the large subunit of Ec Hyd 1 and the small subunit of Ec Hyd 2,⁷⁰ and Sf Hyn.⁵⁵ Arrows show the time of injection of O₂ saturated buffer, to reach a final concentration of 4 μM and 8 μM at time $t = 0$ s and $t = 300$ s in panel (A), and 8 μM at time $t = 0$ s in panel (B). Conditions: in panel (A) $E = 140$ mV; 40 $^{\circ}\text{C}$; pH 5.5; 3000 rpm; 1 atm H₂; in panel (B) $E = 140$ mV; 40 $^{\circ}\text{C}$; pH 6; 3000 rpm; 1 atm H₂; 8 μM O₂ injection. In all cases, the concentration of O₂ increases suddenly after the injection of an aliquot of O₂-saturated solution, and then decreases exponentially as the solution is flushed with H₂.⁵⁵




Fig. 6 Catalytic bias in voltammetry for different hydrogenases. (A) NiFe hydrogenases from *E. coli* Hyd 1 and Hyd 2, the variant C19G/C120G (called GG) and the chimeric dimer L1S2mod.⁷⁰ Conditions: pH 6; 40 °C; 3% H₂. (B) Variants of the NiFe hydrogenase Hyn from *S. fructosivorans*. The curves in this panel were scaled for normalisation. Conditions: pH 5.5; 40 °C; 10% H₂ for all²¹ except H184C,⁴⁷ pH 6; 40 °C; 100% H₂. (C) FeFe hydrogenase CplII WT,²⁷ in red a voltammogram where the potential is reversed before reaching a high value, showing the steady-state catalytic response and no inactivation. In blue a voltammogram recorded over a large potential range showing the oxidative inactivation; pH 7; 5 °C; 100% H₂.

the reason for this is unknown. Replacing the equivalent valine in the oxygen-tolerant hydrogenase *E. coli* Hyd 1 also increases oxygen tolerance further (although at the expense of decreasing activity).⁶⁹

It is somewhat surprising that this valine is very conserved, whereas many of its substitutions improve the resistance to O₂.³⁸ The effects of these substitutions on the catalytic bias (see below) may have put a selective pressure on the conservation of this residue.

O₂-tolerant NiFe hydrogenases from group 1d: the whole small subunit is crucial, not just the proximal cluster

O₂-tolerant NiFe hydrogenases in group 1d have attracted much interest when various investigations showed that they differ from prototypical hydrogenases by the presence of a peculiar [4Fe3S] cluster, attached to the protein by 6 cysteine residues (including C19 and C120, *E. coli* Hyd 1 numbering, Fig. 4D). This cluster is proximal to the active site but embedded in the small accessory subunit, and replaces the standard 4-Cys [4Fe4S] cluster found in many other NiFe hydrogenases and shown in Fig. 4C.

The most familiar hypothesis is that the proximal cluster makes the enzyme O₂-tolerant by providing two electrons to reduce the attacking O₂, favouring the formation of the inactive state NiB, which reactivates quickly, over the more inert inactive state NiA.^{43,71} However, this hypothesis has been challenged by recent investigations based on the engineering of this proximal cluster.

Very recently, Lenz and coworkers showed that replacing with glycine either one or two of the supernumerary cysteines around the proximal cluster of the O₂-tolerant NiFe hydrogenase from *C. necator* (MBH) does make the enzyme more O₂-sensitive (as observed before in *E. coli* Hyd 1 (ref. 72)), but without favouring the formation of the NiA inactive state over

NiB.⁷³ According to the authors, the observation that group 1d NiFe hydrogenases react with O₂ to form only the NiB state is therefore not related to the two-electron transfer capacity of the proximal cluster.⁷³

In our study of the O₂-tolerant NiFe hydrogenase from *E. coli* (Hyd 1), by comparing variants where we replaced either the proximal [4Fe3S] cluster with a standard [4Fe4S] cluster (double substitution C19G/C120G, Fig. 4D), or the entire accessory subunit with that of an O₂-sensitive NiFe hydrogenase (by designing a chimeric dimer, called L1S2mod, made of the large subunit of *E. coli* Hyd 1 and the small subunit of Hyd 2), we concluded that the kinetics of reactivation after oxidative anaerobic inactivation of the enzyme is determined by the large subunit, which bears the NiFe active site, whereas the accessory subunit that houses the electron transfer chain is mostly responsible for O₂ tolerance (pale blue and orange in Fig. 5B).⁷⁰ Most importantly, the variant of *E. coli* Hyd 1 that bears a standard proximal cluster is much less O₂-sensitive than standard hydrogenases (pale blue and green in Fig. 5B), showing that this cluster is not the unique determinant of O₂-resistance.⁷⁰

A SDM study of *S. enterica* Hyd5 (also from group 1d) showed that the substitutions of two residues (E73 and H229, blue in Fig. 4B) negatively impact O₂-tolerance. They are in the large subunit, but H229 is close to the proximal cluster of the small subunit. However these residues are not markers of O₂-tolerance: the histidine is very conserved (including in O₂-sensitive NiFe hydrogenases) and some enzymes from group 1d (e.g. *Aquifex aeolicus* hydrogenase I) have a glutamine instead of the glutamate, as in O₂-sensitive enzymes.⁴²

Other NiFe hydrogenases (those in groups 1h^{74,75} and 2a⁷⁶) appear to be O₂-tolerant despite lacking the proximal [4Fe3S] cluster, but their catalytic activity is very low.⁷⁷ The narrowness of gas channels has been hypothesised to play a role, as mentioned above. However, the O₂-insensitivity of the AH enzyme from *C. necator* (group 1h, formerly group 5) is lost



upon substitution of the aspartate residue that coordinates (in an unusual fashion) the proximal FeS cluster of its electron transfer chain (Fig. 4E); this observation led the authors to suggest again an O₂ tolerance mechanism involving the reductive removal of the attacking O₂.⁴⁶

Protecting the hydrogenase active site by saturating the coordination sphere of the metal ions in FeFe and NiFe hydrogenases

Most FeFe hydrogenases are irreversibly damaged by O₂. Some of them partially recover activity after a short exposure to O₂,^{35,78,79} but this does not prevent irreversible degradation upon continuous exposure. However, Morra and co-workers recently discovered that the FeFe hydrogenase from *C. beijerinckii* (Cb) is “O₂-stable”: it can be exposed to air with minimal damage.^{80,81}

Various observations led to the conclusion that this peculiar protection mechanism results from the binding of the thiolate of a conserved cysteine residue to the distal Fe of the active site, under oxidising conditions. First, the X-ray structure of the Cb enzyme in the Hinact state (pdb 6TTL) is consistent with the formation of a bond between the cysteine and the distal Fe (Fig. 3B).²⁵ Second, the FTIR signature of the oxidised active site of Cb hydrogenase is similar to the so-called “Hinact state”, obtained by oxidising the FeFe hydrogenases from Dd or Cr in the presence of hydrogen sulfide;^{16,82,83} in this state, a sulfido ligand binds the distal iron (“Fe_d” in Fig. 1A) and prevents the binding of H₂ or O₂, hence the enzyme is inactive but protected from O₂.⁸⁴ Third, the replacement of the cysteine with an aspartate suppresses both the stability in air and the formation of the Hinact state under oxidising conditions.²⁵

That the cysteine binds to the oxidized H-cluster in Cb but not in other FeFe hydrogenases is due to a conformational change that involves residues very far away from the active site (up to 20 Å, see below the section on Redox-driven changes in active site structure).^{25,26} It is a clear illustration of how the protein matrix may have a long distance effect on active site chemistry.

Although the inactive state itself (Hinact, cysteine-bound) is protected from O₂, whether or not the formation of this state protects the enzyme depends on the relative kinetics of formation of Hinact and O₂-induced degradation. In the third FeFe hydrogenase from *C. pasteurianum*, CpIII (Fig. 3C), the inactivation (Fig. 2G) that results from the formation of the Hinact state upon binding to Fe_d of a cysteine side chain is too slow to overcome O₂ attack.²⁷

The protection of the H-cluster of Cb and CpIII hydrogenases by the binding under oxidising conditions of a nearby cysteine side chain is reminiscent of the protection mechanism evidenced in the NAD⁺-reducing soluble [NiFe]-hydrogenase (SH) from *Hydrogenophilus thermoluteolus* (Ht). The oxidised active site of this enzyme includes a six-coordinate Ni, with a conserved glutamic acid previously shown to be involved in proton transfer³⁷ bound to the Ni (orange in Fig. 4B). Like in the case of Cb, the formation of this bond inactivates the enzyme and prevents O₂ binding.^{39,85} It is proposed that the reversible

formation of this inactive state under oxidising conditions protects Ht SH from irreversible damage related to metal-assisted ROS production,⁸⁵ but it is unknown which other mechanism allows this enzyme to perform H₂ oxidation (although at a reduced rate) in the presence of O₂.⁸⁶ Replacement of the glutamate residue with alanine or glutamine suppresses the IR signature of the Ni-carboxylate bond in the oxidised state, and greatly decreases the activity (as expected from the substitution of a residue involved in proton transfer),⁸⁵ but the impact of the substitution of this glutamate on O₂-damage has not been evaluated. The reason why this protection mechanism is operational in Ht SH and not in other, homologous NiFe hydrogenases is also unknown.

Catalytic bias

In voltammetry experiments such as those in Fig. 2 and 6, the H₂ oxidation and production activities of hydrogenases are assessed successively as the electrode potential is swept up and down across a large potential window. It immediately appears that some enzymes are more active in one direction of the reaction than the other, or even only active in one direction (see e.g. Fig. 2D and 6A). This property is referred to as the “catalytic bias”, or “catalytic preference”, or “catalytic directionality”,²² although there is no consensus in the hydrogenase literature as to how exactly it should be quantified.

That a particular hydrogenase (or any other catalyst) may be a unidirectional catalyst does not violate the principles of thermodynamics.⁸⁷ Thermodynamics imposes that the current is zero when the electrode potential (*E*) matches the equilibrium potential (*E*_{eq}, or “Nernst potential”) of the H⁺/H₂ couple (as indeed observed in all experiments^{88,89}), and it relates the direction of the reaction (H₂ oxidation or production) to the sign of the overpotential (the difference between *E* and *E*_{eq}), but it does not constrain the magnitudes of the currents observed under oxidising or reducing conditions.

Elucidating why hydrogenases may be biased to catalyse the reaction in a certain direction may help the design of synthetic catalysts, very few of which are active in H₂ oxidation.⁹⁰ This question also has implications in terms of physiology. Indeed, some bacteria produce a number of homologous hydrogenases, which are involved in various metabolic functions: for example, as part of respiration and fermentation pathways, these enzymes are used to either oxidise or produce H₂, respectively. It is tempting to think that a particular function matches the intrinsic property of each isoform, and that in addition to genetic regulation and cellular localisation, whether a hydrogenase is a better catalyst in one of the two directions of the reaction contributes to defining its metabolic contribution.

Evidence that the standard redox potentials of the FeS of the electron transfer chain determine the catalytic bias of hydrogenase is lacking

Twenty five years ago, P. L. Dutton and coworkers proposed that electron transfer chains in redox enzymes are only optimised in terms of distance between the cofactors (allegedly the most



important determinant of the ET rate), whereas the standard redox potentials of the cofactors do not matter. The reasoning was that intramolecular electron transfer was faster than active site chemistry and thus not rate limiting in the catalytic cycle, “endergonic electron transfer steps can still support rapid electron transfer”, and the details of the electron transfer chain (incl. the potentials of the relays) are not important.⁹¹ This paradigm has shifted since, at least in the hydrogenase field, where the potential of the accessory clusters is now often considered a determinant of the catalytic bias (higher potential clusters supposedly accelerating catalysis in the direction of H₂ oxidation, and lower potential clusters favouring H₂ production).

Indeed, recently, the bias in the direction of H₂ oxidation of the group 2a Huc NiFe hydrogenase from *Mycobacterium smegmatis* was tentatively ascribed to the presence of three high redox potential [3Fe4S] clusters in the electron transfer chain of this enzyme,⁹⁶ and the lack of [3Fe4S] clusters in the NiFe-hydrogenase of the Cyanobacterium *Synechocystis* sp. PCC 6803 was tentatively related to the catalytic bias toward proton reduction.⁹² Support for this hypothesis from site-directed mutagenesis experiments has not been provided yet. By explicitly considering the redox relay in the kinetic modelling of the catalytic response, we showed that the rates of intramolecular electron transfer matter, not just the potential of the relay,⁹³ but these rate constants are difficult to measure.⁹⁴

The above-mentioned hydrogenases *E. coli* Hyd 1 (group 1d) and Hyd 2 (group 1c) have also been studied in this context,^{14,70} because, despite their strong homology (43% sequence identity for both the large and the small subunits), the former is strongly biased in the direction of H₂ oxidation whereas the latter is bidirectional (dark blue and green, respectively, in Fig. 6A). In NiFe hydrogenases from group 1d, O₂-tolerance and the presence of an atypical [4Fe3S] proximal cluster (Fig. 4D) seem to correlate with unidirectionality. Another difference between Hyd 1 and Hyd 2 is that the distal cluster of the former has not been detected by EPR,⁹⁵ despite its structure and environment being the same as in Hyd 2. Parkin *et al.* hypothesised that this cluster has higher potential in Hyd 1 than in Hyd 2 and that this may be one of the factors that bias Hyd 1 for H₂ oxidation.⁴⁸

However, investigations of the relation between the redox potential of the clusters of the ET chain and catalytic bias were inconclusive. Decreasing the potential of the distal cluster in Hyd 1 (by substituting a conserved residue, R193, Fig. 4F) has only a small influence on the catalytic bias.⁴⁸ Replacing either the high potential [3Fe4S] middle cluster or the high potential [4Fe3S] proximal cluster with a low potential [4Fe4S] has no impact.^{70,72}

Regarding the structural determinants of the difference in catalytic bias between Hyd 1 and Hyd 2, the effect of assembling the large subunit of the unidirectional enzyme Hyd 1 with the entire electron transfer subunit of the bidirectional enzyme Hyd 2 is unambiguous: this chimeric dimer has no H₂ evolution activity, showing that this catalytic bias is not defined by the electron transfer chain.⁷⁰

Regarding FeFe hydrogenases, attempts to identify the determinants of the catalytic bias by site-directed mutagenesis

have mostly focused on the immediate environment of the H-cluster and the proton transfer chain. Replacing a cysteine ligand of the cubane of the H-cluster (C362 in Cr HydA1, C499 in CpI, see Fig. 3C) with histidine increases the redox potential of the active site and selectively suppressed proton reduction.⁹⁶ The replacement of the same cysteine with aspartate⁹⁷ and certain modifications of residues involved in long range proton transfer also suppress⁹⁸ or favor³⁶ proton reduction. Changing the environment of the CN[−] ligand of the proximal Fe ion (of the dinuclear fragment of the H-cluster) has a mild effect on the catalytic bias.⁹⁹ Regarding the manipulation of the accessory clusters, it was shown that replacing the histidine that binds the surface exposed accessory [4Fe4S] cluster in CpI with a cysteine decreases the redox potential of the cluster (≈ 65 mV), and the bias in solution assays (about two-fold) in the reductive direction.¹⁰⁰

The catalytic bias depends on the rate constants of the steps in the catalytic cycle

The rates of catalysis in either direction depend on the rate constants of the steps in the catalytic cycle, and the discussion of the catalytic bias should be based on the measurement of the rate constants of the steps and on the understanding of which steps define the overall TOF. However, the catalytic cycle of complex metalloenzymes involves steps of very different types (active site chemistry, long range proton and electron transfers, diffusion along substrate channels¹⁰¹) and whose individual rate constants are often impossible to measure. Moreover, the overall turnover frequency may be a very complex function of these rate constants, in particular when all steps in the catalytic cycle are reversible³² (indeed, only when a reaction consists of a sequence of irreversible steps is the rate limiting step easily defined as the slowest of these steps¹⁰²).

Modifying the coordination of the distal cluster of Sf NiFe hydrogenase (Fig. 4F) changes the catalytic bias by impacting the rate of H₂ oxidation in solution assays more than the rate of H₂ evolution; this suggests that the intermolecular ET step contributes to defining the rate of H₂ oxidation.⁴⁷ Furthermore, obstructing the gas channel of the same enzyme by substituting V74 or L122, the residues whose side chains line the gas channel in the large subunit of the enzyme (grey in Fig. 4B), slows H₂ evolution much more than H₂ oxidation, suggesting that H₂ egress is the rate limiting step of the reaction of H₂-production in this series of mutants: indeed, the rates of that diffusion step, measured by interpreting the results of isotope exchange assays, exactly match the overall rates of H₂ evolution measured in solution assays.²¹ The effect of these mutations in the large subunit of NiFe hydrogenase on the catalytic bias is also clear from the voltammetric signatures of this series of variants (*e.g.* red and yellow in Fig. 6B). These are clear examples where the catalytic bias is defined by steps in the catalytic cycle other than active site chemistry.

The discussion of the impact of the mutations on the redox properties of the active site and the catalytic bias (as observed with the C170H mutant of Cr HydA1 (ref. 96)) focussed on how the mutations may affect the standard redox potential of the



active site. A complication in catalytic cycles that are bidirectional is that a thermodynamic constraint applies to the series of steps in the catalytic cycle,^{32,103} and a mutation that tends to accelerate a chemical step in one particular direction of the catalytic cycle must impact the thermodynamics of other steps, possibly redox transitions. There may therefore be cases where the discussion of the rate constants of the chemical steps is intrinsically linked by thermodynamics to the redox potential of some other steps in the catalytic cycle.

Redox-driven changes in the structure of the active site may influence the catalytic bias

A completely distinct discussion of why a catalyst may be better in one particular direction considers the possibility that the structure of the catalyst may change depending on the driving force.

A very classical example of this was provided by John Bockris in his studies of oxygen reduction and evolution on metals: the metal surface that is present at low electrode potential under conditions of oxygen reduction is completely distinct from the metal-oxide or hydroxide surface that is present when a high electrode potential is set to drive O₂-evolution from water.¹⁰⁴ The structure of the active site of many hydrogenases also changes depending on the redox conditions.

The active site of NiFe hydrogenases, for example, is oxidised at high electrode potential (above $E \approx -100$ mV, pH 7 (ref. 105)) to either one or a mixture of inactive states in which hydroxo (or maybe peroxo) ligands bind the active site metal ions, which prevents oxidative turnover.^{53,105,106} Sulfoxylation of the active site¹⁰⁷ or saturation of the Ni coordination sphere with a nearby side chain^{39,85} may also result in oxidative inactivation of NiFe hydrogenases.

Anaerobic oxidative inactivation does not occur in most FeFe hydrogenases that have been characterized to date (unless chloride¹⁷ or sulfide^{16,82} is present), but the group A FeFe hydrogenase from *Clostridium beijerinckii* is unusual in that respect. Under mildly oxidising conditions, the side chain of the conserved cysteine that is near the amine of the dithiolate ligand binds the distal Fe of the dinuclear cluster, which inactivates the enzyme by preventing H₂-binding (resulting in the formation of the above-mentioned Hinact state); this is detected in voltammetry as a decrease of the H₂-oxidation current above ≈ -300 mV (orange in Fig. 2E). This oxidative inactivation disappears when this cysteine (C367) is replaced with an aspartate (red in Fig. 2E, the residual hysteresis in this voltammogram is due to the presence of chloride in the buffer).²⁵ The consequence of this inactivation is that the WT FeFe hydrogenase from Cb is protected from O₂ damage (as discussed above, the saturation of the coordination sphere of the distal Fe prevents O₂ binding) but there is a trade-off: this enzyme is a very poor H₂-oxidation catalyst. This is not because the enzyme is intrinsically unable to oxidise H₂, but because it cannot stay active for long under conditions where H₂ oxidation occurs.

The question remains as to why this conserved cysteine binds the cluster in the FeFe hydrogenase from Cb, and not in

prototypical FeFe hydrogenases from group A. The reason is that the formation of the bond between the distal Fe and the cysteine sulphur requires a significant conformational change, which depends on the side chain of non-conserved residues that are remote from the active site. Winkler *et al.* identified 3 remote residues (at positions 364, 368, and 561, Cb numbering, Fig. 3B), whose bulkier side chains in prototypical FeFe hydrogenase (such as CpI) prevent the conformational change.²⁵ Replacing the amino acids of Cb with those present in CpI slows down the conformational change and prevents oxidative inactivation (Fig. 2F).²⁵ The replacement of the position 382 residue, which is even more distant (18 Å away from the nitrogen of the amine bridge, Fig. 3B), has the opposite effects: it increases the flexibility of the protein and the rate and extent of oxidative inactivation.²⁶ The voltammograms of these variants disclose the intrinsic capability of the active site to oxidise H₂.

The FeFe hydrogenase CpIII is from the phylogenetic group B, which has barely been investigated. Like Cb, CpIII inactivates under mildly oxidising conditions (Fig. 2G)²⁷ and is observed by IR in a state that resembles Hinact.²⁴ The loop that bears the above-mentioned proton transfer cysteine is characterised in CpIII by an unusual motif with three vicinal cysteine residues. According to the alpha-fold structure of CpIII (Fig. 3C),²⁷ we hypothesised that one cysteine binds the [4Fe4S] cluster of the active site (as in prototypical FeFe hydrogenases), another one takes the position of the above mentioned proton transfer cysteine and may bind the distal Fe (to produce the Hinact-like state), and the third one may also be involved in proton transfer.¹⁰⁸

Like the case of Cb, the analysis of the voltammetric responses of CpIII shows that the environment of its active site is not particularly tuned in a way that makes it better at evolving H₂ than oxidising H₂; the reason this enzyme has low H₂ oxidation activity in solution assays is that it inactivates under oxidising conditions.²⁷

Reversibility

We²² and others before us^{90,109} have proposed to call “reversible catalysis” the action of a “reversible catalyst”, which becomes active in one direction or the other in response to a small departure from equilibrium. In contrast, an irreversible redox catalyst is only active at high overpotential.

Irreversibility may be the mere consequence of slow interfacial electron transfer (between the electrode and the enzyme), as observed in the H184C mutant of Sf Hyn (Fig. 4F, and green in Fig. 6B).⁴⁷ In contrast, in the case of the sensory FeFe hydrogenase Tam HydS discussed below, we concluded from the analysis of the waveshape that slow electron transfer is not the reason the catalytic response is irreversible, this is an intrinsic property of the enzyme.²³

Fig. 2 shows reversible and irreversible responses obtained with homologous FeFe hydrogenases. The catalytic response in panel C is from the putative sensory hydrogenase from *Thermoanaerobacter mathranii* (Tam); it is classified as group D, but it resembles the sensory hydrogenase HydS from *Thermotoga maritima* (Tm) which is in group C. In both enzymes, the proton



transfer chain and the environment of the H-cluster differ from that in group A hydrogenases (Fig. 3D).¹⁰⁸ Irreversibility is not specific to hydrogen sensors: the irreversible response in Fig. 2D is that of the 2nd hydrogenase from *C. pasteurianum* (CpII, from group A, prototypical FeFe hydrogenases).²⁴

The separation between the two catalytic waves (the difference, for a given enzyme, between the values of the two catalytic potentials, the midpoint potentials of the H₂ oxidation and H⁺ reduction catalytic waves) is related to the difference between the redox potentials of the two transitions of the active site, although the catalytic potentials are shifted from the thermodynamic values, just like Michaelis constants depart from the true dissociation constants.^{22,32,110} This implies that the more stable the half reduced state of the active site (the so-called H_{red} state), the more irreversible the response; this appears to be the main reason why the enzyme from Tam behaves very irreversibly,³³ although it is still unknown which residues in the environment of the active site make the half reduced state very stable.¹¹¹ That the proton relay near the H-cluster has a lower pK_a in Tam than in Cr also contributes to making the response of the former less reversible.³³

Implications for the performance of hydrogenase-based catalytic devices

The functional variations that are observed among homologous hydrogenases that share the same active site are clear illustrations of very long range effects in these biological inorganic catalysts. They also have immediate implications regarding the design of any device that uses a hydrogenase as a catalyst for oxidation or production. Care must be taken that the properties of the particular hydrogenase that is chosen match the needs and the operational conditions of the device, in terms of activity, directionality, reversibility, resistance to light and inhibitors (including O₂), stability, cost of production, *etc.* Since it appears that no hydrogenase checks all the boxes, the right choice is necessarily a compromise.

Any attempt to evolve H₂ by coupling hydrogenase with a photosensitizer would fail if the enzyme from *D. desulfuricans* is used since that enzyme is destroyed by white light.^{112,113} Other hydrogenases appear more robust in that respect, but the possibility of photodamage¹⁹ should be considered in all sunlight-dependent H₂-evolution systems.¹¹⁴

Since long term stability is required, enzymes produced by hyperthermophilic organisms may be particularly useful.^{115,116} However, some hydrogenases from mesophilic organisms were shown to be very stable: the NiFe hydrogenase from *Desulfovibrio gigas* could be used for oxidising H₂ at 40 °C for weeks without any loss of activity.¹¹⁷

The most important concern regarding the use of hydrogenases is their sensitivity to oxygen. The O₂-tolerant NiFe hydrogenases from group 1d have been extensively used for that reason because all other O₂-tolerant hydrogenases have very low H₂-oxidation activity. The group-1d hydrogenase from *Cupriavidus metallidurans*, which combines O₂-tolerance and high affinity for H₂, was used as the anode catalyst in a membrane-less fuel cell operating on 3% H₂ in air.¹¹⁸

But even oxygen sensitive hydrogenases can be used as electrocatalysts of H₂-oxidation in the presence of O₂, under the condition that the matrix that supports the enzyme blocks O₂. This is the basis of the strategy used by Plummer and coworkers, who have shown that when a hydrogenase is embedded in the thick layer of a redox polymer, a fraction of the incoming H₂ can be redirected to catalytically produce electrons that reduce any molecule of O₂ that may penetrate the film. Long term stability can be achieved under the harsh conditions of a fuel cell,¹¹⁹ even with the most O₂-sensitive FeFe hydrogenase.¹²⁰ The protective effect of the matrix and the intrinsic O₂-resistance of the enzyme can be combined to make the system particularly robust.¹²¹

Reversibility²² is also desirable when a hydrogenase is used as an electrocatalyst because any overpotential needed to trigger catalysis is a waste of energy. Care must also be taken that this useful property is not lost when the electron transfer between the enzyme and an electrode is mediated.¹²²

The active sites of hydrogenases are based on cheap metals, Ni and Fe, but the cost of producing the enzyme in large amounts is rarely discussed in the literature. The heavy biological machinery that the living cells must use to produce the complex active sites shown in Fig. 1 hampers the biological synthesis of these enzymes, and upscaling their production is a challenge. A huge step forward was made by an international collaboration ten years ago when it was demonstrated that some FeFe hydrogenases can be produced in an “apo” form (that lacks the dinuclear fragment of the H-cluster), and then activated by the spontaneous insertion of a synthetic dinuclear cluster.¹²³ This opened the way for the cheap, large-scale production of simple FeFe hydrogenases, and recent results regarding the biological maturation of the NiFe active site¹²⁴ makes us hope that a similar process will one day be available to produce NiFe hydrogenases.

Data availability

The text files of the electrochemical data shown in this paper can be downloaded from the Zenodo repository at <https://zenodo.org/records/10849830>.

Author contributions

AF, VF and CL co-wrote the manuscript.

Conflicts of interest

The authors declare no conflict of interest.

Acknowledgements

We are very grateful to many colleagues and former students who contributed to our work on hydrogenase in Marseille: Abbas Abou Hamdan, Carole Baffert, Aurore Bailly, Melisa del Barrio, Myriam Brugna, Pierre Ceccaldi, Sebastien Dementin, Christina Felbek, Marie-Thérèse Giudici-Orticoni, Chloé Guendon, Pascale Infossi, Arlette Kpebbe, Fanny Leroux, Pierre-Pol



Liebgoth, Matteo Sensi and Jeremy Wozniak. And to our colleagues Gustav Berggren, Luca Bertini, James Birrell, Jochen Blumberger, Maurizio Bruschi, Marc Fontecave, Luca de Gioia, Claudio Greco, Thomas Happe, Adam Kubas, Ines Perreira, Nicolas Plumeré, Sven Stripp, Francesca Valletti, Martin Winkler and their students and colleagues. We also thank Oliver Lenz, Stefan Frielingsdorf and Marius Horch for helpful discussions. This research was funded by the Centre National de la Recherche Scientifique, Aix Marseille Université, Agence Nationale de la Recherche (ANR-12-BS08-0014, ANR-14-CE05-0010, ANR-15-CE05-0020, ANR-21-CE50-0041, and ANR-23-CE50-0016), Région Sud, and the French Government under the France 2030 investment plan, as part of the Initiative d'Excellence d'Aix-Marseille Université – A*MIDEX, AMX-22-RE-AB-097 and ANR-11-IDEX-0001-02. The authors are part of the FrenchBIC CNRS network (<https://www.frenchbic.cnrs.fr/>).

References

- O. Einsle and D. C. Rees, *Chem. Rev.*, 2020, **120**, 4969–5004.
- B. M. Hoffman, D. Lukoyanov, Z.-Y. Yang, D. R. Dean and L. C. Seefeldt, *Chem. Rev.*, 2014, **114**, 4041–4062.
- D. G. Nicholls, *Bioenergetics*, Academic Press, 2013.
- F. Sargent, in *Advances in Microbial Physiology*, ed. R. K. Poole, Academic Press, 2016, vol. 68, pp. 433–507.
- C. Baffert, A. Kpebe, L. Avilan and M. Brugna, in *Advances in Microbial Physiology*, ed. R. K. Poole, Academic Press, 2019, vol. 74, pp. 143–189.
- C. Greening, A. Biswas, C. R. Carere, C. J. Jackson, M. C. Taylor, M. B. Stott, G. M. Cook and S. E. Morales, *ISME J.*, 2016, **10**, 761–777.
- J. Meyer, *Cell. Mol. Life Sci.*, 2007, **64**, 1063–1084.
- M. Stephenson and L. H. Stickland, *Biochem. J.*, 1931, **25**, 205–214.
- S. T. Stripp, B. R. Duffus, V. Fourmond, C. Léger, S. Leimkühler, S. Hirota, Y. Hu, A. Jasniewski, H. Ogata and M. W. Ribbe, *Chem. Rev.*, 2022, **122**, 11900–11973.
- J. A. Birrell, P. Rodríguez-Maciá, E. J. Reijerse, M. A. Martini and W. Lubitz, *Coord. Chem. Rev.*, 2021, **449**, 214191.
- W. Lubitz, H. Ogata, O. Rüdiger and E. Reijerse, *Chem. Rev.*, 2014, **114**, 4081–4148.
- S. Morra, *Front. Microbiol.*, 2022, **13**, 853626.
- J. W. Peters, G. J. Schut, E. S. Boyd, D. W. Mulder, E. M. Shepard, J. B. Broderick, P. W. King and M. W. W. Adams, *Biochim. Biophys. Acta*, 2015, **1853**, 1350–1369.
- M. J. Lukey, A. Parkin, M. M. Roessler, B. J. Murphy, J. Harmer, T. Palmer, F. Sargent and F. A. Armstrong, *J. Biol. Chem.*, 2010, **285**, 3928–3938.
- J. Esselborn, L. Kertess, U.-P. Apfel, E. Hofmann and T. Happe, *J. Am. Chem. Soc.*, 2019, **141**, 17721–17728.
- C. Felbek, F. Arrigoni, D. de Sancho, A. Jacq-Bailly, R. B. Best, V. Fourmond, L. Bertini and C. Léger, *ACS Catal.*, 2021, **11**, 15162–15176.
- M. Del Barrio, M. Sensi, L. Fradale, M. Bruschi, C. Greco, L. de Gioia, L. Bertini, V. Fourmond and C. Léger, *J. Am. Chem. Soc.*, 2018, **140**, 5485–5492.
- S. P. J. Albracht, W. Roseboom and E. C. Hatchikian, *J. Biol. Inorg. Chem.*, 2006, **11**, 88–101.
- M. Sensi, C. Baffert, L. Fradale, C. Gauquelin, P. Soucaille, I. Meynial-Salles, H. Bottin, L. de Gioia, M. Bruschi, V. Fourmond, C. Léger and L. Bertini, *ACS Catal.*, 2017, **7**, 7378–7387.
- A. Ciaccafava, C. Hamon, P. Infossi, V. Marchi, M.-T. Giudici-Orticoni and E. Lojou, *Phys. Chem. Chem. Phys.*, 2013, **15**, 16463–16467.
- A. Abou Hamdan, S. Dementin, P.-P. Liebgoth, O. Gutierrez-Sanz, P. Richaud, A. L. De Lacey, M. Rousset, P. Bertrand, L. Cournac and C. Léger, *J. Appl. Chem. Sci.*, 2012, **134**, 8368–8371.
- V. Fourmond, N. Plumeré and C. Léger, *Nat. Rev. Chem.*, 2021, **5**, 348–360.
- A. Fasano, H. Land, V. Fourmond, G. Berggren and C. Léger, *J. Am. Chem. Soc.*, 2021, **143**, 20320–20325.
- J. H. Artz, O. A. Zadovnyy, D. W. Mulder, S. M. Keable, A. E. Cohen, M. W. Ratzloff, S. G. Williams, B. Ginovska, N. Kumar, J. Song, S. E. McPhillips, C. M. Davidson, A. Y. Lyubimov, N. Pence, G. J. Schut, A. K. Jones, S. M. Soltis, M. W. W. Adams, S. Rauegi, P. W. King and J. W. Peters, *J. Am. Chem. Soc.*, 2020, **142**, 1227–1235.
- M. Winkler, J. Duan, A. Rutz, C. Felbek, L. Scholtyssek, O. Lampret, J. Jaenecke, U.-P. Apfel, G. Gilardi, F. Valetti, V. Fourmond, E. Hofmann, C. Léger and T. Happe, *Nat. Commun.*, 2021, **12**, 756.
- A. Rutz, C. K. Das, A. Fasano, J. Jaenecke, S. Yadav, U.-P. Apfel, V. Engelbrecht, V. Fourmond, C. Léger, L. V. Schäfer and T. Happe, *ACS Catal.*, 2023, **13**, 856–865.
- A. Fasano, A. Bailly, J. Wozniak, V. Fourmond and C. Léger, *bioRxiv*, 2023, preprint, DOI: [10.1101/2023.06.23.541094](https://doi.org/10.1101/2023.06.23.541094), ACS Cat. (2024), in press.
- M. Sensi, M. del Barrio, C. Baffert, V. Fourmond and C. Léger, *Curr. Opin. Electrochem.*, 2017, **5**, 135–145.
- J. N. Butt, L. J. C. Jeuken, H. Zhang, J. A. J. Burton and A. L. Sutton-Cook, *Nat. Rev. Methods Primers*, 2023, **3**, 1–19.
- C. Léger and P. Bertrand, *Chem. Rev.*, 2008, **108**, 2379–2438.
- M. Del Barrio, M. Sensi, C. Orain, C. Baffert, S. Dementin, V. Fourmond and C. Léger, *Acc. Chem. Res.*, 2018, **51**, 769–777.
- V. Fourmond, E. S. Wiedner, W. J. Shaw and C. Léger, *J. Am. Chem. Soc.*, 2019, **141**, 11269–11285.
- A. Fasano, C. Baffert, C. Schumann, G. Berggren, J. A. Birrell, V. Fourmond and C. Léger, *J. Am. Chem. Soc.*, 2024, **146**(2), 1455–1466.
- M. Barrio and V. Fourmond, *ChemElectroChem*, 2019, **6**, 4949–4962.
- A. Kubas, C. Orain, D. De Sancho, L. Saujet, M. Sensi, C. Gauquelin, I. Meynial-Salles, P. Soucaille, H. Bottin, C. Baffert, V. Fourmond, R. B. Best, J. Blumberger and C. Léger, *Nat. Chem.*, 2017, **9**, 88–95.
- P. R. Cabotaje, K. Walter, A. Zamader, P. Huang, F. Ho, H. Land, M. Senger and G. Berggren, *ACS Catal.*, 2023, **13**, 10435–10446.



- 37 S. Dementin, B. Burlat, A. L. De Lacey, A. Pardo, G. Adryanczyk-Perrier, B. Guigliarelli, V. M. Fernandez and M. Rousset, *J. Biol. Chem.*, 2004, **279**, 10508–10513.
- 38 A. Abou Hamdan, P.-P. Liebgott, V. Fourmond, O. Gutiérrez-Sanz, A. L. De Lacey, P. Infossi, M. Rousset, S. Dementin and C. Léger, *Proc. Natl. Acad. Sci. U. S. A.*, 2012, **109**, 19916–19921.
- 39 Y. Shomura, M. Taketa, H. Nakashima, H. Tai, H. Nakagawa, Y. Ikeda, M. Ishii, Y. Igarashi, H. Nishihara, K.-S. Yoon, S. Ogo, S. Hirota and Y. Higuchi, *Science*, 2017, **357**, 928–932.
- 40 T. Buhrke, O. Lenz, N. Krauss and B. Friedrich, *J. Biol. Chem.*, 2005, **280**, 23791–23796.
- 41 O. Duché, S. Elsen, L. Cournac and A. Colbeau, *FEBS J.*, 2005, **272**, 3899–3908.
- 42 L. Bowman, L. Flanagan, P. K. Fyfe, A. Parkin, W. N. Hunter and F. Sargent, *Biochem. J.*, 2014, **458**, 449–458.
- 43 J. Fritsch, P. Scheerer, S. Frielingsdorf, S. Kroschinsky, B. Friedrich, O. Lenz and C. M. T. Spahn, *Nature*, 2011, **479**, 249–252.
- 44 Y. Shomura, K.-S. Yoon, H. Nishihara and Y. Higuchi, *Nature*, 2011, **479**, 253–256.
- 45 A. Volbeda, P. Amara, C. Darnault, J.-M. Mouesca, A. Parkin, M. M. Roessler, F. A. Armstrong and J. C. Fontecilla-Camps, *Proc. Natl. Acad. Sci. U. S. A.*, 2012, **109**, 5305–5310.
- 46 C. Schäfer, M. Bommer, S. E. Hennig, J.-H. Jeoung, H. Dobbek and O. Lenz, *Structure*, 2016, **24**, 285–292.
- 47 S. Dementin, V. Belle, P. Bertrand, B. Guigliarelli, G. Adryanczyk-Perrier, A. L. De Lacey, V. M. Fernandez, M. Rousset and C. Léger, *J. Am. Chem. Soc.*, 2006, **128**, 5209–5218.
- 48 H. Adamson, M. Robinson, J. J. Wright, L. A. Flanagan, J. Walton, D. Elton, D. J. Gavaghan, A. M. Bond, M. M. Roessler and A. Parkin, *J. Am. Chem. Soc.*, 2017, **139**, 10677–10686.
- 49 R. M. Bullock and A. Dey, *Chem. Rev.*, 2022, **122**, 11897–11899.
- 50 K. D. Swanson, M. W. Ratzloff, D. W. Mulder, J. H. Artz, S. Ghose, A. Hoffman, S. White, O. A. Zadovnyy, J. B. Broderick, B. Bothner, P. W. King and J. W. Peters, *J. Am. Chem. Soc.*, 2015, **137**, 1809–1816.
- 51 R. Cammack, V. M. Fernandez and K. Schneider, *Biochimie*, 1986, **68**, 85–91.
- 52 V. M. Fernandez, E. C. Hatchikian and R. Cammack, *Biochim. Biophys. Acta, Protein Struct. Mol. Enzymol.*, 1985, **832**, 69–79.
- 53 A. Abou Hamdan, B. Burlat, O. Gutiérrez-Sanz, P.-P. Liebgott, C. Baffert, A. L. De Lacey, M. Rousset, B. Guigliarelli, C. Léger and S. Dementin, *Nat. Chem. Biol.*, 2013, **9**, 15–17.
- 54 A. Volbeda, Y. Montet, X. Vernède, E. C. Hatchikian and J. C. Fontecilla-Camps, *Int. J. Hydrogen Energy*, 2002, **27**, 1449–1461.
- 55 C. Léger, S. Dementin, P. Bertrand, M. Rousset and B. Guigliarelli, *J. Appl. Chem. Sci.*, 2004, **126**, 12162.
- 56 R. Grinter, A. Kropp, H. Venugopal, M. Senger, J. Badley, P. R. Cabotaje, R. Jia, Z. Duan, P. Huang, S. T. Stripp, C. K. Barlow, M. Belousoff, H. S. Shafaat, G. M. Cook, R. B. Schittenhelm, K. A. Vincent, S. Khalid, G. Berggren and C. Greening, *Nature*, 2023, **615**, 541–554.
- 57 J. Kalms, A. Schmidt, S. Frielingsdorf, P. van der Linden, D. von Stetten, O. Lenz, P. Carpentier and P. Scheerer, *Angew. Chem. Int. Ed. Engl.*, 2016, **55**, 5586–5590.
- 58 T. M. Barbosa, C. S. A. Baltazar, D. R. Cruz, D. Lousa and C. M. Soares, *Sci. Rep.*, 2020, **10**, 10540.
- 59 S. Zacarias, A. Temporão, M. del Barrio, V. Fourmond, C. Léger, P. M. Matias and I. A. C. Pereira, *ACS Catal.*, 2019, **9**, 8509–8519.
- 60 F. Leroux, S. Dementin, B. Burlat, L. Cournac, A. Volbeda, S. Champ, L. Martin, B. Guigliarelli, P. Bertrand, J. Fontecilla-Camps, M. Rousset and C. Léger, *Proc. Natl. Acad. Sci. U. S. A.*, 2008, **105**, 11188–11193.
- 61 P.-P. Liebgott, F. Leroux, B. Burlat, S. Dementin, C. Baffert, T. Lautier, V. Fourmond, P. Ceccaldi, C. Cavazza, I. Meynial-Salles, P. Soucaille, J. C. Fontecilla-Camps, B. Guigliarelli, P. Bertrand, M. Rousset and C. Léger, *Nat. Chem. Biol.*, 2010, **6**, 63–70.
- 62 G. Caserta, C. Papini, A. Adamska-Venkatesh, L. Pecqueur, C. Sommer, E. Reijerse, W. Lubitz, C. Gauquelin, I. Meynial-Salles, D. Pramanik, V. Artero, M. Atta, M. Del Barrio, B. Faivre, V. Fourmond, C. Léger and M. Fontecave, *J. Am. Chem. Soc.*, 2018, **140**, 5516–5526.
- 63 P. Ceccaldi, K. Schuchmann, V. Müller and S. J. Elliott, *Energy Environ. Sci.*, 2017, **10**, 503–508.
- 64 C. Brocks, C. K. Das, J. Duan, S. Yadav, U.-P. Apfel, S. Ghosh, E. Hofmann, M. Winkler, V. Engelbrecht, L. V. Schäfer and T. Happe, *ChemSusChem*, 2023, e202301365.
- 65 J. Koo and J. R. Swartz, *Metab. Eng.*, 2018, **49**, 21–27.
- 66 A. S. Bingham, P. R. Smith and J. R. Swartz, *Int. J. Hydrogen Energy*, 2012, **37**, 2965–2976.
- 67 P.-P. Liebgott, A. L. de Lacey, B. Burlat, L. Cournac, P. Richaud, M. Brugna, V. M. Fernandez, B. Guigliarelli, M. Rousset, C. Léger and S. Dementin, *J. Am. Chem. Soc.*, 2011, **133**, 986–997.
- 68 A. Volbeda, L. Martin, P.-P. Liebgott, A. L. De Lacey and J. C. Fontecilla-Camps, *Metallomics*, 2015, **7**, 710–718.
- 69 M. del Barrio, C. Guendon, A. Kpebe, C. Baffert, V. Fourmond, M. Brugna and C. Léger, *ACS Catal.*, 2019, **9**, 4084–4088.
- 70 A. Fasano, C. Guendon, A. Jacq-Bailly, A. Kpebe, J. Wozniak, C. Baffert, M. D. Barrio, V. Fourmond, M. Brugna and C. Léger, *J. Am. Chem. Soc.*, 2023, **145**, 20021–20030.
- 71 J. A. Cracknell, A. F. Wait, O. Lenz, B. Friedrich and F. A. Armstrong, *Proc. Natl. Acad. Sci. U. S. A.*, 2009, **106**, 20681–20686.
- 72 M. J. Lukey, M. M. Roessler, A. Parkin, R. M. Evans, R. A. Davies, O. Lenz, B. Friedrich, F. Sargent and F. A. Armstrong, *J. Am. Chem. Soc.*, 2011, **133**, 16881–16892.
- 73 A. Schmidt, J. Kalms, C. Lorent, S. Katz, S. Frielingsdorf, R. M. Evans, J. Fritsch, E. Siebert, C. Teutloff, F. A. Armstrong, I. Zebger, O. Lenz and P. Scheerer, *Chem. Sci.*, 2023, **14**, 11105–11120.
- 74 C. Schäfer, B. Friedrich and O. Lenz, *Appl. Environ. Microbiol.*, 2013, **79**, 5137–5145.



- 75 R. A. Schmitz, A. Pol, S. S. Mohammadi, C. Hogendoorn, A. H. van Gelder, M. S. M. Jetten, L. J. Daumann and H. J. M. Op den Camp, *ISME J.*, 2020, **14**, 1223–1232.
- 76 H. Koch, A. Galushko, M. Albertsen, A. Schintlmeister, C. Gruber-Dorninger, S. Lückner, E. Pelletier, D. Le Paslier, E. Spieck, A. Richter, P. H. Nielsen, M. Wagner and H. Daims, *Science*, 2014, **345**, 1052–1054.
- 77 C. Greening and R. Grinter, *Nat. Rev. Microbiol.*, 2022, **20**, 513–528.
- 78 C. Baffert, M. Demuez, L. Cournac, B. Burlat, B. Guigliarelli, P. Bertrand, L. Girbal and C. Léger, *Angew Chem. Int. Ed. Engl.*, 2008, **47**, 2052–2054.
- 79 C. Orain, L. Saujet, C. Gauquelin, P. Soucaille, I. Meynial-Salles, C. Baffert, V. Fourmond, H. Bottin and C. Léger, *J. Am. Chem. Soc.*, 2015, **137**, 12580–12587.
- 80 S. Morra, M. Arizzi, F. Valetti and G. Gilardi, *Biochemistry*, 2016, **55**, 5897–5900.
- 81 F. Valetti, S. Morra, L. Barbieri, S. Dezzani, A. Ratto, G. Catucci, S. J. Sadeghi and G. Gilardi, *Faraday Discuss.*, 2024, DOI: [10.1039/D4FD00010B](https://doi.org/10.1039/D4FD00010B).
- 82 P. Rodríguez-Maciá, E. J. Reijerse, M. van Gastel, S. DeBeer, W. Lubitz, O. Rüdiger and J. A. Birrell, *J. Am. Chem. Soc.*, 2018, **140**, 9346–9350.
- 83 P. Rodríguez-Maciá, L. M. Galle, R. Björnsson, C. Lorent, I. Zebger, Y. Yoda, S. P. Cramer, S. DeBeer, I. Span and J. A. Birrell, *Angew Chem. Int. Ed. Engl.*, 2020, **59**, 16786–16794.
- 84 A. A. Oughli, S. Hardt, O. Rüdiger, J. A. Birrell and N. Plumeré, *Chem. Commun.*, 2020, **56**, 9958–9961.
- 85 C. J. Kulka-Peschke, A.-C. Schulz, C. Lorent, Y. Rippers, S. Wahlefeld, J. Preissler, C. Schulz, C. Wiemann, C. C. M. Bernitzky, C. Karafoulidi-Retsou, S. L. D. Wrathall, B. Procacci, H. Matsuura, G. M. Greetham, C. Teutloff, L. Lauterbach, Y. Higuchi, M. Ishii, N. T. Hunt, O. Lenz, I. Zebger and M. Horch, *J. Am. Chem. Soc.*, 2022, **144**, 17022–17032.
- 86 J. Preissler, S. Wahlefeld, C. Lorent, C. Teutloff, M. Horch, L. Lauterbach, S. P. Cramer, I. Zebger and O. Lenz, *Biochim. Biophys. Acta, Bioenerg.*, 2018, **1859**, 8–18.
- 87 A. Cornish-Bowden, *Fundamentals of Enzyme Kinetics (English Edition)*, Wiley-Blackwell, 4th edn, 2013.
- 88 J. N. Butt, M. Filipiak and W. R. Hagen, *Eur. J. Biochem.*, 1997, **245**, 116–122.
- 89 H. R. Pershad, J. L. Duff, H. A. Heering, E. C. Duin, S. P. Albracht and F. A. Armstrong, *Biochemistry*, 1999, **38**, 8992–8999.
- 90 A. Dutta, A. M. Appel and W. J. Shaw, *Nat. Rev. Chem.*, 2018, **2**, 244–252.
- 91 C. C. Page, C. C. Moser, X. Chen and P. L. Dutton, *Nature*, 1999, **402**, 47–52.
- 92 C. L. McIntosh, F. Germer, R. Schulz, J. Appel and A. K. Jones, *J. Am. Chem. Soc.*, 2011, **133**, 11308–11319.
- 93 V. Fourmond, C. Baffert, K. Sybirna, T. Lautier, A. Abou Hamdan, S. Dementin, P. Soucaille, I. Meynial-Salles, H. Bottin and C. Léger, *J. Am. Chem. Soc.*, 2013, **135**, 3926–3938.
- 94 S. Dementin, B. Burlat, V. Fourmond, F. Leroux, P.-P. Liebgott, A. Abou Hamdan, C. Léger, M. Rousset, B. Guigliarelli and P. Bertrand, *J. Am. Chem. Soc.*, 2011, **133**, 10211–10221.
- 95 M. M. Roessler, R. M. Evans, R. A. Davies, J. Harmer and F. A. Armstrong, *J. Am. Chem. Soc.*, 2012, **134**, 15581–15594.
- 96 P. Rodríguez-Maciá, L. Kertess, J. Burnik, J. A. Birrell, E. Hofmann, W. Lubitz, T. Happe and O. Rüdiger, *J. Am. Chem. Soc.*, 2019, **141**, 472–481.
- 97 L. Kertess, A. Adamska-Venkatesh, P. Rodríguez-Maciá, O. Rüdiger, W. Lubitz and T. Happe, *Chem. Sci.*, 2017, **8**, 8127–8137.
- 98 A. J. Cornish, K. Gärtner, H. Yang, J. W. Peters and E. L. Hegg, *J. Biol. Chem.*, 2011, **286**, 38341–38347.
- 99 O. Lampret, A. Adamska-Venkatesh, H. Konegger, F. Wittkamp, U.-P. Apfel, E. J. Reijerse, W. Lubitz, O. Rüdiger, T. Happe and M. Winkler, *J. Am. Chem. Soc.*, 2017, **139**, 18222–18230.
- 100 C. E. Lubner, J. H. Artz, D. W. Mulder, A. Oza, R. J. Ward, S. G. Williams, A. K. Jones, J. W. Peters, I. I. Smalyukh, V. S. Bharadwaj and P. W. King, *Chem. Sci.*, 2022, **13**, 4581–4588.
- 101 S. M. Kim, S. H. Kang, B. W. Jeon and Y. H. Kim, *Bioresour. Technol.*, 2024, **394**, 130248.
- 102 W. J. Ray Jr, *Biochemistry*, 1983, **22**, 4625–4637.
- 103 A. Fasano, V. Fourmond and C. Léger, The difference bidirectionality makes to the kinetic modeling of molecular catalysis, *Curr. Opin. Electrochem.*, 2024, DOI: [10.1016/j.coelec.2024.101489](https://doi.org/10.1016/j.coelec.2024.101489).
- 104 A. Damjanovic, A. Dey and J. O. M. Bockris, *J. Electrochem. Soc.*, 1966, **113**, 739.
- 105 A. K. Jones, S. E. Lamle, H. R. Pershad, K. A. Vincent, S. P. J. Albracht and F. A. Armstrong, *J. Am. Chem. Soc.*, 2003, **125**, 8505–8514.
- 106 V. Fourmond, P. Infossi, M.-T. Giudici-Orticoni, P. Bertrand and C. Léger, *J. Am. Chem. Soc.*, 2010, **132**, 4848–4857.
- 107 M. Horch, L. Lauterbach, M. A. Mroginiski, P. Hildebrandt, O. Lenz and I. Zebger, *J. Am. Chem. Soc.*, 2015, **137**, 2555–2564.
- 108 J. Duan, M. Senger, J. Esselborn, V. Engelbrecht, F. Wittkamp, U.-P. Apfel, E. Hofmann, S. T. Stripp, T. Happe and M. Winkler, *Nat. Commun.*, 2018, **9**, 4726.
- 109 F. A. Armstrong and J. Hirst, *Proc. Natl. Acad. Sci. U. S. A.*, 2011, **108**, 14049–14054.
- 110 A. Fasano, V. Fourmond and C. Léger, *Bioelectrochemistry*, 2024, **155**, 108511.
- 111 N. Chongdar, P. Rodríguez-Maciá, E. J. Reijerse, W. Lubitz, H. Ogata and J. A. Birrell, *Chem. Sci.*, 2023, **14**, 3682–3692.
- 112 M. Sensi, C. Baffert, V. Fourmond, L. De Gioia, L. Bertini and C. Léger, *Sustainable Energy Fuels*, 2021, **5**, 4248–4260.
- 113 P. Rodríguez-Maciá, J. A. Birrell, W. Lubitz and O. Rüdiger, *ChemPlusChem*, 2017, **82**, 540–545.
- 114 D. Adam, L. Bösche, L. Castañeda-Losada, M. Winkler, U.-P. Apfel and T. Happe, *ChemSusChem*, 2017, **10**, 894–902.
- 115 C. Beauflis, H.-M. Man, A. de Poulpique, I. Mazurenko and E. Lojou, *Catalysts*, 2021, **11**, 497.



- 116 A. de Poulpiquet, A. Ciaccafava, R. Gadiou, S. Gounel, M. T. Giudici-Orticoni, N. Mano and E. Lojou, *Electrochem. Commun.*, 2014, **42**, 72–74.
- 117 M. A. Alonso-Lomillo, O. Rüdiger, A. Maroto-Valiente, M. Velez, I. Rodríguez-Ramos, F. J. Muñoz, V. M. Fernández and A. L. De Lacey, *Nano Lett.*, 2007, **7**, 1603–1608.
- 118 K. A. Vincent, J. A. Cracknell, J. R. Clark, M. Ludwig, O. Lenz, B. Friedrich and F. A. Armstrong, *Chem. Commun.*, 2006, 5033–5035.
- 119 V. Fourmond, S. Stapf, H. Li, D. Buesen, J. Birrell, O. Rüdiger, W. Lubitz, W. Schuhmann, N. Plumeré and C. Léger, *J. Am. Chem. Soc.*, 2015, **137**, 5494–5505.
- 120 A. A. Oughli, F. Conzuelo, M. Winkler, T. Happe, W. Lubitz, W. Schuhmann, O. Rüdiger and N. Plumeré, *Angew Chem. Int. Ed. Engl.*, 2015, **54**, 12329–12333.
- 121 H. Li, D. Buesen, S. Dementin, C. Léger, V. Fourmond and N. Plumeré, *J. Am. Chem. Soc.*, 2019, **141**, 16734–16742.
- 122 S. Hardt, S. Stapf, D. T. Filmon, J. A. Birrell, O. Rüdiger, V. Fourmond, C. Léger and N. Plumeré, *Nat. Catal.*, 2021, **4**, 251.
- 123 J. Esselborn, C. Lambertz, A. Adamska-Venkates, T. Simmons, G. Berggren, J. Noth, J. Siebel, A. Hemschemeier, V. Artero, E. Reijerse, M. Fontecave, W. Lubitz and T. Happe, *Nat. Chem. Biol.*, 2013, **9**, 607–609.
- 124 G. Caserta, S. Hartmann, C. Van Stappen, C. Karafoulidi-Retsou, C. Lorent, S. Yelin, M. Keck, J. Schoknecht, I. Sergueev, Y. Yoda, P. Hildebrandt, C. Limberg, S. DeBeer, I. Zebger, S. Frielingsdorf and O. Lenz, *Nat. Chem. Biol.*, 2023, **19**, 498–506.
- 125 W. Roseboom, A. L. De Lacey, V. M. Fernandez, E. C. Hatchikian and S. P. J. Albracht, *J. Biol. Inorg. Chem.*, 2006, **11**, 102–118.
- 126 K. Zuchan, F. Baymann, C. Baffert, M. Brugna and W. Nitschke, *Biochim. Biophys. Acta, Bioenerg.*, 2021, **1862**, 148401.
- 127 N. Chongdar, J. A. Birrell, K. Pawlak, C. Sommer, E. J. Reijerse, O. Rüdiger, W. Lubitz and H. Ogata, *J. Am. Chem. Soc.*, 2018, **140**, 1057–1068.
- 128 H. Land, A. Sekretareva, P. Huang, H. J. Redman, B. Németh, N. Polidori, L. S. Mészáros, M. Senger, S. T. Stripp and G. Berggren, *Chem. Sci.*, 2020, **11**, 12789–12801.
- 129 C. Léger, A. K. Jones, W. Roseboom, S. P. J. Albracht and F. A. Armstrong, *Biochemistry*, 2002, **41**, 15736–15746.
- 130 L. B. Maia, B. K. Maiti, I. Moura and J. J. G. Moura, *Molecules*, 2024, **29**(1), 120.
- 131 P. Ceccaldi, M. C. Marques, V. Fourmond, I. C. Pereira and C. Léger, *Chem. Commun.*, 2015, **51**, 14223–14226.
- 132 A. Parkin, G. Goldet, C. Cavazza, J. C. Fontecilla-Camps and F. A. Armstrong, *J. Am. Chem. Soc.*, 2008, **130**, 13410–13416.
- 133 M. Ludwig, J. A. Cracknell, K. A. Vincent, F. A. Armstrong and O. Lenz, *J. Biol. Chem.*, 2009, **284**, 465–477.
- 134 J. A. Cracknell, K. A. Vincent, M. Ludwig, O. Lenz, B. Friedrich and F. A. Armstrong, *J. Am. Chem. Soc.*, 2008, **130**, 424–425.
- 135 M.-E. Pandelia, V. Fourmond, P. Tron-Infossi, E. Lojou, P. Bertrand, C. Léger, M.-T. Giudici-Orticoni and W. Lubitz, *J. Am. Chem. Soc.*, 2010, **132**, 6991–7004.
- 136 E. Van der Linden, T. Burgdorf, M. Bernhard, B. Bleijlevens, B. Friedrich and S. P. J. Albracht, *J. Biol. Inorg. Chem.*, 2004, **9**, 616–626.
- 137 M. Horch, L. Lauterbach, M. Saggiu, P. Hildebrandt, F. Lendzian, R. Bittl, O. Lenz and I. Zebger, *Angew Chem. Int. Ed. Engl.*, 2010, **49**, 8026–8029.
- 138 T. Burgdorf, E. van der Linden, M. Bernhard, Q. Y. Yin, J. W. Back, A. F. Hartog, A. O. Muijsers, C. G. de Koster, S. P. J. Albracht and B. Friedrich, *J. Bacteriol.*, 2005, **187**, 3122–3132.
- 139 E. van der Linden, T. Burgdorf, A. L. de Lacey, T. Buhrke, M. Scholte, V. M. Fernandez, B. Friedrich and S. P. J. Albracht, *J. Biol. Inorg. Chem.*, 2006, **11**, 247–260.
- 140 P. Kwan, C. L. McIntosh, D. P. Jennings, R. C. Hopkins, S. K. Chandrayan, C.-H. Wu, M. W. W. Adams and A. K. Jones, *J. Am. Chem. Soc.*, 2015, **137**, 13556–13565.

