

Cite this: *Chem. Sci.*, 2021, 12, 983

All publication charges for this article have been paid for by the Royal Society of Chemistry

Molecular mechanism of inhibiting the SARS-CoV-2 cell entry facilitator TMPRSS2 with camostat and nafamostat†

Tim Hempel,^{ab} Lluís Raich,^{‡a} Simon Olsson,^{ab} Nurit P. Azouz,^{cd} Andrea M. Klingler,^c Markus Hoffmann,^{ef} Stefan Pöhlmann,^{ef} Marc E. Rothenberg,^c and Frank Noé^{*abg}

The entry of the coronavirus SARS-CoV-2 into human lung cells can be inhibited by the approved drugs camostat and nafamostat. Here we elucidate the molecular mechanism of these drugs by combining experiments and simulations. *In vitro* assays confirm that both drugs inhibit the human protein TMPRSS2, a SARS-CoV-2 spike protein activator. As no experimental structure is available, we provide a model of the TMPRSS2 equilibrium structure and its fluctuations by relaxing an initial homology structure with extensive 330 microseconds of all-atom molecular dynamics (MD) and Markov modeling. Through Markov modeling, we describe the binding process of both drugs and a metabolic product of camostat (GBPA) to TMPRSS2, reaching a Michaelis complex (MC) state, which precedes the formation of a long-lived covalent inhibitory state. We find that nafamostat has a higher MC population than camostat and GBPA, suggesting that nafamostat is more readily available to form the stable covalent enzyme–substrate intermediate, effectively explaining its high potency. This model is backed by our *in vitro* experiments and consistent with previous virus cell entry assays. Our TMPRSS2–drug structures are made public to guide the design of more potent and specific inhibitors.

Received 11th September 2020
Accepted 6th November 2020

DOI: 10.1039/d0sc05064d

rsc.li/chemical-science

1 Introduction

In December 2019 several cases of unusual and severe pneumonia were reported in the city of Wuhan, China. These cases were traced back to a new coronavirus, SARS-CoV-2 (severe acute respiratory syndrome coronavirus 2); the disease is called COVID-19.¹ As of October 11, 2020 there are over 37 million confirmed COVID-19 cases and more than 1 million deaths,² with both numbers likely to be severe underestimates. Given

estimates of the infection mortality rate of 0.4 to 1.4% (ref. 3–5) the virus has the potential to kill tens of millions of people unless efficient vaccines or drugs are available.

As other coronaviruses,^{6–9} SARS-CoV-2 exploits host proteins to initiate cell-entry, in particular TMPRSS2 and ACE2, two membrane-bound proteins expressed in the upper and lower respiratory tract.^{10–13} TMPRSS2 contains an extracellular trypsin-like serine-protease domain that can proteolytically activate the spike (S) protein on the surface of SARS-CoV-2 viral particles¹⁴ (Fig. 1). While in certain cell lines, the S-protein can also be activated by the endo/lysosomal pH-dependent cysteine protease cathepsin L,^{14,15} virus entry into human airway cells^{14,16} seems to depend on TMPRSS2 but not cathepsin L. Consistently, epidemiological data of prostate cancer patients

^aFreie Universität Berlin, Department of Mathematics and Computer Science, Berlin, Germany. E-mail: frank.noë@fu-berlin.de

^bFreie Universität Berlin, Department of Physics, Berlin, Germany

^cDivision of Allergy and Immunology, Cincinnati Children's Hospital Medical Center, Department of Pediatrics, University of Cincinnati College of Medicine, Cincinnati, OH, USA

^dDepartment of Pediatrics, University of Cincinnati College of Medicine, Cincinnati, OH, USA

^eInfection Biology Unit, German Primate Center – Leibniz Institute for Primate Research, Göttingen, Germany

^fFaculty of Biology and Psychology, University Göttingen, Göttingen, Germany

^gRice University, Department of Chemistry, Houston, TX, USA

^hChalmers University of Technology,

undergoing androgen-deprivation therapies, which lowers TMPRSS2 levels, indicate a lower risk of contracting the SARS-CoV-2 infection.¹⁷ We further note that low concentration levels of TMPRSS2 are observed in children and infants, possibly explaining lower risks of severe COVID-19 infections in younger age groups.¹⁸

TMPRSS2 is also exploited by other coronaviruses and influenza A viruses for activation of surface glycoproteins, viral spread, and pathogenesis.^{19–25} TMPRSS2 knock-out mice have no phenotype in the absence of infection,²⁶ indicating that inhibiting TMPRSS2 function might not be associated with substantial unwanted side effects. As a result, TMPRSS2 is a promising therapeutic target in the context of influenza A and coronavirus infection, including SARS-CoV-2. Since TMPRSS2 is host encoded and thus genetically stable, treatment should be associated with a low risk of drug resistance.

Here, we study the structural basis and molecular mechanism of TMPRSS2 inhibition by nafamostat, camostat, and its metabolic product 4-(4-guanidinobenzoyloxy)phenylacetic acid (GBPA). These guanidinobenzoyl-containing drugs are approved for human use in Japan and have been demonstrated to inhibit SARS-CoV-2 cell-entry.^{14,27–29} A recent survey of FDA approved drugs further found nafamostat to be an effective inhibitor of SARS-CoV-2 infection in human lung-cell cultures.³⁰ We report experimental measurements demonstrating that nafamostat and camostat inhibit TMPRSS2 activity by using our recently established cell-based assay,³¹ consistent with *in vitro* enzymatic TMPRSS2 activity assays.³²

Despite the hopes associated with TMPRSS2 inhibition, we are, as yet, lacking an experimental structure. We here go beyond the previous dependence on homology models by an extensive 330 microseconds of high-throughput all-atom molecular dynamics (MD) simulations and Markov modeling. This approach provides an ensemble of equilibrium structures of the protein–drug complex and also drug binding kinetics. We show that nafamostat, camostat, and GBPA are covalent inhibitors with an identical covalent complex, but their different inhibitory activity can be explained by different populations of their Michaelis complex preceding the covalent complex. These findings, combined with the simulation structures that we make publicly available, provide an important basis for developing more potent and specific TMPRSS2 inhibitors.

2 Results

Camostat and nafamostat inhibit the catalytic activity of TMPRSS2

First we confirm that camostat and nafamostat are TMPRSS2 inhibitors. To this end, we employ our recently reported activity assay³¹ of the full-length TMPRSS2 protein on the surface of live cells with both inhibitors (Fig. 2A). Briefly, we transfected the human cell-line HEK-293T with a TMPRSS2 expression vector. We then measured the protease activity of the transfected cells using the fluorogenic peptide substrate BOC-QAR-AMC, following incubation of the cell with increasing inhibitor concentrations. Peptide-digestion induced a minimal increase




Fig. 2 (A) Chemical structure of nafam

values to 142 ± 31 nM and 55 ± 7 nM (Fig. 2C). Our results are consistent with the finding that both drugs inhibit cell entry of SARS-CoV-2 and other coronaviruses, and that nafamostat is the most potent inhibitor.^{27,28,32}

Note that in humans, camostat is rapidly processed to 4-(4-guanidinobenzoyloxy)phenylacetic acid (GBPA) (Fig. 2A).³³ It has been recently shown that GBPA also inhibits TMPRSS2 and cell entry of SARS-CoV-2 viruses, although slightly less efficiently than camostat.²⁹ Hence, we subsequently study the molecular interactions between TMPRSS2 and all three compounds: camostat, GBPA, and nafamostat.

Equilibrium structures of TMPRSS2 in complex with camostat and nafamostat


We now set off to investigate the molecular mechanism of TMPRSS2 inhibition by nafamostat, camostat, and its metabolite GBPA. No TMPRSS2 crystal structure is available to date, however it has been shown that all-atom MD simulations can reliably model the equilibrium structures of proteins when (i) a reasonable model is available as starting structure, and (ii) simulations sample extensively, such that deficiencies of the starting structure can be overcome.^{35–39}

Here, we initialize our simulations with recent homology models of the TMPRSS2 protease domain and with camostat/nafamostat docked to them.⁴⁰ Trypsins adopt a common fold and share an active-site charge relay system whose structural

requirements for catalytic activity are well understood;⁴¹ we select our MD model consistent with these structural requirements. In particular, we focus on systems with Asp435 (substrate recognition) deprotonated and His296 (catalytic function) in a neutral form (N_δ protonated), as well as on the interactions of a charged lysine nearby the catalytic Asp345 (Fig. S1 and S2†).

In order to avoid artifacts of the initial structural model and to simulate the equilibrium ensemble of the TMPRSS2–drug complexes, we collected a total of 100 μ s of simulation data for TMPRSS2–camostat, 50 μ s for TMPRSS2–GBPA, and 180 μ s for TMPRSS2–nafamostat. Every drug dataset has converged RMSD distributions (Fig. S5†) and samples various drug poses and multiple association/dissociation events. Using Markov modeling^{42–46} we derive the structures of the long-lived (meta-stable) states and characterize protein–drug binding kinetics and thermodynamics.

We find TMPRSS2 has flexible loops around the binding site but maintains stable structural features shared by other trypsin-like proteases (Fig. 3A and S6†). After formation of a non-covalent substrate–enzyme complex (binding step, Fig. 2B), trypsins cleave peptide-like bonds in two catalytic steps, assisted by a conserved catalytic triad (Asp345, His296, and Ser441 in TMPRSS2). The first step involves the formation of a covalent acyl–enzyme intermediate between the substrate and Ser441.⁴¹ During this step, His296 serves as a general base to deprotonate



the nucleophilic Ser441, and subsequently as a general acid to protonate the leaving group of the substrate. The second step involves the hydrolysis of the acyl-enzyme intermediate, releasing the cleaved substrate and restoring the active form of the enzyme (Fig. 2B).

Along these two steps, the so called “oxyanion hole”, formed by the backbone NHs of Gly439 and Ser441, helps to activate and stabilize the carbonyl of the scissile bond. Another important structural feature is the S1 pocket, which contains a well conserved aspartate (Asp435) that is essential for substrate binding and recognition. At the opposite site of the S1 pocket, a loop containing a hydrophobic patch delimits the binding region of substrates within enzymatic active site. All these structural elements, known to play crucial roles in the function of serine proteases,⁴¹ are generally stable and preserved in our equilibrium structures (*cf.* Fig. S2 and S6†).

Structural basis of TMPRSS2 inhibition by camostat and nafamostat

Drugs with a guanidinobenzoyl moiety can inhibit trypsins by mimicking their natural substrates (Fig. 3B). Indeed, the ester group, resembling a peptide bond, can react with the catalytic serine with rates that are orders of magnitude faster,⁴⁷ forming the acyl-enzyme intermediate. In contrast to peptide catalysis, the drug's guanidinobenzoyl group stays covalently linked to the catalytic serine with a small off-rate, rendering it an effective chemical inhibitor.⁴⁸ Note that in its inhibited state, the TMPRSS2 active site is modified such that protease activity is disabled, preventing SARS-CoV-2 S-protein cleavage.

The present MD simulations sample different conformations of the complex formed by the enzyme with each of the drugs that precede the covalent substrate-enzyme complex. We can, therefore, elucidate their binding and how specific interactions stabilize different modes. However, please note that our simulations do not simulate the covalent complex's formation. All binding modes mimic interactions made between trypsins and their natural substrates, in which lysine heads interact with a conserved aspartate in the S1 pocket (Asp435, Fig. 3B). In camostat, its metabolic product GPBA, and nafamostat, the role of the lysine heads is taken by the guanidinium heads which bind in the S1 pocket and also interact with Asp435 (Fig. 3C–E). However, the guanidinium–Asp435 salt bridge is formed and broken transiently especially for camostat and GPBA (Fig. 3G–I), indicating that these drugs are not optimized for the TMPRSS2 pocket (Fig. S4†).

Nafamostat also binds in a “reverse” orientation where the amidinium head binds into the S1 pocket and interacts with Asp435 (Fig. 3F).⁴⁰ In this orientation, the guanidinium head mainly interacts with Glu299, with the drug reactive center slightly displaced from the oxyanion hole, while the “forward” orientation (Fig. 3E) keeps the amidinium head mainly nearby Val280, with the ester center well positioned for the reaction (Fig. S3†). This observation is in agreement with several crystal structures of acyl-enzyme intermediates between different trypsins and guanidinobenzoyl molecules bound to the S1 pocket (*e.g.* PDBs 2AH4,⁴⁹ 3DFL,⁵⁰ 1GBT⁵¹). There are also

“inverse substrates” known to react with rates comparable to the ones of normal esters, suggesting that the inverted nafamostat orientation may also be reactive.⁴¹

A fraction of the bound-state structures resembles a reactive Michaelis complex (MC) which fulfills the necessary criteria for catalysis of the inhibitory acyl-enzyme complex: small distances of (i) the drug ester carbon to catalytic serine oxygen, and (ii) the catalytic serine hydrogen to catalytic histidine nitrogen (Methods). We observe that besides Asp435 binding to the S1 pocket, drugs in the MC state are particularly stabilized by the oxyanion hole. Our model predicts that nafamostat has the highest MC state population followed by camostat and GPBA (Fig. 4), an order that coincides with the one of experimental drug binding affinities.²⁹ We note that the relative free energies of binding to the MC states are significantly different between nafamostat (2.1 ± 0.1 kcal mol^{−1}) and the other drugs (2.8 ± 0.1 kcal mol^{−1} and 3.1 ± 0.2 kcal mol^{−1} for camostat and GPBA, respectively), with the bootstrap sample




Fig. 4 Binding kinetics model of camostat (top), camostat metabolic product GBPA (middle), and nafamostat (bottom). Inhibition process is depicted from left (apo state) to right (covalent complex). Single representative structures for each intermediate state are shown – note that all states have significant flexibility. Drugs are depicted in yellow, catalytic triad residues in black, leaving groups and covalent group are denoted by LG and CG, respectively. Rates and populations predicted by our model are annotated at reaction arrows and states, respectively. The covalent complex is illustrated using a structure with prostaticin (PDB 3DFL⁵⁰).

arise from either (1) the

Our binding assays provide evidence that both inhibitors directly act on TMPRSS2 and that nafamostat is more potent compared to camostat, and this qualitative difference is in agreement with complementary *in vitro* assays on purified protein construct³² or cell-entry assays.^{27,28} We note that the absolute IC₅₀ values differ between these three assay types, reflecting differences in experimental conditions and which function is being inhibited and measured.

While no crystallographic structure of TMPRSS2 is available, we provide extensive 330 microseconds of all-atom MD simulations starting from a homology model that generate stable equilibrium structure ensembles of the protein–drug complexes. These simulations sample multiple association/dissociation events and various drug poses in the protein active site. Our analyses show that the non-covalent complexes of nafamostat, camostat, and its metabolic product GBPA are relatively short-lived, suggesting that the main inhibitory effect is due to the formation of the long-lived covalent acyl–enzyme complex between the drug's guanidinobenzoyl moiety and the catalytic serine of TMPRSS2.

Although the MC state is not the main cause of inhibition, its population directly translates into the potency of the inhibitor, as higher MC population corresponds to a higher catalytic rate and therefore yields a larger population of inhibited enzyme. Consistently with the higher potency of nafamostat, it is found to have a threefold more stable MC compared to camostat, and sixfold compared to GBPA. A second contribution may be the pK_a of drug leaving groups, affecting the rate of enzyme acylation.

Our detailed models of the thermodynamic and kinetics of inhibitor binding highlight the bound state's heterogeneity, with both drugs adopting multiple distinct poses. We note the importance of residue Asp435 in the conserved S1-pocket, which stabilizes the MC state and helps to orient the reactive molecules in a conformation that is suited for catalysis. Nafamostat has two groups that can potentially bind into the S1 pocket, whereas camostat has only one. However, we find that the population of S1 associated states are similar between nafamostat, camostat, and GBPA, suggesting that non-covalent inhibition is likely a minor contribution to the overall inhibition of TMPRSS2.

We conclude that the design of future TMPRSS2 inhibitors with increased potency and specificity should incorporate the following points:

First, stabilizing the non-covalent complex with the TMPRSS2 active site is beneficial for both, covalent and non-covalent inhibitors. As S1 pocket binding is a major contribution to the stability of the non-covalent complex, effective drugs may contain hydrogen bond donors and positively charged moieties that could interact principally with Asp435, but also with different backbone carbonyls of the loops that compose the cavity (e.g. from Trp461 to Gly464).

Second, for covalent inhibitors, we must consider that the catalytic serine is at a distance of around 1.3 nm from Asp435. Thus, the reactive center of an effective drug and its S1-interacting moieties should be within that distance. We note that, even though all three molecules fit well in the overall active

site, the guanidinobenzoyl moiety is slightly shorter than the ideal size of the TMPRSS2 cavity (Fig. S4†). We further suggest that a drug should be size-compatible to the hydrophobic patch on the S1 distal site (Fig. 3A and S4†). We speculate that drugs with a large end to end distance and high rigidity may not fit well in the described TMPRSS2 scaffold, and in particular, might be significantly less reactive.

Third, optimizing the pK_a of the drug's leaving group might be beneficial for improving covalent TMPRSS2 inhibitors. The first step of the reaction would be faster, and the acetyl–enzyme intermediate would accumulate. We note that the deacetylation off-rate must be very low, ideally on the order of magnitude of guanidinobenzoyl moiety containing drugs.

Finally, we make our simulated equilibrium structures of TMPRSS2 in complex with the simulated drugs available, hoping they will be useful to guide future drug discovery efforts.

4 Materials and methods

TMPRSS2 activity assays

The TMPRSS2 activity assay was described previously.³¹ Briefly, we transfected

plasmid (no TMPRSS2; background noise). We used *scipy*'s⁵⁵ curve fitting algorithms to extract the IC50 with error estimates.

Molecular dynamics simulations

MD simulations were run with OpenMM 7.4.0 (ref. 56) using the CHARMM 36 force field (2019 version).⁵⁷ Camostat and nafamostat structures were taken from PubChem⁵⁸ with PubChem CIDs 4413 (nafamostat) and 2536 (camostat), respectively, and modeled with the CHARMM general force field (CGenFF v. 4.3).⁵⁹ We generated our MD setups with CharmmGUI.⁶⁰ We initiate a simulation box of side length 7.5 nm with a NaCl ion concentration of 0.1 mol l⁻¹ at neutral charge and the TIP3P water model.⁶¹ The setups contain 12038 (camostat), 12030 (GBPA), and 12039 (nafamostat) water molecules, respectively.

We run simulations in the NPT ensemble and keep the temperature at 310 K (physiological temperature) and the pressure at 1 bar. We use a Langevin integrator with 5 fs integration step and heavy hydrogen approximation (4 amu). PME electrostatics, rigid water molecules, and a 1 nm cutoff for non-bonded interactions are used. Simulation times vary between 100 and 500 ns and accumulate to 100 μ s (camostat), 50 μ s (GBPA), 180 μ s (nafamostat), respectively. Structures were visualized using VMD.⁶²

Due to the lack of a crystal structure for TMPRSS2, MD simulations were seeded from a homology model. It is taken from ref. 40, model 3W94 is chosen based on precursive MD analyses that showed that 3W94 has the most stable catalytic triad configuration (Fig. S1 and S2†). The construct includes amino acids 256 to 491 of the full sequence, corresponding to the catalytic chain except for a C-terminal Glycine missing due to homology modeling against a shorter sequence. MD simulations are seeded as follows: equilibrated docking poses (highest scorers of ref. 40) of the ligand were generated in a precursive run using another homology model. We note that the used camostat docking pose resembles the one described by ref. 63. This data set was equilibrated with local energy minimization, 100 ps simulations with 2 fs time steps in NVT and NPT ensemble subsequently. Frames are selected based on a preliminary metastability analysis, protein conformation is constraint to 3W94 homology model using a constraint force minimizing minRMSD. Production run MD simulations are started from these poses, *i.e.* from the same protein configuration and with 77 (nafamostat) and 60 (camostat) ligand docking poses, respectively. To ensure convergence of sampling statistics, we ran multiple adaptive runs of simulations, seeding new simulations with coordinates associated with sparsely sampled states.

We later added the camostat metabolite GBPA by following the same setup procedure. Due to its similarity to camostat, we seeded production simulations from representative structures of the camostat stage 1 Markov model (described below) using 200 representative structures.

Markov modeling

We model the binding and unbinding rates in a two step procedure using Markov state type models.^{42–45,64–66} First, we

describe drug unbound and associated states using a hidden Markov model (HMM).⁶⁷ Second, we define a reactive state by using distance cutoffs.

In detail, in the first stage we define distance features between drug guanidinium group and TMPRSS2 Asp435 (minimal distance), drug amidinium group and TMPRSS2 Asp435 (minimal distance, nafamostat only). We further use a binary “reactive” distance feature defined by drug ester carbon to catalytic Ser441-OG, and catalytic serine (HG) to catalytic histidine (NE2) and a threshold of 0.35 nm. If both last mentioned distances are below the threshold, both nucleophilic attack of the serine-OG to the drug ester group and proton transfer

GBPA, and 6.07 mM (5.55, 6.93) for nafamostat (68% confidence intervals).

Kinetic model

Simplifying the binding kinetics into a three-state model describing the binding to/dissociation from the Michaelis complex (ligand concentration c and rates k_{on} , k_{off}), catalytic rate of entering the covalent complex (k_{cat}) and dissociation to the apo state (k_{dis}), the kinetics are described by the rate matrix:

$$K = \begin{bmatrix} -ck_{\text{on}} & ck_{\text{on}} & 0 \\ k_{\text{off}} & -k_{\text{off}} - k_{\text{cat}} & k_{\text{cat}} \\ k_{\text{dis}} & 0 & -k_{\text{dis}} \end{bmatrix} \quad (2)$$

with the (unnormalized) equilibrium distribution

$$\pi = \begin{bmatrix} \frac{k_{\text{dis}}(k_{\text{off}} + k_{\text{cat}})}{ck_{\text{on}} + k_{\text{cat}}} \\ k_{\text{dis}}/k_{\text{cat}} \\ 1 \end{bmatrix} \quad (3)$$

The overall dissociation constant is then:

$$K_d = \frac{\pi_1}{\pi_2 + \pi_3} = \frac{k_{\text{dis}}(k_{\text{off}} + k_{\text{cat}})}{k_{\text{on}}(k_{\text{dis}} + k_{\text{cat}})} \quad (4)$$

The non-covalent dissociation constant of the Michaelis complex:

$$K_d^M = \frac{\pi_1}{\pi_2} = \frac{k_{\text{off}}}{ck_{\text{on}} + k_{\text{cat}}} \quad (5)$$

The dissociation constant scales as:

$$K_d = K_d^M \frac{k_{\text{dis}}}{k_{\text{cat}} + k_{\text{dis}}} \quad (6)$$

and thus the association constant scales with the stability of the Michaelis complex by a constant factor given by the rates of chemical catalysis and dissociation:

$$K_a = K_a^M \frac{k_{\text{cat}} + k_{\text{dis}}}{k_{\text{dis}}} \quad (7)$$

Software and data availability

Structural ensembles of camostat, GBPA, and nafamostat binding poses are published online at https://github.com/noegroup/tmpRSS2_structures.

Author contributions

M. H., S. P., M. E. R., and F. N. designed the study. T. H., L. R., S. O., N. P. A., and A. M. K. performed research. T. H., L. R., S. O., and N. P. A. analyzed the data. T. H., L. R., S. O., F. N. wrote the manuscript.

Conflicts of interest

M. E. R. is a consultant for Pulm One, Spoon Guru, ClostraBio, Serpin Pharm, Allakos, Celgene, Astra Zeneca, Arena Pharmaceuticals, GlaxoSmith Kline, Guidepoint and Suvretta Capital Management

- 10 C. G. K. Ziegler, *et al.*, SARS-CoV-2 Receptor ACE2 is an Interferon-Stimulated Gene in Human Airway Epithelial Cells and is Detected in Specific Cell Subsets across Tissues, *Cell*, 2020, **181**(5), 1016–1035.
- 11 S. Lukassen, *et al.*, SARS-CoV-2 Receptor ACE2 and TMPRSS2 are Primarily Expressed in Bronchial Transient Secretory Cells, *EMBO J.*, 2020, **39**(10), e105114.
- 12 K. Bilinska, P. Jakubowska, C. S. Von Bartheld and R. Butowt, Expression of the SARS-CoV-2 Entry Proteins, ACE2 and TMPRSS2, in Cells of the Olfactory Epithelium: Identification of Cell Types and Trends with Age, *ACS Chem. Neurosci.*, 2020, **11**(11), 1555–1562.
- 13 Y. J. Hou, *et al.*, SARS-CoV-2 Reverse Genetics Reveals a Variable Infection Gradient in the Respiratory Tract, *Cell*, 2020, **182**(2), 429–446.
- 14 M. Hoffmann, *et al.*, SARS-CoV-2 Cell Entry Depends on ACE2 and TMPRSS2 and is Blocked by a Clinically Proven Protease Inhibitor, *Cell*, 2020, **181**(2), 271–280.
- 15 T. Ou *et al.*, Hydroxychloroquine-mediated inhibition of SARS-CoV-2 entry is attenuated by TMPRSS2, 2020, bioRxiv, DOI: 10.1101/2020.07.22.216150.
- 16 M. Hoffmann, *et al.*, Chloroquine does not inhibit infection of human lung cells with SARS-CoV-2, *Nature*, 2020, **585**, 588–590.
- 17 M. Montopoli, *et al.*, Androgen-Deprivation Therapies for Prostate Cancer and Risk of Infection by SARS-CoV-2: A Population-Based Study (N = 4532), *Ann. Oncol.*, 2020, **31**(8), 1040–1045.
- 18 B. A. Schuler *et al.*, Age-Determined Expression of Priming Protease TMPRSS2 and Localization of SARS-CoV-2 Infection in the Lung Epithelium, 2020, bioRxiv, DOI: 10.1101/2020.05.22.111187.
- 19 Y. Zhou, *et al.*, Protease Inhibitors Targeting Coronavirus and Filovirus Entry, *Antiviral Res.*, 2015, **116**, 76–84.
- 20 N. Iwata-Yoshikawa, *et al.*, TMPRSS2 Contributes to Virus Spread and Immunopathology in the Airways of Murine Models after Coronavirus Infection, *J. Virol.*, 2019, **93**(6), e01815–18.
- 21 B. Hatesuer, *et al.*, Tmprss2 is Essential for Influenza H1N1 Virus Pathogenesis in Mice, *PLoS Pathog.*, 2013, **9**(12), e1003774.
- 22 C. Tarnow, *et al.*, TMPRSS2 is a Host Factor that is Essential for Pneumotropism and Pathogenicity of H7N9 Influenza A Virus in Mice, *J. Virol.*, 2014, **88**(9), 4744–4751.
- 23 K. Sakai, *et al.*, The Host Protease TMPRSS2 Plays a Major Role in in Vivo Replication of Emerging H7N9 and Seasonal Influenza Viruses, *J. Virol.*, 2014, **88**(10), 5608–5616.
- 24 R. L. O. Lambertz, *et al.*, Tmprss2 Knock-out Mice are Resistant to H10 Influenza A Virus Pathogenesis, *J. Gen. Virol.*, 2019, **100**(7), 1073–1078.
- 25 R. L. O. Lambertz, *et al.*, H2 Influenza A Virus is not Pathogenic in Tmprss2 Knock-out Mice, *Virol. J.*, 2020, **17**(1), 56.
- 26 T. S. Kim, C. Heinlein, R. C. Hackman and P. S. Nelson, Phenotypic Analysis of Mice Lacking the Tmprss2-Encoded Protease, *Mol. Cell. Biol.*, 2006, **26**(3), 965–975.
- 27 M. Hoffmann, *et*

- 43 N. Singhal, C. D. Snow and V. S. Pande, Using Path Sampling to Build Better Markovian State Models: Predicting the Folding Rate and Mechanism of a Tryptophan Zipper Beta Hairpin, *J. Chem. Phys.*, 2004, **121**(1), 415–425.
- 44 F. Noé, C. Schütte, E. Vanden-Eijnden, L. Reich and T. R. Weikl, Constructing the Equilibrium Ensemble of Folding Pathways from Short Off-Equilibrium Simulations, *Proc. Natl. Acad. Sci. U. S. A.*, 2009, **106**(45), 19011–19016.
- 45 J.-H. Prinz, *et al.*, Markov Models of Molecular Kinetics: Generation and Validation, *J. Chem. Phys.*, 2011, **134**(17), 174105.
- 46 B. E. Husic and V. S. Pande, Markov State Models: From an Art to a Science, *J. Am. Chem. Soc.*, 2018, **140**(7), 2386–2396.
- 47 B. Zerner, R. P. M. Bond and M. L. Bender, Kinetic Evidence for the Formation of Acyl-Enzyme Intermediates in the α -Chymotrypsin-Catalyzed Hydrolyses of Specific Substrates, *J. Am. Chem. Soc.*, 1964, **86**(18), 3674–3679.
- 48 M. K. Ramjee, I. M. Henderson, S. B. McLoughlin and A. Padova, The Kinetic and Structural Characterization of the Reaction of Nafamostat with Bovine Pancreatic Trypsin, *Thromb. Res.*, 2000, **98**(6), 559–569.
- 49 E. S. Radisky, J. M. Lee, C.-J. K. Lu and D. E. Koshland, Insights into the serine protease mechanism from atomic resolution structures of trypsin reaction intermediates, *Proc. Natl. Acad. Sci. U. S. A.*, 2006, **103**(18), 6835–6840.
- 50 K. W. Rickert, *et al.*, Structure of Human Prostatic, a Target for the Regulation of Hypertension, *J. Biol. Chem.*, 2008, **283**(50), 34864–34872.
- 51 W. F. Mangel, *et al.*, Structure of an acyl-enzyme intermediate during catalysis: (guanidinobenzoyl)trypsin, *Biochemistry*, 1990, **29**(36), 8351–8357.
- 52 Z. Rappoport, *Handbook of Tables for Organic Compound Identification*, Cleveland, Chemical Rubber Co., 1967.
- 53 P. Pearce and R. Simkins, Acid Strengths of Some Substituted Picric Acids, *Can. J. Chem.*, 1968, **46**(2), 241–248.
- 54 C. Ritz, F. Baty, J. C. Streibig and D. Gerhard, Dose-Response Analysis Using R, *PLoS One*, 2015, **10**(12), e0146021.
- 55 P. Virtanen, *et al.*, SciPy 1.0: Fundamental Algorithms for Scientific Computing in Python, *Nat. Methods*, 2020, **17**, 261–272.
- 56 P. Eastman, *et al.*, OpenMM 7: Rapid Development of High Performance Algorithms for Molecular Dynamics, *PLoS Comput. Biol.*, 2017, **13**(7), e1005659.
- 57 R. B. Best, *et al.*, Optimization of the Additive CHARMM All-Atom Protein Force Field Targeting Improved Sampling of the Backbone ϕ , ψ and Side-Chain χ 1 and χ 2 Dihedral Angles, *J. Chem. Theory Comput.*, 2012, **8**(9), 3257–3273.
- 58 S. Kim, *et al.*, PubChem 2019 Update: Improved Access to Chemical Data, *Nucleic Acids Res.*, 2019, **47**(D1), D1102–D1109.
- 59 K. Vanommeslaeghe, *et al.*, CHARMM General Force Field: A Force Field for Drug-like Molecules Compatible with the CHARMM All-Atom Additive Biological Force Fields, *J. Comput. Chem.*, 2009, **31**, 671–690.
- 60 S. Jo, T. Kim, V. G. Iyer