



Cite this: *Org. Biomol. Chem.*, 2024, **22**, 5224

Received 8th April 2024,
Accepted 1st June 2024
DOI: 10.1039/d4ob00573b
rsc.li/obc

Chemodivergent phosphorylation of diazocarboxylates: light-on vs. light-off reactions†

Jalaj Kumar Pathak,^{a,b} Ruchir Kant^c and Namrata Rastogi  ^{a,b}


By tapping into the divergent reactivity of diazocarboxylates under thermal and photocatalytic conditions, we could develop chemodivergent phosphorylation protocols for α -diazocarboxylates with trialkyl phosphites. While the thermal reaction led to N-P bond formation affording phosphorylated hydrazones, the visible light-mediated reaction furnished phosphorylated aryl carboxylates through C-P bond formation. Both reactions are notable for their operational simplicity and mild conditions affording products in good yields without the requirement of a metal, base or photocatalyst.

Introduction

Phosphorus containing organic scaffolds are found extensively among natural as well as synthetic medicinal compounds,¹ agrochemicals² and functional materials.³ Organophosphorus compounds are also valuable as catalysts, ligands, protecting groups and building blocks in organic chemistry.⁴ Therefore, numerous synthetic methods to incorporate phosphorus into organic scaffolds have been invented.⁵ Yet development of more efficient protocols to access various organophosphorus molecules is highly desired.

The diazo compounds are undoubtedly amongst the most versatile organic substrates owing to their tendency to serve as dipoles, nucleophiles, electrophiles, and precursors to carbenes, ketenes and radical intermediates.⁶ This unique reactivity profile of the diazo group offers great opportunity to develop chemodivergent reactions by modulating the reaction conditions.⁷ For instance, aryl diazoacetates form free carbenes upon visible light irradiation⁸ whereas they serve as C-nucleophiles and N-electrophiles under thermal conditions. We envisioned that a nucleophile would undergo both an insertion reaction with the electrophilic diazocarbene and nucleophilic addition reaction with the terminal nitrogen of the diazo group, under suitable conditions. We opted to explore the phosphorus nucleophiles for these reactions con-

sidering the significance of organophosphorus scaffolds, as mentioned above, and also because phosphorus can adopt various oxidation states offering options for further transformation of the products. Notably, several reactions of diazo compounds with $\text{P}(\text{O})\text{H}$ compounds to access organophosphorus scaffolds featuring C-P or N-P linkages have been reported in the recent literature. For instance, phosphine oxides undergo transition metal-mediated P-H insertion or reductive catalytic coupling with diazo substrates (or diazo intermediates *in situ* generated from tosylhydrazones) forming $\text{C}(\text{sp}^3)\text{-P}$ bonds (Scheme 1A).⁹ On the other hand, Zhu and co-workers reported



Scheme 1 Carbon vs. nitrogen phosphorylation of diazo compounds.

^aMedicinal & Process Chemistry Division, CSIR-Central Drug Research Institute, Lucknow-226031, India. E-mail: namrata.rastogi@cdri.res.in

^bAcademy of Scientific and Innovative Research (AcSIR), Ghaziabad-201002, India

^cBiochemistry & Structural Biology Division, CSIR-Central Drug Research Institute, Lucknow-226031, India

† Electronic supplementary information (ESI) available: ^1H , ^{13}C , ^{31}P and ^{19}F NMR spectra of all new compounds. CCDC 2342435. For ESI and crystallographic data in CIF or other electronic format see DOI: <https://doi.org/10.1039/d4ob00573b>

a base-mediated nucleophilic addition of phosphonates with α -diazoesters resulting in *E*-phosphinamides with N-P bonds (Scheme 1B).^{10a} However, during the preparation of this manuscript Nan and Yi published photocatalytic phosphorylation of H-phosphine oxides with α -diazoesters to afford *Z*-phosphinic hydrazone.^{10b} Similar N-P bond formation was reported by Pullarkat and Leung through palladium-catalyzed asymmetric diarylphosphine addition to α -diazoesters leading to the enantioenriched phosphinic hydrazone (Scheme 1B).¹¹

To the best of our knowledge phosphites have never been used as substrates with diazo compounds except for an iron-catalyzed reaction of trimethyl phosphite with phenyldiazomethane and ethyl diazoacetate forming dimethyl benzylphosphonate and ethyl 2-(dimethoxyphosphoryl)acetate, respectively, through a phosphorus ylide intermediate¹² (Scheme 1A). We hereby report diazocarboxylate phosphorylation with phosphites *via* nucleophilic substitution or carbene insertion generating divergent organophosphorus scaffolds featuring N-P or C-P bonds, respectively, by simple “switch-off” or “switch-on” of blue light (Scheme 1; this work).

We first started investigating the nucleophilic substitution reaction with methyl 2-diazo-2-phenylacetate **1a** and trimethyl phosphite **2a** as model substrates (Table 1).

The initial experiments with **1a** and **2a** in a 1:1 ratio employing DMSO or DMF as the solvent at room temperature failed to furnish any identifiable product (entries 1 and 2). Upon changing the solvent to toluene, the expected phosphinamide product **3a** was isolated in 15% yield (entry 3). Further solvent screening revealed moderate and almost similar yields in methanol (40%) and dichloromethane (42%) (entries 4 and 5). However, the yield improved to 50% in acetonitrile with comparable yields in chloroform (48%) and acetone (46%)

Table 1 Optimization of conditions for thermal phosphorylation^{a,b}

Entry	1a : 2a	Solvent	Yield of 3a ^c (%)
1 ^d	1 : 1	DMSO	0
2 ^d	1 : 1	DMF	0
3	1 : 1	PhMe	15
4	1 : 1	MeOH	40
5	1 : 1	CH ₂ Cl ₂	42
6	1 : 1	CHCl ₃	48
7	1 : 1	(Me) ₂ CO	46
8	1 : 1	MeCN	50
9	2 : 1	MeCN	45
10	1 : 2	MeCN	66
11	1 : 3	MeCN	78
12	1 : 4	MeCN	72
13 ^e	1 : 3	MeCN	75

^a Unless otherwise mentioned, the reactions were carried out on a 0.2 mmol scale of **1a** with specified amounts of **2a** in 2 mL solvent at rt under N₂. ^b Time taken for reaction completion: 10–12 h. ^c Isolated yields. ^d Complex mixture. ^e Reaction in 1 mL solvent.

(entries 6–8). Further, the reaction with a 2 : 1 substrate ratio of **1a** and **2a** furnished **3a** in 45% yield (entry 9) but reversing the ratio to 1 : 2 led to the isolation of **3a** in 66% yield (entry 10). Further optimization of the reaction conditions established that a 1 : 3 ratio of **1a** and **2a** in acetonitrile were the best conditions affording **3a** in 78% isolated yield (entries 8–12). Apparently, the yields were largely independent of the reaction concentration since **3a** was isolated in 75% yield in a more concentrated mixture of **1a** and **2a** taken in a 1 : 3 ratio (entry 13). The product was isolated as exclusively the *E*-isomer, as identified by ¹H NMR and single crystal X-ray analysis of **3k**.¹³

After optimizing the reaction conditions, we set out to determine the scope of the thermal phosphorylation reaction. For this purpose, several α -diazo esters **1a**–**1q** and trialkylphosphites **2a**–**2d** were subjected to the optimized reaction conditions (Table 2). Initially, trimethyl phosphite **2a** was used as the reaction partner with various electron-rich and electron-deficient aryl diazo carboxylates to afford the corresponding phosphinamides **3a**–**3l** in high yields ranging from 33–85%. Notably, commonly encountered functional groups such as –OMe, –OCF₃, –Ph, –CF₃, –NO₂ and halogens (Cl, Br, F) on the aryl ring of the diazo substrate were tolerated well in the reaction. The reaction of trimethyl phosphite **2a** also proceeded smoothly with methyl 2-diazo-2-(naphthalen-1-yl)acetate and heteroaryl substrates such as ethyl 2-(benzo[*d*][1,3]dioxol-5-yl)-2-diazoacetate and ethyl 2-diazo-2-(thiophen-3-yl)acetate to

Table 2 Scope of thermal phosphorylation of diazo compounds^{a,b}

Other diazo substrates as electrophiles:			

^a Reaction conditions: **1** (0.2 mmol) and **2** (0.6 mmol) in CH₃CN (2 mL) at rt under N₂. ^b Isolated yields mentioned. ^c Isolated yield for a 5 mmol scale reaction.

furnish the corresponding products **3m**, **3n** and **3o** in high yields. Moreover, a phenyldiazoester with an allyl acetate group **1p** and a propargyl acetate group **1q** also reacted well with **2a** to yield **3p** and **3q** in 63 and 60% yields, respectively. Further, the *N*-phosphonylation of aryl diazocarboxylate **1a** could also be carried out efficiently with other trialkyl phosphites such as triethyl phosphite **2b**, triisopropyl phosphite **2c** and triallyl phosphite **2d** to afford **3r–3t** in 65–72% yields.

Eventually, the *N*-phosphonylation reaction could be successfully extended to other diazo substrates bearing at least one acceptor group including 2-diazo-1,2-diphenylethan-1-one (**1r**), (1-diazo-2,2,2-trifluoroethyl)benzene (**1s**), diethyl 2-diazo-malonate (**1t**) and 2-diazo-1,3-diphenylpropane-1,3-dione (**1u**) furnishing the corresponding products **3u–3x** in excellent yields.

The *E*-configuration was assigned to the products on the basis of analogy with **3k**, the structure of which was confirmed by single X-ray crystallography (Fig. 1).¹³

Next, we set out to establish the suitable conditions for the proposed photochemical phosphonylation reaction between **1a** and **2a** (Table 3).

To begin with, equivalent amounts of **1a** and **2a** were reacted in various solvents under blue light irradiation. The reactions in both dimethyl sulfoxide and dimethyl formamide were too complex to be conclusive (entries 1 and 2). In toluene, although the reaction furnished the anticipated C-P phosphonylation product **4a** in 15% yield, the phosphinamide **3a** was also isolated in 20% yield (entry 3). In other solvents including methanol, dichloromethane, chloroform, acetone and acetonitrile **4a** was obtained in moderate yields (26–35%) along with **3a** (10–20%) (entries 4–8). The reaction when carried out in dichloromethane with a 1:2 ratio of **1a** and **2a** provided **4a** in 38% yield, but **3a** was also isolated in 30% yield (entry 9). Fortunately, formation of **3a** was completely suppressed and **4a** could be isolated in 55% yield upon changing the ratio of **1a** and **2a** to 2:1 (entry 10). Further, a slight reduction in the yield of **4a** was noticed upon changing the substrate ratio to 1:3 (entry 11), therefore the conditions in

Table 3 Optimization of conditions for photochemical phosphonylation^{a,b}

Entry	1a : 2a	Solvent	Yield of 4a ^c (%)		Yield of 3a ^c (%)
			P(OMe) ₃ 2a	4a	
1 ^d	1 : 1	DMSO	0		0
2 ^d	1 : 1	DMF	0		0
3	1 : 1	PhMe	15		20
4	1 : 1	MeOH	28		13
5	1 : 1	CH ₂ Cl ₂	35		10
6	1 : 1	CHCl ₃	26		16
7	1 : 1	(Me) ₂ CO	30		15
8	1 : 1	MeCN	30		20
9	1 : 2	CH ₂ Cl ₂	38		30
10	2 : 1	CH ₂ Cl ₂	55		<5
11	3 : 1	CH ₂ Cl ₂	52		<5

^a Unless otherwise mentioned, the reactions were carried out on a 0.2 mmol scale of **1a** with specified amounts of **2a** in 2 mL solvent at rt irradiated with 455 nm blue LEDs under N₂. ^b Time taken for reaction completion: 5–6 h. ^c Isolated yields. ^d Complex mixture.

entry 10 were finally selected as the optimal reaction conditions.

Subsequently in order to investigate the scope of the photochemical phosphonylation reaction, several aryl α -diazoesters were employed in the blue light-mediated reaction with trimethylphosphite **2a** and triethylphosphite **2b** (Table 4). Evidently, phenyl diazo methylesters bearing substituents with various electronic properties performed well under the photoredox conditions to furnish the corresponding C-phosphonylated products **4a–4f** in good yields. The diazo substrates with an ethyl ester group and an allyl ester group also reacted efficiently with methylphosphite to afford **4g** and **4h** in 45% and 66% yield, respectively. The reaction of methyl

Table 4 Scope of photochemical phosphonylation of diazo compounds^{a,b}

1	P(OR) ₃ 2	CH ₂ Cl ₂ , rt, 6 h, N ₂		4
		Blue LEDs (455 nm, 3W)		
4a : R = H, 55% (53%) ^c	PO(OMe) ₂			PO(OMe) ₂
4b : R = 4-OCH ₃ , 60%				PO(OMe) ₂
4c : R = 4-Ph, 52%				PO(OMe) ₂
4d : R = 4-F, 48%				PO(OMe) ₂
4e : R = 4-NO ₂ , 45%				PO(OMe) ₂
	PO(OMe) ₂			PO(OEt) ₂
				PO(OEt) ₂
				PO(OEt) ₂
				PO(OEt) ₂

^a Reaction conditions: **1** (0.2 mmol) and **2** (0.1 mmol) in CH₂Cl₂ (2 mL) at rt irradiated with 455 nm blue LEDs under N₂. ^b Isolated yields mentioned. ^c Isolated yield for a 5 mmol scale reaction.





Fig. 1 ORTEP diagram drawn with 30% ellipsoid probability for non-H atoms of the crystal structure of compound **3k** determined at 294 K.



Scheme 2 Plausible mechanism.

2-diazo-2-phenylacetate **1a** with ethylphosphite **2b** under the optimized photoredox conditions furnished **4i** in 60% yield.

Mechanistically, the two phosphorylations follow divergent routes, as depicted in Scheme 2. The *N*-phosphonylation proceeds through nucleophilic addition of the phosphite to the terminal nitrogen of the diazo group. The resulting zwitterionic intermediate **A** upon dealkylation during aqueous workup forms intermediate **B** which tautomerizes to the *N*-phosphonylated product **3**. On the other hand, the aryl diazoester upon visible light-mediated photolysis generates free carbene species **X** which undergoes insertion with the phosphite to form ylide intermediate **Y**. Subsequent decomposition of the intermediate **Y** upon workup leads to the *C*-phosphonylated product **4**. The carbene intermediate in the reaction was confirmed by the formation of dimethyl 2,3-diphenylmaleate, the carbene dimerization product, in the visible light-mediated reaction between **1a** and **2a** (confirmed by ESMS of crude **4a**; Fig. S108 in the ESI†).

Conclusions

In conclusion, we developed mild protocols for accessing phosphonylated hydrazones and phosphonylated aryl carboxylates through thermal and photochemical phosphonylation, respectively, of α -diazo carboxylates with trialkyl phosphites. The former reaction is an example of a nucleophilic substitution reaction whereas the latter reaction follows a carbene insertion pathway. Both reactions proceed under mild reaction conditions, show wide substrate scope and afford the corresponding products in good yields.

Conflicts of interest

There are no conflicts to declare.

Acknowledgements

JP thanks DST, New Delhi for an INSPIRE Ph. D. fellowship. We thank the SAIF division of CSIR-CDRI for the analytical support. We also thank Dr T. S. Thakur of the Biochemistry

and Structural Biology Division, CSIR-CDRI for help in X-ray crystal structure determination studies of **3k**. CDRI Communication No: 10806.

References

- (a) U. Pradere, E. C. Garnier-Amblard, S. J. Coats, F. Amblard and R. F. Schinazi, *Chem. Rev.*, 2014, **114**, 9154–9218; (b) S. Demkowicz, J. Rachon, M. Daško and W. Kozak, *RSC Adv.*, 2016, **6**, 7101–7112; (c) J. B. Rodriguez and C. Gallo-Rodriguez, *ChemMedChem*, 2019, **14**, 190–216.
- (a) R. G. Hall, *Phosphorus, Sulfur Silicon Relat. Elem.*, 2008, **183**, 258–265; (b) C. Zhou, X. Luo, N. Chen, L. Zhang and J. Gao, *J. Agric. Food Chem.*, 2020, **68**, 3344–3353.
- (a) T. Baumgartner, *Acc. Chem. Res.*, 2014, **47**, 1613–1622; (b) Z. Wang and T. Baumgartner, *Chem. Rec.*, 2015, **15**, 199–217; (c) M. A. Shameem and A. Orthaber, *Chem. – Eur. J.*, 2016, **22**, 10718–10735; (d) D. Joly, P.-A. Bouit and M. Hissler, *J. Mater. Chem. C*, 2016, **4**, 3686–3698.
- (a) D. Heift, Z. Benkő and H. Grützmacher, *Chem. – Eur. J.*, 2014, **20**, 11326–11330; (b) C. Xie, A. J. Smaligo, X.-R. Song and O. Kwon, *ACS Cent. Sci.*, 2021, **7**, 536–558; (c) T. Gensch, G. P. Gomes, P. Friederich, E. Peters, T. Gaudin, R. Pollice, K. Jorner, A. K. Nigam, M. Lindner-D'Addario, M. S. Sigman and A. Aspuru-Guzik, *J. Am. Chem. Soc.*, 2022, **144**, 1205–1217.
- (a) Y. Gao, G. Tang and Y. Zhao, *Phosphorus, Sulfur Silicon Relat. Elem.*, 2017, **192**, 589–596; (b) K. Neog and P. Gogoi, *Org. Biomol. Chem.*, 2020, **18**, 9549–9561; (c) M. Arisawa, *Synthesis*, 2020, 2795–2806; (d) D. J. Jones, E. M. O'Leary and T. P. O'Sullivan, *Adv. Synth. Catal.*, 2020, **362**, 1825–1830; (e) Y. Mei, Z. Yan and L. L. Liu, *J. Am. Chem. Soc.*, 2022, **144**(4), 1517–1522; (f) Y. Liu, X. Chen and B. Yu, *Chem. – Eur. J.*, 2023, **29**, e202302142; (g) X. Liu, L. Zhou, R. Yang, X.-R. Song and Q. Xiao, *Adv. Synth. Catal.*, 2023, **365**, 2280–2298.
- For selected reviews on diazo compounds, see: (a) A. Ford, H. Miel, A. Ring, C. N. Slattery, A. R. Maguire and M. A. McKervey, *Chem. Rev.*, 2015, **115**, 9981–10080; (b) N. R. Candeias, R. Paterna and P. M. P. Gois, *Chem. Rev.*, 2016, **116**, 2937–2981; (c) Y. Xiang, C. Wang, Q. Ding and Y. Peng, *Adv. Synth. Catal.*, 2019, **361**, 919–944; (d) Z. Yang, M. L. Stivanin, I. D. Jurborg and R. M. Koenigs, *Chem. Soc. Rev.*, 2020, **49**, 6833–6847; (e) P. K. Mykhailiuk, *Chem. Rev.*, 2020, **120**, 12718–12755; (f) J. Durka, J. Turkowska and D. Gryko, *ACS Sustainable Chem. Eng.*, 2021, **9**, 8895–8918; (g) S. Dong, X. Liu and X. Feng, *Acc. Chem. Res.*, 2022, **55**, 415–428; (h) A. Dasgupta, E. Richards and R. L. Melen, *ACS Catal.*, 2022, **12**, 442–452; (i) C. Empel, C. Pei and R. M. Koenigs, *Chem. Commun.*, 2022, **58**, 2788–2798.
- (a) M. Wang, L. Kong, Q. Wu and X. Li, *Org. Lett.*, 2018, **20**(15), 4597–4600; (b) G. G. Faura, T. Nguyen and S. France, *J. Org. Chem.*, 2021, **86**, 10088–10104; (c) Z. Qi and S. Wang, *Org. Lett.*, 2021, **23**(21), 8549–8553;

- (d) C. Empel, C. Pei, F. He, S. Jana and R. M. Koenigs, *Chem. – Eur. J.*, 2022, **28**, e202104397; (e) G. D. Titov, G. I. Antonychev, M. S. Novikov, A. F. Khlebnikov, E. V. Rogacheva, L. A. Kraeva and N. V. Rostovskii, *Org. Lett.*, 2023, **25**(15), 2707–2712.
- 8 I. D. Jurberg and H. M. L. Davies, *Chem. Sci.*, 2018, **9**, 5112–5118.
- 9 (a) A. M. Polozov, N. A. Polezhaeva, A. H. Mustaphin, A. V. Khotinen and B. A. Arbuzov, *Synthesis*, 1990, 515–517; (b) W. Miao, Y. Gao, X. Li, Y. Gao, G. Tang and Y. Zhao, *Adv. Synth. Catal.*, 2012, **354**, 2659–2664; (c) L. Wu, X. Zhang, Q.-Q. Chen and A.-K. Zhou, *Org. Biomol. Chem.*, 2012, **10**, 7859–7862; (d) H. E. Bartrum, D. C. Blakemore, C. J. Moody and C. J. Hayes, *Tetrahedron*, 2013, **69**, 2276–2282; (e) L. Wang, Y. Wu, Y. Liu, H. Yang, X. Liu, J. Wang, X. Li and J. Jiang, *Org. Lett.*, 2017, **19**, 782–785.
- 10 (a) H. Jiang, H. Jin, A. Abdukader, A. Lin, Y. Cheng and C. Zhu, *Org. Biomol. Chem.*, 2013, **11**, 3612–3615; (b) Y. Feng, H. Ding, R. Liu, W. Wei, G. Nan and D. Yi, *Asian J. Org. Chem.*, 2024, **13**, e20240015.
- 11 L. B. Balázs, Y. Huang, J. B. Khalikuzzaman, Y. Li, S. A. Pullarkat and P.-K. Leung, *J. Org. Chem.*, 2020, **85**, 14763–14771.
- 12 V. K. Aggarwal, J. R. Fulton, C. G. Sheldon and J. de Vicente, *J. Am. Chem. Soc.*, 2003, **125**, 6034–6035.
- 13 The crystal structure of compound **3k** has been deposited at the Cambridge Crystallographic Data Centre and the reference no. CCDC 2342436 was allotted.†