

Halogen bonding, chalcogen bonding, pnictogen bonding, tetrel bonding: origins, current status and discussion

Lee Brammer 

Received 26th September 2017, Accepted 27th September 2017

DOI: 10.1039/c7fd00199a

The role of the closing lecture in a Faraday Discussion is to summarise the contributions made to the Discussion over the course of the meeting and in so doing capture the main themes that have arisen. This article is based upon my Closing Remarks Lecture at the 203rd Faraday Discussion meeting on Halogen Bonding in Supramolecular and Solid State Chemistry, held in Ottawa, Canada, on 10–12th July, 2017. The Discussion included papers on fundamentals and applications of halogen bonding in the solid state and solution phase. Analogous interactions involving main group elements outside group 17 were also examined. In the closing lecture and in this article these contributions have been grouped into the four themes: (a) fundamentals, (b) beyond the halogen bond, (c) characterisation, and (d) applications. The lecture and paper also include a short reflection on past work that has a bearing on the Discussion.

1. Introduction: history and foundations

Halogen bonding in its modern incarnation has been the subject of study for approximately 20 years, but is underpinned by many important earlier contributions, sometimes noted in historic introductions to the field and sometimes overlooked. Early papers from the 19th century have more recently been recognised as being observations of halogen bonding, the earliest among these being the association of iodine (I₂) with ammonia.^{1,2} Some of this work pre-dates the development of ideas that led to early definitions of the hydrogen bond,^{3–5} the intermolecular interaction to which halogen bonding is most often compared. Studies by Hassel in the 1950s/60s,^{6–8} which are the subject of his Nobel Prize lecture,⁹ provided a clear characterisation of halogen bonds, not yet named as such, by crystallographic means and notably established their linear geometry. The role of crystallography and solid-state chemistry continues to be important in the study of halogen bonds and their applications, as reflected in the title of this Discussion and many of the papers contributed. These include those by Riley and

Department of Chemistry, University of Sheffield, Brook Hill S3 7HF, UK. E-mail: lee.brammer@sheffield.ac.uk;
Web: https://www.sheffield.ac.uk/chemistry/staff/profiles/lee_brammer



I. Introduction

A. THESIS

A number of recent crystal structure determinations on compounds of the nonmetals have discovered intramolecular distances that are much longer than normal bonds and intermolecular distances that are much shorter than van der Waals distances. In this chapter, these interactions are examined and a qualitative explanation attempted. It will become clear that in most of them an approximately linear arrangement is found,



where $Y-A$ is a normal bond and $A \cdots X$ is a short intermolecular distance.* It is with these approximately linear interactions that we are particularly concerned, and it will be our contention that they are the result of directed forces and that their behavior is sufficiently regular and understandable for the name *secondary bond* to be appropriate.†

Chart 1 Definition of the term *secondary bond* by Alcock. Reproduced from ref. 11 with permission from Elsevier.

It is often difficult to single out individual papers that have made a difference in advancing a field of study, but here I will note two papers that appeared almost 20 years ago, both of which contributed significantly to the renaissance of interest in what is now termed halogen bonding by providing a point of comparison with hydrogen bonding, an interaction that was well-established in the minds of chemists and those working in the molecular sciences. The review in 1999 by Legon²¹ on structural characterisation of gas-phase complexes (Fig. 1) demonstrated the similarities between hydrogen bonding and halogen bonding, which had been systematically developed in experimental spectroscopic studies over a number of years, but was perhaps not well known to those with interests in self-assembly in the condensed phases. The complexes studied provide experimental rigour to underpin the fundamental understanding of these interactions and the extent to which there is parity between the two. The early work of Resnati and Metrangolo beginning around a similar time arose from a background in fluorine chemistry, which led to the development of highly fluorinated organic building blocks that could be deployed in halogen-bonded self-assembly in the solid state. Their report in 2000 showed that halogen bonding could be competitive with or even out-compete hydrogen bonding in the self-assembly of 1 : 1 co-crystals from the solution phase (Fig. 2),²² and brought to the attention of a wider community the possibility that halogen bonding might be used as an alternative to the more well-known hydrogen bonding in self-assembly. Very recently we have returned to this competition between hydrogen bonds and halogen bonds and shown that the choice of solvent affects the outcome (Fig. 3).²³ Generally speaking, less polar solvents favour association *via* hydrogen bonds over halogen bonds in forming






Fig. 1 Experimentally characterised gas-phase hydrogen-bonded and halogen-bonded complexes. Reproduced from ref. 21 with permission from Wiley.

the co-crystals, whereas the reverse is true in polar solvents. The cross-over point is related to the relative strengths of the hydrogen bonds and halogen bonds that are in competition.

2. Fundamentals

A number of the papers presented in the Discussion focused on fundamental aspects of halogen bonding and related interactions. The discussion of the papers centred on understanding the bonding, and being able to use this understanding in a predictive manner. In his opening lecture, and accompanying article (DOI: 10.1039/c7fd00058h), Tim Clark emphasised the fact the chemistry is based on models and proposed a *lex pasimoniae* (Occam's razor) for bonding theory as follows:






Fig. 2 Halogen-bonded self-assembly can out-compete hydrogen-bonded self-assembly in forming co-crystals from solution. Reproduced from ref. 22 with permission from Wiley.

Solvent			
	2a	4a	2a+4a
Toluene	HB	HB	HB
Chloroform	HB/XB	HB	HB
Dichloro- methane	HB/XB	HB/XB	HB
Acetone	XB	XB	HB
MeCN	XB	XB	HB
Nitromethane	XB	XB	HB
<i>iso</i> -Propanol	XB	XB	HB/XB

Fig. 3 Solvent effect on self-assembly of 1 : 1 co-crystals of 4,4'-dipyridylethane with the ditopic hydrogen bond and halogen bond donors shown above each column (HB = hydrogen-bonded co-crystal formed; XB = halogen-bonded co-crystal formed).



- An analysis is adequate when it explains all the known facts.
- Try to stick with physically unique and defined phenomena wherever possible.
- Do not define new interactions where old ones will do.

The paper emphasised an electrostatic description of bonding, which in principle can account for all bonding interactions, but recognised the value of orbital models which have been embedded in the description of bonding in chemistry over many years. The electrostatic model includes:

- The Coulomb interaction, which is the foundation of the description of halogen bonding in terms of the interaction of the σ -hole of the halogen-bond donor and the electron-rich region of the halogen-bond acceptor.
- Polarisation, because it is important for describing the interaction between unperturbed molecules where this is not otherwise obvious.
- Dispersion, because otherwise the attraction of nonpolar species is not explained.

Co-operativity in halogen bonding was demonstrated in calculations by Del Bene and co-workers on a series of binary and ternary complexes (DOI: 10.1039/c7fd00048k). As shown in Fig. 4, the halogen bond between ClF and PH_2F is enhanced slightly by the cooperative polarising effect from the formation of a second halogen bond involving HCl. This polarisation of the central ClF molecule is greater when a stronger halogen-bond donor (a second ClF molecule) is used to form the ternary complex, so much so that it leads to Cl^+ transfer. Co-operativity has long been established in hydrogen bonding, as illustrated in the study by Hermansson of co-operativity in hydrogen-bonded water chains,²⁴ in which the calculated difference electron density of the water molecules clearly shows the most polarised molecule to be the central one of the chain (Fig. 5). Polarisation in halogen bonding was further discussed in the computational study by Riley and Tran on charged halogen bonds between cationic halogen-bond donors and anionic acceptors (DOI: 10.1039/c7fd000106a), indicating a greater polarisation effect in interactions between charged species than neutral ones. Such interactions have also been studied extensively in experimental work involving halides, halometallate ions and other anions in the halogen-bond acceptor role with either cationic or neutral halogen bond donors.^{18,25–34} Kennepohl and co-workers (DOI: 10.1039/c7fd0076f) used X-ray Absorption




Fig. 4 Computational models of complexes $\text{H}_2\text{FP}\cdots\text{ClF}$, $\text{H}_2\text{FP}\cdots\text{ClF}\cdots\text{ClH}$ and $\text{H}_2\text{FP}\cdots\text{ClF}\cdots\text{ClF}$, illustrating co-operativity in halogen bonding interactions. Adapted from DOI: 10.1039/c7fd00048k with permission from The Royal Society of Chemistry.






Fig. 5 Calculated difference electron density of hydrogen-bonded water chains of length n molecules, showing co-operativity in hydrogen bonding, which leads to the central water molecule being the most polarised. Solid contours represent increased electron density and dashed contours decreased electron density relative to the pro-molecule. All electron densities are shown in the plane of the three atoms in each molecule for ease of comparison (note: only the central molecule is shown for $n = \infty$). Adapted from ref. 24 with permission from AIP Publishing.

Spectroscopy (XAS) to demonstrate and quantify the charge-transfer from chloride ions to I_2 halogen-bond donors, by observation of the $Cl_{1s} \rightarrow Cl_{3p}$ transition, which cannot arise in the absence of Cl^- to $I_2(\sigma^*)$ charge-transfer (Fig. 6). Fourmigué and co-workers (DOI: 10.1039/c7fd00067g) further developed the discussion of covalency in halogen bonding in experimental and computational charge density studies of $N-X \cdots N$ halogen bonds ($X = I, Br$), which suggest greater covalency for $X = Br$ than for $X = I$ (Fig. 7), a reverse of what might have been




Fig. 6 Cl_{1s} to Cl_{3p} transition, which can be probed by XAS, arises only if Cl_{3p} to $I_2(\sigma^*)$ charge-transfer is present (A represents a halogen-bond acceptor, here I_2). The intensity of the transition is proportional to the Cl_{3p} character in the ψ^* orbital. Reproduced from DOI: 10.1039/c7fd00076f with permission from The Royal Society of Chemistry.





Fig. 7 Halogen bonding interactions in (a) NBSac·Pic and (b) NISac·Pic (distances in Å; NBSac = *N*-bromosaccharin, NISac = *N*-iodosaccharin, Pic = 4-picoline). Arrows and values correspond to the orientation and the magnitude (in Debye) of the dipolar moments of the adducts calculated at frozen experimental geometries. Geometries indicate the greater asymmetry is in the N–I⋯N halogen bond. Reproduced from DOI: 10.1039/c7fd00067g with permission from The Royal Society of Chemistry.

anticipated. The nature of N–X⋯N halogen bonds have also been extensively studied in solution-phase using NMR spectroscopy by Erdelyi and co-workers.^{35–38}

Alavi and co-workers (DOI: 10.1039/c7fd00064b) described their computational studies of chlorine and bromine hydrates, clathrates based on polyhedral hydrogen-bonded water cages with the dihalogens occupying the cage voids. These materials have been known for some 200 years, but are still revealing structural and bonding details.^{39–42} The study demonstrates the halogen-bonding interaction between the dihalogen and the water molecules, and highlights the bifurcation that occurs involving one of the water molecule lone pairs, which accepts both a hydrogen bond and a halogen bond. Bifurcation of interactions also featured in a number of the calculations presented by Politzer (DOI: 10.1039/c7fd00062f), which focused on σ -hole interactions across the p-block. Specifically, examples were presented in which the site of the most positive electrostatic potential was calculated to be located between atomic sites at which σ -holes might have been expected, *e.g.* S and C in octafluorothiophane and two hydrogens in dimethylurea (see Fig. 8). An alternative to the more conventional description of a bifurcated interaction was proposed, by connecting the sites of most positive and most negative potential on neighbouring molecules *via* a single interaction. Bifurcation remains an appropriate description if presented in terms of an atom–atom interaction framework, and its interpretation is well understood and in common use. Such a bifurcated description has been used previously where sites of maxima or minima in electrostatic potential have been determined to lie between atoms.⁴³

The emphasis so often placed on drawing (dotted/dashed) lines between atoms to represent intermolecular interactions is reflected upon in the paper by Edwards, Spackman and co-workers (DOI: 10.1039/c7fd00072c). Their analysis of intermolecular interactions in crystals is based on the calculated molecular charge densities and enables molecule–molecule interactions to be partitioned into electrostatic and dispersion contributions, which can be visualised in






Fig. 8 Maxima in positive electrostatic potential (shown as black dots) that lie between atoms: (a) between ring atoms S and C, providing the site for interaction with a Lewis base, and (b) between hydrogen atoms providing a site for interaction with the most negative region of the carbonyl oxygen leading to the hydrogen-bonded urea tape. (c) Two alternative descriptions of hydrogen-bonding between urea molecules – the conventional bifurcated interaction in an atom–atom interaction framework (left) and the interaction between sites of electrostatic potential minima and maxima (right). Adapted from DOI: 10.1039/c7fd00062f with permission from The Royal Society of Chemistry.

a quantitative manner by representing the strength of an interaction between two molecules by a line, the thickness of which is proportional to the interaction strength (Fig. 9). Spackman refers to this model of intermolecular interactions as an energy framework.^{44–46} Their analyses, and similar analyses by Gavezzotti, using his PIXEL calculation approach,^{47–49} show that the strongest interactions between molecules are not always those that conform to named interactions such as hydrogen bonds, halogen bonds *etc.*, *i.e.* the interactions which are conveniently depicted as dotted lines between the atoms of the interacting molecules. The paper (DOI: 10.1039/c7fd00072c) also questions the need for so many specific names for many different types of atom–atom intermolecular interactions when intermolecular interactions may be better interpreted in terms of complementary electrostatic interactions between molecular surfaces combined with dispersion forces.





Fig. 9 (a) Crystal structures of C_6Cl_6 and C_6Br_6 , shown with dashed lines representing intermolecular halogen-halogen interactions. (b) Energy frameworks for crystal structures of C_6Cl_6 and C_6Br_6 , showing that the largest contribution to the lattice energy comes from dispersion interactions between the stacked molecules and that halogen-halogen interactions (type I and type II) provide a much smaller contribution to the overall energy, which might be interpreted as providing a steering role for the orientation of adjacent molecular stacks. Adapted from DOI: 10.1039/c7fd00072c with permission from The Royal Society of Chemistry.

3. Beyond the halogen bond

Although the study of halogen bonding has expanded dramatically over the past 15–20 years,^{50–54} analogous interactions involving other main group elements in a Lewis acidic role have seen far less attention over this same period. More recently, with the advent of the σ -hole terminology as a framework for describing such interactions,¹³ there has been a growing interest in the study of such interactions, which like halogen bonding have the potential to lead to a variety of applications in supramolecular chemistry, crystal engineering, materials chemistry and in medical/biological areas. The terms triel bond (group 13), tetrel bond (group 14), pnictogen/pnictogen bond (group 15), chalcogen bond (group 16) and aerogen bond (group 18) have begun to be used by analogy to the term halogen bond for the now well-established interactions involving group 17 elements. As noted in the Introduction, such interactions across the p-block were well established by crystallographic studies in the 1960s and 70s.^{10,11} The interactions were often described within a bonding framework of donor-acceptor interactions, in which a significant covalent or charge-transfer component to the bonding was inferred, rather than *via* an electrostatic (*e.g.* σ -hole) description. Early studies often focused on systems in which the main group element involved in the Lewis acidic role is bound to fluorine (often perfluorinated), leading to strong intermolecular interactions (Fig. 10). There is now scope to develop this palette of






Fig. 10 Examples of secondary bonding interactions involving p-block elements in Lewis acid roles.^{55,56} Equivalent isoelectronic species to secondary bond donors are noted in parentheses. Adapted from ref. 11 with permission from Elsevier.

interactions in a similar manner to the modern development of halogen bonds through use of different electron-withdrawing groups (perfluoroaryl, perfluoroalkyl, alkynyl, cationic, *etc.*) that can be engaged through established synthetic organic and main group chemistry. Such development is necessary to enable opportunities for new applications of supramolecular chemistry across a variety of fields. Examples are known. Indeed in work with Braddock-Wilking and co-workers, we reported, in 1995, examples of intramolecular tetrel bonding, although not then named as such, involving $\text{SiF}_2(2,4,6\text{-C}_6\text{H}_2(\text{CF}_3)_3)_2$ and $\text{SiHF}(2,4,6\text{-C}_6\text{H}_2(\text{CF}_3)_3)_2$ (Fig. 11).⁵⁷ The interactions involving fluorine substituents of the *ortho*- CF_3 groups interact with the Si centres *trans* to the primary bonding (to F, H and C) and are evident from the crystal structures and from solution-phase NMR spectroscopy in which J_{SiF} coupling for the secondary (tetrel) bonds has a magnitude of approximately 3 Hz in comparison with the $^1J_{\text{SiF}}$ coupling of 300 Hz observed for primary Si-F bonds. The tetrel bond is shorter for interactions *trans* to F than *trans* to C, consistent with electrostatic or orbital (charge-transfer) arguments. We noted in our report that secondary $\text{F}\cdots\text{E}$ interactions have also been reported for Ar_2Pb ,⁵⁸ Ar_2Sn ,^{59,60} Ar_3Ga ⁶¹ and for related filled-shell transition metal complexes Ar_2Zn , Ar_2Hg and $\text{Ar}_2\text{Cd}(\text{NCMe})$,⁶² although not all adopt the linear arrangement observed in our examples. Thus, care is needed in interpreting the nature of these interactions.

In an extensive computational study, Scheiner (DOI: 10.1039/c7fd00043j) reported binding energies and geometries for halide ion binding in a series of isostructural ditopic receptors in which hydrogen bonding, tetrel bonding, pnictogen bonding, chalcogen bonding and halogen bonding were compared (Fig. 12). As expected, heavier p-block elements form stronger interactions than their lighter congeners. Of particular interest is the fact that tetrel bonding from RSnH_3 groups to the halides is stronger than that for corresponding halogen





Si–F	1.54–1.56 Å	
Si···F	2.72–2.77 Å	<i>trans</i> to F
Si···F	2.97–3.07 Å	<i>trans</i> to C(Ar)

solution-phase NMR	
J _{SiF}	300 Hz (primary)
J _{SiF}	3 Hz (secondary)




Fig. 11 Crystal structure of SiFHA₂, Ar = C₆H₂(CF₃)₃ (top), with secondary (tetrel) bonding shown as dashed bonds and primary bonding (solid lines) and secondary bonding (dashed lines) at Si in SiFHA₂ and SiF₂Ar₂ (bottom). Secondary (tetrel) Si···F bonds are longer than primary (covalent) Si–F bonds. Secondary bonds *trans* to F are shorter than those *trans* to C. Adapted from ref. 57 with permission from Elsevier.

bond (RI) and at least as strong as the corresponding hydrogen bond (RH). Vargas-Baca and co-workers (DOI: 10.1039/c7fd00075h) presented a series of cyclic tellurazole *N*-oxides, which can form supramolecular assemblies *via* chalcogen bonds involving Te as the chalcogen bond donor. The *N*-oxide has an electron-withdrawing role that enhances the strength of the chalcogen bond. This role can be further amplified by coordination of the oxygen to Lewis acids such as BF₃ (Fig. 13). Southern, Bryce and co-workers (DOI: 10.1039/c7fd00087a) described the use of solid-state ²⁰⁷Pb NMR spectroscopy to explore the formation of tetrel bonds in hemihedrally coordinated Pb(II) complexes. Although not a commonly used NMR nucleus, ²⁰⁷Pb is 22.1% abundant, has *I* = 1/2 and a very large chemical shift range, offering the potential of great sensitivity to the nature of intermolecular interactions involving Pb centres. It is further noted that group 14 elements provide considerable opportunities for the investigation of tetrel bonding by NMR spectroscopy (Table 1).





Fig. 12 Optimized geometries of Cl⁻ anion with ditopic receptors (distances in Å). Binding energies for halide ion binding by receptors shown. Adapted from DOI: 10.1039/c7fd00043j with permission from The Royal Society of Chemistry.




Fig. 13 N–Te⋯P chalcogen bond. Strong chalcogen bonding capability of Te is activated by coordination of *N*-oxide, and further enhanced by Lewis acid coordination of the BF₃ group to the oxygen (blue circle). Phosphine (PPh₃, red ellipse) serves as the (Lewis basic) chalcogen bond acceptor. Adapted from DOI: 10.1039/c7fd00075h with permission from The Royal Society of Chemistry.

4. Characterisation

A number of papers in this Discussion showcased characterisation methods that provide insight into secondary bonding interactions, including halogen bonding. In many examples crystallographic studies were used to establish the geometries of these interactions, including the contributions by Riley and Tran (DOI: 10.1039/c7fd000106a), Alavi *et al.* (DOI: 10.1039/c7fd00064b), Kennepohl *et al.* (DOI: 10.1039/c7fd0076f), Edwards *et al.* (DOI: 10.1039/c7fd00072c), Politzer *et al.*



Table 1 NMR parameters for group 14 elements

NMR-active isotope	I	Natural abundance (%)	Magnetogyric ratio ($10^7 \text{ rad T}^{-1} \text{ s}^{-1}$)	Relative receptivity	Chemical shift range (ppm)
^{13}C	1/2	1.07	6.728284	1.00	250
^{29}Si	1/2	4.6832	-5.3190	2.16	900
^{73}Ge	9/2	7.73	-0.9360303	0.642	1200
^{115}Sn	1/2	0.34	-8.8013	0.711	2600
^{117}Sn	1/2	7.68	-9.58879	20.8	2600
^{119}Sn	1/2	8.59	-10.0317	26.6	2600
^{207}Pb	1/2	22.1	5.58046	11.8	17 000

(DOI: 10.1039/c7fd00062f), Southern *et al.* (DOI: 10.1039/c7fd00087a), Vargas-Baca *et al.* (DOI: 10.1039/c7fd00075h), Guru Row *et al.* (DOI: 10.1039/c7fd00084g), Mosquera *et al.* (DOI: 10.1039/c7fd00079k), Aakerøy *et al.* (DOI: 10.1039/c7fd00080d), Fourmigué *et al.* (DOI: 10.1039/c7fd00067g), Lloyd *et al.* (DOI: 10.1039/c7fd000108h), and Friščić *et al.* (DOI: 10.1039/c7fd00114b). Guru Row and co-workers (DOI: 10.1039/c7fd00084g) studied the structures of 2-bromo-4-chloro-benzoic acid and its isomer 4-bromo-2-chloro-benzoic acid. Both form structures that involve hydrogen-bonded carboxylic acid dimers but differ in the arrangement interactions involving the halogens. Solid solutions of the two compounds could be formed at a variety of different compositions. The resultant crystal structures were found to adopt that of the 4-bromo isomer until high concentrations of the 2-bromo isomer led its structure directing the final arrangement. The intermolecular interactions in crystals of each of the pure compounds were quantified by a topological analysis of the experimental electron density distributions.

Erdelyi and co-workers (DOI: 10.1039/c7fd00107j) investigated natural abundance ^{15}N NMR spectroscopy as a probe for weak halogen bond formation in the solution phase. Nonfluorinated halogen-bond donors involving I, Br and Cl were investigated as were substituent effects and solvent effects (Fig. 14). The study concluded that ^{15}N NMR spectroscopy is suitable for the characterization of strong secondary interactions in solution, but not sufficiently accurate to detect weak halogen-bonded complexes due to the low mole fraction of the complex present. DFT calculations show that the ^{15}N chemical shift does reflect the diamagnetic deshielding associated with the formation of a weak halogen bond, but that experimental detection was unreliable (Fig. 14a). Hydrogen-bonding interactions with the same acceptors show a 10-fold larger change in chemical shift ($\Delta\delta^{15}\text{N}$) and can readily be detected and correlated with the electronic properties of the hydrogen bond donor (Fig. 14b). The effect of solvent upon halogen bond formation is also apparent (Fig. 14c) wherein less polar solvents result in larger deshielding of the ^{15}N nuclei ($\Delta\delta^{15}\text{N} < 0$), the $\Delta\delta^{15}\text{N}$ having the largest magnitude for $\text{I} > \text{Br} > \text{Cl}$. In less polar solvents, the resulting positive $\Delta\delta^{15}\text{N}$ is suggested to arise from the altered polarity of these mixtures as compared to the reference, *i.e.* this effect opposes the very small negative $\Delta\delta^{15}\text{N}$ that results from the formation of the weak halogen bond.

Electrochemical methods were used in conjunction with spectroscopic measurements by Rosokha and by Branca, Schöllhorn and co-workers to






Fig. 14 (a) (Poor) correlation of $\Delta\delta^{15}\text{N}$ with (weak) halogen-bond donor strength expressed in terms of Hammett σ_{para} constant for substituent R. (b) Good correlation of $\Delta\delta^{15}\text{N}$ with hydrogen-bond donor strength expressed in terms of Hammett σ_{para} constant for substituent R. All measurements for (a) and (b) are in toluene- d_8 . Standard error on $\delta^{15}\text{N} < 0.02$ ppm. (c) Average chemical shift changes for (weak) halogen bonds C-I...N (blue), C-Br...N (red), C-Cl...N (green), as a function of solvent. Adapted from DOI: 10.1039/c7fd00107j with permission from The Royal Society of Chemistry.

investigate halogen bond formation in solution. Rosokha (DOI: 10.1039/c7fd00074j) used a combination of electrochemical and UV/Vis spectroscopic measurements to examine charge-transfer in halogen bonds formed by R-CBr₃



halogen-bond donors with tetramethyl-*p*-phenylenediamine (TMPD) or iodide as the acceptor. The work established from kinetics determinations that electron transfer from diamine to R-CBr₃ occurs too rapidly to be *via* an outer-sphere mechanism. The resulting fast redox process implicates transient halogen-bonded complexes in which barriers for electron transfer are lowered by the strong electronic coupling the two molecules. The study also suggests that halogen bonding should be taken into account in the mechanistic analysis of the reactions of halogenated species. In their study of cationic viologen-substituted halide receptors, Branca, Schöllhorn and co-workers (DOI: 10.1039/c7fd00082k) used a combination of electrochemical and NMR titrations to establish binding strengths and binding sites.

5. Applications

In the final section we will focus on applications of halogen bonding, with papers presented on anion binding, supramolecular synthesis, micelle formation in polymers, gel formation, dichroism and light-controlled ion transport. Beer and co-workers have developed rotaxane systems with a goal of selective binding of anions in water.^{63–65} Their use of halogen bonding for anion binding has enabled marked progress towards this aim. In their Discussion paper (DOI: 10.1039/c7fd00077d), they report the tuning of the binding site to enable excellent selectivity of Br[−] over I[−] and Cl[−] in a polar d₆-acetone/D₂O (9 : 1) solution. The anion binding site comprises three C–I halogen bond donors from triazole units (Fig. 15), for which the halogen-bond donor capability is enhanced by cationic charge or ligation to Re. Chelation of the Re centre also influences the spatial arrangement of the two C–I groups on the bistriazole unit improving pre-organisation of the rotaxane for binding.

As part of a continuing programme of study of the competition between intermolecular interactions in supramolecular synthesis of crystals,^{66–69} Aakeröy





Fig. 15 Cationic rotaxane that exhibits selective binding of Br[−] ($K = 436(25) \text{ M}^{-1}$) over Cl[−] and I[−], for which binding is too weak to measure by NMR titration. All measurements are in d₆-acetone/D₂O at 298 K. Adapted from DOI: 10.1039/c7fd00077d with permission from The Royal Society of Chemistry.



and co-workers have designed a series of 1-(pyridylmethyl)-2,2'-biimidazoles, which contain three different sites capable of being hydrogen bond or halogen bond acceptors (Scheme 1). Co-crystallisation with a series of ditopic hydrogen bond donors and halogen bond donors allowed selectivity patterns for intermolecular interaction formation to be examined (DOI: 10.1039/c7fd00080d). Halogen-bond donors consistently led to self-association of the biimidazoles through $R_2^2(10)$ N-H...N hydrogen-bonded motifs, with halogen-bond donors selecting the pyridyl acceptor site over the remaining imidazole site. Carboxylic acid hydrogen-bond donors showed less consistency. In some cases biimidazole self-association occurred with carboxyl hydrogen bonding to pyridyl groups, but in others the carboxyl group disrupted the self-association, leading to a variety of



Scheme 1 Competition between acceptor sites for hydrogen-bond or halogen-bond donors in co-crystal formation. Adapted from DOI: 10.1039/c7fd00080d with permission from The Royal Society of Chemistry.




Fig. 16 Micelles formed *via* halogen bonding between triblock copolymer and halogen-bond donor polymer. Adapted from DOI: 10.1039/c7fd00111h with permission from The Royal Society of Chemistry.



outcomes. Mosquera, Gómez-Sal and co-workers (DOI: 10.1039/c7fd00079k) have carried out a mechanistic investigation that examines in detail the role of halogen bond formation on halide ligand substitution in the octahedral complex $[\text{RuCl}_2(\text{CNR})_4]$. Their study implicates halogen bond formation, initially between dihalogens (I_2 , Br_2) and the chloride ligands, in the mechanism leading to substitution of chloride by iodide or bromide. The rate of substitution depends on the electron-donating ability of the isocyanide (CNR) ligands. A number of intermediate stages of the reaction have been characterised crystallographically and studied by Raman spectroscopy. An unusual observation from these




Fig. 17 Dichroic crystals of $[\text{K}(15\text{-crown-5})][\text{Au}(\text{CN})_2] \cdot (\text{DIFAB}) \cdot (\text{solvent})$ (DIFAB = 4,4'-diiodooctafluoroazobenzene) containing aligned linear halogen-bonded assemblies of DIFAB and $[\text{Au}(\text{CN})_2]^-$. (a) and (b) Crystals presented at different rotation angles exhibit different colours in polarised light. (c) Light intensity as a function of crystal rotation. (d) Crystal structure $[\text{K}(15\text{-crown-5})][\text{Au}(\text{CN})_2] \cdot (\text{DIFAB}) \cdot (\text{solvent})$. Reproduced from DOI: 10.1039/c7fd00114b with permission from The Royal Society of Chemistry.



structures is that formation of $\text{Ru-X}\cdots\text{I}_2$ leads to Ru–X bond lengthening, whereas formation of $\text{Ru-Br}\cdots\text{Br}_2$ leads to Ru–Br bond shortening.

Vanderkooy and Taylor (DOI: 10.1039/c7fd00111h) have appended $\text{C}_6\text{F}_4\text{I}$ halogen-bond donor groups to a polymer backbone enabling the use of halogen bonding in forming multicompartmental micelles from complementary macromolecules. The micelles were able to be imaged by TEM (Fig. 16). Lloyd, Khimyak and co-workers (DOI: 10.1039/c7fd000108h) have examined the effect of halogen bonding, and more generally the influence of intermolecular interactions involving halogen substituents, on a series of phenylalanine (Phe) derivatives with halogen substituents on the aromatic ring. Friščić, Barrett and co-workers (DOI: 10.1039/c7fd00114b) were able to synthesise and demonstrate dichroism in crystals comprising halogen-bonded chains of *trans*-4,4'-diiodooctafluoroazobenzene as chromophore and ditopic halogen-bond donor and dicyanoaurate(i) as ditopic halogen-bond acceptor (Fig. 17). The anionic assembly is charge-balanced by 15-crown-5-encapsulated K^+ ions. The halogen bonding is used to provide alignment of the chromophores. Finally, remaining with the theme of optical properties, Cavallo, Dichiarante and co-workers (DOI: 10.1039/c7fd00120g) have prepared halogen-bonded lamellar liquid crystals (LC), which include an iodide-containing charged region as well as a lipophilic region. The halogen-bond donor unit contains a photo-switchable diazo group. A reversible *cis-trans* photo-isomerization of this group leads to a LC-to-isotropic phase transition, which the authors envision is a first step towards light-induced control over ion transport and conductance in these supramolecular complexes.

6. Conclusions and future outlook

Among the main themes that emerged in the Discussion (DOI: 10.1039/c7fd90061a, DOI: 10.1039/c7fd90062g, DOI: 10.1039/c7fd90063e and DOI: 10.1039/c7fd90064c) were those that involved the bonding model for describing halogen bonds and related interactions. It is clear that the electrostatic model is central to describing halogen bonds and their p-block analogues, but that care is needed in interpreting solid-state arrangements, as while it may be convenient to assign (on the basis of geometry) halogen bonds or hydrogen bonds as structure directing, these may not be the main cohesive intermolecular forces. Experimental measurements were presented to demonstrate charge transfer in halogen bonds indicating the importance that such a description can have in understanding the interactions.

Although computational studies highlighted some differences between halogen bonds, chalcogen bonds, pnictogen bonds and tetrel bonds, more experimental studies are needed to quantify and exploit such differences. A wide variety of characterisation methods were employed in the studies presented, including, X-ray crystallography, multinuclear NMR spectroscopy (solution-phase and solid-state), UV/Vis spectroscopy, IR/Raman spectroscopy, cyclic voltammetry, X-ray absorption spectroscopy (XAS) and electron microscopy. Experimental studies were often complemented by computational studies, which enabled the calculation of structures, spectroscopic parameters, bonding models and energetics. It is also noticeable that studies of halogen bonding spanned a very wide range of molecular sizes from very small, *e.g.* diatomics (I_2 , ClF , *etc.*) to very large, *e.g.* rotaxanes and polymers.



A number of applications of halogen bonding were described, including supramolecular and covalent synthesis, anion binding, micelle formation in polymers, gel formation, dichroism and light-controlled ion transport. These represent examples from among the many areas in which halogen bonding is now being applied, such as medicine and biology, including pharmaceuticals, many aspects of molecule-based materials chemistry, encompassing crystal engineering, and in synthesis and catalysis. It should be anticipated that the recent development of interest in related chalcogen, pnictogen and tetrel bonds will also lead to opportunities for a wide array of applications that are underpinned by an improved understanding and an expanded “toolbox” for supramolecular chemistry.

I will finally return to the question of nomenclature, which was raised in the Introduction. The definition of halogen bonding⁷⁰ (and re-definition of hydrogen bonding⁷¹) has recently been approved through IUPAC projects and the definitions of analogous interactions, commonly referred to as tetrel bonds (group 14), pnictogen bonds (group 15), chalcogen bonds (group 16) and aerogen bonds (group 18) are soon to be determined *via* a similar IUPAC project. Although the names of these interactions convey valuable precision in description and permit distinction to be made between these related interactions, it seems clear that a collective name for such interactions is also needed. Although not formally approved by IUPAC, that collective name already exists. The term *secondary bond*, coined by Alcock,¹¹ fulfils the role and allows for the necessary derivative terms secondary bonding, secondary bond donor, secondary bond acceptor, *etc.* to convey meaning with clarity for this collection of interactions in the same way that more specific terms, *e.g.* halogen bond, chalcogen bonding, pnictogen bond donor and tetrel bond acceptor, can for the more limited subset of cases to which they apply. Some might argue that hydrogen bonds should remain separate, but this seems unnecessary as they can be described in the same manner (either by σ -hole electrostatic model⁷² or charge-transfer orbital model^{73,74}) as the interactions that involve p-block elements in the Lewis acidic role, although forming less directional interactions. Indeed it can be anticipated that analogous classes of interactions involving elements of group 13 (triel) or s-block groups 1 and 2 can be similarly encompassed in the definition of secondary bonding. Thus, it seems appropriate to adopt the term *secondary bonding* as an umbrella term for all classes of intermolecular interaction involving s- and p-block elements in a Lewis acidic role.

Acknowledgements

Thanks are due to the Royal Society of Chemistry, the organising committee and particularly its Chair, David Bryce, who also served as an excellent host for the Discussion meeting. I would like to thank the PhD students and postdocs, Fiorenzo Zordan, Guillermo Mínguez Espallargas, Stefano Libri, Iñigo Vitórica-Yrezábal, Paul Smart, Johnathan Ormond-Prout, Adrián Pérez Redondo, James Wright, Craig Robertson and Tom Roseveare, as well as a number of undergraduate research students, who have worked on halogen-bond related projects in my group since 2001. I also value a number of enriching collaborations that have broadened our research and hastened our progress, particularly those with Robin Perutz (York), Chris Hunter (Cambridge), Paul Sherwood (Daresbury) and



Anthony Meijer (Sheffield). I am grateful to the EPSRC, STFC, CCDC, the White Rose Consortium and University of Sheffield for support of work involving the aforementioned students.

References

- 1 M. Colin, *Ann. Chim.*, 1814, **91**, 252–272.
- 2 F. Guthrie, *J. Chem. Soc.*, 1863, **16**, 239–244.
- 3 For example, see: (a) G. C. Pimental and A. L. McClellan, *The Hydrogen Bond*, W. H. Freeman, San Francisco, 1960; (b) *The Hydrogen Bond: Recent Developments in Theory and Experiment*, ed. P. Schuster, G. Zundeland C. Sandorfy, North-Holland, Amsterdam, 1976.
- 4 G. A. Jeffrey, *An Introduction to Hydrogen Bonding*, Oxford University Press, Oxford, 1997.
- 5 G. R. Desiraju and T. Steiner, *The Weak Hydrogen Bond*, Oxford University Press, Oxford, 1999.
- 6 O. Hassel and J. Hvoslef, *Acta Chem. Scand.*, 1954, **8**, 873.
- 7 O. Hassel and K. O. Stromme, *Acta Chem. Scand.*, 1958, **12**, 1146.
- 8 O. Hassel and K. O. Stromme, *Acta Chem. Scand.*, 1959, **13**, 1781–1786.
- 9 O. Hassel, *Science*, 1970, **170**, 497–502.
- 10 H. A. Bent, *Chem. Rev.*, 1968, **68**, 587–648.
- 11 N. W. Alcock, *Adv. Inorg. Chem. Radiochem.*, 1972, **15**, 1–58.
- 12 T. Clark, M. Hennemann, J. S. Murray and P. Politzer, *J. Mol. Model.*, 2007, **13**, 291–296.
- 13 P. Politzer, J. S. Murray and T. Clark, *Phys. Chem. Chem. Phys.*, 2013, **15**, 11178–11189.
- 14 N. Ramasubbu, R. Parthasarathy and P. Murray-Rust, *J. Am. Chem. Soc.*, 1986, **108**, 4308–4314.
- 15 G. R. Desiraju and R. Parthasarathy, *J. Am. Chem. Soc.*, 1989, **111**, 8725–8726.
- 16 V. R. Pedireddi, D. S. Reddy, B. S. Goud, D. C. Craig, A. D. Raeb and G. R. Desiraju, *J. Chem. Soc., Perkin Trans. 2*, 1994, 2353–2360.
- 17 L. Brammer, E. A. Bruton and P. Sherwood, *Cryst. Growth Des.*, 2001, **1**, 277–290.
- 18 L. Brammer, G. Mínguez Espallargas and H. Adams, *CrystEngComm*, 2003, **5**, 343–345.
- 19 L. Brammer, F. Zordan, G. Mínguez Espallargas, S. L. Purver, L. Arroyo Marín and H. Adams, *Trans. Am. Crystallogr. Assoc.*, 2004, **39**, 114–122.
- 20 F. Zordan, L. Brammer and P. Sherwood, *J. Am. Chem. Soc.*, 2005, **127**, 5979–5989.
- 21 A. C. Legon, *Angew. Chem., Int. Ed.*, 1999, **38**, 2686–2714.
- 22 E. Corradi, S. V. Meille, M. T. Messina, P. Metrangolo and G. Resnati, *Angew. Chem., Int. Ed.*, 2000, **39**, 1782–1786.
- 23 C. C. Robertson, J. S. Wright, E. J. Carrington, R. N. Perutz, C. A. Hunter and L. Brammer, *Chem. Sci.*, 2017, **8**, 5392–5398.
- 24 K. Hermansson and M. Alfredsson, *J. Chem. Phys.*, 1999, **111**, 1993–2000.
- 25 R. D. Willett, F. Awwadi, R. Butcher, S. Haddad and B. Twamley, *Cryst. Growth Des.*, 2003, **3**, 301–311.
- 26 F. Zordan and L. Brammer, *Acta Crystallogr., Sect. B: Struct. Sci.*, 2004, **60**, 512–519.



- 27 G. Mínguez Espallargas, L. Brammer and P. Sherwood, *Angew. Chem., Int. Ed.*, 2006, **45**, 435–440.
- 28 F. Zordan, G. Mínguez Espallargas and L. Brammer, *CrystEngComm*, 2006, **8**, 425–431.
- 29 F. F. Awwadi, R. D. Willett and B. Twamley, *Cryst. Growth Des.*, 2007, **7**, 624–632.
- 30 G. Mínguez Espallargas, F. Zordan, L. Arroyo Marín, H. Adams, K. Shankland, J. van de Streek and L. Brammer, *Chem.–Eur. J.*, 2009, **15**, 7554–7568.
- 31 P. Metrangolo, T. Pilati, G. Terraneo, S. Biella and G. Resnati, *CrystEngComm*, 2009, **11**, 1187–1196.
- 32 P. Cauliez, V. Polo, T. Roisnel, R. Llusar and M. Fourmigué, *CrystEngComm*, 2010, **12**, 558–566.
- 33 A. Abate, J. Marti-Rujas, P. Metrangolo, T. Pilati, G. Resnati and G. Terraneo, *Cryst. Growth Des.*, 2011, **11**, 4220–4226.
- 34 J. E. Ormond-Prout, P. Smart and L. Brammer, *Cryst. Growth Des.*, 2012, **12**, 205–216.
- 35 A.-C. C. Carlsson, J. Grafenstein, J. L. Laurila, J. Bergquist and M. Erdelyi, *Chem. Commun.*, 2012, **48**, 1458–1460.
- 36 A.-C. C. Carlsson, J. Grafenstein, A. Budnjo, J. L. Laurila, J. Bergquist, A. Karim, R. Kleinmaier, U. Brath and M. Erdelyi, *J. Am. Chem. Soc.*, 2012, **134**, 5706–5715.
- 37 A. Karim, M. Reitti, A.-C. C. Carlsson, J. Grafenstein and M. Erdelyi, *Chem. Sci.*, 2014, **5**, 3226–3233.
- 38 A.-C. C. Carlsson, K. Mehmeti, M. Uhrbom, A. Karim, M. Bedin, R. Puttreddy, R. Kleinmaier, A. A. Neverov, B. Nekoueishahraki, J. Grafenstein, K. Rissanen and M. Erdelyi, *J. Am. Chem. Soc.*, 2016, **138**, 9853–9863.
- 39 L. Pauling and R. E. Marsh, *Proc. Natl. Acad. Sci. U. S. A.*, 1952, **38**, 112.
- 40 K. W. Allen and G. A. Jeffrey, *J. Chem. Phys.*, 1963, **38**, 2304.
- 41 K. A. Udachin, G. D. Enright, C. I. Ratcliffe and J. A. Ripmeester, *J. Am. Chem. Soc.*, 1997, **119**, 11481–11486.
- 42 K. A. Udachin, S. Alavi and J. A. Ripmeester, *J. Phys. Chem. C*, 2013, **117**, 14176–14182.
- 43 L. Brammer, J. K. Swearingen, E. A. Bruton and P. Sherwood, *Proc. Natl. Acad. Sci. U. S. A.*, 2002, **99**, 4956–4961.
- 44 J. J. McKinnon, D. Jayatilaka and M. A. Spackman, *Chem. Commun.*, 2007, 3814–3816.
- 45 M. J. Turner, S. P. Thomas, M. W. Shi, D. Jayatilaka and M. A. Spackman, *Chem. Commun.*, 2015, **51**, 3735–3738.
- 46 C. F. Mackenzie, P. R. Spackman, D. Jayatilaka and M. A. Spackman, *IUCrJ*, 2017, **4**, 575–587.
- 47 A. Gavezzotti, *J. Phys. Chem. B*, 2002, **106**, 4145–4154.
- 48 A. Gavezzotti, *J. Phys. Chem. B*, 2003, **107**, 2344–2353.
- 49 J. D. Dunitz and A. Gavezzotti, *Angew. Chem., Int. Ed.*, 2005, **44**, 1766–1787.
- 50 G. Cavallo, P. Metrangolo, R. Milani, T. Pilati, A. Priimagi, G. Resnati and G. Terraneo, *Chem. Rev.*, 2016, **116**, 2478–2601.
- 51 L. C. Gilday, S. W. Robinson, T. A. Barendt, M. J. Langton, B. R. Mullaney and P. D. Beer, *Chem. Rev.*, 2015, **115**, 7118–7195.
- 52 R. K. Rowe and P. S. Ho, *Acta Crystallogr., Sect. B: Struct. Sci., Cryst. Eng. Mater.*, 2017, **73**, 255–264.



- 53 E. Parisini, P. Metrangolo, T. Pilati, G. Resnati and G. Terraneo, *Chem. Soc. Rev.*, 2011, **40**, 2267–2278.
- 54 L. Brammer, G. Mínguez Espallargas and S. Libri, *CrystEngComm*, 2008, **10**, 1712–1727.
- 55 V. M. McRae, R. D. Peacock and D. R. Russell, *Chem. Commun.*, 1969, 62–63.
- 56 A. J. Edwards and R. J. C. Sills, *J. Chem. Soc. A*, 1970, 2697–2699.
- 57 J. Braddock-Wilking, M. Schieser, L. Brammer, J. Huhmann and R. Shaltout, *J. Organomet. Chem.*, 1995, **499**, 89–98.
- 58 S. Brooker, J.-K. Buijink and F. T. Edelmann, *Organometallics*, 1991, **10**, 25–26.
- 59 H. Grützmacher, H. Pritzkow and F. T. Edelmann, *Organometallics*, 1991, **10**, 23–24.
- 60 U. Lay, H. Pritzkow and H. Grützmacher, *J. Chem. Soc., Chem. Commun.*, 1992, 260–262.
- 61 R. D. Schluter, H. S. Isom, A. H. Cowley, D. A. Atwood, R. A. Jones, F. Olbrich, S. Corbelin and R. J. Lagow, *Organometallics*, 1994, **13**, 4058–4063.
- 62 S. Brooker, N. Bertel, D. Stalke, M. Noltemeyer, H. W. Roesky, G. M. Sheldrick and F. T. Edelmann, *Organometallics*, 1992, **11**, 192–195.
- 63 S. W. Robinson, C. L. Mustoe, N. G. White, A. Brown, A. L. Thompson, P. Kennepohl and P. D. Beer, *J. Am. Chem. Soc.*, 2015, **137**, 499–507.
- 64 M. J. Langton, S. W. Robinson, I. Marques, V. Félix and P. D. Beer, *Nat. Chem.*, 2014, **6**, 1039–1043.
- 65 L. C. Gilday and P. D. Beer, *Chem.–Eur. J.*, 2014, **20**, 8379–8385.
- 66 C. B. Aakeröy, M. Fasulo, N. Schultheiss, J. Desper and C. Moore, *J. Am. Chem. Soc.*, 2007, **129**, 13772–13773.
- 67 C. B. Aakeröy, T. K. Wijethunga, M. A. Haj, J. Desper and C. Moore, *CrystEngComm*, 2014, **16**, 7218–7225.
- 68 C. B. Aakeröy, C. L. Spartz, S. Dembowski, S. Dwyre and J. Desper, *IUCrJ*, 2015, **2**, 498–510.
- 69 M. D. Perera, J. Desper, A. S. Sinha and C. B. Aakeröy, *CrystEngComm*, 2016, **18**, 8631–8636.
- 70 G. R. Desiraju, P. S. Ho, L. Kloo, A. C. Legon, R. Marquardt, P. Metrangolo, P. Politzer, G. Resnati and K. Rissanen, *Pure Appl. Chem.*, 2013, **85**, 1711–1713.
- 71 E. Arunan, G. R. Desiraju, R. A. Klein, J. Sadlej, S. Scheiner, I. Alkorta, D. C. Clary, R. H. Crabtree, J. J. Dannenberg, P. Hobza, H. G. Kjaergaard, A. C. Legon, B. Mennucci and D. J. Nesbitt, *Pure Appl. Chem.*, 2011, **83**, 1619–1636.
- 72 P. Politzer, J. S. Murray and T. Clark, *Phys. Chem. Chem. Phys.*, 2010, **12**, 7748–7757.
- 73 H. Ratajczak, *J. Phys. Chem.*, 1972, **76**, 3000–3004.
- 74 L. C. Allen, *J. Am. Chem. Soc.*, 1975, **97**, 6921–6940.

