

Chemical Science

Volume 16
Number 9
7 March 2025
Pages 3761–4178

rsc.li/chemical-science



ISSN 2041-6539

EDGE ARTICLE

Masahiro Terada *et al.*
Chiral Brønsted acid-catalysed enantioselective allylboration of sterically hindered aldehydes enabled by multiple hydrogen bonding interactions

Cite this: *Chem. Sci.*, 2025, 16, 3865

All publication charges for this article have been paid for by the Royal Society of Chemistry

Chiral Brønsted acid-catalysed enantioselective allylboration of sterically hindered aldehydes enabled by multiple hydrogen bonding interactions†

Shigenobu Umemiya,^a Sotaro Osaka,^b Naoya Shinagawa,^b Takumi Hirata^b and Masahiro Terada^{b*}

Chiral phosphoric acids (CPAs) are an important class of chiral Brønsted acid catalysts that can accomplish highly effective enantioselective allylborations of aldehydes. However, traditional CPA-catalysed allylboration has difficulty utilizing sterically hindered aldehydes, where the corresponding products of enantioenriched secondary alcohols adjacent to a quaternary carbon are important moieties in biologically active natural products. To overcome this issue, we employed a chiral phosphoramidate catalyst for allylation and crotylation reactions of the allyl boronic acid pinacol ester with sterically hindered aldehydes to benefit from multiple hydrogen bonding interactions between the chiral phosphoramidate and substrates. As a result, not only the sterically hindered aldehydes, but the less sterically hindered ones, could also be subjected to enantioselective allylboration using the chiral phosphoramidate catalyst by “interaction strategy”. Indeed, conventional CPAs were ineffective for the reactions presented here, resulting in low conversions and enantioselectivities. Computational studies revealed that the most stable transition state comprises weak attractive interactions between phosphoramidate and substrates. These interactions did not exist in traditional allylborations with chiral phosphoric acids. In conclusion, the sum of weak interactions, including S=O...H-C and two C-F...H-C hydrogen bonding interactions, substantially impacts the enantioselectivity of the allylboration of sterically hindered aldehydes.

Received 13th December 2024

Accepted 19th January 2025

DOI: 10.1039/d4sc08443h

rsc.li/chemical-science

Introduction

Enantioenriched secondary alcohols carrying sterically hindered groups, such as a quaternary carbon adjacent to a stereogenic centre, are important moieties in biologically active natural products. For instance, many macrolides have this structural feature in their enantioenriched form (Fig. 1).¹

As these natural products have potent biological activities and unique structures, numerous efforts have been made toward their total synthesis, prompting tremendous advancements in synthetic methods over the past decades.² For example, the Kiyooka group reported an asymmetric aldol reaction of aldehydes with ketene silyl acetal in the presence of stoichiometric valine derivatives and $\text{BH}_3\cdot\text{THF}$; this reaction produced corresponding aldol adducts possessing an enantioenriched secondary hydroxyl group adjacent to a quaternary

carbon in good yields with high enantioselectivities (Scheme 1a).³ Fürstner's group initiated the enzyme-mediated kinetic resolution of a racemic secondary alcohol next to a quaternary carbon centre to furnish the corresponding chiral acetate in the total synthesis of (–)-polycavernoside A (Scheme 1b).⁴




Fig. 1 Natural product possessing enantioenriched secondary alcohols with neighbouring quaternary carbon.

^aResearch and Analytical Center for Giant Molecules, Graduate School of Science, Tohoku University, 6-3 Aramaki Aza Aoba, Aoba-ku, Sendai 980-8578, Japan

^bDepartment of Chemistry, Graduate School of Science, Tohoku University, 6-3 Aramaki Aza Aoba, Aoba-ku, Sendai 980-8578, Japan. E-mail: mterada@tohoku.ac.jp

† Electronic supplementary information (ESI) available. See DOI: <https://doi.org/10.1039/d4sc08443h>



Scheme 1 Some select examples of the synthesis of enantioenriched secondary alcohols with adjacent quaternary carbon.

While these conventional methods are reliable, the interest in catalytic and stereoselective carbon–carbon bond-forming reactions has increased in green and sustainable chemistry.⁵ For instance, Carreira's group described a catalytic and enantioselective allylation reaction of aldehydes with allyltrimethylsilane in the presence of TiF_4 and BINOL complex (Scheme 2a).⁶ This method effectively generates the corresponding allylation product adjacent to a quaternary carbon with good enantioselectivity, despite a hygroscopic and reactive titanium catalyst being required. In 2008, Krische's group published an elegant Ir-catalysed enantioselective allylation reaction to reveal that 3-(benzyloxy)-2,2-dimethylpropanal is an appropriate substrate to obtain the desired product in good yield with excellent enantioselectivity. However, they did not investigate the substrate scope of this methodology (Scheme 2b).⁷

Chiral Brønsted acid catalysts, including chiral phosphoric acids (CPAs),⁸ are valuable tools for enantioselective reactions. These organocatalytic reactions attract a lot of attention in the context of practical and environmentally benign conditions. Recently, we focused on several interactions of chiral Brønsted acid catalysts to develop the enantioselective intramolecular $\text{S}_{\text{N}}2'$ reaction and the intermolecular enantioselective Diels–

Alder reaction; both of these are challenging reactions to achieve *via* conventional methods.⁹ Herein, we report chiral phosphoramidate-catalysed stereoselective allylation and crotylation reactions of allyl boronic acid pinacol ester that contains sterically hindered and less hindered aldehydes, utilizing multiple hydrogen bonding interactions between chiral phosphoramidate and substrates (Scheme 2c).¹⁰ We found that the sum of weak interactions significantly impacts enantioselectivity in the chiral phosphoramidate-catalysed allylation reaction.


Results and discussion

Initial attempts


We investigated the enantioselective allylation reaction using a SPINOL-derived catalyst, (*S*)-**3a** ($\text{TRIP} = 2,4,6\text{-}i\text{-Pr}_3\text{-C}_6\text{H}_2$), which generally features the best results in the conventional enantioselective allylboration reaction, as reported by Hu's group (Scheme 3).¹¹ However, the reaction of **1a** with **2** in the presence of (*S*)-**3a** at -75°C afforded a product with low enantioselectivity (40% ee), even though the chemical yield was good. The reaction catalysed by BINOL-derived CPA (*R*)-**3b**, which also used Antilla's allylboration¹² method, also gave an unsatisfactory result, albeit with a slight improvement in enantioselectivity (67% ee). These results show that conventional CPAs are ineffective for sterically hindered aldehydes. Importantly, when the acidic catalyst, (*R*)-**3c**, rather than (*S*)-**3a** or (*R*)-**3b**, was used, the reaction rate was sluggish, and both chemical yield and selectivity dropped. This result suggests that high catalyst acidity is unrelated to the activation of the allylation reaction. Although various enantioselective allylation reactions using CPA¹³ have been developed due to their importance and utility in organic synthetic chemistry, enantioselective allylation reactions of bulky aldehydes with allylboronic agents have yet to be reported. Therefore, many challenges remain in the use of sterically congested substrates in catalytic and enantioselective allylboration.

The Goodman and Houk groups independently performed intensive theoretical studies of the reaction mechanism in the case of benzaldehyde.¹⁴ The most stable transition state (TS) in the BINOL-derived (*R*)-CPA-catalyzed allylation reaction of benzaldehyde with an allylboronic agent is shown in Fig. 2.

Catalytic allylation reactions using sterically hindered aldehydes



Scheme 2 Representative examples of catalytic and enantioselective allylation reactions using sterically hindered aldehydes and those presented in the Results and discussion section.



Scheme 3 The initial attempt of chiral Brønsted acid-catalyzed allylboration of pivalaldehyde (**1a**) with **2**.




According to this TS model, the CPA catalyst, (*R*)-**3b**, does not activate the carbonyl group of the benzaldehyde by protonation. However, it stabilizes the six-membered TS through two-point hydrogen bonding interactions with the axial oxygen of the boronic ester (P–O–H···O–B) and the formyl hydrogen of the benzaldehyde (P=O···H–C). Notably, DFT calculations have shown that the acidic proton of the CPA catalyst and the axial oxygen atom (P–O–H···O–B) are approximately 1.5 Å apart, indicating that it is not protonation of the oxygen atom, but the hydrogen bonding instead causing stabilization. From these calculation studies, the unfavorable result observed for the CPA (*R*)-**3c** is justifiable because high catalyst acidity is not necessarily a crucial factor stabilizing TS (**Re_A**) during allylboration reactions. Hence, we focused on chiral phosphoramides, rather than CPAs, to stabilize TS in the major reaction pathway using an “interaction strategy”.^{9b}

Development of a new catalytic system through multiple interactions

Through our previous work, we found that the phosphoramidate catalyst with a C₆F₅ group at the sulfonamide moiety yielded favourable results.^{9a} Thus, we first investigated the allylation reactions of **1a** with **2** using (*R*)-**3d** having a C₆F₅SO₂NH moiety (Table 1, entry 1). To our delight, the reaction proceeded smoothly to furnish the desired product in 98% yield with good enantioselectivity (84% ee). When we investigated other substituents on the *N* moiety of the catalyst, we found that the catalyst, (*R*)-**3e**, containing a SO₂CF₃ (Tf) group exhibited better reactivity than (*R*)-**3d**, and thus, it helped obtain the desired product in quantitative yield with higher enantioselectivity (95% ee) than that (84% ee) observed using (*R*)-**3d** (entry 2). The absolute configuration of compound **4a** obtained in this reaction was consistent with those obtained with CPA (*R*)-**3b** having the same chiral backbone as phosphoramidate (*R*)-**3e**. No reaction occurred when the catalyst (*R*)-**3f** containing an SO₂Me group was employed (entry 3). The substituents on the sulfonamide moiety of the catalyst significantly influenced both reactivity and enantioselectivity. The reaction did not proceed when a more acidic thiophosphoramidate catalyst, (*R*)-**3g**, that was previously reported by Yamamoto and co-workers,¹⁵ was used (entry 4). These results suggest that high catalyst acidity is not essential in the allylation reaction of aldehydes with allylboration agents; this can also be recognized from the TS model of Goodman (Fig. 2). SPINOL-derived phosphoramidate catalyst, (*S*)-**3h**, was ineffective in the

Table 1 Optimization of the reaction conditions using **1a** and **2** catalysed by chiral phosphoramidate **3**^a

Entry	Catalyst	Time (h)	Yield of (<i>R</i>)- 4a ^b /%	ee/%
1	(<i>R</i>)- 3d	72	98	84
2	(<i>R</i>)- 3e	24	Quant.	95
3	(<i>R</i>)- 3f	72	No reaction	nd
4	(<i>R</i>)- 3g	72	No reaction	nd
5	(<i>S</i>)- 3h	72	No reaction	nd
6 ^c	(<i>R</i>)- 3e	48	Quant.	95



^a Unless otherwise specified, all reactions were carried out using 0.20 mmol of **1a**, 0.24 mmol (1.2 eq.) of **2**, and 0.010 mmol (5 mol%) of catalyst. ^b Isolated yield. ^c Catalyst loading was 0.5 mol%, and 2 g of pivalaldehyde was employed. TRIP = 2,4,6-*i*-Pr₃-C₆H₂, nd = not determined.

present reaction, probably because of a narrow reaction space (entry 5).¹⁶ After comprehensively investigating reaction conditions, including the solvent, temperature, and concentration, the experimental condition that uses (*R*)-**3e** (entry 2) was found to be the best (see ESI†). Finally, catalyst loading could be reduced to 0.5 mol% in a large-scale experiment. The product was quantitatively formed on a 2 g scale by extending the reaction time without losing enantioselectivity (entry 6).

Transition state analysis

After investigating the optimum conditions of phosphoramidate-catalysed allylboration reaction with pivalaldehyde, we performed DFT computational studies to gain mechanistic insights into the present reaction system. As the present reaction of pivalaldehyde (**1a**) afforded (*R*)-product **4a** in the presence of the catalyst, (*R*)-**3e**, the TS in the *re*-face attack had a lower energy barrier than that in the *si*-face attack.

We calculated four TSs for the *re*-face and the *si*-face attacks, considering the axial and equatorial coordination models, respectively (Fig. 3).¹⁴ DFT calculation was performed using Gaussian 16 in the M06-2X/6-31G(d) level of theory.¹⁷ As a result, TS **Re_A**, namely, *re*-face attack and coordination to the axial oxygen of boronic ester, produced the lowest energy. Resembling CPA-catalysed allylboration, two-point interactions were produced through the hydrogen bonding of the phosphoryl oxygen of the catalyst with the proton of the formyl group (2.43 Å); hydrogen bonding was observed between the axial boronate oxygen and the hydrogen atom (N–H) of the catalyst (1.57 Å) in **Re_A**. We also found that the TS of the major reaction pathway (major-TS) was preferred over the TS of the minor reaction pathway (minor-TS), namely **Si_E**, by 4.2 kcal mol^{−1}. Single-point energies were conducted with the MN15/6-31G(d)+SMD




Fig. 2 Goodman's transition state model.





Fig. 3 Optimized transition states of **Re_A**, **Re_E**, **Si_A**, and **Se_E** at the M06-2X/6-31G(d) level of theory. Bond lengths are given in Å. Values enclosed in parentheses are energies relative to “**Re_A**” calculated by MN15/6-31g(d)+SDM (toluene). Energy differences are given in kcal mol^{−1}. See the ESI† on page S38 in detail.

(toluene) to result in a 3.0 kcal mol^{−1} energy difference.¹⁸ These calculation results are in relatively good agreement with the experimentally observed enantiomeric excess of (*R*)-**3e** (95% ee, theoretically *ca.* 2.2 kcal mol^{−1}).¹⁹ In the major-TS (**Re_A**), the distance for N-H...O-B hydrogen bonding was 1.57 Å, which remained almost unchanged from the calculation result in the CPA-catalysed allylboration.¹⁴ This suggests that the high acidity of the phosphoramidate catalyst does not enhance the N-H...O-B hydrogen bonding in this system because the hydrogen bond is not shortened. Interestingly, not only the hydrogen bonding between the sulfonyl moiety and the olefin C-H (S=O...H-C, 2.47 Å) but also the two weak interactions between a fluorine atom of the CF₃ group and the hydrogen atoms of the pinacol Me group (C-F...H-C, 2.25 and 2.57 Å) were found in addition to the major two-point hydrogen bonding interactions in **Re_A**. Conversely, in the minor-TS (**Si_E**), the P=O...H-C (formyl) interaction and C-F...H-C interactions are not present. Similarly, the C-F...H-C interactions were not observed in the minor-TSs (**Re_E** and **Si_A**). These observations indicate that the minor-TSs were less stable than **Re_A**. This was not due to the

steric congestion but due to the lack of multiple forces of attraction.²⁰ The sum of the multiple hydrogen bonding interactions, including the relatively weak S=O...H-C and C-F...H-C hydrogen bonding interactions, which are not present in conventional CPA-catalysed allylation reactions, is considered to effectively stabilize the major-TS to give the desired product with excellent enantioselectivity. As the catalyst does not occupy the space around the ^tBu group in **Re_A**, we envisioned that various substituents could be introduced to aldehydes. Based on the DFT calculation results, we proceeded to investigate the substrate scope.

Substrate scope

After rationally elucidating the high enantioselectivity of the present allylation reaction under optimal reaction conditions, we explored the substrate scope of the present reaction toward aldehydes with a quaternary centre at the α-position (Scheme 4). Various functional groups at the β-position of the formyl group were found to be suitable for the allylation reaction. They could



effectively furnish the corresponding homoallyl alcohols in excellent yields with good enantioselectivities (**4b–4d**). In contrast, the SPINOL-derived catalyst, (*S*)-**3a**, was ineffective for these substrates, resulting in low conversions and enantioselectivities.^{21a} Aldehydes with sensitive functional groups, such as a Ts group, an α,β -unsaturated ester, and an enone, were also tolerated and they produced the corresponding products with a quantitative yield of up to 95% ee (**4e–4g**). Next, the scope of less hindered aldehydes, such as cyclohexane carboxaldehyde and benzaldehyde, was investigated. The corresponding products were obtained in good yields with high enantioselectivities (**4h**, **4i**). Unfortunately, the allylboration of benzaldehyde was inferior to that of the CPA-catalysed reactions using (*S*)-**3a** and (*R*)-**3b** (99% ee and 98% ee, respectively), whereas the enantioselectivity of the reaction using cyclohexane carboxaldehyde was similar to that with (*S*)-**3a** (91% ee) and superior to that with



Fig. 4 Derivatization of products.

(*R*)-**3b** (73% ee).^{11,12} Next, we applied less sterically hindered alkyl aldehydes to our system, and found that these afforded **4j** and **4k** in good yields and ee values (85% ee and 83% ee, respectively). A similar tendency to reduce enantioselectivities in the case of less hindered aldehydes was observed in our reaction. Diastereoselective allylation reactions of enantioenriched aldehydes with substituents at the α - or β -position under optimized conditions produced the desired products in excellent yields with moderate to high diastereoselectivities (**4l–4o**). In contrast, the conventional CPA-catalysed reactions produced **4n** and **4o** with low diastereoselectivities.^{21b} Subsequently, we extended this protocol to crotylation reactions using diverse aldehydes, including those having a stereogenic centre. Various enantioenriched substrates worked well and afforded products, **4p–4s**, in good yields and moderate to reasonably high diastereomeric ratios (drs). Notably, the *anti*-selective crotylation products, **4p** and **4q**, with three continuous stereogenic centres, commonly found in natural products, such as marine macrocides, were obtained with excellent diastereoselectivities. Not only the enantioenriched aldehydes but also simple aldehydes, such as benzaldehyde and 3-phenylpropionaldehyde, were suitable crotylation, furnishing corresponding *anti*-products in good yields (**4t** and **4u**). In contrast, sterically hindered 3-(benzoyloxy)-2,2-dimethylpropanal did not yield the desired product, and there was no reaction even when the temperature was elevated to 0 °C (**4v**).

Derivatization

A derivatization of products was attempted to show the synthetic utility of the present allylation and crotylation reactions to construct functional chiral building blocks (Fig. 4). The enantioenriched alcohol **4g** was treated with *p*-nitrobenzaldehyde in the presence of catalytic trifluoromethanesulfonic acid (TfOH) to afford sterically congested 1,3-*syn*-dioxane derivative **5** with 87% yield and excellent diastereoselectivity (eqn (1)).²² Alternatively, the esterification of compound **4q** using acryloyl chloride, followed by ring-closing metathesis, produced functionalized α,β -unsaturated lactone **6** in 91% yield in two steps (eqn (2)).

Conclusions

We have developed an enantio- and diastereoselective allylboration of sterically hindered aldehydes that utilize multiple



Scheme 4 ^aUnless otherwise specified, all reactions were carried out using 0.20 mmol of **1a**, 0.24 mmol (1.2 eq.) of **2**, and 0.010 mmol (5 mol%) of catalyst. ^bIsolated yield. ^c(*S*)-**3e** was used instead of (*R*)-**3e**. TRIP = 2,4,6-ⁱPr₃-C₆H₂. Chiral phosphoramidate-catalysed allylation and crotylation of aldehydes.^{a,b}

hydrogen bonding interactions between the chiral phosphoramidate catalyst and substrate to furnish synthetically useful secondary alcohols with a quaternary carbon adjacent to a stereogenic centre. Computational studies suggest that the major-TS is stabilized by weak attractive interactions, including S=O...H-C and two C-F...H-C hydrogen bonding interactions. In contrast, minor-TSs lack some of these multiple interactions. Hence, the sum of weak interactions substantially impacts the enantioselectivity in the allylboration of sterically hindered aldehydes. Using phosphoramidate **3e** as a multiple interaction catalyst, 0.5 mol% catalyst loading is sufficient to complete the allylboration of pivalaldehyde on a gram-scale. The present allylation and crotylation reactions can be applied to both sterically hindered and less hindered aldehydes to obtain the corresponding homoallylic alcohols in excellent yields and high enantioselectivities. Further studies on the efficient synthesis of important natural products using the present methodology are in progress in our laboratory.

Data availability

The exploratory investigation results, experimental procedures, computational data, and characterization data are available.

Author contributions

S. U. contributed to the conceptualization, project administration, and design of the work, as well as data curation, formal analysis, investigation (experimental and theoretical studies), writing – original draft. S. O., N. S., and T. H. contributed to the formal analysis and investigation (experimental). M. T. contributed to the conceptualization, writing – review and editing, supervision, and funding acquisition.

Conflicts of interest

There are no conflicts to declare.

Acknowledgements

The computation was performed at the Research Center for Computational Science, Okazaki, Japan. This work was partially supported by a Grant-in-Aid for Scientific Research on Innovative Areas “Hybrid Catalysis for Enabling Molecular Synthesis on Demand” (JP17H06447) and a Grant-in-Aid for Transformative Research Areas (A) “Green Catalysis Science for Renovating Transformation of Carbon-Based Resources” (JP23H04908) from MEXT, Japan (M. T.), a Grant-in-Aid for Scientific Research (C) (23K04730) from JSPS, Japan (S. U.) and a Grant-in-Aid for Young Scientists (24KJ0444) from JSPS, Japan (N. S.).

Notes and references

- 1 Isolations of these natural products, see: (a) C.-L. Shao, R. G. Linington, M. J. Balunas, A. Centeno, P. Boudreau, C. Zhang, N. Engene, C. Spadafora, T. S. Mutka, D. E. Kyle,

- L. Gerwick, C.-Y. Wang and W. H. Gerwick, *J. Org. Chem.*, 2015, **80**, 7849–7855; (b) K. Gerth, N. Bedorf, G. Hofle, H. Irschik and H. J. Reichenbach, *J. Antibiot.*, 1996, **49**, 560–563; (c) K. Umeda, A. Iwasaki, R. Taguchi, N. Kurisawa, G. Jeelani and T. N. K. Suenaga, *J. Nat. Prod.*, 2023, **86**, 2529–2538; (d) G. R. Pettit, J. F. Day, J. L. Hartwell and H. B. Wood, *Nature*, 1970, **227**, 962–963; (e) G. R. Pettit, C. L. Herald, D. L. Doubek and D. L. Herald, *J. Am. Chem. Soc.*, 1982, **104**, 6846–6848.
- 2 Selected examples of total syntheses of natural products possessing quaternary carbon adjacent to enantioenriched secondary alcohol moieties, see: (a) A. Balog, D. Meng, T. Kamenecka, P. Bertinato, D.-S. Su, E. J. Sorensen and S. J. Danishefsky, *Angew. Chem., Int. Ed.*, 1996, **35**, 2801–2803; (b) K. Ohmori, Y. Ogawa, T. Obitsu, Y. Ishikawa, S. Nishiyama and S. Yamamura, *Angew. Chem., Int. Ed.*, 2000, **39**, 2290–2294; (c) B. M. Trost, Y. Wang, A. K. Buckl, Z. Huang, M. H. Nguyen and O. Kuzmina, *Science*, 2020, **368**, 1007–1011; (d) J. B. Cox, A. A. Kellum, Y. Zhang, B. Li and A. B. Smith III, *Angew. Chem., Int. Ed.*, 2022, **61**, e202204884.
- 3 S. Kiyooka, Y. Kaneko, M. Komura, H. Matsuo and M. Nakano, *J. Org. Chem.*, 1991, **56**, 2276–2278.
- 4 L. Brewitz, J. Llaveria, A. Yada and A. Fürstner, *Chem.-Eur. J.*, 2013, **19**, 4532–4537.
- 5 For selected reviews, see: (a) P. T. Anastas, M. M. Kirchhoff and T. C. Williamson, *Appl. Catal., A*, 2001, **221**, 3; (b) J. F. Jenck, F. Agterberg and M. J. Droesch, *Green Chem.*, 2004, **6**, 544; (c) I. Delidovich and R. Palkovits, *Green Chem.*, 2016, **18**, 590.
- 6 D. R. Gauthier Jr and E. M. Carreira, *Angew. Chem., Int. Ed.*, 1996, **35**, 2363–2365.
- 7 (a) I. S. Kim, M.-Y. Ngai and M. J. Krische, *J. Am. Chem. Soc.*, 2008, **130**, 14891–14899; (b) I. S. Kim, S. B. Han and M. J. Krische, *J. Am. Chem. Soc.*, 2009, **131**, 2514–2520; (c) When more inexpensive BINAP ligand than MeO-BIPHEP ligand was used for the Krische allylation, decrease of enantioselectivity was observed (77% ee) in our investigation. See the ESI† in detail.; (d) Recently, the Krische group reported an enantioselective *tert*-prenylation using chiral ruthenium catalyst, see: *J. Am. Chem. Soc.*, 2023, **145**, 18676–18683.
- 8 For seminal studies of chiral phosphoric acid catalysts, see: (a) T. Akiyama, J. Itoh, K. Yokota and K. Fuchibe, *Angew. Chem., Int. Ed.*, 2004, **43**, 1566–1568; (b) D. Uraguchi and M. Terada, *J. Am. Chem. Soc.*, 2004, **126**, 5356–5357.
- 9 (a) M. Shimizu, J. Kikuchi, A. Kondoh and M. Terada, *Chem. Sci.*, 2018, **9**, 5747–5757; (b) T. Nakanishi and M. Terada, *Chem. Sci.*, 2023, **14**, 5712–5721.
- 10 For seminal study of chiral phosphoramidate catalyst, see: (a) D. Nakashima and H. Yamamoto, *J. Am. Chem. Soc.*, 2006, **128**, 9626–9627. For a recent review, see: (b) G. Caballero-García and J. M. Goodman, *Org. Biomol. Chem.*, 2021, **19**, 9565–9618.
- 11 C.-H. Xing, Y.-X. Liao, Y. Zhang, D. Sabarova, M. Bassous and Q.-S. Hu, *Eur. J. Org. Chem.*, 2012, 1115–1118.



- 12 P. Jain and J. C. Antilla, *J. Am. Chem. Soc.*, 2010, **132**, 11884–11886.
- 13 Selected recent examples of enantioselective allylboration catalyzed by CPA, see: (a) E. Rodríguez, M. N. Grayson, A. Asensio, P. Barrio, K. N. Houk and S. Fustero, *ACS Catal.*, 2016, **6**, 2506–2514; (b) S. Gao, M. Duan, Q. Shao, K. N. Houk and M. Chen, *J. Am. Chem. Soc.*, 2020, **142**, 18355–18368; (c) S. Gao, M. Duan, L. R. Andreola, P. Yu, S. E. Wheeler, K. N. Houk and M. Chen, *Angew. Chem., Int. Ed.*, 2022, **61**, e202208908.
- 14 (a) M. N. Grayson, S. C. Pellegrinet and J. M. Goodman, *J. Am. Chem. Soc.*, 2012, **134**, 2716–2722; (b) H. Wang, P. Jain, J. C. Antilla and K. N. Houk, *J. Org. Chem.*, 2013, **78**, 1208–1215.
- 15 M. Sai and H. Yamamoto, *J. Am. Chem. Soc.*, 2015, **137**, 7091–7094.
- 16 H. Ye, T. Sato, T. Nakanishi, S. Ito, S. Umemiya and M. Terada, *Catal. Sci. Technol.*, 2024, **14**, 6869–6881.
- 17 Whereas Goodman and Houk used the B3LYP density function, we decided to use M06-2X because of the importance of weak attractive interactions in our system. Indeed, after some calculation studies, we found that B3LYP was unsuitable for the present reaction that includes multiple weak interactions. see the ESI†
- 18 H. S. Yu, X. He, S. L. Lia and D. G. Truhlar, *Chem. Sci.*, 2016, **7**, 5032–5051.
- 19 Similar tendency of energy differences was observed by using B3LYP-D3(BJ)/6-311G(d, p) and wB97XD/6-31G(d). See the ESI† in details.
- 20 (a) Interestingly, TS **Si_E** was more stable than TS **Si_A** despite the absence of hydrogen bonding between the phosphoryl oxygen and the hydrogen atom of the formyl group. In TS **Si_E**, weak attractive interactions of S=O...H-C (sp³) in the pinacol moiety of **2** and the 'Bu group of pivalaldehyde were observed in DFT calculations; (b) Goodman's group performed the asymmetric propargylation of cyclohexanecarboxaldehyde with allenylboronic acid pinacol ester in the presence of (*R*)-**3e** to give a product with only 39% ee. They did not mention the multiple hydrogen bonding interactions between phosphoramidate and substrates. See: M. N. Grayson and J. M. Goodman, *J. Am. Chem. Soc.*, 2013, **135**, 6142–6148.
- 21 (a) For instance, the allylation reaction using SPINOL-derived catalyst (*S*)-**3a** afforded product **4b** in 7% yield and 33% ee. See the ESI† for experimental results in detail.; (b) The reaction using (*R*)-**3a** gave 1,3-*syn* product **4n** with dr = 3 : 1, and the reaction using (*S*)-**3a** afforded 1,3-*anti* product **4o** with dr = 2 : 1.
- 22 Selected examples of synthesis of 1,3-*syn* diol derivatives via hemiacetalization/intramolecular oxy-Michael reactions, see: (a) P. A. Evans, A. Grisin and M. J. Lawler, *J. Am. Chem. Soc.*, 2012, **134**, 2856–2859; (b) Y. Hayashi, T. Saitoh, H. Arase, G. Kawauchi, N. Takeda, Y. Shimasaki and I. Sato, *Chem.-Eur. J.*, 2018, **24**, 4909–4915; (c) K. Murata, H. Takeshita, K. Sakamoto and H. Fuwa, *Chem.-Asian. J.*, 2020, **15**, 807–819.

