Chemical Science



COMMENTARY

View Article Online View Journal | View Issue



Cite this: Chem. Sci., 2023, 14, 7608

d All publication charges for this article have been paid for by the Royal Society of Chemistry

DOI: 10.1039/d3sc90114a

rsc.li/chemical-science

A focus on 1-azahomocubane: the new kid on the block

Cecile Elgindy and Mark D. Levin*

Strained hydrocarbons have recently regained interest as potential drug candidates. However, the study of their heteroatom analogs has remained limited, despite differing by only a single atom. The first synthesis of 1-azahomocubane by Williams, Eaton and co-workers (T. Fahrenhorst-Jones et al., Chem. Sci., 2023, 14, 2821-2825, https://doi.org/10.1039/D3SC00001J) is discussed within the context of nitrogen scanning of strained hydrocarbons.

family

providing

related

to

The exchange of a single atom can dramatically alter the properties of molecule. Reactivity, biological activity, stability, selectivity, physical properties, and synthetic tractability of the molecule can all be impacted, simply by the substitution of one atom for another. Such effects have been observed in natural products and drug candidates, exemplified by the "necessary nitrogen effect" in medicinal chemistry whereby the replacement of a carbon atom for a nitrogen atom in an aromatic system can profoundly alter molecule's the pharmacological profile.1,2

Meanwhile, renewed interest in strained hydrocarbons has arisen from their potential as benzene bioisosteres, which increase the parent molecule's three-dimensionality without introducing substantial metabolic hotspots, owing to the high s character of the exocyclic C-H bonds.3 Cubane, bicyclo [1.1.1]pentane, and bicyclo[1.1.2]hexane have all been proposed as attractive substructures for drug candidates along these lines, which has led to a flurry of activity examining the functionalization of these scaffolds, with a particular focus on C-H functionalisation and strain release methods for their synthesis.4-6

compounds. Surprisingly, however, very few of the aza-analogs of these hydrocarbons are known, with two azahomocubanes and a small number of substituted azabicyclohexanes previously reported, mostly of the 2-azabicyclohexane class (Fig. 1).7-9 Williams, Eaton, and co-workers disclose the first synthesis of 1-azahomocubane, an aza-analog of homo-(https://doi.org/10.1039/ cubane D3SC00001J).10 The synthesis and study of this surprisingly stable, strained molecule is a rare foray into the azahomocubane series,

useful

One can imagine, in analogy to the

necessary nitrogen effect for benzenoid

systems, that the corresponding hetero-

cyclic series would be valuable in the

hydrocarbon

The synthesis pinpointed azidyl cubane as a strategic intermediate that enabled the incorporation of nitrogen into the core of the strained ring framework, building upon previous reports of a substituted 9-azahomocubane.11 A sequence of rearrangements beginning from the azidyl cubane was initiated by strong acid followed by Nchlorination, resulting in nitrogen incorporation into the ring system. To complete the synthesis, functional

comparison

this underexplored structural space.

structures and enabling insights into

group manipulation enabled the requisite ring closure via N-alkylation to generate the parent 1-azahomocubane, which was trapped in situ as a salt to simplify its isolation (Fig. 2).

With the target compound in hand, a preliminary survey of chemical and physical properties could be conducted. The free amine was found to be stable as a solution over months, whereas the hydrochloride salt was observed to undergo chloride mediated ringopening. 1-Azahomocubane was calculated to be more stable than homocubane and was observed to be less reactive towards Ag+ mediated isomerisations, suggesting that nitrogen-atom exchange enhances the stability of the strained ring system. Indeed, an X-ray structure of the bis(azahomocubane) silver(1) perchlorate salt was collected. Finally, azahomocubane was found to be meaningfully less basic than related polycyclic tertiary amines, with the strain likely additional inducing increased lone pair s-character.

This work by Williams, Eaton and coworkers highlights an underexplored region of chemical space. The synthesis of strained polycyclic heterocycles conceptually derived from benzene bioisosteres is not without challenges, but this work will hopefully spur further development in this area.

Department of Chemistry, University of Chicago, Chicago, IL 60637, USA. E-mail: marklevin@uchicago. edu

Commentary




Fig. 1 The necessary nitrogen effect in benzenoid systems, imagined for benzene bioisosteres




Fig. 2 Synthesis of 1-azahomocubane

Author contributions

All authors contributed equally.

Conflicts of interest

There are no conflicts to declare.

References

- 1 D. L. Boger, J. Org. Chem., 2017, 82, 11961-11980.
- 2 L. D. Pennington and D. T. Moustakas, J. Med. Chem., 2017, 60, 3552-3579.
- 3 F. Lovering, J. Bikker and C. Humblet, J. Med. Chem., 2009, 52, 6752-6756.
- 4 (a) T. A. Reekie, C. M. Williams, L. M. Rendina and M. Kassiou, J. Med. Chem., 2019, **62**, 1078–1095; (b)

- P. K. Mykhailiuk, Org. Biomol. Chem., 2019, 17, 2839-2849.
- 5 P. S. Baran, et al., J. Am. Chem. Soc., 2017, 139, 3209-3226.
- 6 For some recent examples of BCP and C–H functionalisation: I. F. Yu, J. L. Manske, A. Diéguez-Vázquez, A. Misale, A. E. Pashenko, P. K. Mykhailiuk, S. V. Ryabukhin, D. M. Volochnyuk and J. F. Hartwig, 2023, **15**, 685–693; (b) O. L. Garry, M. Heilmann, J. Chen, Y. Liang, X. Zhang, X. Ma, C. S. Yeung, D. J. Bennett and D. W. C. MacMillan, J. Am. Chem. Soc., 2023, 145, 3092-3100. 7 (a) P. E. Eaton, A. M. Fisher and R. E. Hormann, Synlett, 1990, 12, 737-
- P. Ε. Eaton (b) R. E. Hormann, J. Am. Chem. Soc., 1987, 109, 1268-1269.
- 8 (a) G. R. Krow, Y. B. Lee, W. S. Lester, H. Christian, D. A. Shaw and J. Yuan, J. Org. Chem., 1998, 63, 8558-8560; (b) J. W. Davies, J. R. Malpass and M. P. Walker, J. Chem. Soc., Chem. Commun., 1985, 686-687; G. Mloston and H. Heimgartner, Helv. Chim. Acta, 2006, 89, 442-449; For a review of substituted azabicyclo [2.1.1]hexanes: (d) G. R. Krow and K. C. Cannon, Heterocycles, 2004, 62, 877-898; For more recent examples of substituted aza-bicyclo[2.1.1]hexanes: (e) K. Dhake, K. J. Woelk, J. Becica, A. Un, S. E. Jenny and D. C. Leitch, Angew. Chem., Int. Ed., 2022, 61, e202204719; (f) J. L. Tyler, A. Noble and V. K. Aggarwal, Angew. Chem., Int. Ed.,2022, 61, e202114235; Wright, Matviitsuk,

M. J. Black, P. García-Reynaga, L. E. Hanna, A. T. Herrmann, M. K. Ameriks, R. Sarpong and T. P. Lebold, *J. Am. Chem. Soc.*, 2023, 145, 10960–10966.

Chemical Science

- 9 A 5-aza-bicyclohexane: (a) T. Rigotti and T. Bach, *Org. Lett.*, 2022, 24, 8821–8825;
 A benzo-fused analog: (b) K. Takahashi, K. Suenobu, K. Ogura and H. Iida, *Chem. Lett.*, 1985, 14, 1487–1488;
- Putative characterization in a mixture: (c) A. G. Griesbeck and H. Heckroth, *J. Am. Chem. Soc.*, 2002, **124**, 396–403; Claimed but later reassigned: (d) H. F. Olivo, M. S. Hemenway and M. H. Gezginci, *Tetrahedron Lett.*, 1998, **39**, 1309–1312.
- 10 T. Fahrenhorst-Jones, D. L. Marshall,J. M. Burns, G. K. Pierens,R. E. Hormann, A. M. Fisher,
- P. V. Bernhardt, S. J. Blanksby, G. P. Savage, P. E. Eaton and C. M. Williams, *Chem. Sci.*, 2023, 14, 2821–2825.
- 11 R. Gilardi, C. George, J. Karle, P. E. Eaton and A. M. Fisher, J. Heterocycl. Chem., 1993, 30, 1385–1388.