



INORGANIC CHEMISTRY

FRONTIERS

Accepted Manuscript



This article can be cited before page numbers have been issued, to do this please use: D. W. Leong, Y. Shao, N. Bhuvanesh and O. V. Ozerov, *Inorg. Chem. Front.*, 2024, DOI: 10.1039/D4QI01563K.



This is an Accepted Manuscript, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this Accepted Manuscript with the edited and formatted Advance Article as soon as it is available.

You can find more information about Accepted Manuscripts in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this Accepted Manuscript or any consequences arising from the use of any information it contains.

A Redox-Active Ligand Combines a PCP Pincer Site with a Bidentate N-N Donor in Opposition

Derek W. Leong[†], Yanwu Shao[†], Nattamai Bhuvanesh, and Oleg V. Ozerov*

Department of Chemistry, Texas A&M University, 3255 TAMU, College Station, TX 77842.

ozarov@chem.tamu.edu

[†] These authors contributed equally.

ABSTRACT. A binucleating ligand (**2**) combining a monoanionic PCP pincer cleft with a monoanionic N-N cleft has been prepared on the basis of bis(imidazolyl)methane (**1**). Installation of divalent Pd or Pt into the PCP cleft proceeded smoothly, with the formation of square planar (PCP)MCl complexes (M = Pd or Pt). The “C” in the PCP is the central carbon of the β -diketiminato-like N-N cleft. The N-N cleft in these complexes was obtained in three levels of protonation. In **4-Pd** and **4-Pt**, the N-N cleft carries one proton and is neutral; in **3-Pd** or **3-Pt**, it is additionally protonated by HCl, and in **6-Pd** or **6-Pt**, the proton has been removed by the $(\text{Me}_3\text{Si})_2\text{NLi}$ base and replaced by Li. The chloride in **4-Pd/Pt** was replaced with an iodide to make **6-Pd/Pt** in a metathesis with Me_3SiI . Compounds **4-Pd/Pt** and **6-Pd/Pt** exhibited tautomerism reminiscent of acetylacetonate and β -diketimines, possessing either an NH or a CH bond. Compounds **6-Pd/Pt** possess a lithiated N-N cleft and serve as convenient transmetallation agents. Their reactions with $\text{BF}_3(\text{OEt}_2)$, $\frac{1}{2} \text{ZnCl}_2$, or TaCl_5 produced new compounds **7-Pd/Pt**, **8-Pd/Pt**, and **9-Pd/Pt** that carry a BF_2 , $\frac{1}{2} \text{Zn}$, or TaCl_4 units in the N-N cleft. Compounds **8-Pd/Pt** are trimetallic with a central Zn coordinated by two N-N clefts each of which carries a (PCP)MCl on the opposite side. Structures of **3-Pd**, **7-Pt**, **8-Pd**, and **9-Pd** in the solid state were established by X-ray diffractometry. They demonstrate the remarkable planarity of the extended conjugated organic π -system into which the N-N and the PCP binding sites are incorporated.



Electrochemical studies on **7-9** established that each of these extended π -systems can be oxidized twice quasi-reversibly. Compounds **9-Pd/Pt** could be additionally reduced quasi-reversibly by a single electron, which was ascribed to the reduction of the Ta^V center to Ta^{IV}. They also differed from **7** and **8** in possessing an apparent lower-energy LMCT bands responsible for the blue or purple colors.



Introduction

Pincer ligands have become ubiquitous in the studies of organometallic structure and reactivity.¹⁻³

Pincer-supported transition metal complexes have been studied as catalysts for a large variety of applications,¹⁻³ including alkane dehydrogenation,^{4,5} transfer (de)hydrogenation of compounds with C=N/C-N and C=O/C-O bonds,^{6,7,8} C-C coupling,^{9,10} C-H borylation,^{11,12} and nucleophilic addition to ketones and aldehydes.¹³ The tridentate pincer cleft provides a robust environment for the transition metal, in some cases enabling catalytic performance at a temperature of 200 °C or higher,¹⁴ which is unusual for organometallic transformations. The most obvious way to influence the transition metal center in a pincer environment is through the modification of the pincer donor sites. However, substantial attention has been paid to the more remote influences, including the use of redox non-innocent ligands¹⁵⁻²⁰ and the provision of distal binding sites for the coordination of additional metals.^{21,22} The so-called Janus bis-pincer systems (**A-D**, Figure 1) can be included within this context, with one pincer-metal unit providing a remote influence on the other.²³⁻³⁰ The Janus bis-pincers provide two opposing binding cleft sites of the same kind, naturally leading to symmetric bimetallic complexes. There also exist examples of non-Janus bis(pincer) ligands with a shared atom or unit linking the two pincer domains.³¹⁻³⁴ However, it is synthetically difficult to prepare *heterobimetallic* complexes with symmetric bis(pincer) ligands. Considering ways to introduce two very different metals into the same molecule, we became interested in preparing a “dissymmetric Janus” type of ligand, with one phosphine-anchored pincer site suitable for a late transition metal, and another N-N bidentate site amenable to binding a broad variety of metals and metalloids. This arrangement provides a way to influence the structure and reactivity of the pincer ligated transition metal center through variation of the other element in the N-N cleft, but without changing the immediate coordination environment about the transition metal. We envisaged an N-N site similar to β -diketiminates (also known as “nacnac” ligands)³⁵ that is integrated into a redox-active pincer framework. In considering this design, we were influenced by the binucleating “nindigo” ligands (**E**, Figure 1)



pioneered by the Hicks group,³⁶ and also explored by Caulton and Mindiola,³⁷ and by Kaim and Lahiri with co-workers.³⁸ The “nindigo” ligands are essentially a redox non-innocent Janus bis-“nacnac” system. Herein we report on the chemistry of a binucleating ligand fusing a PCP pincer with a “nacnac”-like N-N site. A similar binucleating ligand combining an NCN pincer with a bidentate N-N site (**F**, Figure 1) was recently explored by Ong, Frenking, Zhao, and coworkers.³⁹ In this work, we report our exploration of how the nature of the element in the N-N cleft influences the coordination environment about the transition metal in the pincer cleft, along with the electrochemical and spectroscopic properties of the molecule as a whole. This synthetic solution for the remote modulation of the properties of a transition metal pincer complex may provide new way for exploring and tuning catalytic activity in the future.





Figure 1. Literature examples of Janus binucleating pincer (**A-D**) and “nacnac” (**E**) ligands, Ong’s compound **F**, and the target of investigation in this work.



Results and discussion.

Synthesis of the Janus ligand. Kamble et al. recently reported a convenient synthesis of benzimidazoles by a thermal or microwave reaction of acid amides with diaminobenzenes in aqueous hydrochloric acid.⁴⁰ Substituting hydrochloric acid with 85% H₃PO₄ allowed us to carry out an analogous synthesis of bis(3-benzimidazolyl)methane (**1**) at a higher temperature without a need for a microwave reactor. We were able to isolate **1** in near-quantitative yield. Treatment of **1** with 2.4 equiv of ClPPrⁱ₂ and 2.4 equiv. of Et₃N in THF overnight at ambient temperature resulted in the formation of the proto-ligand **2**, (Scheme 1) which was isolated in excellent yield upon workup. Compound **2** was characterized by ¹H, ¹³C, and ³¹P NMR spectroscopy, with a singlet resonance at 68.2 ppm in the ³¹P{¹H} NMR spectrum. The protons of the methylene linkage give rise to a singlet at δ 5.21 ppm in the ¹H NMR spectrum and a triplet (*J*_{C-P} = 17 Hz) at δ 31.8 ppm in the ¹³C{¹H} spectrum. The PCP cleft of compound **2** can be seen as a member of a class of proto-pincers with a central CH₂ moiety that connects (hetero)arene rings bearing the two side phosphine arms.⁴¹⁻⁴⁷ On the other hand, the N-N cleft of **2** is related to a number of monoanionic N-N bidentate ligands.⁴⁸



Scheme 1. Synthesis of binucleating ligand **2**.


Synthesis and spectroscopic characterization of bi- and trimetallic compounds. The metalation of the PCP pincer can be carried out with or without the addition of external base, such as Et₃N. In the absence of added Et₃N, compounds **3-Pd** and **3-Pt** were isolated from reactions with (COD)MCl₂ (M = Pd, Pt). The synthesis of **3-Pd** required a rather high temperature (180 °C in *o*-C₆H₄Cl₂) to ensure high conversion, but the Pt reaction proceeded well at 120 °C. The use of Et₃N during the metalation with



(COD)PdCl₂ resulted in the formation of **4-Pd** after 12 h at 120 °C in toluene; however, in the case of Pt, use of NEt₃ resulted in decomposition to multiple unidentified products. Compound **4-Pt** was accessed by deprotonation of isolated **3-Pt** with NEt₃ (Scheme 2). Treatment of **4-Pd** with Me₃SiI in toluene resulted in halogen exchange, allowing for the isolation of **5-Pd** in 82% yield after workup.

Complexes **4-Pd/4-Pt/5-Pd** exist as an equilibrium mixture of two tautomers, in which the proton is located either on the nitrogens of the N-N cleft (**4a-Pd/4a-Pt/5a-Pd**), or on the central carbon (**4b-Pd/4b-Pt/5b-Pd**). The different locations of H are evident from the ¹H and ¹³C NMR spectra. For example, in **4a-Pd**, this N-H proton resonates at δ 10.23 ppm (C₆D₆ solvent) as a broad singlet, while the C-H of **4b-Pd** resonates at δ 5.74 ppm as a sharper singlet. The Pd-bound carbons of the two isomers also have distinct chemical shifts (δ 68.0 ppm, **4a-Pd**; δ 48.9 ppm, **4b-Pd**, CDCl₃ solvent). In **4b-Pt**, the C-H resonance (δ 5.51 in CDCl₃) displays telltale ¹⁹⁵Pt satellites (²J_{H-Pt} = 155 Hz). The two tautomers also possess ³¹P NMR chemical shifts that are different by ca. 15-20 ppm in each pair, with the more upfield resonance belonging to the C-H tautomers (**4b-Pd/4b-Pt/5b-Pd**). The observed equilibrium ratios are solvent-dependent, with chloroform favoring the C-H tautomer to a greater degree than benzene or toluene. Compounds **4-Pt** and **5-Pd** displayed a slightly greater predilection towards the NH tautomer (**4a** or **5a**) than did **4-Pd**, in both arene solvents and CDCl₃. It should be noted that we saw no evidence for tautomerism in compounds **2** or **3**.





Scheme 2. Synthesis of mono-metallic complexes at the PCP-pincer cleft with solvent-dependent tautomer ratios and the synthesis of bi- and tri-metallic complexes.

Deprotonation of **4a/b-Pd** or **4a/b-Pt** with (Me₃Si)₂NLi led to the formation of the Li-Pd/Pt compounds **6-Pd/6-Pt**. Both **6-Pd/Pt** were isolated in high yields as bis-THF adducts, with presumed THF binding



to Li. The N-N cleft of the Janus ligand in **3** can be viewed as accepting both protons of the CH₂ group in **2** upon its metalation (and one of the chlorides from PdCl₂ or PtCl₂). Compounds **4** and **6** can be viewed as products of successive dehydrochlorination and then deprotonation of **3**. In other words, **3**, **4**, and **6** represent three different protonation states with 2, 1, or 0 protons from the original CH₂ group retained. The presence of chloride in the isolated **6-Pd** and **6-Pt** was confirmed by observing formation of one equiv. of Me₃SiCl when each was treated with two equiv. of Me₃SiI.

The Li compounds **6** served as convenient precursors for the synthesis of bi- and trimetallic compounds **7**, **8**, and **9** in decent yields via reactions with BF₃•OEt₂, ZnCl₂ (0.5 equiv), or TaCl₅, respectively. Compounds **7-9** show the maximum possible symmetry (C_{2v} for **7** and **9**, and for each pincer fragment in **8** locally; D_{2d} for **8** as a whole) in their NMR spectra at ambient temperature. The ¹H and ¹³C NMR resonances for the PCP/NN ligand backbone in these compounds are quite similar (Tables S1 and S2). Interestingly, the ³¹P NMR chemical shifts and the ³¹P-¹⁹⁵Pt coupling constant vary considerably in the **3-9** series, Table 1. These differences may be related to the perturbation of the P-Pd-P/P-Pt-P angles and the Pd/Pt-P distances as a function of the size of the moiety in the N-N cleft, as well as the different electronic effects of those moieties. The ¹¹B NMR chemical shifts in **7-Pd** and **7-Pt** of ca. 3 ppm, the ¹¹B-¹⁹F coupling constants of 29-30 Hz, and the ¹⁹F NMR chemical shift of ca. -138 ppm are similar to those observed in BODIPY and the related compounds with a BF₂ unit in an amido/enamine cleft.^{39,49-51}

Table 1. Selected NMR data for palladium and platinum compounds.

	$\delta^{13}\text{C}$ for C-M, ppm	$\delta^{31}\text{P}$, ppm	$^1J_{\text{Pt-P}}$, Hz
3-Pd ^a	70.2	110.7	-
3-Pt ^a	59.4	102.6	3077
4a-Pd	75.2	108.6	-
4b-Pd ^a	45.9	93.1	
4a-Pt	64.0	98.9	3182
4b-Pt ^a	- ^d	90.3	3048
5a-Pd ^a	68.0	114.6	
5b-Pd ^b	- ^d	97.8	
6-Pd	76.6	99.1	-
6-Pt	64.9	92.9	3214
7-Pd	64.7	124.4	-
7-Pt	55.3	115.3	3268



8-Pd	74.3	106.2	-
8-Pt	62.4	99.0	3182
9-Pd	- ^d	114.4	-
9-Pt^c	- ^d	106.8	3074

Spectra were collected in C₆D₆ unless otherwise noted. ^a Spectra collected in CDCl₃. ^b Spectra collected in C₇D₈. ^c Spectra collected in CD₂Cl₂. ^d The Pd/Pt-C resonances were not observed owing to the poor solubility of **9-Pd/Pt** and the low achievable concentration of **4b-Pt** and **5b-Pd**.

Compounds **3-8** are yellow, orange, or red, while **9-Pd** is purple, and **9-Pt** is blue. The properties of **4** and of **7-9** were further studied by UV-Vis spectroscopy. All of these compounds possess strong absorption features in the 300-450 nm range. We surmise that they correspond to the π - π^* transitions in the extended organic π -system of the ligand. The UV-Vis spectra of **4** were collected in toluene where **4a-Pd** and **4a-Pt** are the dominant species in solution that contain an extended organic π -system that is not present in **4b-Pd** and **4b-Pt**. It appears that a switch from Pd to Pt incurs a bathochromic shift for analogous compounds by about 15-20 nm. For the same metal in the PCP cleft, the order of increasing absorption energy for these features is **8** < **4** < **7** < **9**, which can be correlated with the increasing electron-withdrawing power in the series of $\frac{1}{2} \text{Zn}^{2+} < \text{H}^+ < \text{BF}_2^+ < \text{TaCl}_4^+$. In addition to these π - π^* transition features, distinct lower energy broad bands were also observed for **9-Pd** ($\lambda = 581$ nm) and **9-Pt** ($\lambda = 602$ nm). These are presumably responsible for the blue or purple colors of these compounds. We tentatively ascribe LMCT character to them, for the transition from the filled π -orbital of the ligand to an empty d-orbital at the d⁰ Ta center.






Figure 2. A. POV-Ray renditions of ORTEP plots (50% probability ellipsoids) of **3-Pd**, **7-Pt**, **8-Pd**, and **9-Pd** showing selected atom labeling. Hydrogen atoms, solvent molecules, disorder in isopropyl arms are omitted for clarity. An additional molecule of **7-Pt** in the asymmetric unit is also omitted for clarity in **7-Pt** structure. B. Highlights of selected approximate bond angles for **3-Pd**, **7-Pt**, **8-Pd**, and **9-Pd**. Arenes, isopropyl arms, hydrogen atoms, and disorder are omitted for clarity. C. Alternate view of each structure in **B** down the M-Cl axis (M = Pd or Pt) to highlight the planarity or lack thereof.

Table 2. Selected XRD-derived bond distances in compounds **3-Pd**, **7-Pt**, **8-Pd**, and **9-Pd** (M = Pd or Pt, M' = B, Zn, or Ta). that were studied in solid state by X-ray crystallography.

	3-Pd	7-Pt ^a	7-Pt ^a	8-Pd ^b	8-Pd ^b	9-Pd
M - P	2.280(2)	2.2897(12)	2.2974(13)	2.297(3)	2.296(3)	2.2862(10)
M - P	2.271(2)	2.2962(12)	2.3098(13)	2.272(3)	2.299(3)	2.2917(10)
M - C	2.028(2)	1.971(4)	1.964(5)	2.008(10)	2.018(10)	1.991(4)



N - M'	N/A	1.561(6)	1.562(7)	1.996(10)	1.963(9)	2.107(3)
N - M'	N/A	1.572(6)	1.569(7)	1.984(8)	2.008(8)	2.104(3)

^a Data for both independent molecules of **7-Pt** in the unit cell are given. ^b Data for the two crystallographically different Pd-bound ligand frameworks in **8-Pd** are given.

Structural characterization. The structures of **3-Pd**, **7-Pt**, **8-Pd**, and **9-Pd** were determined by single-crystal X-ray diffractometry (Figure 2, Table 2). In all compounds, the geometry about Pd or Pt is approximately square-planar, and that plane is also approximately the plane of the flat, conjugated organic framework. The H₂Cl unit in **3-Pd** and the boron atom in **7-Pt** are also positioned in that plane. The structure of **3-Pd** in the N-N cleft is that of two N-H bonds, with these protons hydrogen-bonded to the chloride. The coordination geometry about boron in **7-Pt** closely mimics BODIPY and other related compounds where a monoanionic, bidentate N-N ligand is bound to the BF₂ unit.^{49,51} The Zn atom in **8-Pd** is approximately in the plane with one “half” of the ligand framework, but is displaced by ca. 0.65 Å from the other half (shown in Figure 2C). It is likely a consequence of the need to minimize the considerable crowding brought upon by the two metallaligands binding to the same Zn atom. This also manifests itself in the twisting distortion of one the conjugated “halves”, in that the two “halves” are not strictly perpendicular to each other, and ultimately in the dissymmetry of the distorted tetrahedral environment about Zn. The coordination environment about Ta in **9-Pd** is quite close to octahedral, but Ta is displaced from the ligand plane by ca. 0.62 Å. We believe this can be partly attributed to that the hypothetical positioning of Ta “in-plane” would result in the steric clash between the pair of “in-plane” chlorides and the hydrogens *ortho* to the nitrogens. In a way, this demonstrates that although the N-N cleft is not “bulky” in the usual three-dimensional sense, it can be thought of as “2D-bulky” in its own plane. Thus, there is little steric congestion with a tetrahedral moiety bound in the N-N cleft, but an octahedral moiety must necessarily contend with this “2D-bulkiness”.

Nonetheless, the series of prepared complexes demonstrate well the ability of the ligand to accommodate a variety of sizes in the N-N cleft of the ligand. The C-C-C bond angle that is highlighted



in Figure 2B serves as a good metric to showcase the flexibility. The angle varies by almost 10° across the series of four structures with H_2Cl in **3-Pd** giving rise to the largest angle, followed by the similar values engendered by Zn in **8-Pd** and TaCl_4 in **9-Pd**, and finally the BF_2 moiety in **7-Pt** giving the smallest C-C-C bond angle. The flexing of the N-N cleft portion of the ligand can be seen to affect the geometry on the opposing side of the ligand, in the PCP portion, as illustrated by the changes in the P-M-P angle (Figure 2B). When the N-N cleft of the ligand has to contract when binding the smaller boron atom in **7-Pt**, this causes the phosphines used in the PCP cleft to be pulled back which contracts the P-M-P angle. The binding of the larger Zn (**8-Pd**) or Ta (**9-Pd**), or especially the more open N-N cleft in **3-Pd**, corresponds to larger P-M-P angles. The changes in the N-N cleft can clearly influence the PCP side without changes in the groups immediately connected to the PCP-bound transition metal. However, the spread of the P-M-P angles is only ca. 5° , and they are all well within the common range^{52,53} for the [5,5-PXP]⁵⁴ pincer complexes. The modest degree of PCP perturbation illustrates that the NN/PCP ligand system is flexible enough to adapt to the significant variation in the N-N cleft without making the binding of the transition metal on the PCP side impossible.

Table 3. Reduction potentials (V) of the events observed by cyclic voltammetry for compounds **7-9**.

Complex N	N^{2+}/N^+	N^+/N^0	N^0/N^-
7-Pd	1.18	0.25	-
7-Pt	1.03	0.11	-
8-Pd	0.92 ^a	0.02 ^b -0.22 ^b	-
8-Pt	0.82 ^a	-0.09 ^b -0.32 ^b	-
9-Pd^c	1.18	0.57	-0.87
9-Pt^c	1.07	0.47	-0.88

Cyclic voltammograms were recorded in CH_2Cl_2 at 1 mM of analyte and referenced to the Fc^+/Fc redox couple. Unless otherwise noted CVs were run with 0.1 M of $[\text{Bu}_4\text{N}][\text{PF}_6]$ as the supporting electrolyte. ^a Corresponds to a two-electron event $\text{N}^{4+}/\text{N}^{2+}$. ^b Two events corresponding to N^{2+}/N^+ and N^+/N^0 because of the presence of two ligands in the molecules. ^c CVs run with 0.1 M of $[\text{Bu}_4\text{N}][\text{BArF}_{24}]$ as the supporting electrolyte.



Electrochemistry. The electrochemical properties of compounds **4**, and **7-9** were studied by means of cyclic voltammetry (CV) experiments (Table 3). A similar picture was observed within each pair of Pd vs Pt analogs, with two quasi-reversible oxidations per ligand. The Pt compounds generally appeared easier to oxidize by ca. 0.1 V. Previously studied oxidations of conventional (POCOP)MCl or (PCP)MCl complexes were found to be irreversible.^{55,56} These considerations suggest that the oxidation events in **7-9** are primarily ligand- and not Pd- or Pt-based. DFT calculations performed on the closely related **F** (Figure 1) indicated that its HOMO is primarily ligand-based.³⁹ We previously studied the reversible oxidation of Pd and Pt complexes supported by diarylamido-based pincer ligands, which was also ligand-based, and where the Pt analogs were also easier to oxidize by several hundredths of a volt.^{57,58} The Pd-Pt difference likely reflects the greater π -interaction of the filled 5d-orbitals of Pt with the π -system of the ligand vs the 4d-orbitals of Pd, driving the energy of the ligand HOMO slightly higher. This notion is also consistent with the slightly lower energy of the LMCT transition in **9-Pt** compared to **9-Pd** (vide supra). Compounds **4-Pd** and **4-Pt** appeared to give rise to two quasi-reversible oxidations; however, the recorded CVs had additional unidentified features (See ESI). They may be owing to the presence of both the **4a** and **4b** tautomers, or to the facile loss of H⁺ after oxidation. With the BF₂ unit in the N-N cleft, the CV waves were well-defined and quasi-reversible for both **7-Pd** and **7-Pt**. The trimetallic species **8-Pd** and **8-Pt** each presented three quasi-reversible waves, one of which was a two-electron wave. We assign the two one-electron waves closer to 0 V as corresponding to the **8⁺/8** and **8²⁺/8⁺** events, with the third wave corresponding to the **8⁴⁺/8²⁺** event. Ostensibly, initial one-electron oxidation of one of the PCP/NN ligands influences the subsequent oxidation of the other PCP/NN ligand. It is interesting that the second oxidation events are not likewise differentiated. It is possible that the second oxidation affects the electron density farther away from the N-N cleft.

In the case of the Ta complexes **9-Pd** and **9-Pt**, while the first oxidation is well-behaved, the second oxidation appeared only partially reversible. It is possible that the higher potential needed for the second oxidation of **9** leads to a less stable dicationic species than in the case of **7** (or the tetracationic in the case of **8**). In addition, both **9-Pd** and **9-Pt** also displayed a quasi-reversible wave corresponding to a **9/9⁺**



event at -0.87 and -0.88 V. We tentatively ascribe it to the reduction of the Ta^V center to Ta^{IV}. These values for **9** fall in between the reduction potentials for the [TaCl₆]⁻/[TaCl₆]²⁻ (-1.11 V) and the [TaCl₅(NCMe)]⁰/[TaCl₅(NCMe)]⁻ (-0.65 V) couples, as reported by Bursten et al.⁵⁹

Two oxidations per ligand in **7-9** can be understood by using simple resonance structures, a few of which are shown in Figure 3 for **7-Pd**. The lone pair localized on the central carbon in resonance structure **II** is delocalized, primarily in the “nacnac” fashion (**I** and **III**), but also beyond, into the six-membered aromatic rings (not shown). The two successive oxidations can be thought to convert this central lone pair first into a radical, and then into an empty orbital site, with extensive delocalization. The importance of the resonance structure similar to **II** in compound **F** (Figure 1) led Ong, Frenking, Zhao, and coworkers to point out the similarity of the central carbon-metal interaction in **F** to the transition metal complexes of “carbodicarbenes”.³⁹ That similarity also exists for **7-9**. The importance of the resonance structure **II** is also related to the PCP pincer complexes featuring a central diarylcarbene, explored by the Piers^{43-45,60} and Iluc⁴⁶ groups.




Figure 3. Main resonance structures for **7-Pd**.



Conclusion.

In summary, a new binucleating ligand system has been presented. It combines a monoanionic PCP pincer on one side with a monoanionic N-N bidentate cleft on the opposite side. The PCP pincer cleft was accommodated with square-planar, d^8 Pd or Pt centers, while the N-N cleft has proven adaptable to binding such divergent set of elements as boron, zinc, or tantalum. Two N-N clefts can bind a single Zn center, resulting in trimetallic ($ZnPd_2$ or $ZnPt_2$) complexes. Structural studies showed that the ligand stays nearly flat in the coordination plane of Pd or Pt, indicating a high degree of conjugation throughout. When bound to metals (or metalloids) via both the N-N and the PCP clefts, the ligand system can undergo two quasi-reversible oxidations, ascribed primarily to the oxidation of the extended organic π -system.



Acknowledgments. We are thankful for the support of this work by the Welch Foundation (grant A-1717 to O.V.O.). We thank Ms. Ruth Ann Gholson for assistance with manuscript preparation.

Data availability statement. Details of experimental procedures, graphical NMR spectra, and details of X-ray structural determination, CV experiments, and graphic UV-Vis spectra have been included as Electronic Supplementary Information. CCDC 2351193-2351196 contain crystallographic data for **3-Pd**, **7-Pt**, **8-Pd**, and **9-Pd**. See <http://www.rsc.org/suppdata/xx/b0/b000000x/>



References


- ¹ *Organometallic Pincer Chemistry*, ed. G. van Koten and D. Milstein, Springer, Heidelberg, 2013.
- ² E. Peris and R. H. Crabtree, Key factors in pincer ligand design, *Chem. Soc. Rev.*, 2018, **47**, 1959-1968.
- ³ M. Vogt and R. Langer, The Pincer Platform Beyond Classical Coordination Patterns, *Eur. J. Inorg. Chem.*, 2020, **41**, 3885-3898.
- ⁴ Kumar, A.; Bhatti, T.; Goldman, A. S. Dehydrogenation of Alkanes and Aliphatic Groups by Pincer-Ligated Metal Complexes. *Chem. Rev.* **2017**, **117**, 12357-12384.
- ⁵ Goldman, A. S.; Roy, A. H.; Huang, Z.; Ahuja, R.; Schinski, W.; Brookhart, M. Catalytic Alkane Metathesis by Tandem Alkane Dehydrogenation-Olefin Metathesis. *Science* **2006**, **312**, 257-261.
- ⁶ Manar, K. K.; Chemg, J.; Yang, Y.; Yang, X.; Ren, P. Promising Catalytic Application by Pincer Metal Complexes: Recent Advances in Hydrogenation of Carbon-Based Molecules. *ChemCatChem* **2023**, **15**, e202300004
- ⁷ Kumar, A.; Daw, P.; Milstein, D. Homogeneous Catalysis for Sustainable Energy: Hydrogen and Methanol Economies, Fuels from Biomass, and Related Topics. *Chem. Rev.* **2022**, **122**, 385-441.
- ⁸ Gunanathan, C.; Milstein, D. Bond Activation and Catalysis by Ruthenium Pincer Complexes. *Chem. Rev.* 2014, **114**, 12024-12087.
- ⁹ Arora, V.; Narjinari, H.; Nandi, P. G.; Kumar, A. Recent advances in pincer-nickel catalyzed reactions. *Dalton Trans.* **2021**, **50**, 3394-3428.
- ¹⁰ Timpa, S. D.; Pell, C. J.; Ozerov, O. V. [A Well-Defined \(POCOP\)Rh Catalyst for the Coupling of Aryl Halides with Thiols](#), *J. Am. Chem. Soc.* **2014**, **136**, 14772-14779.
- ¹¹ Press, L. P.; Kosanovich, A. J.; McCulloch, B. J.; Ozerov, O. V. [High-Turnover Aromatic C-H Borylation Catalyzed by POCOP-type Pincer Complexes of Iridium](#), *J. Am. Chem. Soc.* **2016**, **138**, 9487-9497.




- ¹² Foley, B. J.; Bhuvanesh, N. S.; Zhou J.; Ozerov, O. V. [Combined Experimental and Computational Studies of the Mechanism of Dehydrogenative Borylation of Terminal Alkynes \(DHBTA\) Catalyzed by PNP Complexes of Iridium](#), *ACS Catal.* **2020**, *10*, 9824-9836.
- ¹³ Selander, N.; Szabó, K. J. Catalysis by Palladium Pincer Complexes. *Chem. Rev.* **2011**, *111*, 2048–2076.
- ¹⁴ Bézier, D.; Brookhart, M. Applications of PC(sp³)P Iridium Complexes in Transfer Dehydrogenation of Alkanes. *ACS Catal.* **2014**, *4*, 3411–3420.
- ¹⁵ V. Lyaskovskyy and B. de Bruin, Redox Non-Innocent Ligands: Versatile New Tools to Control Catalytic Reactions, *ACS Catal.*, 2012, **2**, 270-279.
- ¹⁶ P. J. Chirik, Preface: Forum on Redox-Active Ligands, *Inorg. Chem.*, 2011, **50**, 9737-9740.
- ¹⁷ B. de Bruin, P. Gualco and N. D. Paul, in *Ligand Design in Metal Chemistry*, ed. John Wiley & Sons, Ltd, Hoboken, NJ, 2016, pp. 176-204.
- ¹⁸ A. I. Nguyen, R. A. Zarkesh, D. C. Lacy, M. K. Thorson and A. F. Heyduk, Catalytic nitrene transfer by a zirconium(iv) redox-active ligand complex, *Chem. Sci.*, 2011, **2**, 166-169.
- ¹⁹ J. J. Davidson, J. C. DeMott, C. Douvris, C. M. Fafard, N. Bhuvanesh, C.-H. Chen, D. E. Herbert, C.-I. Lee, B. J. McCulloch, B. M. Foxman and O. V. Ozerov, Comparison of the Electronic Properties of Diarylamido-Based PNZ Pincer Ligands: Redox Activity at the Ligand and Donor Ability Toward the Metal, *Inorg. Chem.*, 2015, **54**, 2916- 2935.
- ²⁰ D. M. Beagan, N. A. Maciulis, M. Pink, V. Carta, I. J. Huerfano, C.-H. Chen, and K. G. Caulton, A Redox-Active Tetrazine-Based Pincer Ligand for the Reduction of N-Oxyanions Using a Redox-Inert Metal, *Chem. Eur. J.*, 2021, **27**, 11676-11681.
- ²¹ S. Acosta-Calle and A.J.M. Miller, Tunable and Switchable Catalysis Enabled by Cation-Controlled Gating with Crown Ether Ligands, *Acc. Chem. Res.*, 2023, **56**, 8, 971–98.



- ²² H. Valdés, J.M. Germán-Acacio, G. van Koten and D. Morales-Morales, Bimetallic complexes that merge metallocene and pincer-metal building blocks: synthesis, stereochemistry and catalytic reactivity, *Dalton Trans.*, 2022, **51**, 1724-1744.
- ²³ C.-H. Yu, C. Zhu, X. Ji, W. Hu, H. Xie, N. Bhuvanesh, L. Fang, and O. V. Ozerov, Palladium bis-pincer complexes with controlled rigidity and inter-metal distance, *Inorg. Chem. Front.*, 2020, **7**, 4357-4366.
- ²⁴ D. W. Leong, Y. Shao, N. Zhen, N. Bhuvanesh and O. V. Ozerov, A bis(PCN) palladium pincer complex with a remarkably planar 2,5-diarylpyrazine core, *Dalton Trans.*, 2024, **53**, 6520-6523.
- ²⁵ P. Steenwinkel, H. Kooijman, W. J. J. Smeets, A. L. Spek, D. M. Grove and G. van Koten, Intramolecularly Stabilized 1,4-Phenylene-Bridged Homo- and Heterodinuclear Palladium and Platinum Organometallic Complexes Containing N,C,N-Coordination Motifs; η^1 -SO₂ Coordination and Formation of an Organometallic Arenium Ion Complex with Two Pt-C σ -Bonds, *Organometallics*, 1998, **17**, 5411–5426.
- ²⁶ S. L. Jeon, D. M. Loveless, W. C. Yount and S. L. Craig, Thermodynamics of Pyridine Coordination in 1,4-Phenylene Bridged Bimetallic (Pd, Pt) Complexes Containing Two N,C,N' Motifs, 1,4-M₂-[C₆(CH₂NR₂)_{4-2,3,5,6}], *Inorg. Chem.*, 2006, **45**, 11060–11068.
- ²⁷ S. J. Loeb and G. K. H. Shimizu, Dimetallated thioether complexes as building blocks for organometallic coordination polymers and aggregates, *J. Chem. Soc. Chem. Commun.*, 1993, **18**, 1395–1397.
- ²⁸ S. J. Loeb, G. K. H. Shimizu and J. A. Wisner, Mono- versus Dipalladation of the Durene-Based Tetrathioether Ligand 1,2,4,5-(^tBuSCH₂)₄C₆H₂. Structures of [PdCl(^tBuSCH₂)₄C₆H)] and [Pd₂(^tBuSCH₂)₄C₆](MeCN)₂][BF₄]₂, *Organometallics*, 1998, **17**, 2324–2327.
- ²⁹ D. Das, P. Singh, M. Singh and A. K. Singh, Tetradentate selenium ligand as a building block for homodinuclear complexes of Pd(II) and Ru(II) having seven membered rings or bis-pincer



- coordination mode: high catalytic activity of Pd-complexes for Heck reaction, *Dalton Trans.*, 2010, **39**, 10876–10882.
- ³⁰ N. P. N. Wellala, H. T. Dong, J. A. Krause and H. Guan, Janus POCOP Pincer Complexes of Nickel, *Organometallics*, 2018, **37**, 4031–4039.
- ³¹ D. Wang, S. V. Lindeman, and A.T. Fiedler, Bimetallic Complexes Supported by a Redox-Active Ligand with Fused Pincer-Type Coordination Sites, *Inorg. Chem.*, 2015, **54**, 8744–8754.
- ³² J. I. van der Vlugt, S. Demeshko, S. Dechert and F. Meyer, Tetranuclear Co^{II}, Mn^{II}, and Cu^{II} Complexes of a Novel Binucleating Pyrazolate Ligand Preorganized for the Self-Assembly of Compact [2 × 2]-Grid Structures, *Inorg. Chem.*, 2008, **47**, 1576–1585.
- ³³ M. Li, S. K. Gupta, S. Dechert, S. Demeshko, and F. Meyer, Merging Pincer Motifs and Potential Metal–Metal Cooperativity in Cobalt Dinitrogen Chemistry: Efficient Catalytic Silylation of N₂ to N(SiMe₃)₃, *Angew. Chem. Intl. Ed.*, 2021, **60**, 14480–14487.
- ³⁴ C-W. Chan, D. M. P. Mingos, A. J. P. White and D. J. Williams, Multi-domain hydrogen-bond forming metal chelates: X-ray crystal structures of dicyclopalladated 2,3-bis[6-(2-amino-4-phenylamino-1,3,5-triazinyl)]pyrazine (H₂L)[Pd₂Br₂L] and 2,6-bis[6-(2-amino-4-phenylamino-1,3,5-triazinyl)]pyridine dichloride, *Chem. Commun.*, 1996, 81–86.
- ³⁵ a) L. Bourget-Merle, M. F. Lappert and J. R. Severn, The Chemistry of β-Diketiminatometal Complexes, *Chem. Rev.*, 2002, **102**, 3031–3065. b) P. L. Holland, Electronic Structure and Reactivity of Three-Coordinate Iron Complexes, *Acc. Chem. Res.*, 2008, **41**, 905–914. c) D. J. Mindiola, Oxidatively Induced Abstraction Reactions. A Synthetic Approach to Low-Coordinate and Reactive Early Transition Metal Complexes Containing Metal–Ligand Multiple Bonds, *Acc. Chem. Res.*, 2006, **39**, 813–821. d) D. J. Mindiola, Nacnac ... □ Are You Still There? The Evolution of β-Diketimate Complexes of Nickel, *Angew. Chem., Int. Ed.*, 2009, **48**, 6198 —6200. e) Y.-C. Tsai, The chemistry of univalent metal β-diketiminates, *Coord. Chem. Rev.*, 2012, **256**, 722–758. f) R. L. Webster, β-



- Diketiminato complexes of the first row transition metals: applications in catalysis, *Dalton Trans.*, 2017, 4483-4498.
- ³⁶ S. R. Oakley, G. Nawn, K. M. Waldie, T. D. MacInnis, B. O. Patrick and R. G. Hicks, “Nindigo”: synthesis, coordination chemistry, and properties of indigo diimines as a new class of functional bridging ligands, *Chem. Commun.*, 2010, **46**, 6753-6755.
- ³⁷ S. Fortier, O. González-del Moral, C.H. Chen, M. Pink, J. J. LeRoy, M. Murugesu, D. J. Mindiola and K. G. Caulton, Probing the redox non-innocence of dinuclear, three-coordinate Co(ii) nindigo complexes: not simply β -diketiminato variants, *Chem. Commun.*, 2012, **48**, 11082-11084.
- ³⁸ P. Mondal, F. Ehret, M. Bubrin, A. Das, S. M. Mobin, W. Kaim and G. K. Lahiri, A Diruthenium Complex of a “Nindigo” Ligand, *Inorg. Chem.* 2013, **52**, 8467–8475.
- ³⁹ C.-H. Yu, K. C. AuYeung, R. Liu, C.-H. Lee, D. Jiang, B. S. Aweke, C.-H. Wu, Y.-J. Wang, T.-H. Wang, K. V. Kong, G. P. A. Tap, W.-C. Chen, G. Frenking, L. Zhao and T.-G. Ong, Diversification of the Carbodicarbene Class by Embedding an Anionic Component in its Scaffold, *Chem. Eur. J.*, 2023, **29**, e202302886.
- ⁴⁰ P. P. Kattimani, R. R. Kamble and G. Y. Meti, Expedient synthesis of benzimidazoles using amides, *RSC Adv.*, 2015, **5**, 29447-29455.
- ⁴¹ W. Weng, S. Parkin and O. V. Ozerov, Double C–H Activation Results in Ruthenium Complexes of a Neutral PCP Ligand with a Central Carbene Moiety, *Organometallics*, 2006, **25**, 5345-5354.
- ⁴² W. Weng, C.-H. Chen, B. M. Foxman and O. V. Ozerov, Palladium Complexes of a $P_2C=$ Ligand Containing a Central Carbene Moiety, *Organometallics*, 2007, **26**, 3315-3320.
- ⁴³ M. L. Clapson, J. K. Kirkland, W. E. Piers, D. H. Ess, B. Gelfand and J.-B. Lin, Carbene Character in a Series of Neutral $PC_{\text{carbene}}P$ Cobalt(I) Complexes: Radical Carbenes versus Nucleophilic Carbenes, *Organometallics*, 2022, **41**, 235–245.



- ⁴⁴ J. D. Smith, G. Durrant, D. H. Ess, B. S. Gelfand and W. E. Piers, Carbene Character in a Series of Neutral PC_{carbene}P Cobalt(I) Complexes: Radical Carbenes versus Nucleophilic Carbenes, *Chem. Sci.*, 2020, **11**, 10705-10717.
- ⁴⁵ E. A. LaPierre, W. E. Piers and C. Gendy, Redox-state dependent activation of silanes and ammonia with reverse polarity (PC_{carbene}P)Ni complexes: electrophilic vs. nucleophilic carbenes, *Dalton Trans.*, 2018, **47**, 16789-16797.
- ⁴⁶ P. Cui and V. M. Iluc, Redox-induced umpolung of transition metal carbenes, *Chem. Sci.*, 2015, **6**, 7343-7354.
- ⁴⁷ M. R. Hoffbauer, C. C. Comanescu, B. J. Dymm and V. M. Iluc, C–H Activation Reactions of a Nucleophilic Palladium Carbene, *Organometallics* 2018, **37**, 2086–2094.
- ⁴⁸ J. Kretsch, I. Koehne, M. Lökov, I. Leito and D. Stalke, Bis(benzoxazol-2-yl)methanes Hounding NacNac: Varieties and Applications in Main Group Metal Coordination, *Eur. J. Inorg. Chem.*, 2019, **28**, 3256-3351.
- ⁴⁹ K. Perumal, J. A. Garg, O. Blacque, R. Saiganesh, S. Kabilan, K. K. Balasubramanian and K. Venkatesan, β -Iminoamine-BF₂ Complexes: Aggregation-Induced Emission and Pronounced Effects of Aliphatic Rings on Radiationless Deactivation, *Chem. Asian J.*, 2012, **7**, 2670-2677.
- ⁵⁰ B. R. Grove, S. M. Crawford, T. Lundrigan, C. F. Matta, S. Sowlati-Hashijin and A. Thompson, Synthesis and characterisation of the unsubstituted dipyrin and 4,4-dichloro-4-bora-3a,4a-diaza-s-indacene: improved synthesis and functionalisation of the simplest BODIPY framework, *Chem. Commun.*, 2013, **49**, 816-818.
- ⁵¹ S. Madhu, M. R. Rao, M. S. Shaikh and M. Ravikanth, 3,5-Diformylboron Dipyrromethenes as Fluorescent pH Sensors, *Inorg. Chem.*, 2011, **50**, 4392–4400.
- ⁵² J.-F. Gong, Y.-H. Zhang, M.-P. Song and C. Xu, New PCN and PCP Pincer Palladium(II) Complexes: Convenient Synthesis via Facile One-Pot Phosphorylation/Palladation Reaction and Structural Characterization, *Organometallics*, 2007, **26**, 6487–6492.



- ⁵³ L. González-Sebastián and D. Morales-Morales, Cross-coupling reactions catalysed by palladium pincer complexes. A review of recent advances, *J. Organomet. Chem.*, 2019, **893**, 39-51.
- ⁵⁴ Fryzuk, M. D.; MacNeil, P. A., Hybrid multidentate ligands. Tridentate amidophosphine complexes of nickel(II) and palladium(II), *J. Am. Chem. Soc.*, 1981, **103**, 3592-3593.
- ⁵⁵ N. Á. Espinosa-Jalapa, S. Hernández-Ortega, D. Morales-Morales and R. Le Lagadec, Facile synthesis of heterobimetallic compounds from the cyclopentadienyl-ruthenium moiety and group 10 POCOP pincer complexes, *J. Organomet. Chem.*, 2012, **716**, 103-109.
- ⁵⁶ S. Bonnet, M. Lutzs, A. L. Spek, G. van Koten and R. J. M. Klien Gebbink, Bimetallic η^6, η^1 SCS- and PCP-Pincer Ruthenium Palladium Complexes: Synthesis, Structure, and Catalytic Activity, *Organometallics*, 2010, **29**, 1157–1167.
- ⁵⁷ J. J. Davidson, J. C. DeMott, C. Douvris, C. M. Fafard, N. Bhuvanesh, C.-H. Chen, D. E. Herbert, C. - I. Lee, B. J. McCulloch, B. M. Foxman and O. V. Ozerov, Comparison of the Electronic Properties of Diarylamido-Based PNZ Pincer Ligands: Redox Activity at the Ligand and Donor Ability Towards the Metal, *Inorg. Chem.* 2015, **54**, 2916-2935.
- ⁵⁸ C.-H. Yu, X. Yang, X. Ji, C.-H. Wang, Q. Lai, N. Bhuvanesh and O. V. Ozerov, Redox Communication between Two Diarylamido/Bis(phosphine) (PNP)M Moieties Bridged by Ynediyl Linkers (M = Ni, Pd, Pt), *Inorg. Chem.*, 2020, **59**, 10153-10162.
- ⁵⁹ B. E. Bursten, M. R. Green, V. Katovic, J. R. Kirk and D. Lightner Jr., Electrochemistry of niobium(IV) and tantalum(IV) complexes: ligand additivity in d1 octahedral complexes, *Inorg. Chem.*, 1986, **25**, 831–834.
- ⁶⁰ D. V. Gutsulyak, W. E. Piers, J. Borau-Garcia and M. Parvez, Activation of Water, Ammonia, and Other Small Molecules by PC_{carbene}P Nickel Pincer Complexes, *J. Am. Chem. Soc.*, 2013, **135**, 11776–11779.



Data availability statement. Details of experimental procedures, graphical NMR spectra, and details of X-ray structural determination, CV experiments, and graphic UV-Vis spectra have been included as Electronic Supplementary Information. CCDC 2351193-2351196 contain crystallographic data for **3-Pd**, **7-Pt**, **8-Pd**, and **9-Pd**. See <http://www.rsc.org/suppdata/xx/b0/b000000x/>

