




Cite this: *Chem. Sci.*, 2026, 17, 2199

All publication charges for this article have been paid for by the Royal Society of Chemistry

UV-vis-NIR magnetic linear dichroism: a powerful complement to MCD for f-block electronic structure

Sydney M. Giles,  Kevin O'Neil, Ian E. Ramsier, Gina Angelo, Xin Gui  and Wesley J. Transue *

The ability to synthesize next-generation lanthanide and actinide molecular materials with designer photophysical properties rests squarely on our ability to predict, control, and measure their electronic structure. This is especially true of the crystal field (CF) interactions of the metal, which are the only interactions that can be appreciably tuned by ligand design. Herein we present ultraviolet-visible-near infrared magnetic linear dichroism (MLD) spectroscopy as an underutilized magneto-optical technique that holds immense promise in the elucidation of f-block electronic structure. We use a Pr^{III} polyoxometalate complex with pseudo-*D*_{4d} symmetry, [*n*-Bu₄N]₃[Pr{Mo₅O₁₃(OMe)₄(NO)}₂] (1·Pr), to demonstrate that acquisition of both magnetic circular dichroism (MCD) and MLD spectra allows definitive assignment of the observed CF levels through the complementary selection rules of these techniques. We provide general MCD and MLD sign patterns that can be applied to any (pseudo)-*D*_{4d} Pr^{III} complex to facilitate the assignment of fine structure. Our assignments for 1·Pr allow us to fit its transitions with a phenomenological Hamiltonian, providing insight into its CF splitting and solution geometry along with entirely experimentally-derived wavefunctions for its states without use of density functional theory or multireference computational techniques.

Received 4th August 2025
Accepted 20th November 2025

DOI: 10.1039/d5sc05890b

rsc.li/chemical-science

1 Introduction

A primary focus of contemporary f-block chemistry is the development of bespoke magnetic, optical, chiroptical, and magneto-optical properties in molecules, including ultranarrow transitions,¹ circularly polarized luminescence,² magnetochiral dichroism,³ spin–electric coupling,⁴ and more.⁵ Crystal field (CF) interactions control all of these properties. The design of next-generation f-block molecular materials requires that synthetic chemists can precisely tune CF interactions about the metal,⁶ and this importance has led to the development of several experimental techniques to characterize the CF. The most popular are ground state (GS) techniques such as magnetometry,⁷ electron paramagnetic resonance,⁸ far-infrared magnetospectroscopy,⁹ and inelastic neutron scattering.¹⁰ However, there is a complication: strong spin–orbit coupling (SOC) intrinsic to f-block elements causes interstate mixing of $|M_j\rangle$ levels between energetically proximal states.¹¹ This means correlation of GS CF measurements with excited state (ES) properties is not straightforward. It is even known that CF splitting can vary from state to state in ways that cannot be modeled with the usual one-electron CF operators.^{12–15} As f-block chemists increasingly focus on precision engineering of

ES properties, it is important to develop new methods to characterize the CF through direct ES observation.

Ultraviolet-visible-near infrared (UV-vis-NIR) magneto-optical techniques like magnetic circular dichroism (MCD) spectroscopy and its sister spectroscopy, magnetic linear dichroism (MLD), are perfectly suited to this task. They offer three main advantages: (1) the signed nature of MCD/MLD features provides greater insight into overlapping absorption bands,¹⁶ (2) their SOC-driven intensity mechanisms preferentially highlight metal-centered transitions,¹⁷ and (3) their complementary selection rules to absorption spectroscopy assist in assignments.¹⁸ MCD in particular has been used in the evaluation of lanthanide electronic structure, both on its own¹⁷ and in combination with a battery of other experimental techniques.¹¹ This is not the case for MLD, which has been almost completely unreported for lanthanide coordination complexes.^{19–22} In fact, MLD is rarely encountered outside of X-ray synchrotron experiments, and we are aware of only a few molecular examples of UV-vis-NIR MLD studies.^{22–27}

Combined acquisition of MCD and MLD spectra has the potential to provide great insight into the identities of the GS/ES levels split by the CF due to the different positive/negative intensity patterns between the two techniques. Herein, we demonstrate the power of these combined spectroscopies using an example praseodymium(III) polyoxometalate (POM) complex [ⁿBu₄N]₃[Pr{Mo₅O₁₃(OMe)₄(NO)}₂] (1·Pr) (Fig. 1a). We outline

Department of Chemistry, University of Pittsburgh, Pittsburgh, Pennsylvania, USA.
E-mail: wtransue@pitt.edu





Fig. 1 (a) A 50% thermal ellipsoid plot of $1\cdot\text{Pr}$ with the cations and hydrogens omitted for clarity. (b) The energies of all $f-f$ transitions of a free atomic Pr^{III} ion are determined primarily by interelectron repulsion and SOC.^{30,31} There are further CF splittings when Pr^{III} is in a molecule, here shown only for the ${}^3\text{H}_4$ and ${}^1\text{D}_2$ levels (CF splitting not to scale).

MCD and MLD selection rules that we have derived for D_{4d} -symmetric $4f^2$ complexes without the use of Judd–Ofelt theory, which cannot be used at low temperature and is also known to describe Pr^{III} poorly.^{28,29} These selection rules allow unambiguous assignment of the observed fine structure of $1\cdot\text{Pr}$, and fitting the average CF splitting among the states provides a calculated model that closely agrees with the observed experimental transition energies. This model yields experimentally-determined wavefunctions for the system without need for computational methods like density functional theory (DFT) or multireference computational techniques. Interpretation of these CF parameters using the angular overlap model (AOM) shows that $1\cdot\text{Pr}$ retains a very similar pseudo- D_{4d} geometry in solution as in its X-ray crystal structure. The success in modeling the electronic structure of $1\cdot\text{Pr}$ highlights the power of joint acquisition of MCD and MLD spectra to assist in future design of lanthanide and actinide optical materials.

2 Theoretical background

2.1 Magneto-optical spectroscopies

Due to the shielding of $4f$ orbitals from metal–ligand interactions and the Laporte selection rule, $f-f$ transitions are characteristically narrow in lineshape and weak in intensity.²⁹ Molecules in non-centrosymmetric point groups (like D_{4d} and D_4) can display increased $f-f$ intensities through the ‘induced electric dipole’ mechanism arising from CF-driven mixing with Laporte-allowed $f-d$, $f-g$, and charge transfer transitions.^{32,33} Through this mechanism, $f-f$ transitions that satisfy the electric dipole selection rule $\Gamma_i \otimes \Gamma_m \otimes \Gamma_f \ni A_1$ are more likely to be observed in the absorption spectrum, where Γ_i/Γ_f are the irreducible representations (irreps) of the initial/final states and Γ_m is the (possibly reducible) representation for the transition dipole moment operator. Unfortunately, the absorption spectrum does not offer many other ways to discriminate between the identities of $f-f$ transitions beyond this selection rule.

Here, dichroic spectroscopies like MCD and MLD offer a distinct advantage. MCD and MLD spectroscopies are techniques that are nominally similar to but physically distinct from their natural circular dichroism (CD) and linear dichroism (LD) counterparts. Natural CD and LD are limited to chiral and/or anisotropic materials, but the addition of a magnetic field induces MCD and MLD signals in all materials, making them more broadly useful.¹⁶ It is also worth mentioning that MLD is distinct from the Cotton–Mouton effect, in which an applied magnetic field causes molecular reorientation in solution and the development of LD. The great utility of MCD and MLD in f -block spectroscopy comes from the deep connection between these dichroic responses and the molecular symmetry.³⁴

MCD and MLD spectroscopies are closely related but differ in the polarization of light and the orientation of the external magnetic field: MCD orients the field parallel with the direction of light and uses left/right-circular polarization ($\Delta\epsilon_{\text{MCD}} = \epsilon_{\text{LCP}} - \epsilon_{\text{RCP}}$), whereas MLD orients the field perpendicular to the direction of light and uses linear polarization either parallel or perpendicular to the magnetic field direction ($\Delta\epsilon_{\text{MLD}} = \epsilon_{\parallel} - \epsilon_{\perp}$). The intensities of the MCD/MLD signals for a transition $A \rightarrow J$ are³⁵

$$\frac{\Delta\epsilon_{\text{MCD}}}{E} = \gamma \sum_{\substack{a \in A \\ j \in J}} (N_a - N_j) \left(|\langle a | \hat{m}_{-} | j \rangle|^2 - |\langle a | \hat{m}_{+} | j \rangle|^2 \right) f(E) \quad (1)$$

$$\frac{\Delta\epsilon_{\text{MLD}}}{E} = \gamma \sum_{\substack{a \in A \\ j \in J}} (N_a - N_j) \left(|\langle a | \hat{m}_{\parallel} | j \rangle|^2 - |\langle a | \hat{m}_{\perp} | j \rangle|^2 \right) f(E) \quad (2)$$

where E is energy, γ a proportionality constant, N_X the fractional population of level X , and f a lineshape function. The sums run over all thermally accessible sublevels a within GS A and all sublevels j within ES J . The response to left-circular, right-circular, parallel, and perpendicular polarizations of light are calculated from transition dipole moment operators \hat{m}_{-} , \hat{m}_{+} , \hat{m}_{\parallel} , and \hat{m}_{\perp} , respectively. Use of eqn (1) and (2) requires the ability to construct a spin Hamiltonian to describe the GS magnetic response,^{36–39} and complicated nonlinear behavior is often



encountered when the system experiences magnetic saturation.

Taylor expansion can be used to greatly simplify these equations when the system is far from magnetic saturation ($\mu_B B/k_B T \ll 1$, Boltzmann constant k_B , temperature T). At sufficiently weak fields or sufficiently high temperatures, Taylor expansion of eqn (1) predicts a linear MCD response,

$$\frac{\Delta\epsilon_{\text{MCD}}}{E} = \gamma\mu_B B \left\{ -\overline{\mathcal{A}}_1 \frac{\partial f}{\partial E} + \left(\overline{\mathcal{B}}_0 + \frac{\overline{\mathcal{C}}_0}{k_B T} \right) f \right\}. \quad (3)$$

The $\overline{\mathcal{A}}_1$, $\overline{\mathcal{B}}_0$, and $\overline{\mathcal{C}}_0$ constants are called ‘‘Faraday’’ parameters, and their signs can be inferred from symmetry in favorable point groups.³⁴ It should be noted that the $\overline{\mathcal{A}}_1$ term contributes derivative-shaped features to the spectrum, while $\overline{\mathcal{B}}_0$ and $\overline{\mathcal{C}}_0$ terms contribute typical absorption-shaped features (Fig. 2a). The relative orders of magnitude of these parameters for an f-f transition are roughly³⁵

$$\begin{aligned} \overline{\mathcal{A}}_1 : \overline{\mathcal{B}}_0 : \overline{\mathcal{C}}_0 &\sim \frac{1}{I} : \frac{1}{\Delta W} : \frac{1}{k_B T} \\ &\rightarrow 10^2 : 1 : 10^3 \quad \text{at } T = 3 \text{ K} \\ &\rightarrow 10^2 : 1 : 10 \quad \text{at } T = 300 \text{ K} \end{aligned} \quad (4)$$

where we have used values typical of 1·Pr: bandwidth at half-maximum $I = 50 \text{ cm}^{-1}$ and energy differences between states $\Delta W = 2500 \text{ cm}^{-1}$. It is always possible to use a weaker magnetic field in order to remain in a linear MCD response region, even at cryogenic temperatures.

Analogous Taylor expansion of eqn (2) gives no MLD intensity at first order. Instead, the first nonzero term is at second order in the applied field, yielding a quadratic MLD response,

$$\begin{aligned} \frac{\Delta\epsilon_{\text{MLD}}}{E} = \gamma\mu_B^2 B^2 \left\{ \overline{\mathcal{A}}_2 \left(\frac{1}{2} \frac{\partial^2 f}{\partial E^2} \right) - \left(\overline{\mathcal{B}}_1 + \frac{\overline{\mathcal{C}}_1}{k_B T} \right) \frac{\partial f}{\partial E} \right. \\ \left. + \left(\overline{\mathcal{E}}_0 + \frac{\overline{\mathcal{F}}_0}{k_B T} + \frac{\overline{\mathcal{G}}_0}{(k_B T)^2} \right) f \right\}. \end{aligned} \quad (5)$$

Eqn (5) shows that six Faraday parameters ($\overline{\mathcal{A}}_2$, $\overline{\mathcal{B}}_1$, $\overline{\mathcal{C}}_1$, $\overline{\mathcal{E}}_0$, $\overline{\mathcal{F}}_0$, and $\overline{\mathcal{G}}_0$) are required to model non-saturating MLD intensity, and features can appear as zeroth-, first-, or second-order derivatives of the absorption lineshape (Fig. 2b).³⁵ The relative orders of magnitude of these parameters for lanthanides generally vary as⁴⁰

$$\begin{aligned} \overline{\mathcal{A}}_2 : \overline{\mathcal{B}}_1 : \overline{\mathcal{C}}_1 : \overline{\mathcal{E}}_0 : \overline{\mathcal{F}}_0 : \overline{\mathcal{G}}_0 \\ \sim \frac{5}{I^2} : \frac{1}{\Delta W T} : \frac{1}{k_B T T} : \frac{1}{\Delta W^2} : \frac{1}{k_B T \Delta W} : \frac{1}{(k_B T)^2} \quad (6) \\ \rightarrow 10^4 : 10^2 : 10^5 : 1 : 10^3 : (10^6 \text{ or } 0) \text{ at } T = 3 \text{ K}. \end{aligned}$$

It has been shown that isolated doublets (or effective doublets) are unable to produce any $\overline{\mathcal{G}}_0$ MLD intensity,^{25,41} and this is consistent with our observations of $\overline{\mathcal{C}}_1$ as the dominant intensity pattern for 1·Pr (*vide infra*). As temperatures rise, any $\overline{\mathcal{C}}_1$ and $\overline{\mathcal{G}}_0$ MLD intensity should decrease as T^{-1} and T^{-2} , respectively, meaning that $\overline{\mathcal{A}}_2$ features may dominate the MLD spectra of lanthanides under warmer conditions than those explored here.

Every transition within an MCD or MLD spectrum has its own unique set of Faraday parameters associated with it. These MCD and MLD Faraday parameters can be positive, zero, or negative, and their signs vary depending on the identities and symmetries (irreps) of the initial and final states involved in the transition. We have predicted the signs of MCD $\overline{\mathcal{A}}_1$ and $\overline{\mathcal{C}}_0$ signals and MLD $\overline{\mathcal{C}}_1$ signals for a generic $4f^N$ ion using the Wigner-Eckart theorem in the D_{4d} double group,^{34,42} and these signs are summarized in Table 1 (see SI Section S3 for derivation). Our prediction of Faraday parameter signs extends previous analyses of MCD $\overline{\mathcal{A}}_1$ intensity patterns that were developed using Judd-Ofelt theory.^{17,43} Judd-Ofelt theory is a common model of f-f intensity based on mixing of $4f^N$ states with $4f^{N-1}5d^1$ and $4f^{N-1}5g^1$ states, and its most well-known form fits absorptivities using only three parameters (Ω_2 , Ω_4 , Ω_6).²⁸ Several of its core assumptions, however, break down at cryogenic temperatures and for lanthanides with low-lying $4f^{N-1}5d^1$ states (such as Pr^{III}).²⁹ Our predictions in Table 1 are thus more broadly applicable since they rely only on symmetry. Nonetheless, it is worth pointing out that deviations from these expectations can still occur if mixing between M_J levels appreciably alters the effective g values of the GS or ES away from the Landé g values, or if the geometry strays from ideal D_{4d} symmetry. Additionally, $\overline{\mathcal{G}}_0$ intensity can begin to grow as the second-lowest CF level approaches the energy of the lowest level.



Fig. 2 Observed (a) MCD and (b) MLD features for an $A \rightarrow J$ transition are typically modeled as sums of zeroth-, first-, and second-order derivative lineshapes.²⁴ These lineshapes are caused by the close energetic spacing of Zeeman-split M_J levels, their uneven Boltzmann population, and their different interactions with polarized light, among other considerations (see SI Section S6.1).

2.2 Electronic structure and molecular symmetry

Lanthanide electronic structure is usually modeled as a sum of atomic contributions and crystal field contributions: $\hat{H} = \hat{H}_{\text{atom}} + \hat{H}_{\text{CF}}$. We will describe each in turn.



Table 1 Signs of MCD ($\overline{\mathcal{A}}_1$ and $\overline{\mathcal{C}}_0$) and MLD ($\overline{\mathcal{C}}_1$) Faraday parameters from the $^3\text{H}_4$ ground state of Pr^{III} assuming perfect D_{4d} symmetry and ideal Landé g factors^a

$^{2S+1}L_J$ ES	$^3\text{H}_4$ GS M_J and irrep Γ														
	0 A_1 ^a			$\pm 1 E_3$			$\pm 2 E_2$			$\pm 3 E_1$			± 4 ^b $B_1 + B_2$		
$M_J \Gamma$	$\overline{\mathcal{A}}_1$	$\overline{\mathcal{C}}_0$	$\overline{\mathcal{C}}_1$	$\overline{\mathcal{A}}_1$	$\overline{\mathcal{C}}_0$	$\overline{\mathcal{C}}_1$	$\overline{\mathcal{A}}_1$	$\overline{\mathcal{C}}_0$	$\overline{\mathcal{C}}_1$	$\overline{\mathcal{A}}_1$	$\overline{\mathcal{C}}_0$	$\overline{\mathcal{C}}_1$	$\overline{\mathcal{A}}_1$	$\overline{\mathcal{C}}_0$	$\overline{\mathcal{C}}_1$
0 $A_{1,2}$ ^a	0	0	0	0	0	0	0	0	0	–	–	–	0	0	0
$\pm 1 E_3$	0	0	0	0	0	0	–	–	–	0	0	+	+	0	0
$\pm 2 E_2$	0	0	0	–	–	–	0	0	+	+	+	–	0	0	0
$\pm 3 E_1$	–	0	0	0	0	+	+	+	–	0	0	0	0	0	0
± 4 ^c $B_1 + B_2$	0	0	0	+	+	–	0	0	0	0	0	0	0	0	0
$\pm 5 E_1$	+	0	0	0	0	–	–	+	+	0	0	0	0	0	0
$\pm 6 E_2$	0	0	0	+	–	+	0	0	–	–	+	+	0	0	0

^a This table can be used to predict the signs of MCD or MLD Faraday coefficients for a given transition originating from the $^3\text{H}_4$ ground state. For a given initial M_J or irrep (columns), the expected sign of Faraday coefficients for a transition to the final M_J or irrep (rows) is given. ^a The $M_J = 0$ levels transform as A_1 for even J and A_2 for odd J . ^b Treatment of GS $M_J = \pm 4$ levels was performed by assuming the B_1 combination was lowest; the same results are obtained if B_2 is lowest. ^c Treatment of ES $M_J = \pm 4$ assumed the splitting between B_1 and B_2 levels was less than the linewidth.

The atomic Hamiltonian \hat{H}_{atom} characterizes the energy levels of a $4f^N$ ion in the absence of any ligands, and it takes the form

$$\hat{H}_{\text{atom}} = E_0 + \sum_k F^{(k)} \hat{f}_k + \zeta \hat{A}_{\text{SOC}} + \alpha \hat{L}^2 + \beta \hat{G}(G_2) + \gamma \hat{G}(\text{SO}(7)) + \sum_h T^{(h)} \hat{t}_h + \sum_i M^{(i)} \hat{m}_i + \sum_j P^{(j)} \hat{p}_j \quad (7)$$

This equation parametrizes the effects of interelectronic repulsion ($F^{(k)}$), SOC (ζ), configuration interaction ($\alpha, \beta, \gamma, T^{(h)}$), and more (see SI for more detailed definitions of the terms). The strongest of these effects is interelectronic repulsion. This interaction breaks the $4f^N$ levels into multiple states characterized by spin S and orbital L quantum numbers, also known as Russell–Saunders or LS coupling. Within the f block, SOC is the next strongest effect and it couples the spin and orbital angular momenta into a total angular momentum J . This splits each ^{2S+1}L state into a series of $^{2S+1}L_J$ states, which are shown for a free atomic Pr^{III} ion in Fig. 1b.

The introduction of a crystal field (CF) disrupts the spherical symmetry of an atom or atomic ion, lowering the system into one of the molecular point groups. This descent in symmetry lifts the degeneracy of the M_J levels within each $^{2S+1}L_J$ state, and the resulting splitting is typically modeled using one-electron operators as

$$\hat{H}_{\text{CF}} = \sum_{\substack{k=2,4,6 \\ -k \leq q \leq k}} B_q^{(k)} \hat{C}_q^{(k)}, \quad (8)$$

where $B_q^{(k)}$ are the CF parameters and $\hat{C}_q^{(k)}$ are the spherical tensor operators of rank k and component q (see SI Section S2.2). Here, we have used the Wybourne convention,⁴⁴ but the reader should be aware that there are multiple equivalent conventions to describe f -block CF interactions.⁴⁵ Even within a single convention, different choices of xyz axis orientations

can lead to multiple equivalent sets of $B_q^{(k)}$ values, so care must be taken when making comparisons.

Eqn (8) appears to require 27 $B_q^{(k)}$ parameters, but many $B_q^{(k)}$ are necessarily zero due to molecular symmetry. Only $\hat{C}_q^{(k)}$ (or linear combinations thereof) that transform as the totally symmetric irrep of the point group may have nonzero $B_q^{(k)}$ parameters.⁴⁶ There are only three valid (k, q) pairs in a D_{4d} -symmetric system,⁴²

$$\hat{H}_{\text{CF}} = B_0^{(2)} \hat{C}_0^{(2)} + B_0^{(4)} \hat{C}_0^{(4)} + B_0^{(6)} \hat{C}_0^{(6)}. \quad (9)$$

The pattern of M_J splitting caused by this \hat{H}_{CF} perturbation can be straightforwardly predicted using the D_{4d} double group. For a $4f^2 \text{Pr}^{\text{III}}$ ion in D_{4d} symmetry, its M_J levels are expected to split as shown in the leftmost column of Table 1; for example, the $^3\text{H}_4$ GS should split into an A_1 level ($M_J = 0$), an E_3 level (degenerate $M_J = \pm 1$ pair), an E_2 level ($M_J = \pm 2$), an E_1 level ($M_J = \pm 3$), and a $B_1 + B_2$ level (quasi-degenerate $M_J = \pm 4$ pair). This process of mixing into levels that correspond to irreps of the double group generally causes M_J to cease to be a good quantum number; however, we have tried whenever possible to continue to associate M_J values to levels according to the largest component of the wavefunction.

While the double group reveals the pattern of CF splitting, prediction of their energetic ordering requires a model. A popular heuristic uses charge density distributions of the M_J levels to predict how electrostatic repulsion of point-like ligands around the f ion will influence the energies.^{47,48} These distributions (Fig. 3a) suggest the ± 3 levels of the $^3\text{H}_4$ GS experience the least destabilization by the pseudo- D_{4d} CF of $1 \cdot \text{Pr}$. More quantitative predictions were found using the angular overlap model (AOM) to estimate the CF splitting of the GS (Fig. 3b), which agreed that the E_1 ($M_J = \pm 3$) levels are expected to lie lowest in energy for ligands with typical π interactions ($e_\pi < e_\sigma$). Twisting of the two polyoxometallate ligands away from the ideal 45° angle causes a $D_{4d} \supset D_4$ descent in symmetry and further splits the levels (Fig. 3c), so accurate assignment of MCD






Fig. 4 (a) Absorption, (b) and (c) MCD, and (d) MLD spectra of **1-Pr** are shown at various temperatures. Regions of interest are shown in blue and labeled, and an absorption spectrum of **1-La** is included for comparison. There are no data within region 4 in the room temperature spectra due to the strong absorption of the solvent.

$\overline{\mathcal{E}}_0$ intensity at low temperatures and revealed a negative $\overline{\mathcal{A}}_1$ intensity after warming to 80 K (SI Fig. S6a). Additionally, the MLD intensity was best modeled with both negative $\overline{\mathcal{E}}_1$ and $\overline{\mathcal{A}}_2$ Faraday parameters (SI Fig. S6c). The strong intensity of the MCD feature and the uniformly negative MCD and MLD Faraday parameters (Table 1) are consistent with this feature arising from an $E_1 \rightarrow A_1$ transition. This interpretation agrees with the AOM prediction of an E_1 ($M_J = \pm 3$) level lowest (Fig. 3b).

Identification of an E_1 ($M_J = \pm 3$) 3H_4 GS enabled the assignment of the remainder of the features. We will demonstrate by focusing on region 2, which shows ${}^3H_4 \rightarrow {}^1D_2$ transitions that we have labeled with Roman numerals in Fig. 5. A D_{4d} CF will split the 1D_2 state into an A_1 ($M_J = 0$) level, an E_3 ($M_J = \pm 1$) level, and an E_2 ($M_J = \pm 2$) level according to Table 1, and this trifurcation is seemingly confirmed by the observation of three transitions: I (17 085 cm^{-1}), II (16 975 cm^{-1}), and III (16 576 cm^{-1}). These transitions display two positive (I, II) and one

negative (III) MCD $\overline{\mathcal{E}}_0$ term features at 1.7 K. Warming the sample to 80 K revealed negative and positive $\overline{\mathcal{A}}_1$ intensities associated with transitions I and II, respectively. The negative $\overline{\mathcal{E}}_0$ and $\overline{\mathcal{A}}_1$ MCD intensities associated with transition III clearly implicate an A_1 level; however, the Table 1 suggests that only one positive $\overline{\mathcal{E}}_0$ MCD feature should be observed. It may thus be suspected that the small feature for transition I could be due to the MCD-forbidden $E_1 \rightarrow E_3$ transition, its intensity coming from some slight deviation from D_{4d} symmetry. MCD sign and lineshape unfortunately give no further insight. Here, the utility of simultaneous MLD analysis comes into relief. Transition II with positive MCD $\overline{\mathcal{E}}_0$ and $\overline{\mathcal{A}}_1$ intensity has negative MLD $\overline{\mathcal{E}}_1$ intensity; whereas, the weaker MCD $\overline{\mathcal{E}}_0$ feature of transition I has strong positive MLD $\overline{\mathcal{E}}_1$ intensity. These patterns of MCD and MLD intensities demand that transition II is assigned to the ${}^1D_2 E_2$ ($M_J = \pm 2$) level, and transition I is indeed due to the ${}^1D_2 E_3$ ($M_J = \pm 1$) level. Explanations of the assignments in other regions (Table 2) follow similar logic and can be found in the SI





Fig. 5 The MCD and MLD spectra of the three ${}^3\text{H}_4 \rightarrow {}^1\text{D}_2$ features in region 2 are labeled with Roman numerals I, II, and III. These features together allow for confident assignment of each transition through comparison of the \mathcal{C}_0 , \mathcal{A}_1 , and \mathcal{C}_1 intensities.

(Section S4.1). Satisfyingly, our assignments were found to match in MCD \mathcal{C}_0 sign to those from CASSCF(2,7)/RI-NEVPT2 calculations using ORCA 6.0.1 (ref. 56–62) (SI Section S5).

When performing this analysis, we want to emphasize caution in interpretation of MCD/MLD signs in the presence of saturation because MCD and MLD features are able to vary in both strength and sign as a function of field and temperature. If analyzing a cryogenic MCD spectrum collected at strong field, it is crucial to ensure that the sign of the MCD feature is the same at weak fields. For our MCD analysis, we have generally done so by estimating the derivative of MCD intensity with respect to field at zero field, $(\partial\Delta\epsilon_{\text{MCD}}/\partial B)|_{B=0}$. For our MLD analysis, it was not as easy to estimate second derivatives at weak field, so we have collected spectra at a large number of weak and intermediate field strengths to ensure no flips in sign were apparent.

3.3 Crystal field splittings

The determination of the identities of these transitions allowed us to fit the electronic structure of $1\cdot\text{Pr}$ to a D_{4d} Hamiltonian $\hat{H} = \hat{H}_{\text{atom}} + \hat{H}_{\text{CF}}$ using eqn (7) and (9). Fitting was done in a stepwise manner, gradually adding increasing numbers of off-diagonal matrix elements to the Hamiltonian (see SI Section S4.3) until we arrived at the parameters listed in Table 3. Comparison between the experimental and calculated values in Table 2 shows close agreement, generally falling within a few dozen wavenumbers of the experimental value. Most notably,

Table 2 Comparison between experimental energies (cm^{-1}) of $1\cdot\text{Pr}$ $4f^2$ levels and calculated values derived from the D_{4d} fitted parameters

Level $\Gamma (M_J)$	E_{exp}	E_{calc}	ΔE	
${}^1\text{S}_0$ $A_1 (0)$	—	47 301.2	—	
${}^3\text{P}_2$	$E_3 (\pm 1)$	22 328.0	22 332.7	+4.7
	$E_2 (\pm 2)$	22 285.0	22 255.9	−29.1
	$A_1 (0)$	22 222.9	22 261.1	+38.2
${}^1\text{I}_6$	$B_1 + B_2 (\pm 4)$	—	22 052.7	—
	$E_1 (\pm 5)$	—	21 978.1	—
	$E_1 (\pm 3)$	—	21 845.1	—
	$E_2 (\pm 2)$	—	21 604.8	—
	$E_3 (\pm 1)$	—	21 431.6	—
	$E_2 (\pm 6)$	—	21 399.0	—
	$A_1 (0)$	—	21 369.2	—
	$E_3 (\pm 1)$	21 140.7	21 121.2	−19.5
${}^3\text{P}_1$	$A_2 (0)$	—	21 084.3	—
	$A_1 (0)$	20 527.6	20 534.1	+6.5
${}^3\text{P}_0$	$E_3 (\pm 1)$	17 084.9	17 074.9	−10.0
	$E_2 (\pm 2)$	16 794.8	16 788.3	−6.5
${}^1\text{D}_2$	$A_1 (0)$	16 575.6	16 590.0	+14.4
	$A_1 (0)$	—	10 221.1	—
${}^1\text{G}_4$	$B_1 + B_2 (\pm 4)$	—	10 097.1	—
	$E_3 (\pm 1)$	—	10 047.7	—
	$E_2 (\pm 2)$	—	9 731.9	—
	$E_1 (\pm 3)$	—	9 703.4	—
${}^3\text{F}_4$	$A_1 (0)$	7136.3	7140.8	+4.5
	$B_1 + B_2 (\pm 4)$	—	7086.5	—
	$E_3 (\pm 1)$	7073.2	7067.9	−5.3
	$E_2 (\pm 2)$	6894.6	6879.3	−15.3
${}^3\text{F}_3$	$E_1 (\pm 3)$	—	6849.9	—
	$A_2 (0)$	6581.3	6552.2	−29.1
	$E_3 (\pm 1)$	6520.2	6501.9	−18.3
${}^3\text{F}_2$	$E_2 (\pm 2)$	6480.7	6497.3	+16.6
	$E_1 (\pm 3)$	—	6486.9	—
	$A_1 (0)$	5171.2	5178.6	+7.4
	$E_2 (\pm 2)$	5092.4	5119.0	+26.6
${}^3\text{H}_6$	$E_3 (\pm 1)$	5090.5	5088.2	−2.3
	$E_2 (\pm 6)$	4894.1	4886.0	−8.1
	$A_1 (0)$	—	4662.8	—
	$E_3 (\pm 1)$	—	4652.4	—
	$E_2 (\pm 2)$	4587.0	4599.8	+12.8
	$E_1 (\pm 3)$	—	4486.6	—
${}^3\text{H}_5$	$E_1 (\pm 5)$	—	4421.4	—
	$B_1 + B_2 (\pm 4)$	—	4358.5	—
	$E_1 (\pm 5)$	—	2544.9	—
	$A_2 (0)$	—	2475.9	—
	$E_3 (\pm 1)$	—	2417.8	—
${}^3\text{H}_4$	$E_2 (\pm 2)$	—	2336.7	—
	$B_1 + B_2 (\pm 4)$	—	2225.2	—
	$E_1 (\pm 3)$	—	2216.5	—
	$B_1 + B_2 (\pm 4)$	—	422.7	—
	$A_1 (0)$	—	371.9	—
	$E_3 (\pm 1)$	—	323.9	—
${}^3\text{H}_3$	$E_2 (\pm 2)$	—	142.8	—
	$E_1 (\pm 3)$	0	−5.2	−5.2

our fitting procedure yields entirely experimentally-determined wavefunctions for the states of $1\cdot\text{Pr}$. This is a significant achievement, as the wavefunctions are rich with information about the system that is valuable in the understanding and optimization of optical and magnetic properties. It is also worth emphasizing that these wavefunctions were obtained without



Table 3 Fitted parameters from D_{4d} model^a

Parameter	Atomic ^b	1·Pr
E_{avg}	10 201(40)	10 109(13)
$F^{(2)}$	71 761(211)	68 819(180)
$F^{(4)}$	51 721(558)	50 926(503)
$F^{(6)}$	33 675(629)	33 930(289)
α	24.04(5)	27.4(1.3)
β	-626(5)	-873(65)
γ	1476(238)	1343 ^c
ζ	763.025(266)	744(9)
$M^{(0)}$	1.663(61)	0.0(1.6) ^d
$P^{(2)}$	235(6)	409(149) ^d
$B_0^{(2)}$		-109(38)
$B_0^{(4)}$		-1927(111)
$B_0^{(6)}$		424(287)

^a All values in cm^{-1} and parentheses show one standard uncertainty in the final digits. ^b Values were converted from the orthogonal convention in ref. 31 using definitions in ref. 63. ^c Fixed to the indicated value taken from aqueous Pr(III).⁶⁴ ^d Ratios were fixed using the values from Carnall, which were informed from Hartree-Fock calculations: $M^{(0)} : M^{(2)} : M^{(4)}$ was taken as 1 : 0.56 : 0.31 and $P^{(2)} : P^{(4)} : P^{(6)}$ was taken as 1 : 0.5 : 0.1.⁶⁵

need for computational methods like DFT or multireference calculations.

Our best-fit parameters also provided useful information about the molecular geometry in solution. The AOM was used to correlate the $B_q^{(k)}$ CF parameters with metal-ligand interaction strengths and the geometry of the ligands about the metal.⁶⁶⁻⁶⁸ If we assume cylindrically symmetric π -type interactions for each Pr-O bond, a D_{4d} arrangement gives the following equations for $B_q^{(k)}$ parameters:

$$B_0^{(2)} = \frac{10}{7} (3 \cos^2 \theta - 1) (2e_\sigma + 3e_\pi)$$

$$B_0^{(4)} = \frac{3}{7} (35 \cos^4 \theta - 30 \cos^2 \theta + 3) (3e_\sigma + e_\pi)$$

$$B_0^{(6)} = \frac{13}{28} (231 \cos^6 \theta - 315 \cos^4 \theta + 105 \cos^2 \theta - 5) (2e_\sigma - 3e_\pi),$$
(10)

where θ is the angle from the fourfold (z) axis to the O atoms of the POM ligands, e_σ is the strength of the σ -type interaction, and e_π is the individual strength of each π -type interaction. Using an average angle of $\theta = 55^\circ$ from the crystal structure allows fitting of $e_\sigma = 421(28) \text{ cm}^{-1}$ and $e_\pi = 198(48) \text{ cm}^{-1}$. When the angle θ is allowed to vary, it is found that the best-fit $B_q^{(k)}$ values correspond to an angle of $\theta = 55.8(4)^\circ$ and strengths of $e_\sigma = 431(28) \text{ cm}^{-1}$ and $e_\pi = 211(44) \text{ cm}^{-1}$. The closeness between this fitted θ value and the crystallographic one shows that 1·Pr retains its pseudo- D_{4d} geometry in solution.

3.4 Magnetic saturation behavior

A natural question arises whether 1·Pr should truly be treated as D_{4d} in symmetry, or whether its twisting angle causes appreciable changes in electronic structure. Variable temperature variable field (VT VH) MCD analysis gives us insight. A truly D_{4d} system with its E_1 ($M_J = \pm 3$) levels lowest should experience magnetic saturation with an effective g value of $g_{\text{eff}} = 3 \times (4/5) =$

12/5 (or 2.4), where 4/5 is the Landé g value for the 3H_4 GS. Structural distortions away from ideal symmetry will cause deviations in the observed g value, and fitting a saturation curve to the \mathcal{E}_0 intensity of the 3P_0 at 1.7 K shows a decreased value of $g_{\text{eff}} = 1.79(5)$. Clearly, the average molecule of 1·Pr in solution has some degree of distortion. There are many ways that the molecule could distort in solution, but the most obvious distortion from the crystal structure is a change in the twist angle ϕ between POM ligands. Twisting to the angle in the structure ($\phi = 40^\circ$) causes a $D_{4d} \supset D_4$ descent in symmetry, mixing the $M_J = \pm 3$ levels with the $M_J = \mp 1$ levels and lowering the g_{eff} value. This twisting also splits the low-lying E_2 state into $B_1 = (|+2\rangle + |-2\rangle)/\sqrt{2}$ and $B_2 = (|+2\rangle - |-2\rangle)/\sqrt{2}$ states (Fig. 3c).

The VT VH MCD data show nesting, symptomatic of a low-lying ES (Fig. 6). We were able to model the VT VH MCD saturation curves through construction of an effective spin Hamiltonian,³⁶⁻³⁹ giving $g_{\text{eff}} = 2.34(3)$ and the presence of the low-lying $|B_1\rangle$ ES at $E = 92(4) \text{ cm}^{-1}$. This g_{eff} value could be obtained through introduction of $B_{\pm 4}^{(4)}$ and $B_{\pm 4}^{(6)}$ terms in eqn (9) (see SI Section S2.2) and would correspond to a twist angle of $37(2)^\circ$ according to the AOM (Fig. 3c). At this twisting angle, the GS would be 97% $|M_J = \pm 3\rangle$ in composition, suggesting the D_{4d} CF model is adequate for modeling the ES CF splitting of 1·Pr. This angle is also remarkably similar to the one in the crystal structure, again underscoring the power of the AOM in providing geometric insight into f-block elements in solution.

It is interesting to note that a different set of CF parameters is obtained by fitting T vs. $\chi_m T$ and B vs. M data from vibrating sample magnetometry (VSM) of 1·Pr (Fig. 7a and b) than those obtained through fitting of ES splittings from MCD/MLD spectra. Fitting the VSM data using EasySpin⁶⁹ gave $B_0^{(2)} =$




Fig. 6 Nesting of the VT VH MCD saturation curves indicate the presence of a low-lying ES. The best-fit curves from an effective spin Hamiltonian fitting procedure are indicated, and they correspond to a low-lying ES at 92(4) K.






Fig. 7 Vibrating sample magnetometry was used to measure the (a) T vs. $\chi_m T$ and (b) B vs. M response of 1-Pr. Fitting these curves gave slightly different CF parameters than fitting the ES CF splittings. Preliminary VTVH saturation curves of ${}^3\text{H}_4 \rightarrow {}^1\text{D}_2$ MLD intensity for the ${}^3\text{H}_4 \rightarrow {}^1\text{D}_2$ feature are also shown for two of its transitions (c) and (d).

-393 , $B_0^{(4)} = -1373$, $B_0^{(6)} = -1154$, $B_{\pm 4}^{(4)} = -344$, and $B_{\pm 4}^{(6)} = -1054 \text{ cm}^{-1}$. This set of $B_0^{(k)}$ CF parameters is similar in sign and magnitude to those in Table 3, but it deviates more than three standard uncertainties from the ES MCD/MLD fit. The differences between GS and ES fits highlight an important albeit inconvenient fact of f-block CF splittings: CF parameters are known to vary from state to state.^{12–15} If one is interested in the ES photophysical properties of a molecule, joint MCD/MLD analysis of ES splittings can be expected to provide more useful insight into CF interactions. If instead one is interested in the magnetic response of a molecule, GS magnetometry through VTVH MCD saturation curves, VSM, or SQUID (superconducting quantum interference device) measurements are likely to be more relevant.

Despite the differences between ES and GS fits, the two GS magnetometry techniques agree closely: the VSM CF parameters predict the lowest-lying excited state to be at $E = 89 \text{ cm}^{-1}$, which is within one standard uncertainty of our effective Hamiltonian treatment of VTVH MCD saturation curves. More accurate insight would have to come from higher resolution techniques than magnetometry, such as far-infrared magnetospectroscopy or high-field electron paramagnetic resonance spectroscopy. For any interested readers, we also include two plots of VTVH MLD data for the ${}^3\text{H}_4 \rightarrow {}^1\text{D}_2$ transition (Fig. 7c and d). We have not incorporated these data in our fitting routines, but VTVH MLD measurements clearly hold promise for future studies of MLD magnetometry.

4 Conclusions

This work has demonstrated the power of UV-vis-NIR MLD spectroscopy as a complement to MCD spectroscopy for studying f-block electronic structure. Together, these two magneto-optical techniques have provided conclusive assignments of much of the fine structure observed for 1-Pr and allowed fitting of the transition energies with the Hamiltonian $\hat{H} = \hat{H}_{\text{atom}} + \hat{H}_{\text{CF}}$. Our fit gave detailed structural information about the geometry of 1-Pr in solution and, perhaps most excitingly, yielded wavefunctions derived strictly from experimental observables. Because such $4f^N$ wavefunctions determine the molecular optical and magnetic properties, experimentally derived wavefunctions are rich with information that is valuable in future tuning of f-block molecular materials and nanomaterials.⁷⁰

We have also highlighted the deep relation between molecular symmetry and these magneto-optical spectroscopies. Symmetry is known to control many desirable properties like circularly polarized luminescence (CPL),⁷¹ magnetic CPL,^{34,72} magnetochiral dichroism,⁷³ spin–electric coupling,⁷⁴ and ultra-narrow optical transitions;⁷⁵ thus, the utility of symmetry-based insight into electronic structure from joint MCD–MLD analysis is difficult to overstate. Our lab is continuing to explore the implementation of MLD spectroscopy in the understanding of f-block complexes, both moving downwards into the actinides and rightwards into ions with higher f^N counts, especially Kramers systems. Many research groups with MCD spectroscopy instrumentation may already be equipped to acquire MLD spectra, and we hope this work encourages broader adoption of this information-dense technique.

Author contributions

WJT conceived the project. IER and KO synthesized 1-Pr, and KO developed its final preparative route. GA and XG collected vibrating sample magnetometry data. SMG collected most of the MCD, MLD, and absorption spectra. WJT and SMG performed all analysis and wrote the manuscript.

Conflicts of interest

There are no conflicts of interest to declare.

Data availability

The absorption, MCD, and MLD data supporting this article have been included as part of the supplementary information (SI). Supplementary information is available. See DOI: <https://doi.org/10.1039/d5sc05890b>.

CCDC 2457433 contains the supplementary crystallographic data for this paper.⁷⁶

Acknowledgements

Thanks to Carolina Fernandes for collection of the crystal structure of 1-Pr, and to Dr James Peterson for helpful suggestions in the practical acquisition of MLD data. WJT



thanks the University of Pittsburgh, the Spectroscopy Society of Pittsburgh, and the Pitt Momentum Funds for funding, and SMG thanks funding by OUSD/R&E (The Under Secretary of Defense-Research and Engineering), National Defense Education Program (NDEP)/BA-1, Basic Research in the form of a SMART scholarship. This research was supported in part by the University of Pittsburgh Center for Research Computing and Data (RRID:SCR_022735) through the resources provided. Specifically, this work used the H2P cluster, which is supported by NSF award number OAC-2117681. The translating magnet stand used to support the MCD/MLD magnet was constructed by the University of Pittsburgh Dietrich School Machine Shop Core Facility. Work performed in the University of Pittsburgh Dietrich School Machine Shop Core Facility (RRID:SCR_023720) and services and instruments used in this project were graciously supported, in part, by the University of Pittsburgh.

References

- D. Serrano, S. K. Kuppusamy, B. Heinrich, O. Fuhr, D. Hunger, M. Ruben and P. Goldner, *Nature*, 2022, **603**, 241–246.
- F. Zinna and L. D. Bari, *Chirality*, 2015, **27**, 1–13.
- K. Taniguchi, M. Nishio, S. Kishiue, P.-J. Huang, S. Kimura and H. Miyasaka, *Phys. Rev. Mater.*, 2019, **3**, 045202.
- J. Liu, J. Mrozek, A. Ullah, Y. Duan, J. J. Baldoví, E. Coronado, A. Gaita-Ariño and A. Ardavan, Quantum coherent spin–electric control in a molecular nanomagnet at clock transitions, *Nat. Phys.*, 2021, **17**(11), 1205–1209.
- N. Mavragani, D. A. Gálico, A. A. Kitos and M. Murugesu, *J. Am. Chem. Soc.*, 2025, **147**, 1387–1391.
- C.-K. Duan and P. A. Tanner, *J. Phys. Chem. A*, 2010, **114**, 6055–6062.
- J.-L. Liu, Y.-C. Chen and M.-L. Tong, *Chem. Soc. Rev.*, 2018, **47**, 2431–2453.
- J. E. McPeak, S. S. Eaton and G. R. Eaton, in *Methods in Enzymology*, ed. J. A. Cotruvo, Academic Press, 2021, vol. 651 of Rare-Earth Element Biochemistry: Characterization and Applications of Lanthanide-Binding Biomolecules, pp. 63–101.
- J. G. C. Kragoskow, J. Marbey, C. D. Buch, J. Nehr Korn, M. Ozerov, S. Piligkos, S. Hill and N. F. Chilton, *Nat. Commun.*, 2022, **13**, 825.
- M. A. Dunstan, R. A. Mole and C. Boskovic, *Eur. J. Inorg. Chem.*, 2019, **2019**, 1090–1105.
- M. Slota, S.-D. Jiang, E. Heintze, Y. Rechkemmer, M. Dressel, J. van Slageren and L. Bogani, *Phys. Rev. B*, 2019, **99**, 134410.
- B. R. Judd, *Phys. Rev. Lett.*, 1977, **39**, 242–244.
- B. R. Judd, *J. Chem. Phys.*, 1977, **66**, 3163–3170.
- M. F. Reid, *J. Chem. Phys.*, 1987, **87**, 2875–2884.
- C. K. Jayasankar, F. S. Richardson and M. F. Reid, *J. Less-Common Met.*, 1989, **148**, 289–296.
- J. Peterson and E. L. Bominaar, *Anal. Chem.*, 2002, **74**, 527A–533A.
- C. Görrler-Walrand and L. Fluyt, in *Handbook on the Physics and Chemistry of Rare Earths*, Elsevier, 2010, vol. 40, pp. 1–107.
- J. C. Deaton, M. S. Gebhard and E. I. Solomon, *Inorg. Chem.*, 1989, **28**, 877–889.
- Other than ref. 20–22, the only other molecular lanthanide MLD spectra that we know of are unpublished results shown in Chapter 9 of the textbook ‘A Practical Guide to Magnetic Circular Dichroism Spectroscopy’ by Dr W. R. Mason; see ref. 35.
- B. E. Argyle, J. C. Suits and M. J. Freiser, *Phys. Rev. Lett.*, 1965, **15**, 822–825.
- A. C. Boccara and N. Moreau, *Phys. Status Solidi B*, 1971, **45**, 573–581.
- A. C. Boccara, J. Ferre, B. Briat, M. Billardon and J. P. Badoz, *J. Chem. Phys.*, 1969, **50**, 2716–2718.
- J. Rivoal, B. Briat and M. Vala, *Mol. Phys.*, 1979, **38**, 1829–1854.
- J. Peterson, L. L. Pearce and E. L. Bominaar, *J. Am. Chem. Soc.*, 1999, **121**, 5972–5980.
- E. L. Bominaar and J. Peterson, *J. Chem. Phys.*, 1999, **111**, 7512–7518.
- L. L. Pearce, E. L. Bominaar and J. Peterson, *Biochem. Biophys. Res. Commun.*, 2002, **297**, 220–223.
- J. Peterson, T. J. Collins, E. Münck and E. L. Bominaar, *Chem. Phys. Lett.*, 2002, **365**, 164–169.
- M. P. Hehlen, M. G. Brik and K. W. Krämer, *J. Lumin.*, 2013, **136**, 221–239.
- C. Görrler-Walrand and K. Binnemans, in *Handbook on the Physics and Chemistry of Rare Earths*, Elsevier, 1998, vol. 25, pp. 101–264.
- J. Sugar, *J. Opt. Soc. Am.*, 1965, **55**, 1058–1059.
- J.-F. Wyart, A. Meftah, J. Sinzelle, W.-Ü. L. Tchchang-Brillet, N. Spector and B. R. Judd, *J. Phys. B: At., Mol. Opt. Phys.*, 2008, **41**, 085001.
- G. S. Ofelt, *J. Chem. Phys.*, 1962, **37**, 511–520.
- B. R. Judd, *Phys. Rev.*, 1962, **127**, 750–761.
- S. B. Piepho and P. N. Schatz, *Group Theory in Spectroscopy with Applications to Magnetic Circular Dichroism*, John Wiley & Sons, New York, 1983.
- W. R. Mason, *A Practical Guide to Magnetic Circular Dichroism Spectroscopy*, John Wiley & Sons, Inc., Hoboken, NJ, 1st edn, 2007.
- F. Neese and E. I. Solomon, *Inorg. Chem.*, 1999, **38**, 1847–1865.
- V. S. Oganessian, S. J. George, M. R. Cheesman and A. J. Thomson, *J. Chem. Phys.*, 1999, **110**, 762–777.
- V. S. Oganessian and A. J. Thomson, *J. Chem. Phys.*, 2000, **113**, 5003–5017.
- W. J. Transue, R. A. Snyder, J. D. Caranto, D. M. J. Kurtz and E. I. Solomon, *Inorg. Chem.*, 2022, **61**, 16520–16527.
- M. Kreglewski and M. Vala, *J. Chem. Phys.*, 1981, **74**, 5411–5419.
- E. L. Bominaar, C. Achim and J. Peterson, *J. Chem. Phys.*, 1998, **109**, 942–950.
- S. L. Altmann and P. Herzog, *Point-Group Theory Tables*, University of Vienna PHAIDRA, Wien, 2nd edn, 2011.
- C. Görrler-Walrand and L. Fluyt-Adriaens, *J. Less-Common Met.*, 1985, **112**, 175–191.



- 44 B. G. Wybourne, *Spectroscopic Properties of the Rare Earths*, John Wiley & Sons (Interscience Publishers), New York, 1965.
- 45 O. Duros, A. Juhin, H. Elnaggar, G. S. Chiuzbăian and C. Brouder, *J. Phys. A: Math. Theor.*, 2024, **58**, 025207.
- 46 G. Liu and B. Jacquier, *Spectroscopic Properties of the Rare Earths in Optical Materials*, Springer-Verlag, 2005.
- 47 J. Sievers, *Z. Phys. B: Condens. Matter*, 1982, **45**, 289–296.
- 48 J. D. Rinehart and J. R. Long, *Chem. Sci.*, 2011, **2**, 2078–2085.
- 49 C. M. Granadeiro, D. Julião, S. O. Ribeiro, L. Cunha-Silva and S. S. Balula, *Coord. Chem. Rev.*, 2023, **476**, 214914.
- 50 S. Giménez-Santamarina, S. Cardona-Serra, J. M. Clemente-Juan, A. Gaita-Ariño and E. Coronado, *Chem. Sci.*, 2020, **11**, 10718–10728.
- 51 Z.-X. Yang, F. Gong, D. Lin and Y. Huo, *Coord. Chem. Rev.*, 2023, **492**, 215205.
- 52 S. She, C. Gao, K. Chen, A. Bayaguud, Y. Huang, B.-W. Wang, S. Gao and Y. Wei, *Inorg. Chem.*, 2018, **57**, 963–969.
- 53 P. Gouzerh, Y. Jeannin, A. Proust and F. Robert, *Angew. Chem., Int. Ed. Engl.*, 1989, **28**, 1363–1364.
- 54 A. Proust, P. Gouzerh and F. Robert, *Inorg. Chem.*, 1993, **32**, 5291–5298.
- 55 G. H. Dieke and H. M. Crosswhite, *Appl. Opt.*, 1963, **2**, 675–686.
- 56 F. Neese, *Wiley Interdiscip. Rev.: Comput. Mol. Sci.*, 2018, **8**, e1327.
- 57 D. Ganyushin and F. Neese, *J. Chem. Phys.*, 2006, **125**, 024103.
- 58 G. L. Stoychev, A. A. Auer and F. Neese, *J. Chem. Theory Comput.*, 2017, **13**, 554–562.
- 59 C. Kollmar, K. Sivalingam, B. Helmich-Paris, C. Angeli and F. Neese, *J. Comput. Chem.*, 2019, **40**, 1463–1470.
- 60 L. Lang and F. Neese, *J. Chem. Phys.*, 2019, **150**, 104104.
- 61 C. Kollmar, K. Sivalingam, Y. Guo and F. Neese, *J. Chem. Phys.*, 2021, **155**, 234104.
- 62 Y. Guo, K. Sivalingam and F. Neese, *J. Chem. Phys.*, 2021, **154**, 214111.
- 63 B. R. Judd and H. Crosswhite, *J. Opt. Soc. Am. B*, 1984, **1**, 255–260.
- 64 W. T. Carnall, in *Handbook on the Physics and Chemistry of Rare Earths*, Elsevier, 1979, vol. 3 of Non-Metallic Compounds - I, pp. 171–208.
- 65 W. T. Carnall, *J. Chem. Phys.*, 1992, **96**, 8713–8726.
- 66 M. Suta, F. Cimpoesu and W. Urland, *Coord. Chem. Rev.*, 2021, **441**, 213981.
- 67 H. Ramanantoanina, W. Urland, F. Cimpoesu and C. Daul, *Phys. Chem. Chem. Phys.*, 2014, **16**, 12282–12290.
- 68 D. J. Newman and B. Ng, *Rep. Prog. Phys.*, 1989, **52**, 699.
- 69 S. Stoll and A. Schweiger, *J. Magn. Reson.*, 2006, **178**, 42–55.
- 70 Y. Luo, Z. Chen, S. Wen, Q. Han, L. Fu, L. Yan, D. Jin, J.-C. G. Bünzli and G. Bao, *Coord. Chem. Rev.*, 2022, **469**, 214653.
- 71 D. Thakur and S. Vaidyanathan, *J. Mater. Chem. C*, 2025, **13**, 9410–9452.
- 72 J. P. Riehl and F. S. Richardson, *J. Chem. Phys.*, 1977, **66**, 1988–1998.
- 73 L. Cahya Adi, O. G. Willis, A. Gabbani, G. L. J. A. Rikken, L. Di Bari, C. Train, F. Pineider, F. Zinna and M. Atzori, *Angew. Chem., Int. Ed.*, 2024, **63**, e202412521.
- 74 W. T. Morrillo, H. I. J. Cumming, A. Mattioni, J. K. Staab and N. F. Chilton, *J. Am. Chem. Soc.*, 2024, **146**, 25841–25851.
- 75 B. Y. Li, C. E. Dickerson, A. J. Shin, C. Zhao, Y. Shen, Y. He, P. L. Diaconescu, A. N. Alexandrova and J. R. Caram, *Chem. Sci.*, 2024, **15**, 12451–12458.
- 76 CCDC 2457433: Experimental Crystal Structure Determination, 2025, DOI: [10.5517/ccdc.csd.cc2nh51d](https://doi.org/10.5517/ccdc.csd.cc2nh51d).

