



ChemComm

Effect of copper salts on hydrothermal oxidative decarboxylation: A study of phenylacetic acid

Journal:	<i>ChemComm</i>
Manuscript ID	CC-COM-12-2019-009825.R2
Article Type:	Communication

SCHOLARONE™
Manuscripts

COMMUNICATION

Effect of copper salts on hydrothermal oxidative decarboxylation: A study of phenylacetic acid

Received 00th January 20xx,
Accepted 00th January 20xx

Xuan Fu,^a Megan Jamison,^a Aaron M. Jubb,^b Yiju Liao,^a Alexandria Aspin,^a Kyle Hayes,^a Christopher R. Glein,^c and Ziming Yang ^{*a}

DOI: 10.1039/x0xx00000x

Decarboxylation of carboxylic acids is favored under hydrothermal conditions, and can be influenced by dissolved metals. Here, we use phenylacetic acid as a model compound to study its hydrothermal decarboxylation in the presence of copper (II) salts but no O₂. Our results showed a strong oxidizing role of copper in facilitating oxidative decarboxylation.

Organic compound transformations in hydrothermal systems are of great importance to a wide range of chemical and geological processes, including degradation of sedimentary organic matter, metabolism in the subsurface biosphere, and the deep carbon cycle.¹⁻⁴ Unlike at ambient conditions, water at elevated temperatures and pressures assumes key properties that promote organic reactions. For example, hydrothermal water has a considerably lower dielectric constant but a higher dissociation constant (K_w) than ambient water, making it not only a good solvent for organic molecules but also providing Brønsted acid and base catalysts in the form of increased concentrations of H⁺ and OH⁻.^{5, 6} The chemistry of a variety of organic functional groups has been examined under hydrothermal conditions, and their reaction pathways and mechanisms have been elucidated.⁷⁻¹² In particular, reversible and irreversible interconversions between organic functional groups have been proposed in sedimentary basins,¹ and experimental data were collected showing the interconversions between alkanes, alkenes, alcohols, aldehydes/ketones, and carboxylic acids.^{7, 13} However, the hydrothermal chemistry of organic compounds can be influenced by the surrounding rocks and minerals, and water-rock interactions can release dissolved metal species that potentially dominate hydrothermal organic reactivity. Transformations of organic molecules could be promoted by the presence of minerals, and indeed some remarkable hydrothermal redox reactions have been observed.¹⁴⁻¹⁷ For example, reduction of

ketones to alcohols can be facilitated by iron sulfide minerals,¹⁵ oxidation of alcohols to aldehydes can be efficient in the presence of copper (II),¹⁴ and selective reduction of alkenes to alkanes is feasible using Fe/Ni under hydrothermal conditions.¹⁶

Organic-inorganic hydrothermal reactions are also of growing interest to green chemistry. The concept of “geomimicry” has been proposed, which applies relatively benign and Earth-abundant materials (e.g., iron, copper, zinc) to organic synthesis, rather than the use of toxic or rare metal catalysts.^{14, 16} For example, copper salts have been used as effective catalysts for organic oxidation in green chemistry,^{18, 19} but the reactions usually require O₂ to be present as the oxidant.^{20, 21} In a previous study, we found that under hydrothermal conditions copper (II) chloride was an efficient oxidant for alcohols, even in the absence of O₂.¹⁴ The oxidizing role of copper could create novel redox pathways for organic compounds in both natural and human-made hydrothermal systems. For example, deep-sea mining indicates that copper-bearing minerals were deposited at the seafloor from hydrothermal fluids,^{22, 23} and these minerals could release dissolved copper species that interact with organic carbon. Here we investigate the scope of hydrothermal organic oxidation using a suite of copper (II) salts, and we describe results that compare effects of different cupric salts in oxidative decarboxylation of carboxylic acids. Carboxylic acids are known to undergo decarboxylation under hydrothermal conditions,^{12, 24, 25} but studies of hydrothermal interactions between carboxylic acids and inorganic species are few, and in particular, the effect of dissolved metal ions on acid decarboxylation is largely unknown. In this study, we use phenylacetic acid (PAA) as a model compound to examine the effects of cupric salts, based on its known hydrothermal chemistry and predictable decarboxylation pathways.^{12, 14} Results from this work may provide new insights into the roles of dissolved metal species in organic transformations under hydrothermal conditions.

Hydrothermal experiments were performed in fused silica glass tubes using a previously developed method.²⁶ In short, 0.03 mmol of PAA was weighed and carefully transferred into the tube through a Pasteur pipette. A 0.3 mL aliquot of metal salts solution (0.2 molal in

^a Department of Chemistry, Oakland University, Rochester, MI 48309, USA. E-mail: zimingyang@oakland.edu

^b U.S. Geological Survey, 12201 Sunrise Valley Drive, Reston, VA 20192, USA.

^c Space Science and Engineering Division, Southwest Research Institute, San Antonio, TX 78228, USA.

† Electronic Supplementary Information (ESI) available: Detailed experimental setups and experimental materials. See DOI: 10.1039/x0xx00000x





Fig. 1 Hydrothermal decarboxylation of phenylacetic acid (PAA) in the presence of copper or sodium salts at 200 °C and 15 bar (P_{sat}) for up to 6 hours.

degassed and deionized water) was subsequently added using a gas-tight micro-syringe, to reach a molar ratio of 1:2 between PAA and the divalent metal ions. The 1:2 ratio was used to ensure enough organic-metal interactions but also to avoid high metal concentrations that may exceed the “geomimicry” concept. In sodium salts experiments, the anions concentrations were kept the same as those in copper salts experiments, with a molar ratio of 1:4 between PAA and the sodium ions, except for sodium sulfate (1:2). The headspace air in silica tubes was evacuated through three pump-freeze-thaw cycles and the tubes were sealed with an oxyhydrogen flame under vacuum. The sealed tubes were then heated in a gas chromatography (GC) oven at 200 °C for up to 6 hours. The pressure inside the tube was calculated to be ca. 15 bar (P_{sat} at 200 °C) based on thermodynamics calculations using SUPCRT92.²⁷

Experiments were conducted under a hydrothermal condition (200 °C and 15 bar) in the absence of O_2 , and the effects of four copper (II) salts including $CuCl_2$, copper sulfate ($CuSO_4$), copper acetate [$Cu(OAc)_2$], and copper nitrate [$Cu(NO_3)_2$] were compared. At 200 °C, PAA reacted very slowly in pure hydrothermal water, with a < 2% decomposition after 6 hours. In the presence of the copper salts, however, more than 40% of the acid was degraded after 6 hours (Fig. 1). Time-series experiments showed that the decomposition of PAA mostly occurred in the first 2 hours and then reached a steady state (Fig. 1). Among the four copper (II) salts with the same concentration of copper (i.e., 0.2 molal), the trend in

decomposing PAA followed as copper nitrate \gg copper chloride \approx copper acetate $>$ copper sulfate, with copper nitrate exhibiting the most significant effect (e.g., ~70% acid decomposition after 2 hours). Experiments with a lower molar ratio of 1:4 between PAA and copper (II) were also conducted, in which even higher decomposition of PAA was observed when larger amounts of copper (II) were used (Table S1). In contrast, hydrothermal experiments with sodium salts with the same counterions showed negligible PAA decomposition (all below 2% after 4 hours) (Fig. 1), suggesting the reactions were not driven by the anions or nitrate alone. These results demonstrate different effects of copper salts on carboxylic acid decomposition, which could be attributed to the formation of different copper (II) complexes under the hydrothermal condition (see below).

Copper (II)-facilitated PAA hydrothermal transformation was proposed to undergo a series of oxidative steps starting with oxidative decarboxylation to form benzyl alcohol, then benzaldehyde, and benzoic acid (Scheme 1).



Scheme 1. Proposed hydrothermal oxidative decarboxylation pathway from phenylacetic acid (PAA) to benzyl alcohol, benzaldehyde, and benzoic acid in the presence of copper (II).

In these steps, copper ions were proposed to serve as a reactive oxidant to promote acid decarboxylation and the following oxidation of primary products. UV-Vis spectroscopy analysis showed that the dissolved copper (II) was mostly consumed after the reaction (Fig. S3), which verifies the oxidative role of copper (II) in this reaction scheme. Benzaldehyde and benzoic acid were formed as the major products of PAA, but their distributions were dependent on the type of copper (II) salts. As shown in Fig. 2, all copper experiments showed trends of increasing concentrations of benzaldehyde and benzoic acid over time, except for copper nitrate which showed a decreasing concentration of benzaldehyde that implies a second derivative after 0.5 hours. To compare the oxidizing effects between the copper salts, the ratio of benzoic acid to benzaldehyde was calculated. Fig. 2c shows that the acid:aldehyde ratios were between 0.3 and 2.3 in the presence of copper chloride/sulfate/acetate, whereas the ratio in the copper nitrate experiment substantially increased to a value of 47.6 after 6 hours. This high acid:aldehyde ratio indicates that copper




Fig. 2 Concentrations (mmolal) of major product benzaldehyde (a) and benzoic acid (b), and ratios of benzoic acid/benzaldehyde vs. time in hydrothermal experiments of PAA with copper chloride (blue circles), copper sulfate (orange squares), copper acetate (green diamonds), and copper nitrate (red triangles) at 200 °C and 15 bar (P_{sat}).

(II) nitrate was a much stronger oxidant than the other copper salts in hydrothermal transformation of carboxylic acids.

Because sodium salts did not show any oxidizing effects under the same hydrothermal condition (Fig. 1), we hypothesize that the main driver of oxidative decarboxylation could be complexes between copper (II) and the anions, rather than freely solvated copper (II). Indeed, thermodynamic calculations on copper speciation under similar hydrothermal conditions (e.g., 250 °C and 40 bar) suggested that the dominant forms of aqueous copper species were copper complexes (e.g., with chloride or acetate) instead of uncomplexed copper (II).¹⁴ To investigate the effect of the nitrate complex, other divalent but non-oxidizing metal ions including magnesium (Mg^{2+}), zinc (Zn^{2+}), and nickel (Ni^{2+}) were studied in the form of nitrate salts to react with PAA at 200 °C for 4 hours, respectively. Compared to sodium nitrate, magnesium nitrate, zinc nitrate, and nickel nitrate all increased the PAA conversion by 10–15%, with benzaldehyde and benzoic acid formed as the major products (Fig. 3). Since these metals are non-oxidizing, forming the oxidation products implies a potential oxidizing role of divalent metal-nitrate complexes in the hydrothermal system. Interestingly, all three metal nitrates yielded more benzoic acid than benzaldehyde (acid/aldehyde ratio between 3.6 and 4.5), indicating the oxidation step from benzaldehyde to benzoic acid was promoted by the presence of nitrate. However, the PAA decomposition with non-oxidizing metal nitrates was not comparable to that with copper nitrate (Fig. 3), suggesting that the oxidative decarboxylation was mainly driven by the copper (II) complexes, whereas the oxidation of benzaldehyde was facilitated by the nitrate (Scheme 1).

To further study the copper complexation, Raman spectroscopy analysis was also performed for hydrothermal experiments with PAA and with PAA + $CuCl_2$, respectively. Raman spectra were collected every 30 minutes during a 2-hour hydrothermal reaction of PAA at 200 °C, and the peaks were assigned according to Lin-Vien et al (1991).²⁸ No obvious changes were observed in the PAA alone experiment (Fig. 4a), which is consistent with the low reactivity of PAA under the hydrothermal condition. In the presence of $CuCl_2$, however, several significant changes are observed in the spectra (Fig. 4b). The C=O group became more abundant through the hydrothermal process, which is consistent with the production of benzaldehyde and benzoic acid from PAA. The intensity of a Raman response is equal to the product of the input laser energy, the number of oscillators, and the Raman cross-section of the vibration;²⁸ the Raman cross-section is proportional to the change in polarizability induced during the vibration. The increased C=O and C=C (from the benzene ring) intensities in Fig. 4b could be explained

by a higher cross-section of the vibration in benzaldehyde and benzoic acid than that in PAA, since an increased proximity of the




Fig. 4 Time-series Raman spectroscopy analysis (every 30 minutes) on hydrothermal decarboxylation of PAA (a) and PAA with copper chloride (b) at 200 °C and 15 bar (P_{sat}) for up to 2 hours. Dashed lines indicate major peaks with assignments taken from Lin-Vien et al. (1991). Aro = aromatic, sym = symmetric.

aromatic ring to the C=O moiety may induce a greater change in polarizability during vibration. More interestingly, an increasing fluorescence was detected over the reaction time, which may suggest the formation of copper-coordinated complexes in the system.^{29–31} These Raman results provided additional evidence for the presence of copper complexes in the PAA experiments.

Copper (II) nitrate is a readily available cupric salt and has emerged as a versatile reagent for organic synthesis.³² It has been used as an oxidizing agent and also a Lewis acid catalyst. For example, an organic synthesis study showed that copper nitrate allows facile oxidative decarboxylation of benzoyl formic acid to form benzonitrile in the presence of aqueous ammonium and O_2 ,³³ although the reaction conditions were quite different from the current work. To investigate whether copper nitrate played a role of Lewis acid in this decarboxylation, we chose another common Lewis acid – copper (II) triflate [$Cu(OTf)_2$] to study. The triflate ion is weakly nucleophilic, which makes the cupric ion more cationic and $Cu(OTf)_2$ a stronger Lewis acid than $Cu(NO_3)_2$.³⁴ At the same experimental conditions at 200 °C, however, we found the conversions of PAA after 1 and 2 hours were $4.4 \pm 0.2\%$ and $14.6 \pm 0.3\%$, respectively, which were much lower than those in the $Cu(NO_3)_2$ experiments (Fig. 1). In contrast, experiment of PAA with a mixture of $Cu(OTf)_2$ and $NaNO_3$ yielded similar conversion to that with $Cu(NO_3)_2$ (Table S1). These results




Fig. 3 Hydrothermal decarboxylation of PAA (a) and its product distribution (b) in the presence of magnesium nitrate, zinc nitrate, nickel nitrate, and copper nitrate at 200 °C and 15 bar (P_{sat}) after 4 hours.




Fig. 5 Hydrothermal transformation of benzyl alcohol (a) and its product distribution (b) in the presence of the copper salts at 200 °C and 15 bar (P_{sat}) after 4 hours.

further suggest that the complexation, instead of Lewis acidity, is more important to enhance the oxidative ability of copper (II).

Copper nitrate experiments with substituted phenylacetic acids, including *p*-tolylacetic acid and *p*-fluorophenylacetic acid, were also conducted to further clarify the decarboxylation mechanism. At 200 °C, the acid conversions after 1 hour were $42.1 \pm 0.3\%$ and $13.6 \pm 0.2\%$ for *p*-tolylacetic acid and *p*-fluorophenylacetic acid, respectively, while the conversion for parent PAA was $36.1 \pm 0.1\%$. This conversion trend (*p*-tolylacetic acid > PAA > *p*-fluorophenylacetic acid) thus suggests a carbocation intermediate (in Scheme 1), since the electron-donating substituent ($-\text{CH}_3$) on the benzene ring accelerated the reaction, whereas the electron-withdrawing substituent ($-\text{F}$) slowed down the reaction. In addition, experiments starting with benzyl alcohol were also performed, and results showed that benzaldehyde and benzoic acids were formed as the major products (Fig. 5), which further support the reaction steps proposed in Scheme 1. Specifically, in the presence of CuCl_2 , CuSO_4 , and $\text{Cu}(\text{OAc})_2$, the benzyl alcohol conversions were below 40% with an acid:aldehyde ratio < 0.1 , while in the presence of $\text{Cu}(\text{NO}_3)_2$ the conversion was nearly 100%, with a ratio increased to 15.1. These observations are also consistent with the PAA experiments, in which copper nitrate behaved as the strongest oxidant.

In conclusion, the fact that the organic reactivity and product distribution were dependent on the type of copper salts highlights a selective role of metal ions/complexes in hydrothermal organic redox transformations. The unique properties of water at high temperatures and pressures not only allow organic reactions to occur in aqueous medium, but also influence the speciation and complexation of dissolved metals, which can selectively control organic redox transformations in hydrothermal environments. While the majority of hydrothermal organic geochemistry research have been focusing on organic-mineral interactions, this work provides new insights into the hydrothermal interactions between organic and dissolved metal species, which are important to further our understanding of hydrothermal systems, as well as to discover more "geomimicry" approaches in green chemistry.^{14, 16}


ZY acknowledges the support from the Michigan Space Grant Consortium and startup fund from Oakland University. AMJ was supported by the USGS Energy Resources Program. The authors thank Bill Orem (USGS) for his review that improved this manuscript. Use of trade, product, or firm names is for descriptive purposes only and does not imply endorsement by the U.S. Government.

Conflicts of interest

There are no conflicts to declare.

Notes and references

- J. S. Seewald, *Nature*, 2003, **426**, 327-333.
- I. M. Head, D. M. Jones and S. R. Larter, *Nature*, 2003, **426**, 344-352.
- T. M. McCollom and J. S. Seewald, *Chemical Reviews*, 2007, **107**, 382-401.
- D. A. Sverjensky, V. Stagno and F. Huang, *Nature Geosci.*, 2014, **7**, 909-913.
- N. Akiya and P. E. Savage, *Chemical Reviews*, 2002, **102**, 2725-2750.
- M. Watanabe, T. Sato, H. Inomata, R. L. Smith, K. Arai, A. Kruse and E. Dinjus, *Chemical Reviews*, 2004, **104**, 5803-5822.
- Z. Yang, I. R. Gould, L. B. Williams, H. E. Hartnett and E. L. Shock, *Geochimica et Cosmochimica Acta*, 2012, **98**, 48-65.
- C. Bockisch, E. D. Lorange, H. E. Hartnett, E. L. Shock and I. R. Gould, *ACS Earth and Space Chemistry*, 2018, **2**, 821-832.
- K. M. Fecteau, I. R. Gould, C. R. Glein, L. B. Williams, H. E. Hartnett and E. L. Shock, *ACS Earth and Space Chemistry*, 2019, **3**, 170-191.
- K. J. Robinson, I. R. Gould, K. M. Fecteau, H. E. Hartnett, L. B. Williams and E. L. Shock, *Geochimica et Cosmochimica Acta*, 2019, **244**, 113-128.
- Z. Yang, E. D. Lorange, C. Bockisch, L. B. Williams, H. E. Hartnett, E. L. Shock and I. R. Gould, *The Journal of Organic Chemistry*, 2014, **79**, 7861-7871.
- C. R. Glein, I. R. Gould, E. D. Lorange, H. E. Hartnett and E. L. Shock, *Geochimica et Cosmochimica Acta*, 2020, **269**, 597-621.
- E. L. Shock, P. Canovas, Z. Yang, G. Boyer, K. Johnson, K. Robinson, K. Fecteau, T. Windman and A. Cox, *Reviews in Mineralogy and Geochemistry*, 2013, **76**, 311-350.
- Z. Yang, H. E. Hartnett, E. L. Shock and I. R. Gould, *The Journal of Organic Chemistry*, 2015, **80**, 12159-12165.
- Z. Yang, I. R. Gould, L. B. Williams, H. E. Hartnett and E. L. Shock, *Geochimica et Cosmochimica Acta*, 2018, **223**, 107-126.
- C. Bockisch, E. D. Lorange, G. Shaver, L. B. Williams, H. E. Hartnett, E. L. Shock and I. R. Gould, *Green Chemistry*, 2019, **21**, 4159-4168.
- J. A. Shipp, I. R. Gould, E. L. Shock, L. B. Williams and H. E. Hartnett, *Proceedings of the National Academy of Sciences*, 2014, **111**, 11642-11645.
- P. Gamez, P. G. Aubel, W. L. Driessen and J. Reedijk, *Chemical Society Reviews*, 2001, **30**, 376-385.
- J. Liu, X. Zhang, H. Yi, C. Liu, R. Liu, H. Zhang, K. Zhuo and A. Lei, *Angewandte Chemie International Edition*, 2015, **54**, 1261-1265.
- S. E. Allen, R. R. Walvoord, R. Padilla-Salinas and M. C. Kozlowski, *Chemical Reviews*, 2013, **113**, 6234-6458.
- S. D. McCann and S. S. Stahl, *Accounts of Chemical Research*, 2015, **48**, 1756-1766.
- J. R. Hein, K. Mizell, A. Koschinsky and T. A. Conrad, *Ore Geology Reviews*, 2013, **51**, 1-14.
- K. A. Miller, K. F. Thompson, P. Johnston and D. Santillo, *Frontiers in Marine Science*, 2018, **4**, 418.
- J. B. Dunn, M. L. Burns, S. E. Hunter and P. E. Savage, *The Journal of Supercritical Fluids*, 2003, **27**, 263-274.
- T. M. McCollom and J. S. Seewald, *Geochimica et Cosmochimica Acta*, 2003, **67**, 3625-3644.
- Z. Yang and X. Fu, *JoVE*, 2018, e58230.
- J. W. Johnson, E. H. Oelkers and H. C. Helgeson, *Computers & Geosciences*, 1992, **18**, 899-947.
- D. Lin-Vien, N. B. Colthup, W. G. Fateley and J. G. Grasselli, *The handbook of infrared and Raman characteristic frequencies of organic molecules*, Academic Press, 1991.
- M. H. Lim and S. J. Lippard, *Journal of the American Chemical Society*, 2005, **127**, 12170-12171.
- R. Krämer, *Angewandte Chemie International Edition*, 1998, **37**, 772-773.
- X.-M. Chen and M.-L. Tong, *Accounts of Chemical Research*, 2007, **40**, 162-170.
- M. Gao, R. Ye, W. Shen and B. Xu, *Organic & Biomolecular Chemistry*, 2018, **16**, 2602-2618.
- P. K. Arora and L. M. Sayre, *Tetrahedron Letters*, 1991, **32**, 1007-1010.
- C. Hertweck, *Journal für praktische Chemie*, 2000, **342**, 316-321.

Table of Content:

Efficient and selective effects of copper salts on hydrothermal oxidative decarboxylation have been discovered and studied.