



Cite this: RSC Adv., 2018, 8, 23757

Characterization of key aroma compounds in Meilanchun sesame flavor style baijiu by application of aroma extract dilution analysis, quantitative measurements, aroma recombination, and omission/addition experiments

Jinyuan Sun,^{ab} Qinya Li,^{ab} Siqi Luo,^{ab} Jinglin Zhang, ^{ab} Mingquan Huang, ^{*ab} Feng Chen, ^c Fuping Zheng,^{ab} Xiaotao Sun^{ab} and Hehe Li^{ab}

The aroma components in Meilanchun sesame flavor style baijiu were identified by aroma extract dilution analysis (AEDA), quantitative analysis, aroma active compound recombination, and omission/addition experiments. 92 components in Meilanchun were identified. Among them, 47 odor active compounds were further confirmed by GC-MS/O with aroma extract dilution analysis (AEDA). Ethyl pentanoate, 3-methyl-1-butanol, methional, ethyl 3-phenylpropanoate, phenethyl alcohol had the highest flavor dilution (FD) factors (FD = 2187). Among the 88 components that were determined in terms of their odor activity values (OAVs), 35 compounds showed OAVs ≥ 1 . Furthermore, a reconstitution model was prepared by mixing the above mentioned 35 compounds, lactic acid, phenethyl alcohol and 2-methoxy-1,3-dioxolane, and showed a good similarity to the aroma of Meilanchun baijiu. Omission/addition experiments further confirmed that ethyl hexanoate (OAV 1945), ethyl butanoate (OAV 838), 3-methyl-1-butanol (OAV 2), 3-methylbutanal (OAV 618), methional (OAV 59), and dimethyl trisulfide (OAV 44), might be the most important compounds for the unique flavor of Meilanchun baijiu. In addition, phenols and acetoin (OAV 66) were confirmed to be the key odorants and the odorless compound lactic acid played a significant role in the roasted sesame flavor baijiu by the omission experiments.

Received 30th March 2018
Accepted 13th June 2018

DOI: 10.1039/c8ra02727g

rsc.li/rsc-advances

Introduction

Baijiu is a kind of traditional alcoholic beverage in China, which has a long history, and is regarded as a pearl of ancient Chinese wisdom. Since the introduction of classification of baijiu flavor style in 1979, 12 major flavor styles of Chinese baijiu have been recognized. Among them, Jiang (soy sauce), Nong (strong), Mild (light), and Mi (rice) flavor styles are considered as the four basic flavor styles, while other flavor styles were considered to be derived from the aforementioned 4 basic flavor styles. For instance, sesame flavor style baijiu was derived from Jiang flavor style baijiu. The product possesses a unique style and complex aroma with roasted sesame flavor.¹⁻³

There are some representative baijius with a sesame flavor, including Jingzhi, Bandaojing, Meilanchun, Baotuquan, etc.

However, the whole flavor profiles of these baijius are different from each other. Liu *et al.* studied 12 sesame flavor style baijius by quantitative descriptive analysis, which included 8 aroma descriptions (*i.e.*, sesame, baked, Chen-aroma, Jiang, strong, light, distilled grain, Qu-aroma), 3 taste descriptions (*i.e.*, sweet, sour, bitter) and 5 mouthfeel descriptions (*i.e.*, rich, soft, harmony, clean, aftertaste) to describe the sensory characteristics of these baijius. By the principal component analysis, the sesame flavor style baijius were further divided into 4 styles (pure sesame, partial strong, partial light, and partial Jiang flavor style), and the flavor profile expressed the quality difference of different sesame flavor baijius.⁴

Meilanchun baijiu with sesame flavor (Meilanchun) was produced successfully by Ge Chongkai and Shen Yifang in 1985 according to a modified production process from Moutai liquor, one of the world's top three wines. The baijiu has a typical roasted sesame aroma and stable quality.⁵ In 1993, Jin Peizhang studied the components in Meilanchun and compared its characteristic components with those of other flavor style baijius, and the results showed that the contents of furfural, furfuralcohol, benzyl alcohol and 2,3-butanediol were high, which reflected some characteristics of soy sauce flavor style baijiu.⁶ The level of ethyl hexanoate was low, but had higher alcohols

^aBeijing Advanced Innovation Center for Food Nutrition and Human Health, Beijing Technology and Business University, Beijing, China 100048. E-mail: huangmq@btbu.edu.cn; Fax: +86-10-68984890; Tel: +86-10-68984613

^bBeijing Key Laboratory of Flavor Chemistry, Beijing Technology and Business University, Beijing, China, 100048

^cDepartment of Food Science and Human Nutrition, Clemson University, Clemson, SC 29634, USA



and isoamyl acetate. In 2012, Zhu Shuangliang compared the components of two Meilanchuns with other four famous baijius, and the results showed the amounts of 1,1-diethoxyethane, ethyl linoleate, ethyl oleate, ethyl palmitate, lactic acid and acetic acid were higher in Meilanchun than those of other baijius. In addition, its tetramethylpyrazine was 1.5–4.4 times of the same compound in other baijius with sesame flavor, but the content of butanoic acid was lower.⁷ Zhu Mengxu found that 1,1-diethoxymethane was not detected in all fresh liquors with the sesame flavor, but its content increased along with the aging process, as well as the great decrease of methanethiol, and both of them showed good correlations with the aging course.⁸ However, to the best of our knowledge, there is no report on the aroma-active compounds in Meilanchun up to now.

Sha Sha *et al.* investigated the aroma-active compounds in Jingzhi commercial baijiu with sesame flavor, and found 36 odorants had concentrations higher than their corresponding odor thresholds. For instance, 2-furfurylthiol (OAV 1182), dimethyl trisulfide (odor activity value, OAV 220), β -damascenone (OAV 116), and methional (OAV 99) could be responsible for the unique aroma of roasted sesame like flavor type liquor. Particularly, it was proposed that 2-furfurylthiol was the key typical potent odorant in the roasted sesame-like flavor type liquor by an omission test.⁹ Yang Zheng also studied the aroma-active compounds in two Jingzhi sesame flavor style baijius, and found 26 aroma compounds were further confirmed as the important odorants due to their OAVs ≥ 1 , and the omission experiments further corroborated the importance of ethyl hexanoate, 3-methylbutanal, ethyl pentanoate, methional and dimethyl trisulfide for the overall aroma of Jingzhi commercial baijiu.¹⁰ However, 2-furfurylthiol were neither found in the commercial baijiu nor in the base distillate. In addition, by direct injection combined with GC-MS, Sun *et al.* identified a total of 125 volatile compounds in 36 baijiu samples with the sesame flavor from 6 leading manufacturers. Among them, 30 volatiles, including two sulfur-containing compounds dimethyl disulfide and dimethyl trisulfide, were further identified as the active aroma compounds of Guojing sesame flavor style baijiu by the aroma extract dilution analysis (AEDA).¹¹ Yet, 2-furfurylthiol was not found either. Therefore, it is questionable on the function of 2-furfurylthiol for the sesame flavor of Jingzhi baijiu, although 2-furfurylthiol has a typical sesame aroma and is often used as a food flavoring ingredient. Nevertheless, Meilanchun baijiu with a sesame flavor was selected and studied for this research according to the means of sensomics approach.¹²

The aims of the present study were (i) to identify important odorants in Meilanchun sesame flavor style baijiu by GC-MS/O with AEDA, OAVs, recombination and omission/addition experiments, and (ii) to confirm the contribution of lactic acid for sesame flavor style baijiu.

Materials and methods

Chemicals

All chemicals were of analytical reagent grade, with at least 97% purity. Ethyl acetate, ethyl 2-methylpropanoate, ethyl butanoate, ethyl 2-methylbutanoate, ethyl 3-methylbutanoate,

ethyl pentanoate, ethyl hexanoate, ethyl lactate, diethyl butanedioate, ethyl propanoate, ethyl 4-methylpentanoate, 2-methyl-1-propanol, 1-butanol, 3-methyl-1-butanol, 1-hexanol, acetic acid, butyric acid, 3-methylbutanoic acid, pentanoic acid, hexanoic acid, octanoic acid, decanoic acid, 3-methylbutanal, acetoin, phenethyl alcohol, phenyl acetaldehyde, ethyl phenylacetate, ethyl 3-phenylpropanoate, vanillin, 2,6-dimethylpyrazine, 2,3,5-trimethyl pyrazine, guaiacol, 4-methylphenol, 4-ethylguaiacol, dimethyl trisulfide, methional, methionol, 1,1-diethoxyethane, ethyl nicotinate, 3-phenylpyridine, ethyl octanoate, hexyl acetate, propyl hexanoate, butyl hexanoate, hexyl hexanoate, ethyl nonanoate, ethyl decanoate, pentylhexanoate, ethyl 2-hydroxy-4-methylvalerate, ethyl heptanoate, isobutyl hexanoate, isopentyl hexanoate, ethyl dodecanoate, ethyl myristate, butyl butyrate, hexyl butyrate, diethyl octanedioate, ethyl hexadecanoate, 1-propanol, 1-heptanol, 1-octanol, propionic acid, pentanoic acid, heptanoic acid, nonanoic acid, ethyl benzoate, benzyl alcohol, acetophenone, benzaldehyde, 2-phenylethyl acetate, 2-furfural, furfuryl alcohol, 2-acetyl furan, 5-methylfurfural, 2-acetyl-5-methylfuran, tetramethylpyrazine, phenol, 4-ethylphenol, 2,4-di-*tert*-butylphenol, ethyl 3-methylthiopropionate, 1,1,3-triethoxypropane, γ -valerolactone, 3-(2-furyl)-2-propenal, α -terpenol, nonanal, ethyl 2-furoate, lactic acid, 3-phenylpyridine, 1,1-diethoxy-2-methylpropane, 2,3-diethyl-5-methylpyrazine, 2-methoxy-1,3-dioxolane, ethyl acrylate, 2,3-butandione, C₆–C₃₀ *n*-alkane mixture, were purchased from Sigma-Aldrich Co., Ltd. (Beijing, China), J&K Scientific Ltd. (Beijing, China), and Tokyo Chemical Industry Co., Ltd. (Shanghai, China). Dichloromethane, sodium chloride, anhydrous sodium sulfate, anhydrous ethanol, hydrochloric acid, sodium hydroxide were purchased from Sinopharm Chemical Reagent Co., Ltd. (Beijing, China). High purity nitrogen and high-purity helium (purity $\geq 99.999\%$), were purchased from Beijing AP BAIF Gases Industry Co., Ltd (Beijing, China).

Baijiu samples

The commercial baijiu with sesame flavor product (37.0% vol, 500 mL), provided by Meilanchun Company Limited, was stored at 4 °C prior to analysis. It is worthy of mention that the brand name did neither imply any of our research contact with the baijiu manufacturer, nor for advertising purpose.

Isolation of the volatiles

A total of 25 mL of Meilanchun commercial baijiu was diluted to 14% ethanol by volume with room-temperature boiled ultrapure water. The diluted baijiu sample was saturated with NaCl, and extracted 3 times with freshly distilled dichloromethane (50 mL each time), then the organic phase extracts were merged to get the organic phase O₁. Next, the organic phase O₁ was extracted 3 times with an alkaline solution (pH = 10.0, 50 mL each time), and separated to get the organic phase O₂ and the merged water phase extracts W₂. Then the W₂ was adjusted to pH = 2.0 with hydrochloric acid solution of 2 mol L⁻¹ and 1 mol L⁻¹, and the solution was extracted 3 times with freshly redistilled dichloromethane (50 mL each



time) after being saturated with sodium chloride, and then the combined organic phase O_3 was obtained. Organic phase O_2 and O_3 were dried over anhydrous Na_2SO_4 (30 g) in $-20^\circ C$, then filtered, and the filtrate were concentrated to $\sim 500 \mu L$ with using Vigreux columns followed by gentle stream of nitrogen. Finally the neutral/basic fraction (NBF) and acidic fraction (AF) of components in Meilanchun were obtained, respectively.

Gas chromatography & mass spectrometric/olfactometric (GC-MS/O) analysis

GC-MS/O analysis was performed on an Agilent 7890 gas chromatograph, which was equipped with an Agilent 5975 mass detector and an olfactometer (ODP2, Gerstel, Germany). Samples were analyzed on two different fused silica capillaries, including DB-WAX (60 m \times 0.25 mm i.d., 0.25 μm film thickness, J&W Scientific) and HP-5MS (30 m \times 0.25 mm i.d., 0.25 μm film thickness, J&W Scientific). The injector temperature was $250^\circ C$, and the splitless injection mode was used. The temperature of the olfactory port was kept at $220^\circ C$. Ultra-high purity helium (purity $> 99.999\%$) was used as the carrier gas at a constant rate of 1.5 mL min^{-1} . An aliquot of $2.0 \mu L$ of the concentrated samples (NBF and AF) was analyzed by GC-MS/O.

DB-WAX column heating program: the oven temperature was held at $40^\circ C$ firstly, then raised to $50^\circ C$ at $10^\circ C \text{ min}^{-1}$ and held for 10 min, then ramped to $80^\circ C$ at $3^\circ C \text{ min}^{-1}$ and held for 10 min, and finally increased at $5^\circ C \text{ min}^{-1}$ to $240^\circ C$, held for 7 min.

HP-5MS column heating program: the oven temperature was held at $35^\circ C$ for 0.5 min firstly, then raised to $70^\circ C$ at $0.6^\circ C \text{ min}^{-1}$ and held for 5 min, then ramped to $230^\circ C$ at $5^\circ C \text{ min}^{-1}$, and finally increased to $280^\circ C$ at $20^\circ C \text{ min}^{-1}$ and held for 7 min.

The mass spectrometer was operated in electron ionization (EI) mode at 70 eV, and the ion source temperature was $230^\circ C$. The mass-to-charge (m/z) ranged from 35 to 450 in full scan acquisition mode.

The sniffing experiments were performed by 3 experienced panelists (two females and one male). Before the experiments, they were trained by sniffing about 40 reference compounds in their concentrations 10 times above their odor thresholds in water or air.¹³ The identification of a compound was confirmed by comparing its mass spectra, retention time, odor quality and retention index (RI) with that of the corresponding reference compounds. All analyses were repeated in triplicate by each panelist.

Aroma extract dilution analysis (AEDA)

The flavor dilution factors of the active aroma compounds in the NBF and AF were determined by GC-MS/O on a DB-WAX capillary column. The two concentrated extracts of Meilanchun, NBF and AF, were diluted stepwise with dichloromethane in a series of 1 : 3, 1 : 9, 1 : 27, ..., until no scent was perceived. The flavor dilution factor of each compound was the maximum dilution in which the compound could be perceived.

Quantitative analysis of aroma compounds

The quantitation analysis of the volatile components was carried out on DB-WAX column, and their signal-to-noise ratios were required bigger than 10.

Quantitative analysis by GC-MS. Baijiu sample was diluted by ethanol to 80.0% vol alcohol solution in order to reduced moisture content, and then 1.0 mL of the diluted sample added with 10.0 μL internal standards solution was used for GC-MS analysis. The internal standard solutions were prepared as follows. The standard stock solutions were prepared with 80.0% vol ethanol solution. Then the different concentration standard solutions were obtained by diluting the stock solution step by step. Finally, 1.0 mL aliquots of these standard solutions were determined by GC-MS with selected-ion monitoring (SIM) mode. The standard curves were carried out by plotting the response ratios of standard compounds and internal standards against their concentration ratios.

The above diluted samples and different level standard solutions were spiked with the pivalic acid (IS1, 4.67 mg L^{-1} final concentration) and 2-ethyl butyric acid (IS2, 52.87 mg L^{-1} final concentration) as internal standards for acids, with methyl octanoate (IS3, 117.14 mg L^{-1} final concentration), methyl hexanoate (IS4, 1.1238 mg L^{-1} final concentration) and octyl propionate (IS5, 1.1429 mg L^{-1} final concentration) as internal standards for esters and other compounds. The internal standards with high concentrations were selected for the high concentrations of compounds in baijiu sample.

Quantitative analysis by headspace solid-phase microextraction-gas chromatography-mass spectrometry (HS-SPME-GC-MS). The baijiu sample was diluted by saturated sodium chloride solution to 10% ethanol solution in volume. Then, 8 mL of the diluted solution with 10 μL internal standards (IS4-2, methyl hexanoate, 73.75 $\mu g L^{-1}$ final concentration; IS5-2, octyl propionate, 75 $\mu g L^{-1}$ final concentration) was poured into a 20 mL screw-capped vial. An automatic headspace sampling system (Multi Purpose Sample MPS2 with a solid-phase microextraction (SPME) adapter, from Gerstel Inc., Mülheim, Ruhr, Germany) with a carboxen/polydimethylsiloxane (CAR/PDMS 75 μm , Supelco, Inc., Bellefonte, PA, USA) was used to extract the volatile components. After the sample was preheated at $45^\circ C$ for 5 min, the SPME fiber was inserted into the headspace and adsorbed for 40 min at $45^\circ C$. Then the SPME fiber was desorbed at $250^\circ C$ for 5 min in the injection port of GC-MS with a splitless mode. The flow rate of helium was 1.5 mL min^{-1} , and the heating procedure was the same as the DB-WAX column analyzed by GC-MS/O. The GC-MS was used in selected-ion monitoring (SIM) mode.

The standard solutions on different levels were prepared by 10% vol alcohol solution, which was formulated with pure alcohol and the saturated sodium chloride solution, and then followed by an adjustment of their pH values with a dilute hydrochloric acid to the same pH of the corresponding baijiu sample. Then these standard solutions were analyzed by GC-MS as the above baijiu sample.

Quantitative analysis of lactic acid by liquid chromatography (LC). Because the content of lactic acid is very high in many



baijius and probably has some contributions to the flavor of baijiu, it needs to be quantitated. However, lactic acid can be decomposed easily at high temperature, its content can not be measured by GC or GC/MS. Therefore, a high performance liquid chromatography (HPLC) (1260, Agilent Technologies, Inc., Santa Clara, CA) was used in our experiment.

The baijiu sample in a volume of 10.0 mL was evaporated and concentrated to 3–4 mL at 40 °C by a rotatory evaporator, and then the concentrated solution was added by ultra-pure water to 10.0 mL in total volume. Then the solution and a series of standard solutions of lactic acid were simultaneously analyzed by the same HPLC mentioned above. The standard solutions of lactic acid were prepared by ultra-pure water.

The LC conditions were as follows. Venusil XBP C₁₈ (4.6 mm × 250 mm, 5 µm) was used and hold at 25 °C. The variable wavelength detector worked at 205 nm. The mobile phase was the mixture of methanol and potassium dihydrogen phosphate solution at the constant ratio 5 : 95 by volume. The concentration of potassium dihydrogen phosphate solution was 0.01 mol L⁻¹ in water (pH = 2.8). The flow rate was 1.0 mL min⁻¹. The injection volume was 80.0 µL. Each samples was tested 3 times.

Determination of methionol and methional. The content of methionol and methional was measured by an external standard method and a liquid–liquid extraction followed by a GC-MS analysis, with the organic phase O₁ from the above part of isolation of the volatiles as the analysis sample. The standard solutions of two sulfur-containing compounds were prepared with 37.0% vol ethanol solution. These different standard solutions were prepared as the above baijiu samples.

Odor threshold determination and odor activity value (OAV) analysis

The threshold values of components were mainly quoted from literatures, and other threshold values were determined in this study according to the literature.

Method of threshold determination. Based on the described method,^{13,14} the orthonasal thresholds of the odorants were determined by a forced-choice test at seven concentration steps. The proper high concentration of every odorant was firstly prepared with hydroalcoholic solution at 46% ethanol in volume, and then the solution was stepwise diluted (1 : 3 by volume) with the same matrix to get 7 different concentration solutions for each odorant. Seven triangular series were prepared for every odorant, and each of them comprised of one glass of the odorant dilution and two glasses of hydroalcoholic solution. All of the series were labeled with random four-digit numbers and presented in decreasing concentrations. A sensory panel consisting of 25 panelists was asked to sniff and select a different one in each group of the triangular series. Then the two concentrations for every odorant were recorded, *i.e.* the minimum concentration that the assessors correctly selected and the maximum concentration incorrectly selected. Then, the odor threshold of each odorant was calculated by the following formula described in ref. 15.

$$OT_i = \sqrt{C_x \times C_{x+1}}$$

OT_i: the individual recognition/detection odor threshold of each assessor; C_x: the lowest concentration of the odorant, which was correctly selected by the assessor; C_{x+1}: the highest concentration of the odorant, which was incorrectly selected by the assessor.

The recognition/detection odour threshold (OT_n) of the compound tested by the panel was calculated by the following formula¹⁵

$$OT_n = \sqrt[n]{\prod_{i=1}^n OT_i}$$

OT_n: the recognition/detection odour threshold of the compound tested by a panel; n: number of assessors; $\prod_{i=1}^n OT_i$: the product of individual recognition/detection odor threshold.

Based on the quantitative results, odor activity value (OAV) of each substance was calculated as follows: OAV = the concentration of a compound/the odor threshold of the compound.

Descriptive profile test

The sensory evaluation of the Meilanchun sample was performed by 10 trained panelists (5 males and 5 females, 24 years old on average). The Meilanchun sample in 30 mL was poured in an odorless tasting glass of Chinese baijiu, and the descriptive profile test was analyzed under 20 ± 1 °C.¹⁵ The assessors were asked to rank the intensities of 8 characteristic aroma attributes on a seven-point scale (0, 0.5, 1.0, 1.5, 2.0, 2.5, 3.0) with 0 indicating not perceivable, 1 indicating weak, 2 indicating significant, 3 indicating strong. Eight characteristic aroma attributes and their responding references were as follows: roasted sesame seed for baked sesame note, acetic acid for acidic note, steamed grains including sorghum and rice husk for grain-like note, old pit mud for pit mud-like note, ethyl pentanoate for fruity note, 3-methylbutanal for malty note, 2-methoxy-1,3-dioxolane for sweet note, ethanol for alcoholic note. After being trained, the overall aroma was evaluated for the Meilanchun baijiu by the panel.

Aroma recombination experiments

By using the odorants with OAVs ≥ 1, phenethyl alcohol, 2-methoxy-1,3-dioxolane and lactic acid, the complete aroma recombination models was prepared in 37% ethanol solution in pure water matrix (by volume), and their pH were adjusted to be same with Meilanchun. The aroma profiles of the recombination model were analyzed with the same method for Meilanchun.

Omission experiments and addition experiments

As the above odor threshold determination, a triangle test was performed to determine the significance of one odorant. A glass (20 mL) of the mixture was prepared by omitting or adding one or a group of selected components from the complete recombination models, and then presented with two glasses of



Table 1 Aroma compounds identified by gas chromatography–mass spectrometry/olfactometry in Meilanchun sesame flavor style baijiu

No.	Compound	Odor quality	Fraction ^a	Base of Id ^b	RI		
					DB-WAX	HP-5MS	log ₃ FD
Esters							
1	Ethyl acetate	Fruity, nail polish-like	NBF	MS,S,O,RI	910	614	1
2	Ethyl 2-methylpropanoate	Fruity	NBF	MS,S,O,RI	954	754	5
3	Ethyl propanoate	Fruity, nail polish-like	NBF	MS,S,O,RI	946	711	2
4	Ethyl acrylate	Plastic-like	NBF	MS,S,O	976	—	0
5	Ethyl butanoate	Fruity, apple-like, strawberry-like	NBF/AF	MS,S,O,RI	1018	803	6
6	Ethyl 2-methylbutanoate	Fruity, apple-like, strawberry-like	NBF	MS,S,O,RI	1036	846	5
7	Ethyl 3-methylbutanoate	Fruity, apple-like, strawberry-like	NBF	MS,S,O,RI	1053	850	5
8	Ethyl pentanoate	Fruity, apple-like, strawberry-like	NBF	MS,S,O,RI	1118	903	7
9	Ethyl 4-methylpentanoate	Fruity	NBF	MS,S,O,RI	1170	970	4
10	Ethyl hexanoate	Fruity, fermented pear-like	NBF/AF	MS,S,O,RI	1215	1005	4
11	Ethyl lactate	Fruity	NBF	MS,S,O,RI	1309	820	0
12	Diethyl butanedioate	Sweet	NBF	MS,S,O,RI	1661	1185	3
13	Ethyl phenylacetate	Flowerly, honey	NBF	MS,S,O,RI	1777	1247	2
14	Ethyl nicotinate	Honey, sweet	NBF	MS,S,O,RI	1819	1213	3
15	Ethyl 3-phenylpropanoate	Sweet	NBF	MS,S,O,RI	1882	1351	7
Alcohols							
16	2-Methyl-1-propanol	Malty, roast nuts-like	NBF/AF	MS,S,O,RI	1076	623	1
17	1-Butanol	Malty, roast nuts-like	NBF/AF	MS,S,O,RI	1128	658	1
18	3-Methyl-1-butanol	Malty, roast nuts-like	NBF/AF	MS,S,O,RI	1191	733	7
19	1-Hexanol	Green, flowery	NBF	MS,S,O,RI	1315	865	0
20	Phenethyl alcohol	Flowery	NBF/AF	MS,S,O,RI	1906	1112	7
Acetals							
21	1,1-Diethoxyethane	Fruity	NBF	MS,S,O,RI	916	726	5
22	2-Methoxy-1,3-dioxolane	Sweet, cake-like	NBF	MS,S,O,RI	1207	927	6
23	2,4,6-Trimethyl-1,3-dioxane	Sweet, fruity	NBF	MS	1402	—	3
24	1,1-Diethoxy-2-methylpropane	Sweet, fruity	NBF	MS,S,O	970	—	3
Pyrazines							
25	2,6-Dimethylpyrazine	Woody, roast nuts-like	AF	MS,S,O	1289	—	1
26	2,3,5-Trimethyl pyrazine	Nutty, almond-like, buttery	NBF	MS,S,O	1386	—	1
27	2,3-Diethyl-5-methylpyrazine	Smoky, woody	AF/NBF	MS,S,O	1475	—	4
Carboxylic acids							
28	Acetic acid	Sour, vinegar-like	AF	MS,S,O,RI	1428	600	1
29	Butanoic acid	Sweaty, rancid	AF	MS,S,O	1610	—	4
30	3-Methylbutanoic acid	Sweaty	AF	MS,S,O	1657	—	4
31	Pentanoic acid	Sweaty	AF	MS,S,O	1723	—	2
32	Hexanoic acid	Sour, vinegar-like	AF	MS,S,O	1830	—	3
33	Octanoic acid	Sweaty	AF	MS,S,O	2049	—	1
34	Decanoic acid	Sour	AF	MS,S,O	2271	—	3
Sulfur-containing odorants							
35	Dimethyl trisulfide	Stir-fried vegetable-like, sulfur	NBF	MS,S,O,RI	1353	968	3
36	Methional	Cooked potato	NBF/AF	MS,S,O,RI	1434	905	7
37	Methionol	Cooked potato	NBF	MS,S,O,RI	1704	980	1
Phenols							
38	Guaiacol	Smoky, woody	NBF	MS,S,O,RI	1849	1089	4
39	4-Ethylguaiacol	Smoky	NBF	MS,S,O,RI	2024	1280	5
40	4-Methylphenol	Fecal, horse stable-like	NBF/AF	MS,S,O,RI	2071	1082	5
Carbonyl odorants							
41	2,3-Butandione	Butter-like	NBF/AF	MS,S,O,RI	960	604	0
42	3-Methylbutanal	Malty, roast nuts-like	NBF/AF	MS,S,O,RI	921	648	6
43	Phenylacetaldehyde	Honey	NBF	MS,S,O,RI	1629	1041	0
44	Vanillin	Caramel-like	AF	MS,S,O	2582	—	3



Table 1 (Contd.)

No.	Compound	Odor quality	Fraction ^a	Base of Id ^b	RI		
					DB-WAX	HP-5MS	\log_3 FD
Other odorants							
45	3-Phenylpyridine	Sour	AF	MS,O	2261	—	3
46	uk1-M ^c	Cooked potato	NBF/AF	O	1430	—	3
47	uk2-M ^c	Sour	AF	O	2245	—	3

^a Fraction(s) in which odorant was detected by GC-MS/O after fractionation. AF, acidic/water-soluble fraction; NBF, neutral and basic fraction. ^b MS, compounds were identified by NIST MS spectra; S, compounds were identified by standards; O, compounds were identified by comparison to reference standards by GC-MS/O; RI, compounds were identified on DB-WAX and HP-5MS by comparison to reference standards. ^c Unknown compound that could not be identified.

complete recombination models to the sensory panel in a triangle test.¹⁶ The sensory panel was the same as descriptive profile experiments. The significance α was calculated according to ref. 17. The sensory data were analyzed by one-way analysis of variance (ANOVA) by use of SPSS 20.0 (SPSS Inc., Chicago, IL, USA).

Results and discussion

Analysis of aroma-active compounds in Meilanchun baijiu

Ninety-two compounds were identified from NBF and AF, which were extracted from Meilanchun commercial baijiu. With GC-MS/O, 47 odor-active peaks were detected, and 45 odor-active components were confirmed further, although two could not be identified yet. Besides, 35 odorants showed FD factors ≥ 9 , which were shown in Table 1. The highest FD factors were determined for the ethyl pentanoate (8, fruity, apple-like, strawberry-like odor), ethyl 3-phenylpropanoate (15, sweet odor), 3-methyl-1-butanol (18, malty, roast nuts-like odor), phenylethyl alcohol (20, flowery odor), and methional (36, cooked potato odor) (FD = 2187), followed by 3-methylbutanal (42, malty, roast nuts-like odor), ethyl butanoate (5, fruity, apple-like, strawberry-like), and 2-methoxy-1,3-dioxolane (22, sweet, cake-like odor) with FD 729. A majority of these odorants in Table 1 have also been reported as aroma compounds in other flavor type of Chinese baijiu.^{13,18}

Among the 45 odorants shown in Table 1, there were 15 esters, 5 alcohols, 4 acetals, 3 pyrazines, 7 carboxylic acids, 3 sulfur-containing compounds, 3 phenols, 4 carbonyl odorants and 1 pyridine. The esters and acetals mainly contributed the fruity and sweet flavors to the Meilanchun baijiu, while the alcohols and 3-methylbutanal offered malty, roasted nut-like, the phenethyl alcohol contributed the flowery, the acids provided sour and sweaty, the phenols contributed the smoky, and methional offered cooked potato aromas. Pyrazines contributed some nutty and woody to the flavor of Meilanchun baijiu. The FDs of uk1-M (cooked potato) and uk2-M (sour) were 9 and 27 respectively.

Concentration of compounds in Meilanchun baijiu

In order to get deeper insight into the Meilanchun baijiu, a total of 90 detected compounds were quantitated (Table 2) by GC-MS

or HPLC. Among these compounds, the content of ethyl acetate (808.66 mg L^{-1}) was the highest, followed by ethyl lactate (751.75 mg L^{-1}), acetic acid (711.08 mg L^{-1}), 3-methyl-1-butanol (249.72 mg L^{-1}), 2-methyl-1-propanol (158.23 mg L^{-1}), furfural (131.36 mg L^{-1}), hexanoic acid (115.83 mg L^{-1}) and ethyl hexanoate (107.62 mg L^{-1}). These 8 compounds were present at levels above 100 mg L^{-1} , and were taken as the skeleton component (SC) of Meilanchun. Besides, the following compounds had higher concentrations, such as butyric acid (69.01 mg L^{-1}), ethyl butanoate (68.32 mg L^{-1}), 1-butanol (31.99 mg L^{-1}), pentanoic acid (23.98 mg L^{-1}), phenethyl alcohol (21.94 mg L^{-1}), 1,1-diethoxyethane (21.09 mg L^{-1}). The employed quantitative methods were able to detect all of the identified compounds in Meilanchun. The obtained calibration curves had good linearity with correlation coefficient ($R^2 \geq 0.99$); RSDs in triplicate of samples were $\leq 15\%$, which revealed that the good precision of the quantitative methods.

Four kinds of carboxylic acids (lactic acid, acetic acid, hexanoic acid and butyric acid) are important to flavor in baijiu. All acids are volatiles except lactic acid which may usually ignored the contribution to baijiu aroma. Lactic acid in Meilanchun was quantitated by HPLC. Noteworthy the content of lactic acid was high ($1754.67 \text{ mg L}^{-1}$) which may influence the aroma of Meilanchun.

Odor activity values (OAVs)

To overcome the shortcomings of GC-MS/O analysis with AEDA,¹⁹ and in view of ethanol concentration on the volatility of an odorant,²⁰ the OAVs (ratio of concentration to its odor threshold) of the odorants were calculated to evaluate their contributions to the flavor of Meilanchun (Table 2). Besides, 13 odor thresholds determined in a mixture of ethanol/water (46/54 by vol) were used in this study.

The results in Table 2 showed there were 35 components with OAVs ≥ 1.0 in Meilanchun, including 6 compounds undetected by GC-MS/O. They were ethyl octanoate (OAV 104), butyl hexanoate (OAV 1), 1-propanol (OAV 2), 4-methyl pentanoic acid (OAV 33), acetoin (OAV 66), 2-furfural (OAV 3). This showed that the influence of ethanol concentration on the volatility of an odorant.²⁰ The OAVs of 7 compounds was greater than 100, which was ethyl hexanoate (OAV 1957), ethyl butanoate (OAV 833), 3-methylbutanal (OAV 638), ethyl 3-



Table 2 Compounds' concentrations of in Meilanchun sesame flavor style baijiu

Count	Compound	Meilanchun		OAVs		
		AV ^b ± RSD (%)	Odor threshold ^c (μg L ⁻¹)			
Aroma-active compounds in Meilanchun						
Esters						
1	Ethyl acetate ^a	808.66 ± 6.3	32552 (ref. 21)	25		
2	Ethyl 2-methylpropanoate ^a	15.61 ± 12.5	58 (ref. 21)	269		
3	Ethyl butanoate ^a	68.32 ± 4.5	82 (ref. 21)	833		
4	Ethyl 2-methylbutanoate ^a	0.80 ± 8.1	18 (ref. 23)	44		
5	Ethyl 3-methylbutanoate ^a	2.78 ± 6.1	6.9 (ref. 21)	403		
6	Ethyl pentanoate ^a	6.75 ± 4.3	27 (ref. 21)	250		
7	Ethyl hexanoate ^a	107.62 ± 9.1	55 (ref. 21)	1957		
8	Ethyl lactate ^a	751.75 ± 6.4	128 084 (ref. 21)	6		
9	Diethyl butanedioate ^a	1.33 ± 7.5	353 193 (ref. 21)	<1		
10	Ethyl propanoate	1351.71 ± 10.3	19 019 (ref. 21)	<1		
11	Ethyl 4-methylpentanoate	74.38 ± 1.5	1409	<1		
12	Ethyl phenylacetate	497.78 ± 6.8	407 (ref. 21)	1		
13	Ethyl 3-phenylpropanoate	651.00 ± 8.4	125 (ref. 21)	5		
14	Ethyl nicotinate	95.44 ± 5.4	7781	<1		
15	Ethyl acrylate	8.34 ± 7.6	0.2 (ref. 22)	42		
Alcohols						
1	2-Methyl-1-propanol ^a	158.23 ± 5.0	28 300 (ref. 13)	6		
2	1-Butanol ^a	31.99 ± 5.9	2730 (ref. 21)	12		
3	3-Methyl-1-butanol ^a	249.72 ± 4.0	179 191 (ref. 21)	1		
4	1-Hexanol ^a	2.04 ± 11.0	5370 (ref. 13)	<1		
5	Phenethyl alcohol ^a	21.94 ± 6.8	28 923 (ref. 21)	<1		
Carboxylic acids						
1	Acetic acid ^a	711.08 ± 4.6	160 000 (ref. 13)	4		
2	Butyric acid ^a	69.01 ± 4.4	965 (ref. 21)	72		
3	3-Methylbutanoic acid ^a	6.89 ± 5.8	1045 (ref. 21)	7		
4	Pentanoic acid ^a	23.98 ± 4.3	389 (ref. 21)	62		
5	Hexanoic acid ^a	115.83 ± 4.6	2517 (ref. 21)	46		
6	Octanoic acid ^a	7.48 ± 5.4	2701 (ref. 21)	3		
7	Decanoic acid	5571.81 ± 9.9	13736 (ref. 21)	<1		
Aldehydes and ketone						
1	3-Methylbutanal ^a	10.21 ± 4.5	16 (ref. 21)	638		
2	Phenyl acetaldehyde	61.33 ± 15.6	262 (ref. 13)	<1		
3	Vanillin	421.69 ± 13.7	438 (ref. 21)	1		
4	2,3-Butanedione	8.67 ± 12.03	5 (ref. 22)	2		
Pyrazines						
1	2,6-Dimethylpyrazine	58.02 ± 12.2	790 (ref. 21)	<1		
2	2,3,5-Trimethylpyrazine	91.10 ± 6.1	729 (ref. 21)	<1		
3	2,3-Diethyl-5-methylpyrazine	—	—	—		
Phenols						
1	Guaiacol	20.93 ± 15.6	13 (ref. 21)	2		
2	4-Methylphenol	475.38 ± 5.9	167 (ref. 21)	3		
3	4-Ethyl guaiacol	250.75 ± 6.4	123 (ref. 21)	2		
Sulfur-containing odorants						
1	Dimethyl trisulfide	15.95 ± 15.1	0.36 (ref. 21)	44		
2	Methional	422.20 ± 10	7.12 (ref. 23)	59		
3	Methionol	280.23 ± 5.7	2110 (ref. 21)	<1		
Carbonyl odorants						
1	1,1-Diethoxyethane ^a	21.09 ± 6.7	2090 (ref. 13)	10		
2	Isobutyraldehyde diethyl acetal	8.13 ± 10.23	—	—		
3	2,4,6-Trimethyl-1,3-dioxane	—	—	—		
4	2-Methoxy-1,3-dioxolane	21.46 ± 8.6	—	—		
Other odorants						
1	3-Phenylpyridine	59.23 ± 14.2	19 138	<1		
2	uk1-M	—	—	—		
3	uk2-M	—	—	—		
Other compounds in Meilanchun						
1	Ethyl octanoate ^a	1.33 ± 7.2	12.87 (ref. 21)	104		
2	Hexyl acetate	79.19 ± 2.8	5560 (ref. 24)	<1		



Table 2 (Contd.)

Count	Compound	Meilanchun		OAVs
		AV ^b ± RSD (%)	Odor threshold ^c (μg L ⁻¹)	
3	Propyl hexanoate	148.59 ± 1.2	12 783 (ref. 21)	<1
4	Butyl hexanoate	658.11 ± 1.1	678 (ref. 24)	1
5	Hexyl hexanoate	1166.35 ± 2.1	1890 (ref. 24)	<1
6	Ethyl nonanoate	41.05 ± 12.1	3150 (ref. 21)	<1
7	Ethyl decanoate	652.13 ± 6.3	1122 (ref. 21)	<1
8	Pentylhexanoate	158.99 ± 4.6	13 802 (ref. 25)	<1
9	Ethyl 2-hydroxy-4-methylvalerate ^a	12.73 ± 6.8	—	—
10	Ethyl heptanoate ^a	2.08 ± 3.3	13 153 (ref. 21)	<1
11	Isobutyl hexanoate	309.86 ± 9.1	5250 (ref. 25)	<1
12	Isopentylhexanoate	166.66 ± 6.6	1400 (ref. 24)	<1
13	Ethyl dodecanoate	354.51 ± 12.8	25 619	<1
14	Ethyl myristate	221.93 ± 7.5	46 606	<1
15	Butyl butyrate	672.24 ± 5.2	14 066	<1
16	Hexyl butyrate	73.03 ± 12	30 466	<1
17	Diethyl octanedioate	18.37 ± 1.9	641 000 (ref. 23)	<1
18	Ethyl hexadecanoate	123.77 ± 11.5	39 299	<1
19	1-Propanol^a	96.47 ± 8.3	53 952 (ref. 21)	2
20	1- Heptanol	138.19 ± 12.1	26 600 (ref. 23)	<1
21	1-Octanol	119.53 ± 5.6	1100 (ref. 23)	<1
22	Propionic acid ^a	15.28 ± 10.2	18 100 (ref. 23)	<1
23	4-Methyl pentanoic acid^a	4.83 ± 1.1	144 (ref. 24)	33
24	Heptanoic acid ^a	6.18 ± 3.0	13 821 (ref. 21)	<1
25	Nonanoic acid	3583.2 ± 1.6	3559 (ref. 21)	<1
26	Acetoin^a	17.00 ± 2.0	259 (ref. 23)	66
27	Ethyl benzoate	127.56 ± 13.5	1433 (ref. 21)	<1
28	Benzyl alcohol	61.42 ± 14.4	40 927 (ref. 21)	<1
29	Acetophenone	8.65 ± 16.6	9474	<1
30	Benzaldehyde ^a	2.09 ± 7.6	4203 (ref. 21)	<1
31	2-Phenylethyl acetate	168.64 ± 8.6	908 (ref. 21)	<1
32	2-Furfural^a	131.36 ± 7.7	44 029 (ref. 21)	3
33	Furfuryl alcohol	381.89 ± 9.9	12 323	<1
34	2-Acetyl furan	41.91 ± 14.9	58 504 (ref. 21)	<1
35	5-Methylfurfural	124.80 ± 13.6	466321 (ref. 21)	<1
36	2-Acetyl-5-methylfuran	53.16 ± 14.6	40 870 (ref. 21)	<1
37	Tetramethylpyrazine	84.76 ± 3.4	80 073 (ref. 21)	<1
38	Phenol	64.30 ± 9.5	18 900 (ref. 21)	<1
39	4-Ethylphenol	367.80 ± 2.5	617 (ref. 21)	<1
40	2,4-Di- <i>tert</i> -butylphenol	438.40 ± 13.1	36 373	<1
41	Ethyl 3-methylthiopropionate	78.65 ± 7.7	3080 (ref. 25)	<1
42	1,1,3-Triethoxypropane	492.35 ± 0.8	3700 (ref. 23)	<1
43	γ-Valerolactone	304.86 ± 9.6	25 982	<1
44	3-(2-furyl)-2-propenal	130.59 ± 7.6	13 128	<1
45	α-Terpenol	10.63 ± 16.1	1960 (ref. 25)	<1
46	Nonanal	39.83 ± 13.0	122 (ref. 21)	<1
47	Ethyl 2-furoate	131.23 ± 5.5	132 000 (ref. 25)	<1
48	Lactic acid ^a	1754.67 ± 1.3	—	—

^a The unit of concentration was mg L⁻¹, the unit of other compounds was μg L⁻¹. ^b Average concentration of triplicates. ^c Odor thresholds were determined in 46% ethanol/water solution according to our laboratory or references.^{13,21,22,23,24,25} reference numbers.

methylbutanoate (OAV 403), ethyl 2-methylpropionate (OAV 269), ethyl pentanoate (OAV 250) and ethyl octanoate (OAV 104). These components with OAVs ≥ 1.0 should be important to the flavor of Meilanchun. There were 12 odorants with the OAVs between 10 and 100, such as ethyl hexanoate, ethyl butanoate and 3-methylbutanal. However, methionol, pyrazines and phenethyl alcohol had little effect on the aroma of Meilanchun because of their OAVs smaller than 1.

In consideration of both FD values and the OAVs, it was found that some compounds with high FD value had relatively smaller OAVs. For example, the FD factors of phenethyl alcohol, 3-methyl-1-butanol and ethyl 3-phenylpropanoate possessed the largest FD values 2187, but their OAVs were low which indicated that the influence of the food matrix on odorant binding.¹⁹



Based on the OAVs in Table 2, esters (especially ethyl hexanoate and ethyl butanoate), alcohols, acids, 3-methylbutanal, phenols, dimethyl trisulfide, methional, and 1,1-diethoxyethane were considered to have important influence on the aroma of Meilanchun, which was basically in accordance with FD values and the reported results by Zheng *et al.*¹⁰ and Sha *et al.*,⁹ especially the contributions of ethyl hexanoate, ethyl butanoate, 3-methylbutanal, dimethyl trisulfide, and methional to the aroma of roasted sesame flavor baijiu.

However, 2-furfurylthiol with a roasted sesame aroma reported by Sha *et al.*⁹ was not found in the study. According to the previous work,^{26,27} the odor of roasted sesame seeds was characterized by sulfurous, roasty, nutty, and meaty notes. The authors had not found a compound with a sesame-like flavor. The sesame-like flavor might be a composite flavor, and so there was probably no compound with sesame-like flavor.

Descriptive profile and aroma recombination of Meilanchun

To validate the quantitative results, the descriptive profile experiments were performed for the aroma recombine in comparison to Meilanchun by rating the intensity of 7 odor attributes, as well as the overall similarity. The aroma recombine was prepared in an ethanolic solution (37%, ethanol by volume), and contained all important odorants with OAVs ≥ 1.0 , phenethyl alcohol and 2-methoxy-1,3-dioxolane because of their high FD values, as well as lactic acid due to its high concentration. Besides, the pH in the aroma recombine was adjusted to 3.9. The aroma descriptive profiles for Meilanchun and its simulation sample (M1) are shown in Fig. 1.

As shown in Fig. 1, a distinct sweet aroma quality was detected in Meilanchun, followed by acid, fruity, grain aroma, baked sesame-like, malty. The aroma of the recombine had good similarities for sweet, fruity, grain-like, acidic, ethanol and pit mud-like, although the roasted sesame had some deficiencies. In addition, the floral note was very weak either in Meilanchun or in the recombine, although a very strong

floral aroma could be smelt during the GC-MS/O analysis. This implied that there were interactions among the odorants.¹⁹ The overall aroma similarities between the recombination model and Meilanchun were judged to be 2.7 to 3.0 points and 2.8 to 3.0 points, respectively. These results indicated the successes in simulating the typical flavor of Meilanchun. This is the first time to study the flavor profile of Chinese baijiu.

Omission and addition experiments

To confirm the contributions of some key odorants to the flavor of Meilanchun, a total of 17 aroma omission and addition experiments were carried out, and each model was evaluated by a triangle test with the above complete recombination model. The results of omission and addition experiments were shown in Table 3.

The data in Table 3 showed there were very highly significant differences ($\alpha \leq 0.001$) in the aroma of these omission models with comparison to the complete recombine, such as esters, ethyl hexanoate, phenols, methional, and highly significant differences ($\alpha \leq 0.01$) in the omission models of ethyl butanoate, acids, and dimethyl trisulfide. However, no significant difference was observed when ethyl lactate was omitted, or methionol was added. The results revealed that esters, phenols and acids were very important for the aroma of Meilanchun, especially ethyl hexanoate, methional, ethyl butanoate, and dimethyl trisulfide. The esters were mainly responsible for the typical fruity note of Meilanchun, but ethyl lactate has a little influence in spite of its relatively high level in Meilanchun, which were traditionally regarded as the most key aroma compounds because of their particularly high concentrations.^{28,29} The acids provided the acidic odor, and methional had an important contribution to the roasted note, and methionol contributed little to the Meilanchun aroma, which were in agreement with the reported by Zheng *et al.*¹⁰ These phenols, such as guaiacol, 4-methylphenol and 4-ethyl guaiacol, were identified as key odorants at the first time for the roasted sesame flavor style baijiu.

The two model mixtures without 3-methylbutanal or 3-methyl-1-butanol were also evaluated with significant differences ($\alpha \leq 0.05$) in comparison to the complete reconstitution model. Besides, when the two compounds were omitted, the intensity of malty aroma decreased significantly, while the roasted note decreased slightly. Therefore, they were responsible for the malty aroma and part of roasted note in the entire model mixture.

As shown in Table 3, the omission of lactic acid or acetoin resulted in a significant difference ($\alpha \leq 0.05$). When the mixture model was lack of lactic acid or acetoin, their aroma profiles had some changes in the softness or harmony. Therefore, lactic acid and acetoin (a mild creamy odor) could be as two blenders for the aroma of Meilanchun, since the creamy odor has good compatible with other odor note. This is the first time to confirm the effects of lactic acid and acetoin on the flavor of roasted sesame flavor baijiu.

In addition, the omission of phenethyl alcohol or compounds without odor threshold resulted in an




Fig. 1 Aroma descriptive profiles of Meilanchun sesame flavor style baijiu and its simulation sample with compounds of which OAVs ≥ 1 (M1).



Table 3 Omission and addition experiments from complete recombine

No.	Compounds	Category ^a	n ^b	Significance ^c
1-1	Esters	—	10	***
1-2	Ethyl hexanoate	—	9	***
1-3	Ethyl butanoate	—	8	**
1-4	Ethyl 2-methylpropanoate, ethyl 3-methylbutanoate, ethyl pentanoate, ethyl octanoate	—	7	**
1-5	Ethyl lactate	—	3	
2-1	Acids	—	8	**
2-2	Acetic acid	—	4	
2-3	Lactic acid	—	7	*
3	3-Methyl-1-butanol	—	8	*
4	3-Methylbutanal	—	7	*
5	Phenols	—	9	***
6	Dimethyl trisulfide	—	8	**
7	Methional	—	9	***
8	Acetoin	—	7	*
9	Methionol	+	5	
10	Phenethyl alcohol	—	5	
11	Compounds (odor threshold unavailable)	—	3	

^a “—”, The compounds were omitted; “+”, the compounds were added. ^b Number of correct judgments from 10 assessors evaluating the aroma difference by triangle test. ^c Significance: *, significant ($\alpha \leq 0.05$); **, highly significant ($\alpha \leq 0.01$); ***, very highly significant ($\alpha \leq 0.001$).

insignificant difference from the complete reconstitution. Although phenethyl alcohol had a very high FD value, it had a little contribution to the aroma of Meilanchun because of its low concentration. This was different from the results of the roasted sesame flavor baijiu reported by Zheng *et al.*¹⁰ and Sha *et al.*⁹

In conclusion, 92 compounds were identified from Meilanchun, 47 aroma-active compounds were detected by GC-MS/O, including 45 identified aroma-active compounds and 2 unknown peaks, but 2-furfurylthiol with the roasted sesame aroma was not found. In addition, 43 active odorants and 47 components without being smelt by GC-MS/O were accurately quantified by GC-MS. Meanwhile lactic acid was quantified by HPLC. There were 35 important odorants with OAVs more than 1. The aroma profile of Meilanchun sesame flavor style baijiu were successfully reconstituted by mixing 35 odorants with OAVs ≥ 1.0 , phenethyl alcohol, 2-methoxy-1,3-dioxolane, as well as lactic acid. The omission experiments further confirmed that ethyl hexanoate, phenols, methional were the key odorants making contributions to the overall aroma of Meilanchun baijiu, and ethyl butanoate, 3-methyl-1-butanol, 3-methylbutanal, acetoin and lactic acid also were important compounds to aroma. Methionol and phenethyl alcohol were not the characteristic odorants for Meilanchun sesame aroma-type commercial baijiu. Further researches are needed to identify the function of 2-furfurylthiol for the sesame flavor style baijiu, and explore whether nonvolatile compounds have effects on its aroma, which may explain why the aroma simulation was weak in the roasted sesame aroma in ethanol/water.

Conflicts of interest

The authors declare no competing financial interest.

Abbreviations and nomenclature

AEDA	Aroma extract dilution analysis
OAVs	Odor activity values
GC-MS/	Gas chromatography-mass spectrometry/
O	olfactometry
GC-MS	Gas chromatography-mass spectrometry
IS	Internal standard
AF	Acidic fraction
NBF	Neutral/basic fraction
FD factor	The flavor dilution factor

References

- Q. Yu, The production processes of various flavor baijiu, in *The Brewing Technology of Traditional Baijiu (Liquor)*, China Light Industry Press, Beijing, China, 2013, vol. 17, p. 69.
- B. Sun, J. Wu, M. Huang, J. Sun and F. Zheng, Recent Advances of Flavor Chemistry in Chinese Liquor (Baijiu), *J. Chin. Inst. Food Sci. Technol.*, 2015, **15**, 1–8.
- H. L. Liu and B. G. Sun, Effect of Fermentation Processing on the Flavor of Baijiu, *J. Agric. Food Chem.*, 2018, **66**, 5425–5432.
- C. Liu, M. Liu, Q. Zhong, Z. Xiong, Z. Meng, L. Liu, Z. Lv and X. Li, Study on Typical Sensory Characteristics of Zhimaxiang Baijiu (Sesame-flavor Liquor) by Quantitative Description Analysis, *Liquor-making Science & Technology*, 2014, (06), 10–15.
- C. Gao, Formation and Innovation of Typical Styles of Meilanchun Sesame-flavor Liquor, *Liquor-making Science & Technology*, 2010, (06), 101–107.
- P. Jin, Analysis of aroma components from high quality Meilanchun Baijiu, *Liquor-making Science & Technology*, 1993, (01), 62–65.



7 S. Zhu, C. Gao and G. Cui, Analysis of Trace Compositions of Meilanchun Sesame-flavor Liquor, *Liquor-making Science & Technology*, 2012, (06), 106–110.

8 M. Zhu, W. Fan, Y. Xu and Q. Zhou, 1,1-Diethoxymethane and methanethiol as age markers in Chinese roasted-sesame-like aroma and flavour type liquor, *Eur. Food Res. Technol.*, 2016, 242, 1985–1992.

9 S. Sha, S. Chen, M. Qian, C. Wang and Y. Xu, Characterization of the typical potent odorants in Chinese roasted sesame-like flavor type liquor by headspace solid phase microextraction–aroma extract dilution analysis, with special emphasis on sulfur-containing odorants, *J. Agric. Food Chem.*, 2017, 65, 123–131.

10 Y. Zheng, B. Sun, M. Zhao, F. Zheng, M. Huang, J. Sun, X. Sun and H. Li, Characterization of the key odorants in Chinese Zhima aroma-style Baijiu by gas chromatography–olfactometry, quantitative measurements, aroma recombination, and omission studies, *J. Agric. Food Chem.*, 2016, 64, 5367–5374.

11 J. Sun, D. Zhao, F. Zhang, B. Sun, F. Zheng, M. Huang, X. Sun and H. Li, Joint direct injection and GC–MS chemometric approach for chemical profile and sulfur compounds of sesame-flavor Chinese Baijiu (Chinese liquor), *Eur. Food Res. Technol.*, 2018, 244, 145–160.

12 L. Franitz, M. Granvogl and P. Schieberle, Characterization of the key aroma compounds in two commercial rums by means of the sensomics approach, *J. Agric. Food Chem.*, 2016, 64, 637–645.

13 W. Gao, W. Fan and Y. Xu, Characterization of the key odorants in light aroma type Chinese liquor by gas chromatography–olfactometry, quantitative measurements, aroma recombination, and omission studies, *J. Agric. Food Chem.*, 2014, 62, 5796–5804.

14 M. Czerny, M. Christlbauer, M. Christlbauer, A. Fischer, M. Granvogl, M. Hammer, C. Hartl, N. M. Hernandez and P. Schieberle, Re-investigation on odour thresholds of key food aroma compounds and development of an aroma language based on odour qualities of defined aqueous odorant solutions, *Eur. Food Res. Technol.*, 2008, 228, 265–273.

15 L. Poisson and P. Schieberle, Characterization of the key aroma compounds in an American Bourbon whisky by quantitative measurements, aroma recombination, and omission studies, *J. Agric. Food Chem.*, 2008, 56, 5820–5826.

16 GB/T 10345-2007. Method of analysis for Chinese spirits.

17 G. Jellinek, *Sensory Evaluation of Food. Theory and Practice*, Ellis Horwood Ltd., Chichester, UK, 1985.

18 P. P. Wang, Z. Li, T. T. Qi, X. J. Li and S. Y. Pan, Development of a method for identification and accurate quantitation of aroma compounds in Chinese Daohuaxiang liquors based on SPME using a sol-gel fibre, *Food Chem.*, 2015, 169, 230–240.

19 P. Schieberle, New Developments in Methods for Analysis of Volatile Flavor Compounds and Their Precursors, in *Characterization of Food: Emerging Methods*, ed. A. G. Goankar, Elsevier, Amsterdam, The Netherlands, 1995, pp. 403–431.

20 W. Grosch, Aroma compounds, in *Handbook of Food Analysis*, ed. L. M. L. Nollet, edn 2nd, Marcel Dekker Inc., New York, 2004, vol. 1, pp. 717–746.

21 W. L. Fan and Y. Xu, Determination of Odor Thresholds of Volatile Aroma Compounds in Baijiu by A Forced-choice Ascending Concentration Series Method of Limits, *Liquor Making*, 2011, 38(04), 80–84.

22 G. A. Burdock, *Fenaroli's Handbook of Flavor Ingredients*, edn 5th, CRC Press, New York, Washington, D.C, 2005: vol. 411, p. 543.

23 H. Fan, W. Fan and Y. Xu, Characterization of key odorants in Chinese Chixiang aroma-type liquor by gas chromatography–olfactometry, quantitative measurements, aroma recombination, and omission studies, *J. Agric. Food Chem.*, 2015, 63, 3660–3668.

24 X. Wang, W. Fan and Y. Xu, Comparison on aroma compounds in Chinese soy sauce and strong aroma type liquors by gas chromatography–olfactometry, chemical quantitative and odor activity values analysis, *Eur. Food Res. Technol.*, 2014, 239, 813–825.

25 Q. Zhou, *Study on flavor compounds of sesame flavor liquor*, Jiangnan University, Wuxi, Jiangsu Province, 2015.

26 H. Tamura, A. Fujita, M. Steinhaus, E. Takahisa, H. Watanabe and P. Schieberle, Identification of Novel Aroma-Active Thiols in Pan-Roasted White Sesame Seeds, *J. Agric. Food Chem.*, 2010, 58, 7368–7375.

27 H. Tamura, A. Fujita, M. Steinhaus, E. Takahisa, H. Watanabe and P. Schieberle, Assessment of the Aroma Impact of Major Odor-Active Thiols in Pan-roasted White Sesame Seeds by Calculation of Odor Activity Values, *J. Agric. Food Chem.*, 2011, 59, 10211–10218.

28 Y. Wang, Effects of the main trace components and its quantity relative ratio relationship of Fen-flavor liquor on the sensory quality, *Liquor-making Science & Technology*, 2004, 3, 27–29.

29 Y. Shen, *Manual of Chinese Liquor Manufactures Technology*, Light Industry Publishing House of China, Beijing, China, 1996.

