# Chemical Science



# **EDGE ARTICLE**

View Article Online
View Journal | View Issue



Cite this: Chem. Sci., 2020, 11, 3672

d All publication charges for this article have been paid for by the Royal Society of Chemistry

Received 20th November 2019 Accepted 5th March 2020

DOI: 10.1039/c9sc05883d

rsc.li/chemical-science

# Diazaphosphinanes as hydride, hydrogen atom, proton or electron donors under transition-metal-free conditions: thermodynamics, kinetics, and synthetic applications†

Jingjing Zhang, a Jin-Dong Yang (10 \*\* and Jin-Pei Cheng \*\* and Jin-Pei

Exploration of new hydrogen donors is in large demand in hydrogenation chemistry. Herein, we developed a new 1,3,2-diazaphosphinane  ${\bf 1a}$ , which can serve as a hydride, hydrogen atom or proton donor without transition-metal mediation. The thermodynamics and kinetics of these three pathways of  ${\bf 1a}$ , together with those of its analog  ${\bf 1b}$ , were investigated in acetonitrile. It is noteworthy that, the reduction potentials ( $E_{\rm red}$ ) of the phosphenium cations  ${\bf 1a}$ -[P]<sup>+</sup> and  ${\bf 1b}$ -[P]<sup>+</sup> are extremely low, being -1.94 and -2.39 V (vs.  ${\rm Fc}^{+/0}$ ), respectively, enabling corresponding phosphinyl radicals to function as neutral superelectron-donors. Kinetic studies revealed an extraordinarily large kinetic isotope effect KIE( ${\bf 1a}$ ) of 31.3 for the hydrogen atom transfer from  ${\bf 1a}$  to the 2,4,6-tri-( ${\it tert}$ -butyl)-phenoxyl radical, implying a tunneling effect. Furthermore, successful applications of these diverse P–H bond energetic parameters in organic syntheses were exemplified, shedding light on more exploitations of these versatile and powerful diazaphosphinane reagents in organic chemistry.

# Introduction

Hydrogen transfer plays a very important role in chemical science and related fields. This process occurs through the cleavage of the targeted R-H bond in three possible pathways (Scheme 1), i.e., hydride (H<sup>-</sup>), hydrogen atom (H<sup>\*</sup>)<sup>2</sup> and proton (H<sup>+</sup>)<sup>3</sup> transfers, which, from a thermodynamic aspect, depends on the R-H bond strength and the nature of the atom bound to the transferred hydrogen atom. Accordingly, knowledge of the relevant thermodynamics of hydricity  $\Delta G_{H^-}$ , 1b,c,4 bonddissociation free-energy, BDFE<sup>5</sup> and acidity,  $pK_a^{3c,6}$  as well as the kinetics<sup>7</sup> relevant to these processes would facilitate rational exploitations of new transformations. In recent years, considerable attention has been paid to the development of new hydrogen sources with versatile hydrogen donor reactivities, with particular interests in transition-metal hydrides M-H,1c such as hydrogenase enzyme analogs8 and hydrogenation catalysts,9 and main-group organic hydrides X-H,1b such as nicotinamide coenzyme models10 and Hantzsch esters,11 and so on.

It is well known that metal hydrides (M-H) could serve as H<sup>-</sup>,

H' and H<sup>+</sup> donors (Scheme 1a). 1c,12 Their diverse hydrogen reactivities originate from the readily modulated polarity of

M-H bonds through electronic communication between the

(b) For cobalt coordinated N-H bonds (transition-metal system)

$$Cp \qquad I^{2^{-}}$$

$$Cp \qquad N^{-}Ph \qquad H^{-} \qquad \Delta G_{H^{-}} = 88.7 \text{ kcal/mol}$$

$$Cp \qquad N^{-}Ph \qquad H^{+} \qquad DK_{a} = 12.1$$

$$Cp \qquad I^{-}$$

$$Cp \qquad N^{-}Ph \qquad H^{+} \qquad DK_{a} = 12.1$$

$$Cp \qquad I^{-}$$

$$Cp \qquad N^{-}Ph \qquad H^{+} \qquad DK_{a} = 12.1$$

$$Cp \qquad I^{-}$$

$$Cp \qquad N^{-}Ph \qquad H^{+} \qquad DK_{a} = 12.1$$

(c) For C-H bonds in triarylmethanes (transition-metal-free system)

$$Ph_{3}C\text{-H} \begin{cases} -HT \rightarrow Ph_{3}C^{+} + H^{-} & \Delta G_{H^{-}} = \sim 100 \text{ kcal/mol} \\ -PT \rightarrow Ph_{3}C^{-} + H^{+} & pK_{a} = \sim 40 \\ -HAT \rightarrow Ph_{3}C^{-} + H^{-} & BDFE = \sim 80 \text{ kcal/mol} \end{cases}$$

Scheme 1 Possible pathways of hydrogen transfers for transition-metal and transition-metal-free systems as well as corresponding thermodynamic driving forces in acetonitrile.

<sup>(</sup>a) For M-H bonds (*transition-metal system*)

M-H  $\begin{cases} \frac{\Delta G_{H^-}}{pK_a} & M^+ + H^- & \text{hydride transfer (HT)} \\ \frac{pK_a}{m} & M^- + H^+ & \text{proton transfer (PT)} \\ \frac{BDE}{m} & M^+ + H^- & \text{hydrogen-atom transfer (HAT)} \end{cases}$ 

<sup>&</sup>lt;sup>a</sup>Center of Basic Molecular Science, Department of Chemistry, Tsinghua University, Beijing 100084, China. E-mail: jdyang@mail.tsinghua.edu.cn; jinpei\_cheng@mail.tsinghua.edu.cn

<sup>&</sup>lt;sup>b</sup>State Key Laboratory of Elemento-Organic Chemistry, College of Chemistry, Nankai University, Tianjin 300071, China

<sup>†</sup> Electronic supplementary information (ESI) available: Details of syntheses and experimental determinations. CCDC 1947293. For ESI and crystallographic data in CIF or other electronic format see DOI: 10.1039/c9sc05883d

Edge Article Chemical Science

metal centers and coordinated ligands. Compared to an M–H system, the X–H hydride manifested some advantages in mild reaction conditions, good functional-group compatibility, easy modification, *etc.* Integration of these three dissociation possibilities into one X–H covalent bond is a substantial challenge, due to the great disparity in the electronegativity of the X and H fragments. Nevertheless, Waymouth *et al.* very recently reported an excellent new example of such a X–H bond where the N–H bond of the phenylazopyridine ligand in the Co(I) complex can serve as either a H<sup>-</sup>, H<sup>+</sup> or a H<sup>+</sup> donor (Scheme 1b).<sup>13</sup> It is noted, however, that the diverse reactivity of this N–H bond may require latent electronic communication between the azopyridine ligand and cobalt atom within the five-membered ring, so it may be viewed as a quasi-M–H system.

For a true transition-metal-free system, till now only triaryl-methane analogs, where there exist steric as well as resonance stabilization effects on the incipient radical or charges, were reported to have  $\alpha$ -C–H bond-cleavage energies determined for all three pathways (Scheme 1c). However, their poor hydrogen donability and severe steric hindrance make these dissociation processes not very useful for synthesis. This, in turn, stimulated our interest to find more useful transition-metal-free X–H systems that promise both the desired energetic measurement and new synthetic applications.

Recently, diazaphospholenium hydrides (P–H) have attracted substantial research interest due to their superior hydricity endowed by the unique diazaphospholene skeleton. Many diazaphospholenes were developed and successfully used in various hydridic reductions. However, the superior hydricity resulted in their other promising applications, such as serving as hydrogen atom and proton donors, as well as the precursors of strong electron donors, being greatly overlooked. Considering the similar electronegativity of the hydrogen ( $\chi^{AR}=2.20$ ) and phosphorus ( $\chi^{AR}=2.06$ ) atoms, the precursors of strong electron donors, as well as the precursors of strong electron donors, being greatly overlooked. Considering the similar electronegativity of the hydrogen ( $\chi^{AR}=2.20$ ) and phosphorus ( $\chi^{AR}=2.06$ ) atoms, the precursors of the P–H bond, beside being a good hydride donor, where the potential to release proton and hydrogen atoms as well, by fine-tuning the electrical properties of the P fragment. And indeed luckily enough, this was realized in the present work.

Herein, we reported a new N-heterocyclic phosphine 1a, which could act as a H<sup>-</sup>, H<sup>+</sup> and H<sup>+</sup> donor (Scheme 2). Thermodynamics and kinetics pertinent to these three processes were examined in detail. For comparison, 1b was investigated as well. Based on the P-H bond energetic studies, we also exploited their versatile applications in the reduction of pyridines, synthesis of bisphosphines, H/D exchange, and activation of carbon-halogen bonds as original examples.

Scheme 2 P–H reagents 1a and 1b serving as diverse hydrogen donors.  $A^+$ ,  $B^-$  and  $C^-$  are the hydride, proton and hydrogen atom acceptors, respectively.

# Results and discussion

#### **Synthesis**

The substrate 1a was prepared via a H/Cl exchange between the corresponding phosphoric chloride and LiAlH4 in THF according to the literature 15,17 (see the ESI† for details). LiAlD4 was used for deriving the deuterated substrates 1a-D and 1b-D. The crystal structure of 1a was shown in Fig. 1 and features an envelope configuration with the P atom standing out of the naphthalene plane and the P-H bond adopting a flagpole position, similar to the five-membered ring analogs disclosed previously. 15a And a P-H distance of 1.42(2) Å is slightly shorter than that of the five-membered ring analog.15 31P NMR spectra of 1a in CD<sub>3</sub>CN show a dt peak at 25.66 ppm that is split by the hydrogen atoms on the P atom and isopropyl groups. It moves to the higher magnetic field in comparison with other Nheterocyclic phosphines.<sup>17</sup> Under an inert atmosphere, 1a is stable enough for over 12 hours in tetrahydrofuran, acetonitrile, alcohols, toluene, dimethylsulfoxide and chloroform. On the other hand, 1b is stable in these solvents as well except in alcohols and chlorinated hydrocarbons, where it decomposes rapidly to yield phosphonites (RO)2PH19 or 2-chloro-1,3,2diazaphosphinane [P]-Cl (Scheme 3), showing that 1b is more air- and moisture-sensitive than 1a.

#### Redox properties

The oxidation potentials of **1a** and **1b** were examined by cyclic voltammetry (CV), exhibiting a partially reversible oxidation peak of **1a** to **1a**<sup>\*\*</sup> at 0.23 V (vs. Fc<sup>\*\*</sup>), Fig. S1a†) and an irreversible oxidation wave of **1b** at 0.47 V (vs. Fc<sup>\*\*</sup>), Fig. S1b†) in acetonitrile. The low oxidation potentials of **1a** and **1b** indicate their good reducing capacity. It is interesting to find that **1a** is a better electron donor, in spite of being a far poorer hydride donor than **1b** (vide infra). Besides, the redox behaviors of the respective phosphinyl radicals (**1a**-[**P**]\* and **1b**-[**P**]\*) were also




Fig. 1 The crystal structure of  $\mathbf{1a}$  (50% probability thermal ellipsoids).<sup>20</sup> Selected bond lengths (in Å): P1-H1 1.42(2), P1-N1 1.747(6), N1-C1 1.329(9), N1-C7 1.440(8), and C1-C6 1.413(8).

$$(RO)_2PH$$

ROH

 $N$ 
 $P-H$ 
 $OH_2Cl_2$ 
 $OHCH_2Cl_3$ 
 $O$ 

Scheme 3 Decomposition of 1b in alcohols and chlorinated hydrocarbons. [P] =  $(CH_2)_3(N^tBu)_2P$ , R = Me or Et.

examined through the reduction of the corresponding phosphenium cations ( $1a-[P]^{+}$  and  $1b-[P]^{+}$ ), resulting in the CV peaks at -1.94 V and -2.39 V (vs. Fc<sup>+/0</sup>) in acetonitrile, respectively (Fig. S1c and 1d†). The extremely low potentials, especially for  $1b-[P]^{+}$ , indicate the very high reducing capacity of the corresponding  $1a-[P]^{+}$  and  $1b-[P]^{+}$  radicals. This suggests that they both have a very promising potential to serve as a super electron donor in organic synthesis to reduce aryl halides.<sup>22</sup> And indeed, this was verified in the tentative application of the present work in radical hydrodebromination of bromobenzene (see Application part, *vide infra*).

#### Thermodynamic driving force and reactivity

**Hydride transfer (HT).** Hydride transfer from **1a** or **1b** <sup>17</sup> to *N*-methylacridinium iodide **A1**<sup>+</sup> yielded equimolar products (Fig. S3†). To exclude the suspected interruption of the hydridic isopropyl α-C-H, the deuterated substrate **1a-D** was employed, and over 95% deuterated **A1D** was derived (eqn (1) and Fig. S4†). This confirmed that the P-H hydride of **1a** is a stronger donor than that of the isopropyl α-C-H.

The hydricity ( $\Delta G_{H^-}$ , defined by eqn (2)) of **1a** and **1b** in acetonitrile was determined by measuring their equilibrium constants with the acceptors of known hydricities (eqn (3)). For example, the measured hydride transfer equilibrium constant  $K_{\rm eq}(1a)$  of 0.48 (equivalent to 0.43 kcal mol<sup>-1</sup>) between 1a and phenanthridinuim trifluoromethanesulfonate  $A2^+$  ( $\Delta G_{H^-}(A2H)$ = 61.4 kcal mol<sup>-1</sup>)<sup>1b</sup> provided the hydricity  $\Delta G_{\rm H^-}$  of 1a to be 61.8 kcal mol<sup>-1</sup> (see Fig. S5 in the ESI† for details) and other equilibria (in Fig. S6 and S7†) verified this value, giving a mean hydricity of  $62.2 \pm 1.0 \text{ kcal mol}^{-1}$ . This demonstrates **1a** to be a moderate hydride donor, similar to the hydride-donating ability of the NADH analogs.16 Thus, it should be capable of reducing commonly used hydride acceptors, such as N-methylacridinium cations ( $\Delta G_{H^-} = 76 \text{ kcal mol}^{-1}$ ), 9-phenylxanthylium cations (89 kcal mol<sup>-1</sup>) and trityl cations (99 kcal mol<sup>-1</sup>). Similarly, a much lower equilibrium constant  $K_{\rm eq}$  (1b) of 1.6  $\times$  10<sup>-3</sup> (equivalent to 3.8 kcal mol<sup>-1</sup>, eqn (5) and Fig. S8†) established between 1b and benzimidazolium perchlorate  $A3^+$  ( $\Delta G_{H^-}(A3H) = 45.0 \text{ kcal mol}^{-1})^{4b}$  resulted in  $\Delta G_{\rm H}$  (1b) being 48.8  $\pm$  1.0 kcal mol<sup>-1</sup>, which is comparable to that of the commonly used good hydride donor, 2,3-dihydrobenzo[d]imidazoles.16 The good hydridic reactivity of 1b was illustrated in our recent work in the reduction of various unsaturated compounds.<sup>17</sup> It is interesting to note that **1b** is about 13 kcal mol<sup>-1</sup> stronger than **1a** in hydricity, but is 0.24 V weaker in electron-donating ability.

$$P - H \xrightarrow{\Delta G_{H^-}} P^+ + H^- \tag{2}$$

$$\Delta G_{\text{rxn}} = -RT \ln K_{\text{eq}} = \Delta G_{\text{H}} - (1) - \Delta G_{\text{H}} - (AH)$$
 (3)

The hydricity can be used to rationalize the disparate reactivity between 1a and some reported diazaphospholenium hydrides. The latter compounds were reported to react with strong acid HBF<sub>4</sub> to give H<sub>2</sub> and the corresponding phosphenium cations. 15b However, when **1a** was treated with HBF<sub>4</sub> (or HOTf), no gaseous product was generated. Instead, a new phosphorus species with a doublet peak at about 80 ppm was detected in the <sup>31</sup>P NMR spectrum (Fig. S9a and S9b†). To identify this newly formed species, a base pyridine was added to the reaction mixture, which rendered 1a the primary product with a small amount of unidentified impurities (Fig. S9c†). Consequently, this mysterious P species was speculated to be protonated 1a (1aH<sup>+</sup>, Scheme 4). Based on the doublet peak of the phosphorus atom in <sup>31</sup>P NMR spectra (if the extraneous proton was connected to the P atom, a triplet 31P peak would be observed) and the unsymmetrical peaks of naphthyl and isopropyl hydrogen atoms in <sup>1</sup>H NMR spectra (Fig. S10†), the protonated site was thus assigned to the N atom (Scheme 4). We thought that their dissimilarity in reactivity between 1a and some reported diazaphospholenium hydrides should originate from the relatively weak hydricity of **1a** ( $\Delta G_{H^-}$  = 62.2 kcal  $\text{mol}^{-1}$ ). As a consequence, it makes the  $\text{H}_2$  release unable to compete with its protonation even under the condition of heating (80 °C for 5 hours). The moderate hydricity of 1a endows it with good tolerance to even strong acids, avoiding direct elimination of dihydrogen. This in turn offers a possibility for 1a to act as a hydride donor in acid-catalyzed reduction. Lots of examples in this connection a can be found, especially in the hydrogenation of imines by hydride donors with comparable hydricity to 1a (~60-70 kcal mol<sup>-1</sup>, like Hantzsch ester,<sup>23</sup> benzothiazoline24 and cyclohexadiene25) under the catalysis of Brønsted acids (for examples BINOL-phosphoric acid, trifluoroacetic acid and Tf<sub>2</sub>NH).

The kinetics of the hydride transfer from 1a to  $A1^+$  was conducted by following the decay of the absorption of  $A1^+$  in

Scheme 4 Protonation of 1a by HBF<sub>4</sub> and recovery of it by adding pyridine in CD<sub>3</sub>CN.

Edge Article Chemical Science

CH<sub>3</sub>CN at 430 nm with a stopped-flow spectrophotometer (Fig. 2, see the ESI† for details). The second-order rate constant  $k_{\rm HT}(1a)$  was found to be 5.94 M<sup>-1</sup> s<sup>-1</sup> (Inset in Fig. 2 and Table S1†). Moreover, the nucleophilicity of 1a can be estimated by the second-order rate constant of the reaction of 1a with A1+. Because 1a has a similar structure with 1b, based on the nucleophile-specific sensitivity parameter  $s_{\rm N}$  of  ${\bf 1b}$  (0.52) and the electrophilicity of A1<sup>+</sup> in our recent work, <sup>17</sup> nucleophilicity N, of 1a is estimated to be 8.64. Similarly, the kinetics of **1a-D** with  $A1^+$   $k_{HT}$ (**1a-D**) was determined to be 2.34 M<sup>-1</sup> s<sup>-1</sup> (Table S2†). These gave a primary kinetic isotope effect KIE  $[k_{\rm HT}(1a)/k_{\rm HT}(1a-D)]$  of 2.5. A comparison with the previously determined  $k_{\rm HT}(1\mathbf{b})$  of  $5.41 \times 10^2 \, {\rm M}^{-1} \, {\rm s}^{-1}$  for the reaction of  $1\mathbf{b}$ with A1+ in our earlier work17 indicates that 1b is about 100 times more reactive than 1a. This is exactly in accordance with the order of their hydricities ( $\Delta G_{H^-}$ ) of **1a** vs. **1b** found here.

**Hydrogen atom transfer (HAT).** To examine the possibility of a HAT reaction from diazaphosphinane substrates **1**, the reactions of **1a** and **1b** with the 2,4,6-tri-*tert*-butyl-phenoxyl radical **O** were conducted, and yielded 2,4,6-tri-*tert*-butylphenol **OH** and the corresponding phosphorus species (eqn (6)) without generation




Fig. 2 Monoexponential decay of the absorbance Abs (at 430 nm) with the time t (s) for the reaction of 1a (3.03  $\times$  10<sup>-3</sup> M) with A1<sup>+</sup> (5.00  $\times$  10<sup>-5</sup> M) in CH<sub>3</sub>CN at 20 °C. Inset: Correlation of  $k_{\rm obs}$  with [1a].

of bisphosphines (compared to the reaction of **1a** or **1b** with AIBN, Table 1). A stoichiometric study found a [1]/[0] ratio of 1:2, consistent with the product analysis (Fig. S11 and S12†).

To evaluate the hydrogen atom donability of diazaphosphinanes (i.e., P–H BDFE), the thermodynamic cycle was applied on the basis of the  $\Delta G_{\rm H^-}$  and  $E_{\rm red}([{\bf P}]^+)$  in hand (Scheme 5 and eqn (7);  $FE_{\rm ox}({\bf H}^-)$  is -26.0 kcal  ${\rm mol}^{-1}{}^{1c}$  in acetonitrile vs.  ${\rm Fc}^{0/+}$ ). <sup>26</sup> Substituting the known values into eqn (7) <sup>26a,27</sup> gives P–H BDFEs of  $80.9 \pm 2.0$  and  $77.9 \pm 2.0$  kcal  ${\rm mol}^{-1}$  for  ${\bf 1a}$  and  ${\bf 1b}$ , respectively, being comparable with those of phenol and thiophenol antioxidants. Therefore, they could transfer hydrogen atoms smoothly to acceptors with X–H BDFEs around or larger than 78 kcal  ${\rm mol}^{-1}$ , such as 2,4,6-tri-tert-butylphenoxyl  ${\bf O}^{\cdot}$  (77.1 kcal  ${\rm mol}^{-1}$ ),  ${\rm CN}({\rm CH}_3)_2{\rm C}^{\cdot}$  (about 88 kcal  ${\rm mol}^{-1}$ ) and phenyl radicals (104.7 kcal  ${\rm mol}^{-1}$ ). <sup>5b,28</sup> Compared to  ${\bf 1b}$ , though the fused aryl groups attenuate the hydricity of  ${\bf 1a}$  by 13 kcal  ${\rm mol}^{-1}$  thermodynamically, they show a negligible effect on the HAT reactivity.

BDFE = 
$$\Delta G_{\text{H}^-} - F(E_{\text{red}}([\mathbf{P}]^+) - E_{\text{ox}}(\mathbf{H}^-)) = \Delta G_{\text{H}^-} - 23.06E_{\text{red}}([\mathbf{P}]^+) - 26.0 \text{ (kcal mol}^{-1})$$
 (7)

The reaction mechanism of **1a** with radical **O'** (eqn (6)) is proposed in Scheme 6 based on the derived thermodynamic data (the reaction of **1b** with **O'** is similar). Since hydrogen atom transfer from **1a** to **O'** is slightly endothermic ( $\sim$ 4 kcal mol<sup>-1</sup>), it implies the possibility for a reversible HAT. This reversible HAT

Table 1 Applications of 1a and 1b in organic syntheses

| Entry | Reactant    | Condition  | Product            | Yield               | Туре   |
|-------|-------------|--|--------------------|---------------------|--------|
| (1)   | CN          | <b>1a-[P]</b> <sup>+</sup> (15 mol%), HBpin (1.5 equiv.), CH₃CN, 80 °C, 36 h | CN<br>N<br>Bpin    | $40\%^a$            | НТ     |
| (2)   | P-H<br>N/Pr | <sup>t</sup> BuOK, CD <sub>3</sub> CN, 20 °C, 10 min                         | P-D<br>N-D<br>1a-D | Quant. <sup>b</sup> | PT     |
| (3)   | P-H         | AIBN (1.5 equiv.), $C_6D_6$ , 80 °C, 3 h                                     | Npp                | >90% <sup>b</sup>   | НАТ    |
| (4)   | Br          | <b>1b</b> (1.5 equiv.), AIBN 15 mol%, toluene-d <sub>8</sub> , 90 °C, 5 h    |                    | >90% <sup>c</sup>   | HAT&ET |

<sup>&</sup>lt;sup>a</sup> Isolated yield. <sup>b</sup> NMR yields determined by the amount of phosphorus species. <sup>c</sup> NMR yields with 1,3,5-trimethoxybenzene as the internal standard.

1a or 1b BDFE 
$$[P]' + H'$$

$$AG_{H}$$

$$F[E_{red}[P]'] \cdot E_{ox}[H']$$

Scheme 5 Thermodynamic cycle for deriving the P-H BDFEs of 1a and 1b.

Scheme 6 The possible mechanism for the reaction of 1a with  $0^{\circ}$ . Ar = 2,4,6-tri-tert-butyl-phenyl group.

could be further driven to the far right by the largely exothermic electron transfer, ET of  $\mathbf{1a}$ -[P]' radical  $(E_{\rm ox}(\mathbf{1a}$ -[P]') = -1.94 V) or a follow-up radical coupling. The observed stoichiometric ratio of 1:2 for the reactants [1] vs. [O'] is in exact accordance with the HAT mechanism proposed. On the other hand, a suspected proton transfer between  $\mathbf{1a}$  and the  $\mathbf{O}$ ' radical can be easily ruled out due to the very weak acidity of  $\mathbf{1a}$  (vide~infra) and strong acidity ( $pK_a$  of  $\sim -3$ )<sup>5b</sup> reported for the phenol radical cations (eqn (8)). The other suspected possibility, *i.e.* the ET path between  $\mathbf{1a}$  and the  $\mathbf{O}$ ' radical ( $E_{\rm red} = -0.70$  V  $^{5b}$ ) can also be excluded by a similar strategy on the basis of their respective thermodynamics determined, that indicates that the ET from  $\mathbf{1a}$  to  $\mathbf{O}$ ' is remarkably uphill ( $\Delta G_{\rm ET} = 21.4$  kcal  $\mathrm{mol}^{-1}$ , eqn (9)).  $^{13,29}$  It is thus safe to conclude that the initial HAT should be the rate-determining step (RDS) for the reaction expressed in eqn (6).

$$o' + 1a = \frac{ET}{21.4 \text{ kcal/mol}} o^- + 1a'^+$$
 (9)

The rate constants of this hydrogen atom transfer from  ${\bf 1a}$  and  ${\bf 1a}$ - ${\bf D}$  to radical  ${\bf O}$  were measured to be a  $k_{\rm HAT}({\bf 1a})$  of  $0.70~{\rm M}^{-1}~{\rm s}^{-1}$  and a  $k_{\rm HAT}({\bf 1a}$ - ${\bf D})$  of  $2.24\times 10^{-2}~{\rm M}^{-1}~{\rm s}^{-1}$ , respectively (Tables S3 and S4†). This intermediately points out an extremely large KIE( ${\bf 1a}$ ) of 31.3, implying a tunneling effect in this HAT process. Such a large KIE has been found in biochemical on and transition-metal systems,  $^{31}$  but seldom observed in organic HAT processes. Similarly, the rate of the analogous HAT of  ${\bf 1b}~k_{\rm HAT}({\bf 1b})$  was determined to be  $0.74~{\rm M}^{-1}~{\rm s}^{-1}$  (Table S5†). The quite close  $k_{\rm HAT}$  for  ${\bf 1a}$  and  ${\bf 1b}$  agrees well with their comparable P–H BDFEs (80.9)

vs. 77.9 kcal mol<sup>-1</sup>, *vide supra*), verifying their similar HAT reactivity. However, the same kinetic study for **1b-D** gave a  $k_{\text{HAT}}(\mathbf{1b-D})$  of 0.12 M<sup>-1</sup> s<sup>-1</sup> (Table S6†), resulting in a primary KIE(**1b**) of 6.2 with no tunneling phenomenon detected.

To understand this discrepancy, temperature-dependent kinetics were further performed (Tables S7–S10†). The ratio of Arrhenius pre-factors  $[A(\mathbf{1a})/A(\mathbf{1a-D})]$  was obtained to be 0.021, and the activation energy difference  $[E_a(\mathbf{1a}) - E_a(\mathbf{1a-D})]$  was -5.53 kcal  $\mathrm{mol}^{-1}$  (Fig. S2 and Table S11†). These kinetic characteristics  $[A(\mathbf{1a})/A(\mathbf{1a-D}) << 1$  and  $E_a(\mathbf{1a}) - E_a(\mathbf{1a-D}) << 0]$  are in fact in line with the tunneling model illustrated by Klinman,  $^{30a}$  which features a "through-the-barrier" transition state near the top of the classical barrier. In the  $\mathbf{1b}$  system, on the other hand, the normal primary KIE of 6.2,  $A(\mathbf{1b})/A(\mathbf{1b-D})$  of 3.39 and  $[E_a(\mathbf{1b}) - E_a(\mathbf{1b-D})]$  of -0.38 kcal  $\mathrm{mol}^{-1}$  all support a classical "over-thebarrier" transition state. A more comprehensive investigation of the tunneling issue is presently underway.

**Proton transfer (PT).** It's known that a good hydride donor usually exhibits poor acidity. This explains why the acidity of widely-used X-H type hydride donors, e.g. 2,3-dihydrobenzo[d]imidazoles, has never been determined in the absence of metalcoordination. The hydricity of the present N-heterocyclic phosphines is also quite good (vide supra), hence an investigation of their acidic reactivity cannot be expected to be too straightforward. Nevertheless, as we learned from the initial failures in deprotonating 1a or 1b with some ordinary strong bases, such as 1,8-diazabicyclo[5,4,0]-7-undecene (DBU, p $K_a$  = 24.34 in CH<sub>3</sub>CN), 1,3,4,6,7,8-hexahydro-2*H*-pyrimido[1,2-a] pyrimidine (TBD,  $pK_a = 26.03$ ) and (tert-butylimino) tris(pyrrolidino)-phosphorane (BTPP,  $pK_a = 28.42$ )<sup>28</sup> (see the ESI† for details), we finally found that the H/D exchange of the 1a P-H proton in the presence of stoichiometric <sup>t</sup>BuOK in CD<sub>3</sub>CN (eqn (10) and Fig. S13†) was accomplished in about 10 minutes at room temperature,32 whereas the same operation for 1a-D in CH<sub>3</sub>CN was finished in 8 hours at room temperature (eqn (11) and Fig. S14†). The reversible H/D exchange of 1a in acetonitrile under the promotion of <sup>t</sup>BuOK suggested that the acidity of 1a reaches the measurable acidity limitation in acetonitrile (see the ESI† for details).33 As for 1b with much stronger hydricity, its acidity is, unfortunately, too weak to examine in solution by any existing experiment. However, it is worth noting that the acidity of a P-H bond can be significantly enhanced upon activation by coordination with transition metals34 or other Lewis acids.35

Edge Article Chemical Science

#### Applications in syntheses

Based on the above thermodynamic and kinetic investigations, the potential of applying **1a** and **1b** in organic syntheses, encompassing the diverse hydrogen (H<sup>+</sup>, H<sup>+</sup> and H<sup>-</sup>) or electron transfer, was preliminarily explored and showcased herein.

Catalytic reduction of pyridines is a prevailing strategy for the synthesis of dihydropyridines. Recently, Kinjo *et al.* <sup>16g</sup> *and* Speed *et al.* <sup>36</sup> exploited 1,3,2-diazaphosphenium-catalyzed hydroboration of pyridines with good regio- and chemo-selectivity. <sup>16g</sup> However, in their system, the substrate 3-CN-pyridine gave a mixture of three dihydropyridine isomers. To our delight, we found that under the catalysis of 15 mol% of our 1a-[P]<sup>+</sup>, <sup>37</sup> the sole 1,4-hydroborated isomer of this reaction was selectively produced in a moderate yield of 40% (entry 1 in Table 1 and Fig. S15†). The good regio-selectivity may be ascribed to the steric factors of the isopropyl group on the nitrogen atom of 1a based on Kinjo's DFT calculations. <sup>16g</sup> This good regio-selectivity exhibits the catalytic potential of 1a in the reduction of unsaturated compounds.

Moreover, the acidic character of 1a ( $vide\ supra$ ) offers a more economic method for the synthesis of deuterated reagent 1a-D (entry 2 in Table 1) by employing the much cheaper deuterated solvent  $CD_3CN$  instead of expensive  $LiAlD_4$  in an almost quantitative yield (see the ESI† for details about the synthetic method). Meanwhile, it also provides a new approach for isotope labeling of many other hydridic species. The generation of the hydridic hydrogen from a "protic" solvent as illustrated here may serve as an example of Umpolung which was also proposed by Gudat  $et\ al$ . for the P–H bond.  $^{15a}$ 

Since bisphosphines can easily decompose into phosphinyl radicals,  $^{38}$  they could serve as radical reservoirs for relevant studies and applications. Based on thermodynamic analysis,  $CN(CH_3)_2C^*$  (C–H BDFE  $\approx 85$  kcal  $mol^{-1}$ ), generated from the homolysis of azo-bis-isobutyronitrile (AIBN), could abstract the hydrogen atoms from  $\mathbf{1a}$  (P–H BDFE = 80.9 kcal  $mol^{-1}$ ). Then, the generated phosphinyl radical  $\mathbf{1a}$ -[P]\* coupled quickly with each other to yield bisphosphines (entry 3 in Table 1, Fig. S16† for  $\mathbf{1a}$  and Fig. S17† for  $\mathbf{1b}$ ). The targeted dimers  $^{39}$  are readily extracted from reaction mixtures in a high yield. This transformation, together with Gudat's photochemical dehydrocoupling,  $^{18}$  offers easy access to bisphosphines, avoiding complicated and harsh operations and workups required in other synthetic processes using metals such as sodium  $^{38a-c}$ .

Furthermore, we attempted to design a new radical transformation by an integrated use of hydrogen atoms and electron transfers from N-heterocyclic phosphines. This was inspired by the reactions of AIBN with  ${\bf 1a}$  and  ${\bf 1b}$ , where phosphinyl radicals can be easily generated from their P–H precursors through hydrogen atom release. The highly negative oxidation potentials of phosphinyl radicals imply their competency in the reduction of challenging substrates. If a substrate in the reaction system is sufficiently oxidative, its single-electron capture from phosphinyl radicals may be able to compete with radical coupling. In such cases, subsequent transformations may be triggered. Based on the redox potentials of  ${\bf 1b}$ - $[{\bf P}]$  ( $E_{\rm ox}=-2.39~{\rm V}~vs.~{\rm Fc}^{+/0},$ 

the same with the  $E_{\rm red}({\bf 1b} - [{\bf P}]^+))$  and bromobenzene ( $E_{\rm red} = -2.8$  V), <sup>40</sup> hydrodehalogenation of bromobenzene by **1b** was performed in toluene with 15 mol% AIBN as the initiator (entry 4 in Table 1, Fig. S18†). After 5 hours, the expected reduction product, benzene was obtained in over 90% yield. Due to the poor reduction ability of **1a**-[**P**] $^+$ , the reaction could not proceed when **1a** was used. <sup>41</sup> Therefore, it could be anticipated that these N-heterocyclic phosphines may open up a promising avenue for the development of super electron donors.

# Conclusions

Hydrogen and electron transfers from phosphines 1a and 1b were thermodynamically and kinetically investigated in acetonitrile. The P-H bond of 1a can participate in reactions involving formal transfer of a hydride/hydrogen atom/proton to a substrate under respective conditions in the absence of the usually required transition-metal- or Lewis acid-mediation. It was observed that 1a is a moderate hydride and hydrogen atom donor, but a poor Brønsted acid. In comparison, 1b is a good hydride donor and a moderate hydrogen atom donor; but did not show a detectable acidic reactivity even under the strongest basic solution. With regard to electron donability,  $\mathbf{1a}$  ( $E_{ox}(\mathbf{1a}) =$ 0.23 V) is superior to **1b** ( $E_{ox}(\mathbf{1b}) = 0.47$  V), contrasting with the order of their hydridic reactivity. Their corresponding phosphinyl radicals are very strong electron donors as reflected by their extremely negative redox potentials. Kinetic studies revealed a tunneling contribution (KIE = 31.3) to the hydrogen atom transfer from 1a to the phenoxy radical. Based on the derived energetic parameters, their synthetic applications were tentatively exploited. Preliminary results from the synthesis attempts by using the new reagents 1a and 1b confirmed their competence as effective hydrogen and electron donors in various redox transformations. The effectiveness of applying the relevant physicochemical parameters in the analysis and design of hydrogen transfer reactions was also shown.

### Conflicts of interest

There are no conflicts to declare.

# Acknowledgements

We are grateful for the financial grants from the National Natural Science Foundation of China (No. 21973052, 21933008, 21602116, and 91745101), the National Science & Technology Fundamental Resource Investigation Program of China (No. 2018FY201200), and the Tsinghua University Initiative Scientific Research Program (No. 20181080083).

## Notes and references

(a) J.-D. Yang, J. Xue and J.-P. Cheng, *Chem. Soc. Rev.*, 2019,
 48, 2913–2926; (b) S. Ilic, A. Alherz, C. B. Musgrave and K. D. Glusac, *Chem. Soc. Rev.*, 2018, 47, 2809–2836; (c) E. S. Wiedner, M. B. Chambers, C. L. Pitman,

- R. M. Bullock, A. J. M. Miller and A. M. Appel, *Chem. Rev.*, 2016, **116**, 8655–8692.
- 2 (a) A. Wu and J. M. Mayer, *J. Am. Chem. Soc.*, 2008, **130**, 14745–14754; (b) T. Matsuo and J. M. Mayer, *Inorg. Chem.*, 2005, **44**, 2150–2158; (c) J. M. Mayer, *Acc. Chem. Res.*, 2011, **44**, 36–46.
- 3 (a) Z. Wang, F. Gao, P. Ji and J.-P. Cheng, Chem. Sci., 2018, 9, 3538–3543; (b) K. Kaupmees, N. Tolstoluzhsky, S. Raja, M. Rueping and I. Leito, Angew. Chem., Int. Ed., 2013, 52, 11569–11572; (c) J.-D. Yang, P. Ji, X.-S. Xue and J.-P. Cheng, J. Am. Chem. Soc., 2018, 140, 8611–8623.
- 4 (a) S. Ilic, U. P. Kadel, Y. Basdogan, J. A. Keith and K. D. Glusac, *J. Am. Chem. Soc.*, 2018, 140, 4569–4579; (b)
  X.-Q. Zhu, M.-T. Zhang, A. Yu, C.-H. Wang and J.-P. Cheng, *J. Am. Chem. Soc.*, 2008, 130, 2501–2516.
- (a) M. J. Bezdek and P. J. Chirik, Angew. Chem., Int. Ed., 2018, 57, 2224–2228; (b) J. J. Warren, T. A. Tronic and J. M. Mayer, Chem. Rev., 2010, 110, 6961–7001; (c) G. Ali, P. E. VanNatta, D. A. Ramirez, K. M. Light and M. T. Kieber-Emmons, J. Am. Chem. Soc., 2017, 139, 18448–18451.
- 6 (a) Z. Li, X. Li and J.-P. Cheng, J. Org. Chem., 2017, 82, 9675–9681; (b) C. Mao, Z. Wang, Z. Wang, P. Ji and J.-P. Cheng, J. Am. Chem. Soc., 2016, 138, 5523–5526.
- 7 (a) D. Richter and H. Mayr, Angew. Chem., Int. Ed., 2009, 48, 1958–1961; (b) V. Morozova, P. Mayer and G. Berionni, Angew. Chem., Int. Ed., 2015, 54, 14508–14512; (c) M. Horn, L. H. Schappele, G. Lang-Wittkowski, H. Mayr and A. R. Ofial, Chem.–Eur. J., 2013, 19, 249–263; (d) H. Mayr, G. Lang and A. R. Ofial, J. Am. Chem. Soc., 2002, 124, 4076–4083.
- 8 (a) R. H. Morris, Chem. Rev., 2016, 116, 8588-8654; (b)
  M. E. Ahmed, S. Chattopadhyay, L. Wang, D. Brazzolotto,
  D. Pramanik, D. Aldakov, J. Fize, A. Morozan, M. Gennari,
  C. Duboc, A. Dey and V. Artero, Angew. Chem., Int. Ed.,
  2018, 57, 16001-16004.
- 9 (a) R. Zhong, Z. Wei, W. Zhang, S. Liu and Q. Liu, Chem, 2019, 5, 1552–1566; (b) W. Ai, R. Zhong, X. Liu and Q. Liu, Chem. Rev., 2019, 119, 2876–2953.
- (a) Y. Zu, R. J. Shannon and J. Hirst, *J. Am. Chem. Soc.*, 2003,
   6020–6021; (b) A. Bucci, S. Dunn, G. Bellachioma,
   G. M. Rodriguez, C. Zuccaccia, C. Nervi and A. Macchioni,
   ACS Catal., 2017, 7, 7788–7796.
- (a) Q.-A. Chen, M.-W. Chen, C.-B. Yu, L. Shi, D.-S. Wang, Y. Yang and Y.-G. Zhou, J. Am. Chem. Soc., 2011, 133, 16432–16435;
   (b) V. N. Wakchaure, P. S. J. Kaib, M. Leutzsch and B. List, Angew. Chem., Int. Ed., 2015, 54, 11852–11856.
- 12 (a) D. E. Berning, B. C. Noll and D. L. DuBois, J. Am. Chem. Soc., 1999, 121, 11432–11447; (b) J. R. Norton and J. Sowa, Chem. Rev., 2016, 116, 8315–8317.
- 13 E. A. McLoughlin, K. M. Waldie, S. Ramakrishnan and R. M. Waymouth, *J. Am. Chem. Soc.*, 2018, **140**, 13233–13241.
- 14 (a) W. S. Matthews, J. E. Bares, J. E. Bartmess, F. G. Bordwell,
  F. J. Cornforth, G. E. Drucker, Z. Margolin, R. J. McCallum,
  G. J. McCollum and N. R. Vanier, *J. Am. Chem. Soc.*, 1975,
  97, 7006–7014; (b) X.-M. Zhang, J. W. Bruno and
  E. Enyinnaya, *J. Org. Chem.*, 1998, 63, 4671–4678; (c)

- M. Finn, R. Friedline, N. K. Suleman, C. J. Wohl and J. M. Tanko, *J. Am. Chem. Soc.*, 2004, **126**, 7578–7584; (*d*) K. Xia, G.-B. Shen and X.-Q. Zhu, *Org. Biomol. Chem.*, 2015, **13**, 6255–6268.
- 15 (a) D. Gudat, A. Haghverdi and M. Nieger, Angew. Chem., Int. Ed., 2000, 39, 3084–3086; (b) S. Burck, D. Gudat, M. Nieger and W.-W. Du Mont, J. Am. Chem. Soc., 2006, 128, 3946–3955.
- 16 (a) C. C. Chong, B. Rao and R. Kinjo, ACS Catal., 2017, 7, Adams, C. 5814-5819; (b) M. R. H. B. S. N. Huchenski, M. J. Ferguson and A. W. H. Speed, Angew. Chem., Int. Ed., 2017, 56, 6268-6271; (c) C. C. Chong, H. Hirao and R. Kinjo, Angew. Chem., Int. Ed., 2014, 53, 3342-3346; (d) C. C. Chong, H. Hirao and R. Kinjo, Angew. Chem., Int. Ed., 2015, 54, 190-194; (e) C. C. Chong and R. Kinjo, Angew. Chem., Int. Ed., 2015, 54, 12116-12120; (f) M. R. Adams, C. T. Tien, R. McDonald and A. W. H. Speed, Angew. Chem., Int. Ed., 2017, 56, 16660-16663; (g) B. Rao, C. C. Chong and R. Kinjo, J. Am. Chem. Soc., 2018, 140, 652-656; (h) T. Lundrigan, E. N. Welsh, T. Hynes, C.-H. Tien, M. R. Adams, K. R. Roy, K. N. Robertson and A. W. H. Speed, J. Am. Chem. Soc., 2019, 141, 14083-14088.
- 17 J. Zhang, J.-D. Yang and J.-P. Cheng, *Angew. Chem., Int. Ed.*, 2019, **58**, 5983–5987.
- 18 O. Puntigam, L. Könczöl, L. Nyulászi and D. Gudat, *Angew. Chem.*, *Int. Ed.*, 2015, **54**, 11567–11571.
- 19 R. B. King and P. M. Sundaram, J. Org. Chem., 1984, 49, 1784– 1789.
- 20 A single crystal of **1a** was obtained by volatilization of solution of **1a** in hexane at −30 °C. Due to the air- and moisture- sensitivity of this compound, the single crystal could not be fine sealed when X-ray analyzing in the cool air and the minor compound **1a** (about 10%) was oxidized. And in case of the 10% by product, the obtained crystals are of insufficient quality despite several attempts. Thus the quality of the acquired diffraction data was slightly below the average level (see the attached CIF file for details). Despite this, the present X-ray diffraction data confirmed the chemical structure of **1a** unambiguously.
- 21 H. A. Spinney, I. Korobkov, G. A. DiLabio, G. P. A. Yap and D. S. Richeson, *Organometallics*, 2007, 26, 4972–4982.
- 22 (a) S. S. Hanson, E. Doni, K. T. Traboulsee, G. Coulthard, J. A. Murphy and C. A. Dyker, *Angew. Chem., Int. Ed.*, 2015, 54, 11236–11239; (b) J. A. Murphy, T. A. Khan, S.-z. Zhou, D. W. Thomson and M. Mahesh, *Angew. Chem., Int. Ed.*, 2005, 44, 1356–1360; (c) J. A. Murphy, S.-z. Zhou, D. W. Thomson, F. Schoenebeck, M. Mahesh, S. R. Park, T. Tuttle and L. E. A. Berlouis, *Angew. Chem., Int. Ed.*, 2007, 46, 5178–5183; (d) L. Zhang and L. Jiao, *Chem. Sci.*, 2018, 9, 2711–2722.
- 23 (a) S. Hoffmann, A. M. Seayad and B. List, *Angew. Chem., Int. Ed.*, 2005, 44, 7424–7427; (b) R. I. Storer, D. E. Carrera, Y. Ni and D. W. C. MacMillan, *J. Am. Chem. Soc.*, 2006, 128, 84–86; (c) Q.-A. Chen, K. Gao, Y. Duan, Z.-S. Ye, L. Shi, Y. Yang and Y.-G. Zhou, *J. Am. Chem. Soc.*, 2012, 134, 2442–2448.
- 24 C. Zhu, K. Saito, M. Yamanaka and T. Akiyama, *Acc. Chem. Res.*, 2015, **48**, 388–398.

- 25 (a) R. P. Thummel, W. E. Cravey and D. B. Cantu, J. Org. Chem., 1980, 45, 1633–1637; (b) I. Chatterjee and M. Oestreich, Org. Lett., 2016, 18, 2463–2466.
- 26 (a) V. D. Parker, K. L. Handoo, F. Roness and M. Tilset, J. Am. Chem. Soc., 1991, 113, 7493-7498; (b) M. Tilset and V. D. Parker, J. Am. Chem. Soc., 1989, 111, 6711-6717.
- 27 F. G. Bordwell, J.-P. Cheng and J. A. Harrelson, J. Am. Chem. Soc., 1988, 110, 1229–1231.
- 28 J.-P. Cheng, et al., Internet Bond-energy Databank (iBonD) Home Page, http://ibond.nankai.edu.cn, accessed at Oct. 2019.
- 29 (a) The energetics of the electron transfer (ET) from  $\mathbf{1a}$  ( $E_{\rm ox} = 0.23$  V) to the  $\mathbf{O}$  radical ( $E_{\rm red} = -0.70$  V) shows that its thermodynamic driving force is remarkably uphill ( $\Delta G_{\rm ET} = 21.4$  kcal mol<sup>-1</sup>), rendering its equilibrium constant close to 0. Such an unfavorable thermodynamic driving force prohibits electron transfer to be a feasible path; (b) J.-P. Cheng, Y. Lu, X.-Q. Zhu and L. Mu, *J. Org. Chem.*, 1998, **63**, 6108–6114.
- 30 (a) J. P. Klinman and A. R. Offenbacher, Acc. Chem. Res., 2018, 51, 1966–1974; (b) J. C. Price, E. W. Barr, T. E. Glass, C. Krebs and J. M. Bollinger, J. Am. Chem. Soc., 2003, 125, 13008–13009; (c) J. P. Layfield and S. Hammes-Schiffer, Chem. Rev., 2014, 114, 3466–3494; (d) M. A. Ehudin, D. A. Quist and K. D. Karlin, J. Am. Chem. Soc., 2019, 141, 12558–12569.
- 31 (a) E. J. Klinker, S. Shaik, H. Hirao and L. Que Jr, Angew. Chem., Int. Ed., 2009, 48, 1291–1295; (b) S. H. Bae, X.-X. Li, M. S. Seo, Y.-M. Lee, S. Fukuzumi and W. Nam, J. Am. Chem. Soc., 2019, 141, 7675–7679.
- 32 When the same reaction was performed in MeOH and toluene solutions (with other conditions identical), none of the H/D exchange was observed.
- 33 A. Kütt, S. Selberg, I. Kaljurand, S. Tshepelevitsh, A. Heering, A. Darnell, K. Kaupmees, M. Piirsalu and I. Leito, *Tetrahedron Lett.*, 2018, 59, 3738–3748.
- 34 (a) J.-J. Brunet, R. Chauvin, O. Diallo, B. Donnadieu, J. Jaffart and D. Neibecker, *J. Organomet. Chem.*, 1998, 570, 195–200;
  (b) P. K. Majhi, A. W. Kyri, A. Schmer, G. Schnakenburg and R. Streubel, *Chem.–Eur. J.*, 2016, 22, 15413–15419.
- 35 (a) A. Longeau and P. Knochel, *Tetrahedron Lett.*, 1996, 37, 6099–6102; (b) M. Blum, J. Kappler, S. H. Schlindwein, M. Nieger and D. Gudat, *Dalton Trans.*, 2018, 47, 112–119.
- 36 T. Hynes, E. N. Welsh, R. McDonald, M. J. Ferguson and A. W. H. Speed, *Organometallics*, 2018, 37, 841–844.

- 37 Although phosphenium triflate **1a-[P]**<sup>+</sup> (as hydride acceptor) is used as a catalyst in the hydride transfer reaction (entry 1 in Table 1), generation of P–H **1a** from **1a-[P]**<sup>+</sup> is the initial step of the catalytic cycle. Then, the pyridine is reduced by *in situ* generated **1a**. The hydride donor **1a** and acceptor **1a-[P]**<sup>+</sup> coexist in the reaction system. Their hydride-donating/accepting abilities (hydricity) are crucial in the catalytic cycle.
- 38 (a) O. Puntigam, D. Förster, N. A. Giffin, S. Burck, J. Bender, F. Ehret, A. D. Hendsbee, M. Nieger, J. D. Masuda and D. Gudat, Eur. J. Inorg. Chem., 2013, 2013, 2041-2050; (b) R. Edge, R. J. Less, E. J. L. McInnes, K. Müther, V. Naseri, J. M. Rawson and D. S. Wright, Chem. Commun., 2009, 1691-1693; (c) D. Förster, H. Dilger, F. Ehret, M. Nieger and D. Gudat, Eur. J. Inorg. Chem., 2012, 2012, 3989-3994; (d) N. A. Giffin, A. D. Hendsbee, T. L. Roemmele, M. D. Lumsden, C. C. Pye and J. D. Masuda, Inorg. Chem., 2012, 51, 11837–11850; (e) J.-P. Bezombes, K. B. Borisenko, P. B. Hitchcock, M. F. Lappert, J. E. Nycz, D. W. H. Rankin and H. E. Robertson, Dalton Trans., 2004, 1980-1988; (f) A. Dumitrescu, V. L. Rudzevich, V. D. Romanenko, A. Mari, W. W. Schoeller, D. Bourissou and G. Bertrand, Inorg. Chem., 2004, 43, 6546-6548; (g) M. Blum, O. Puntigam, S. Plebst, F. Ehret, J. Bender, M. Nieger and D. Gudat, Dalton Trans., 2016, 45, 1987-1997.
- 39 E. E. Nifantiev, N. S. Vyazankin, S. F. Sorokina, L. A. Vorobieva, O. A. Vyazankina, D. A. Bravo-Zhivotovsky and A. R. Bekker, J. Organomet. Chem., 1984, 277, 211–225.
- 40 (a) C. Costentin, M. Robert and J.-M. Savéant, J. Am. Chem. Soc., 2004, 126, 16051–16057; (b) L. Pause, M. Robert and J.-M. Savéant, J. Am. Chem. Soc., 1999, 121, 7158–7159.
- 41 DFT calculations showed that  ${\bf 1a}$ -[ ${\bf P}$ ] and  ${\bf 1b}$ -[ ${\bf P}$ ] should have a comparable ability (with an energy difference of 1.3 kcal mol<sup>-1</sup>, see the ESI† for details) in abstracting the bromine atom. This failed to explain the disparate yields of <10% for  ${\bf 1a}$ -[ ${\bf P}$ ] and 90% for  ${\bf 1b}$ -[ ${\bf P}$ ]. Besides, according to the redox potentials of  ${\bf 1b}$ -[ ${\bf P}$ ]  $(E_{\rm ox}=-2.39~{\rm V}~vs.~{\rm Fc}~{\rm in}~{\rm MeCN})$  and bromobenzene ( $E_{\rm red}=-2.8~{\rm V}$ ), the electron transfer from  ${\bf 1b}$ -[ ${\bf P}$ ] to bromobenzene is a feasible reversible process, while that for  ${\bf 1a}$ -[ ${\bf P}$ ]  $(E_{\rm ox}=-1.94~{\rm V})$  is thermodynamically prohibited. These findings preferentially support an ET-initiated mechanism rather than a direct bromine abstraction.