

**An Electrochemically Controlled Release of NHCs Using Iron
Bis(dithiolene) N-heterocyclic Carbene Complexes**

Journal:	<i>Inorganic Chemistry Frontiers</i>
Manuscript ID	QI-RES-06-2020-000638.R2
Article Type:	Research Article
Date Submitted by the Author:	02-Sep-2020
Complete List of Authors:	Selvakumar, Jayaraman ; Wright State University, Department of Chemistry Simpson, Scott; St. Bonaventure University, Chemistry Zurek, Eva; SUNY at Buffalo, Chemistry Arumugam, Kuppuswamy; Wright State University, Department of Chemistry

ARTICLE

An Electrochemically Controlled Release of NHCs Using Iron Bis(dithiolene) N-heterocyclic Carbene Complexes

Jayaraman Selvakumar,^a Scott M. Simpson,^b Eva Zurek,^c and Kuppaswamy Arumugam^{a*}

Received 00th January 20xx,
Accepted 00th January 20xx

DOI: 10.1039/x0xx00000x

A series of five coordinated iron bis(dithiolene) complexes $[\text{Fe}(\text{NHC})(\text{S}_2\text{C}_2\text{R}_2)_2]$ ($\text{R} = \text{C}_6\text{H}_5$ or $\text{C}_6\text{H}_4\text{-}p\text{-OCH}_3$) containing N-heterocyclic carbene (NHC) (NHC = 1,3-bis(2,4,6-trimethylphenyl)imidazol-2-ylidene or 1,3-bis(2,4,6-trimethylphenyl)-4,5-dihydroimidazol-2-ylidene) were isolated in high yield (84–92%). The iron complexes were characterized by NMR spectroscopy and confirmed by single crystal X-ray diffraction studies. The combination of cyclic voltammetry and spectroelectrochemical analysis revealed that iron complexes undergoes $\text{Fe} \rightarrow \text{C}_{\text{NHC}}$ bond cleavage and releases NHC upon subjection to electrochemical reduction. The electrochemically released NHC was trapped using 1-naphthylisothiocyanate and the adduct was isolated in nearly quantitative yield (~99%). As a proof of concept, the electrochemically released NHC was subsequently used as a catalyst for synthesis of γ -butyrolactones from commercially available cinnamaldehydes.

Introduction

Since their discovery,¹ transition metal dithiolene complexes have been an active area of research. Intriguing structural characteristics² coupled with interesting electrochemical properties² makes this class of metal complexes attractive for a variety of potential applications,³ including sensing,^{3h} photophysical,^{3a, 3d} superconducting,^{3c, 3f} and hydrogen evolution catalysts.^{3b, 4} In general, a dithiolene ligand can exist in three different oxidation states, a fully reduced ene-1,2-dithiolate ($\text{S}_2\text{C}_2\text{R}_2^{2-}$) or a radical monoanion ($\text{S}_2\text{C}_2\text{R}_2^{\cdot-}$) or a α -dithione ($\text{S}_2\text{C}_2\text{R}_2$) (Figure 1). When coordinated to a transition metal atom, the ligand may exist in either one of these oxidation states, thus leaving the metal oxidation state ambiguous.^{1b, 2a} The redox non-innocent character of the dithiolene ligand allows it to adopt multiple redox states without affecting the oxidation state of the central metal atom.^{2a, 2b} This distinctive feature assists the metal complexes in ligand based reactivity, i.e., reactions with alkenes.^{3g, 5}

Of all metal dithiolene complexes, dimeric iron bis(dithiolene) complexes $[\text{Fe}(\text{S}_2\text{C}_2\text{R}_2)_2]_2$ are among the most studied.^{2, 6} Dimeric iron bis(dithiolene) complexes readily react with Lewis bases such as tri-alkyl/aryl phosphines,⁷ phosphites,⁸ nitrosyl,⁹ pyridines,¹⁰ and halides,¹¹ to yield the corresponding five

coordinate square pyramidal adducts $[\text{Fe}(\text{X})(\text{S}_2\text{C}_2\text{R}_2)_2]$ ($\text{X} = \text{PR}_3,$




Figure 1. Redox states of dithiolene ligands.

$\text{P}(\text{OR})_3$, NO, pyridine, and I). The appealing feature of five coordinated metal complexes, especially, the phosphine and the phosphite adducts is that they undergo reversible release and binding of phosphine/phosphite when subject to electrochemical reduction and oxidation cycle. Donahue and coworkers have elegantly demonstrated an electrochemical release and recovery of triphenylphosphine with $[\text{Fe}(\text{PPh}_3)(\text{S}_2\text{C}_2(\text{C}_6\text{H}_4\text{-}p\text{-OCH}_3)_2)_2]$ (Figure 2a).^{7b} Similarly, Wiegardt and coworkers has also observed an electrochemical release of phosphite with five coordinated




Figure 2. Electrochemically controlled release and binding of phosphine, phosphite and NHC ligands to iron bis(dithiolene) complexes.


^a Department of Chemistry, Wright State University, 3640 Colonel Glenn Hwy. Dayton, OH 45435 USA.

^b Department of Chemistry, St. Bonaventure University, 3261 West State Road, St. Bonaventure, NY 14778 USA.

^c Department of Chemistry, State University of New York at Buffalo, Buffalo, NY 14260, USA.

† Footnotes relating to the title and/or authors should appear here.

Electronic Supplementary Information (ESI) available: See DOI: 10.1039/x0xx00000x



Scheme 1. Synthesis of iron bis(dithiolene) N-heterocyclic carbene adducts **5-8**.

[Fe(P(OPh)₃)(S₂C₂Ph₂)₂] when subjected to electrochemical reduction.^{8d} The binding characteristics of an exogenous ligand (PPh₃ or P(OPh)₃) as a function of the redox state of the host ([Fe(S₂C₂R₂)₂]) could potentially be exploited to design an “on-off” switch. If the exogenous ligand is a catalyst, the strategy may be used to control the activity of an organic transformation. Relevant to this context, we envisioned to develop [Fe(NHC)(S₂C₂R₂)₂] type iron complexes to facilitate the release of NHC as a function of the redox state of the metal bis(dithiolene) units (Figure 2b).

N-heterocyclic carbenes¹² are one of the most attractive classes of molecules with a variety of applications.¹³ The prime use of this ligand is in homogeneous catalysis, such as olefin metathesis,¹⁴ C–C cross-coupling reactions,¹⁵ transfer hydrogenation,¹⁶ and C–H activation reactions.¹⁷ A number of transition metal complexes containing NHC ligands also exhibit potential therapeutic value.¹⁸ In addition, free NHCs are also used as organocatalysts for a variety of organic transformations,^{13a, 19} including benzoin condensation reactions,²⁰ Stetter reactions,²¹ umpolung acyanion catalysis,²² annulation reactions,²³ catalysis involving acyl azolium intermediates,²⁴ and zwitterionic ring-opening polymerization reactions.²⁵

Free NHCs are usually generated *in situ* by treatment of imidazolium, pyrazolium or thiazolium salts with strong bases such as NaN(SiMe₃)₂, KNa(SiMe₃)₂, NaH, and KO^tBu.^{13a} Considerable developments have been made to generate NHCs from masked carbene sources such as NHC-carboxylates,²⁶ imidazolidines-pentafluorophenyl,²⁷ NHC-isothiocyanates,²⁸ imidazolidines-trichloromethyl,²⁹ NHC-alcohol adducts and NHC-metal systems.^{14d, 30} Alternatively, NHCs can also be generated via electrochemical reduction of imidazolium salts.^{31, 32} Careful analysis reveals that the usefulness of NHCs in organocatalysis is inherently tied to its ease of generation and recovery from the reaction media. Unfortunately, the methods that hinge on these principles are not common with NHCs. As a proof of concept, herein, we disclose a system that uses a less hazardous approach than traditional chemical methods for the release of NHC from [Fe(NHC)(S₂C₂Ph₂)₂] adducts and its subsequent use in organocatalysis.

Results and discussion

Synthesis and Characterization. Since NHCs are excellent σ-donors with π-back bonding capability, the metal complexes they form are expected to be stable to oxygen and moisture.³³

Hence, we envisaged that like PPh₃, NHCs (**3** and **4**) would readily react with [Fe(S₂C₂R₂)₂] (**1** and **2**) to yield air and moisture stable square pyramidal adducts. The resulting adducts can be subsequently subjected to electrochemical release and recovery studies. The results of these studies are discussed below.

The neutral iron bis(dithiolene) dimers (**1** and **2**)^{1b} and NHCs (**3** and **4**)^{12, 34} were readily prepared according to the known literature procedure. Using the dimers and the NHCs, metal complexes **5-8** were prepared (Scheme 1). Two equiv. of *in situ* generated **3** was treated with 1 equiv. of [Fe(S₂C₂Ph₂)₂] in THF. The resulting mixture was allowed to stir for 2 h at 25 °C to yield **5** in 90% yield. Similarly, two equiv. of **4** was treated with [Fe(S₂C₂Ph₂)₂] to yield **6** in 84% yield. By following the same procedure, **7** and **8** were isolated in 92% and 87% yield, respectively. Compounds **1** and **2** are sparingly soluble in polar organic solvents, while compounds **5-8** were readily soluble in common organic solvents such as THF and CH₂Cl₂.

Compounds **5-8** displayed characteristic ¹H and ¹³C signals corresponding to bis(dithiolene) and NHC units in the NMR spectroscopy. For example, in ¹H NMR, **5** displayed characteristic phenyl hydrogens (20H) as a multiplet between 7.32–7.18 ppm, aromatic C–H (mesityl) hydrogens (4H) as a singlet at 6.96 ppm, and olefinic (imidazole) hydrogens (2H) as a singlet at 6.84 ppm. Two sets of singlets corresponding to mesityl-CH₃ hydrogens (6H and 12H) were observed at 2.36 and 2.00 ppm, respectively. Likewise, compounds **6-8** were all characterized using ¹H NMR spectroscopy. In ¹³C NMR, all four metal complexes displayed a characteristic C_{NHC} peak between 180–183 ppm, confirming the presence of a Fe–C_{NHC} bond.³⁵




Figure 3. Thermal ellipsoid plots of **5-8** drawn at the 50% probability level, hydrogen atoms are omitted for clarity. The plots were rendered using POV-Ray. Selected distances and angles are presented in Table 1.

Crystal Structures. Molecular structure of compounds **5-8** were unambiguously assigned using single-crystal X-ray diffraction studies. The thermal ellipsoid plots of **5-8** are presented in Figure 3 and selected bond parameters are presented in Table 1. Compounds **5**, **6** and **8** crystallized in monoclinic ($C2/c$ or $P2_1/c$) space group, while **7** crystallized in a triclinic ($P-1$) space group. All four compounds **5-8** displayed iron centered distorted square pyramidal geometries with four sulfur atoms occupying the base of a pyramid and the apex occupied by a C_{NHC} atom. The average $Fe-C_{NHC}$ distance for **5-8** was observed to be $1.958[2]_{ave}$ Å, which is consistent with values previously reported for $Fe-C_{NHC}$ ($1.982 - 2.148$ Å).^{35b, 36} To better understand the nature of bonding in **5-8**, the refinement parameters were compared with those of previously reported phosphine iron complexes, $[Fe(X)(S_2C_2Ph_2)_2]$ ($X = PPh_3, P(OPh)_3$).⁷ The average $C-S$ and $S-Fe-S$ angles (deg) for compounds **5**, **6**, **7** and **8**.

respectively, which is consistent with the phosphine/phosphite iron complexes ($C-S = 1.710[2]_{ave}$ Å; $S-C-C-S = 1.3825[2]_{ave}$ Å). Moreover, the direct comparison with the phosphine iron complex strongly suggests that the central iron is divalent and the electronic structure for dithiolene units can be best described as radical monoanion ($S_2C_2Ar_2$)^{•-}.^{7b, 8c, 8d, 37} For all four compounds **5-8**, the iron atom was observed to be ~ 0.46 Å above the mean plane defined by the four sulfur atoms and all four compounds displayed a distorted square pyramidal geometry. The magnitude of distortion from an ideal square pyramidal geometry can be deduced by computing Addison's τ value.³⁸ These values are used to determine whether five-coordinate iron complexes adopt a perfect square pyramidal or a trigonal bipyramidal geometry. For example, for an ideal square pyramidal geometry $\tau = 0$, while for an ideal trigonal bipyramidal geometry $\tau = 1$. The

	5	6	7	8	$[Fe(PPh_3)(S_2C_2(C_6H_4-p-OCH_3)_2)_2]$	$[Fe(P(OPh)_3)(S_2C_2Ph_2)_2]$
$Fe-C_{carbene}$	1.963(3)	1.962(3)	1.968(3)	1.940(3)		
$N-C=N^a$	1.327[8]	1.512[6]	1.333[4]	1.496[6]		
$S-C-C-S^a$	1.381[3]	1.383[3]	1.3845[4]	1.382[5]	1.3825[2]	1.3955[4]
$C-S^a$	1.717[3]	1.718[2]	1.728[3]	1.718[3]	1.710[2]	1.70475[3]
$Fe-S^a$	2.1683[5]	2.1688[5]	2.1712[8]	2.1650[9]	2.1567[8]	2.1576[11]
$S-Fe-S^a$	88.25[2]	87.025[19]	87.825[3]	87.81[3]	88.13[10]	88.46[4]
$N-C-N^a$	103.7[3]	107.4[2]	103.4[2]	107.1[3]		
τ^b	0.22	0.41	0.63	0.60	0.19	0.11

^aAverage values are reported when two or more identical bond parameters are present. ^b τ values are computed as described in $S-C-C-S$ distances in **5-8** are $1.720[3]_{ave}$ Å and $1.381[2]_{ave}$ Å, calculated τ values for **5-8** are presented in Table 1.³⁹ Of all

ARTICLE

the complexes, **5** was the least distorted with $\tau_{\text{M-L}} = 0.22$ and compound **7** was the most distorted with $\tau_{\text{M-L}} = 0.33$. This distortion may be attributed to steric and electronic effects exerted by an NHC unit.⁴⁰

Electrochemistry. To assign formal reduction and oxidation potentials for compounds **5-8**, cyclic voltammetry (CV) was recorded in THF with $[\text{N}(\text{nBu})_4]\text{PF}_6$ as the supporting electrolyte at 25°C using a platinum disc working electrode, platinum wire working electrode and silver wire quasi-reference electrode under N_2 atmosphere. The CVs of **5-8** were very similar and the key electrochemical potentials are summarized in Table 2. As a representative example only compound **5** is discussed here. Compound **5** sustained one oxidation and two reduction waves in the cyclic voltammetry (Figure 4). A reversible oxidation wave was observed at $E_{1/2}^{\text{ox}} = +0.51$ V, yielding 5^{1+} .^{8d} A reversible reduction wave was observed at $E_{1/2}^{\text{red}} = -0.67$ V, ascribed to $\text{5} \rightarrow \text{5}^{1-}$ and an irreversible reduction at $E^3 = -1.97$ V attributed to $\text{5} \rightarrow \text{5}^{2-}$. The irreversibility of the second reduction (E^3) signifies dissociation of compound **5**, which is consistent with breaking of $\text{Fe} \rightarrow \text{C}_{\text{NHC}}$ bond and formation of $[\text{Fe}(\text{S}_2\text{C}_2\text{Ph}_2)_2]^{2-}$.^{7b, 8d, 41} Similar behaviour was reported by Donahue and coworkers and Weighardt and coworkers with $[\text{Fe}(\text{PPh}_3)(\text{S}_2\text{C}_2(\text{C}_6\text{H}_4\text{-}p\text{-OCH}_3)_2)_2]$ and $[\text{Fe}(\text{P}(\text{OPh})_3)(\text{S}_2\text{C}_2\text{Ph}_2)_2]$ in CH_2Cl_2 , respectively. Surprisingly for phosphine/phosphite iron complexes, the groups observed only one irreversible reduction wave in CH_2Cl_2 . However, in a separate report, McCleverty has reported the existence of two reductions for $[\text{Fe}(\text{PPh}_3)(\text{S}_2\text{C}_2(\text{C}_6\text{H}_4\text{-}p\text{-OCH}_3)_2)_2]$ in DMF solvent. This clearly suggests that the solvent plays a critical role in stabilizing the chemical species that are generated during the course of the reduction processes. Moreover, Donahue and Weighardt studies with $[\text{Fe}(\text{PPh}_3)(\text{S}_2\text{C}_2(\text{C}_6\text{H}_4\text{-}p\text{-OCH}_3)_2)_2]$ and




Figure 4. Cyclic voltammogram of **5** with 0.1M $[\text{N}(\text{nBu})_4]\text{PF}_6$ in THF (1 mM), decamethylferrocene was added as an internal standard and the potentials were adjusted to SCE.

$[\text{Fe}(\text{P}(\text{OPh})_3)(\text{S}_2\text{C}_2\text{Ph}_2)_2]$ suggests that the reduction processes is predominantly ligand based, rather metal based. A similar behaviour was observed with compounds **5-8**. Molecular gas-phase calculations (*vide infra*) is in close accordance with ligand-based reduction processes.

Table 2. Electrochemical redox potentials of **5-8** as determined by CV at 25°C (V vs. SCE)

Journal Name

Compound	$E_{1/2}^{\text{ox}}$	$E_{1/2}^{\text{red}}$	E^3
5	0.51 (r)	-0.67 (r)	-1.97 (ir)
6	0.55 (r)	-0.73 (r)	-1.96 (ir)
7	0.38 (r)	-0.74 (r)	-2.04 (ir)
8	0.43 (r)	-0.79 (r)	-2.02 (ir)

Potentials obtained from CV in THF with 0.10 M $[\text{N}(\text{nBu})_4]\text{PF}_6$ and 0.1 mM analyte. The obtained redox potentials were referenced to SCE. See Figure S6, S9-S11 for corresponding cyclic voltammograms of **5-8**. Abbreviations: r = reversible, ir = irreversible, ox = oxidation, red = reduction.

Just like compound **5**, compounds **6-8** displayed analogous electrochemical behaviour with $[\text{N}(\text{nBu})_4]\text{PF}_6$ in THF. Close examination of the oxidation potentials for **5-8** revealed that **7** and **8** are easier to oxidize than **5** and **6** (Table 2). This characteristic is attributed to the electron rich nature of anisyl-substituted dithiolene ligands in **7** and **8**. Subsequently, compounds **5** and **6** were easier to reduce relative to **7** and **8**. Hence, these finding suggests that the electrochemical potentials are highly regulated by the nature of the ligand (Figure S6, S8-S10). Again, these assignments are consistent with those reported for $[\text{Fe}(\text{PPh}_3)(\text{S}_2\text{C}_2(\text{C}_6\text{H}_4\text{-}p\text{-OCH}_3)_2)_2]$ and $[\text{Fe}(\text{P}(\text{OPh})_3)(\text{S}_2\text{C}_2\text{Ph}_2)_2]$ complexes.^{7b, 8d}

During the course of reduction, small peaks were observed upon scanning to negative potentials (around -0.1 and -1 V). These peaks may be attributed to the electrochemical response generated from the formation of $[\text{Fe}(\text{S}_2\text{C}_2\text{Ph}_2)_2]^{2-}$, as a consequence of $\text{Fe} \rightarrow \text{C}_{\text{NHC}}$ bond dissociation.⁴¹ To corroborate this, cyclic voltammetry was performed with a narrow electrochemical window that curtails the second reduction (Figure S6). The absence of small peaks with the newly defined window clearly suggests that these peaks were observed upon scanning to the potential pertaining to the second reduction. To probe the reversible nature of first reduction and confirm that the bond cleavage exclusively occurs after the second reduction, a scan rate dependent study was undertaken (see Figure S7). Either at low scan rate or at higher scan rate, the reversibility of $E_{1/2}^{\text{red}}$ was maintained, indicative of stability of **5** after the first reduction.




Figure 5. Stick models of the (bottom; blue/red) HOMO and (top; orange/green) the LUMO of compound **5**. The geometry displayed on the right/left are aligned with the x-axis/y-axis, respectively. Isovalue of ± 0.05 au.

Density Functional Theory Calculations. Molecular gas phase density functional theory (DFT) calculations were performed on compound **5**. The starting coordinates in each geometry optimization were taken from those experimentally determined for the crystalline compound, and all parameters were relaxed to the nearest stationary point. The experimental NMR spectra of **5** clearly show that the iron complex is diamagnetic ($S = 0$), however there are two ways in which such a state may be achieved: one in which the dithiolene ligand radicals are Pauli coupled, and another in which they are strongly antiferromagnetically coupled localized ligand radicals. We refer to the latter as the $S = 0^*$ state, and it was studied utilizing the broken-symmetry approach and methodology developed by Noodleman.⁴¹⁻⁴² We also considered the $S = 1, 2$ spin states, since their energies relative to the $S = 0/0^*$ states, may be able to help determine the most appropriate density functional to model our system. To these ends the ground state geometry of **5** was optimized with a variety of density functionals, and the resulting distances were compared with those measured for the

Table 3. The binding energy (BE) and Fe-C_{NHC} distance for **5** and its reduced forms as calculated by the BP86, B3LYP*, and TPSS functionals in kcal/mol.

Charge	BP86		B3LYP*		TPSS	
	BE (kcal/mol)	d(Fe-C _{NHC}) Å	BE (kcal/mol)	d(Fe-C _{NHC}) Å	BE (kcal/mol)	d(Fe-C _{NHC}) Å
0	-28.44	1.960	-12.93	2.019	-29.24	1.963
-1	-12.29	1.936	-6.33	2.335	-12.47	1.932
-2	-2.27	1.870	-2.61	2.309	-2.74	1.859

crystal. Detailed results are provided in the Supporting Information (SI). The functionals that were tested include those that employ the

generalized gradient approximation, GGA, (BP86, PBE, PBEsol, revPBE), hybrid functionals (B3LYP, B3LYP*, PBE0), and the TPSS meta-GGA.⁴³ Hartree-Fock type theories systematically favor the high-spin state in Fe(II) transition-metal complexes. As a result, the admixture of exact exchange in standard hybrid functionals, such as B3LYP, tends to over-stabilize the high-spin state. It has therefore been proposed that reducing the coefficient of exact exchange can yield more reliable results, and the resulting functional has been dubbed B3LYP*.^{43h} Nonetheless, Jacobsen and Donahue have found that the BP86 functional provides a more appropriate description of iron bis(dithiolene) complexes than either B3LYP or B3LYP*.⁴¹ We have arrived to a similar conclusion based upon the rationalization contained in the supporting information, as we report our results of our BP86 calculations. We also choose to report our results for the B3LYP* and TPSS functionals, for comparison purposes.

A symmetrized fragment orbital (SFO) analysis was carried out on the $S = 0$ state, using the BP86 functional, to determine the composition of the frontier molecular orbitals (Figure 5). The highest occupied molecular orbital (HOMO) was comprised primarily of the iron's d_{yz} -orbital ($\sim 59\%$) and occupied/unoccupied orbitals localized on the dithiolene ligand (31%). The major contribution to the lowest unoccupied molecular orbital (LUMO) of the iron complex was from the unoccupied LUMO+1 of the distorted dithiolene ligands (66%) and a contribution of 28% from the iron's d_{xz} -orbital. In the previously reported phosphine compound, [Fe(P(CH₃)₃)(S₂C₂(C₆H₄-*p*-OCH₃)₂)₂], the doubly degenerate π^* orbitals of the dithiolene ligand contributed to the LUMO.^{7c} However, in the case of **5**, the degeneracy of the π^* orbitals of the dithiolene is broken because of the distortion of the dithiolene ligand from an ideal square pyramidal geometry.

The Fe-C_{NHC} binding energy (BE) of **5**, along with the reduced forms of **5** ([**5**]⁻¹ & [**5**]⁻²) was calculated with the BP86, B3LYP*, and TPSS functionals. The BE was calculated by:

$$BE = E(\mathbf{5}^{\ominus q}) - E(\mathbf{3}) - E(\text{bis(dithiolene)}^{\ominus q})$$

where $E(\mathbf{5}^{\ominus q})$ is the energy of molecule **5**, $E(\mathbf{3})$ is the energy of the isolated NHC molecule, and $E(\text{bis(dithiolene)}^{\ominus q})$ is the energy of the isolated bis(dithiolene) ligand. The results for each functional are contained in Table 3. In the case of all functionals, the binding energy decreases as the charge on the

iron complex increases. Again, the BE results of the BP86 functional essentially mirror the TPSS results, while the BEs as

calculated with B3LYP* are approximately half of the value obtained using BP86/TPSS for the neutral and the anionic species. It is interesting to note that the Fe-C_{NHC} distance decreases as the bond weakens. This has also been observed for iron bis(dithiolene) phosphine complexes.⁴¹ It is clear from our calculations that Fe-C_{NHC} bond significantly weakens upon first and second reduction, this phenomenon is consistent with our experimental observation.

In order to investigate the nature of the reduction of **5** to [5]¹⁻ to [5]²⁻, we shall discuss their electronic structure in terms of atomic charges. We elect to use the Hirschfeld charges (q) as our charge models over the widely used Mulliken charge analysis as Mulliken charges have been shown to be highly basis set dependent. The results of our analysis are shown in Table S3. The SI contains the iron spin state derived for Mulliken populations for **5**, [5]¹⁻, and [5]²⁻ using the BP86, B3LYP*, and TPSS functionals.

It is apparent comparing the Hirshfeld charges of the reduction of **5** to [5]¹⁻ that the charge of the iron does not significantly change for the BP86 and TPSS functionals. While there is an appreciable amount of change in the charge for the Fe in the B3LYP* calculations, it is overmasked by the change in charge on the bis(dithiolene) ligand. The BP86 and TPSS functionals also find that the change in charge accumulation is localized on the sulfur containing ligand. This trend appears to continue for the dianionic species. It is clear from our calculations that it is not only the iron center that is involved in the reduction, but also the ligands surrounding the iron center. Predominantly, the bis(thiolene) ligand gains a substantial amount of charge during the reduction. One may conclude that this reduction is a ligand-based event.





Figure 6. Electronic absorption spectra recorded during potential bulk electrolysis (-2.25 V vs AgCl) of **5** (**5** → **5**²⁻) in THF with 0.1M [N(nBu)₄][PF₆] as a supporting electrolyte with platinum mesh working electrode. The arrows indicate spectra change during the process.

Electronic Absorption Spectra. To better understand electronic excitations in compounds **5-8**, UV-vis absorption

measurements were carried out in THF. All four compounds **5-8** displayed distinctive absorptions at λ = 681, 672, 697 and 704 nm, respectively (Figure S11), these absorptions were attributed to intra ligand charge transfer transitions (ILCT).^{7b, 8c} Such absorption is completely absent for reduced iron bis(dithiolene) [Fe(S₂C₂R₂)₂]ⁿ⁻ complexes (n = 1 or 2).^{7b} Hence, electrochemically reducing compound **5** followed by concomitant monitoring of its electronic excitations *in situ* will help to verify the breaking of the Fe-C_{NHC} bond. Independently, we treated compound **5** with 1 and 2 equiv. of cobaltocene and analysed its UV-vis absorption spectra (Figure S13). The spectra revealed absence of characteristic ILCT transition at 681 nm for the doubly reduced compound ([Fe(S₂C₂R₂)₂]²⁻).

To ascertain the electronic absorption properties of [5]²⁻, the adduct was subjected to UV-vis spectroelectrochemical measurements (Figure 6). An electrochemical quartz cell fitted with Pt mesh working electrode, Pt wire counter electrode and quasi reference Ag wire was used for the measurement. The potentiostat was held at a constant potential (-2.25 V) and the absorption measurements (400–900 nm) were made using a UV-Vis absorption spectrometer. Compound **5** displayed a strong absorption feature at 681 nm attributed to ILCT. Upon electrochemical reduction, an exponential decrease in the concentration of the 681 nm peak was observed, consistent with a decrease in the concentration of **5**. As evident from Figure 6, during pre-electrolysis, the solution contained only **5**, while during electrolysis, the concentration of **5** gets depleted and the concentration of [5]²⁻ increases. After 300 seconds of electrolysis, significant suppression of the absorbance peak corresponding to ILCT was observed, this indicates disappearance of **5** and appearance of [5]²⁻.⁴¹

Having documented spectroscopic evidence for two-electron reduction of **5** and concomitant dissociation of [5]²⁻ to generate **3**, we turned our focus towards trapping the generated carbene (**3**). We hypothesized that bulk electrolytic reduction of **5** → **5**²⁻ is expected to release free carbenes that can be chemically trapped using isothiocyanates.²⁸ By quantifying the amount of isothiocyanate-**3** adduct formation, the amount of free carbene released during the process can be computed. Bulk electrolysis of **5** was conducted in a two-compartment cell separated by a glass frit at 25°C in 20 mL of THF with 0.1M [N(nBu)₄][PF₆] as the supporting electrolyte using a Pt mesh working electrode, Pt mesh counter electrode, and Ag/AgCl non-aqueous reference electrode (see the SI for more information). Adduct **5** (0.02 mmol) was taken in the working electrode compartment and the reference electrode and counter electrode were immersed in the other compartment. Bulk reduction was performed by holding a constant potential at -2.25 V (vs Ag/AgCl). The initial current during the beginning of the experiment was read at 1mA, as the experiment progressed, the current continue to steadily drop and read near zero as the concentration of **5** completely exhausted. After the passage of 3.81 coulombs of electrons (0.0395 mmol), the relative current read near zero, signifying the endpoint of electrolysis. The resulting electrolytic solution on




Scheme 2. Trapping of electrochemically generated IMES carbene

the cathodic compartment was transferred to a 20 mL vial containing 2 equivalent of 1-naphthyl isothiocyanate and the resulting reaction mixture was stirred for 1 h. Further purification of the reaction mixture yielded the expected adduct **9** in near quantitative yield (Scheme 2).

Organocatalysis with Electrochemically Released NHC. After successfully trapping the electro-generated carbene, we turned our focus towards electrochemically-controlled organocatalysis. Carbene catalyzed conjugate umpolung reaction of an unsaturated aldehyde was recently disclosed by Glorius, such reaction has great utility in synthesizing γ -butyrolactones.^{23, 44} Electrochemically released carbene (**3**) via bulk electrolysis of **5** was transferred into a 20 mL vial containing *trans*-cinnamaldehyde (0.4 mmol) and the resulting mixture was stirred at 25 °C for 24 h (Scheme 3). The progress of the reaction was monitored using thin layer chromatography. At the end of 24 h, the crude mixture was subjected to column chromatography to yield γ -butyrolactone **10** in 34% yield (1:0.15/*cis*:*trans* isomers). Control experiments were conducted without **5** and with **5** and no electrolysis, in either case, the formation of γ -butyrolactone was not observed. Moreover, heating **5** with *trans*-cinnamaldehyde to reflux temperature in THF resulted with no product. These control experiments further confirm that suitable electrochemical potential has to be applied for the release **3**.

To increase the scope of electro-released NHC catalysis, the methodology was expanded to other substituted cinnamaldehydes. Varying substituents on aryl include electron-donating functionalities such as methyl and methoxy and moderately electron withdrawing group such as bromine



Scheme 3. Electrochemically generated carbene catalyzed synthesis of γ -butyrolactones.

were subjected to similar catalytic conditions. Methyl and methoxy substituted aldehydes furnished the respective γ -

butyrolactones **11** in 32% and **12** in 21% yield. Bromo substituted cinnamaldehyde yielded **13** in 24% yield. γ -Butyrolactones **10-12** were isolated as *cis* and *trans* mixture, while **13** was isolated in pure *cis* form. Electrochemical methods employed here use a less hazardous approach than traditional reagent-based chemistry. The benefits of such system include mild reaction conditions, functional group tolerance and scalability. However, such systems also present challenges in product isolation and purification.

Conclusions

To the best of our knowledge, these are the first five coordinated $[\text{Fe}(\text{NHC})(\text{S}_2\text{C}_2\text{R}_2)]$ systems that are prepared by cleaving the $[\text{Fe}(\text{S}_2\text{C}_2\text{Ar}_2)_2]_2$ dimer with N-Heterocyclic carbenes (**3** and **4**). The iron complexes **5-8** were characterized by NMR spectroscopy and confirmed by X-ray crystallography. DFT calculations show that the reduction of **5** weakens the Fe-C_{NHC} bond. The calculations of the neutral and reduced forms of the iron complex demonstrate that the reduction is primarily ligand based. Electrochemical and spectroelectrochemical measurements revealed that $[\text{Fe}(\text{NHC})(\text{S}_2\text{C}_2\text{R}_2)]$ undergo facile electrochemical reduction that leads to dissociation of the adduct to release free NHCs. This behaviour was further confirmed by a trapping **3** with 1-naphthylisothiocyanate. The *in situ* electro-released free NHC (**3**) was used as a catalyst in organocatalysis to convert cinnamaldehydes to γ -butyrolactones. Efforts are underway to fine tune the electronics of the dithiolene system to achieve reversible binding of NHC so that activity of an organic transformation can be effectively controlled and the pre-catalyst can be actively restored. Furthermore, efforts are under way to explore a variety of redox-active NHC systems to expand the scope of $[\text{Fe}(\text{NHC})(\text{S}_2\text{C}_2\text{R}_2)]$ systems for other potential applications, including electronic and material applications.

Conflicts of interest

There are no conflicts to declare.

Experimental Section

Materials and Methods. All synthetic manipulations were done in a nitrogen filled glove box (Inert Purelab) unless otherwise noted. All glassware were oven dried at 120 °C for 12 hours prior to use. The compounds 1,3-bis(2,4,6-trimethylphenyl)imidazolium chloride (**3**·HCl) and 1,3-bis(2,4,6-trimethylphenyl)imidazolium chloride (**4**·HCl) were synthesized according to the literature procedure.^{12, 34} Iron bis(dithiolene) dimers $[\text{Fe}(\text{S}_2\text{C}_2\text{Ph}_2)_2]_2$ and $[\text{Fe}(\text{S}_2\text{C}_2(\text{C}_6\text{H}_4\text{-}p\text{-OCH}_3)_2)]_2$ were prepared following the procedure reported by Schrauzer et. al.^{1b} All other reagents were purchased from commercial sources and used as received, such as sodium bis(trimethylsilyl)amide, *trans*-cinnamaldehyde, and substituted *trans*-cinnamaldehyde. $[\text{N}(n\text{Bu})_4][\text{PF}_6]$ was recrystallized two times from ethanol and dried at 70 °C for 12 hours prior to use. Solvents were dried with a solvent

purification system from Inert Pure Company (THF, CH₂Cl₂, Et₂O and toluene) and degassed using three freeze–pump–thaw cycles prior to use.⁴⁵ All solvents were stored over 4 Å molecular sieves in a nitrogen filled glove box. CDCl₃ (99.9%) was purchased from Acros Laboratories and dried over 4 Å molecular sieves prior to use.

Instrumentation. UV-Vis spectra were obtained at 25 °C with a Varian Cary 50 Bio UV-vis spectrophotometer and molar absorptivities were reported in M⁻¹ cm⁻¹. ¹H and ¹³C NMR spectra were recorded on a Bruker 300 MHz NMR spectrometer. Spectra were referenced to the residual solvent as an internal standard, for ¹H NMR: CDCl₃, 7.26 ppm; for ¹³C NMR: CDCl₃, 77.16. Coupling constants (J) are expressed in hertz (Hz). Elemental analyses were performed by Midwest Microlab, LLC in Indianapolis, IN. Electrochemical measurements were performed on a CHI620E electrochemical workstation using a silver wire quasi-reference electrode, a platinum disk working electrode, and a Pt wire auxiliary electrode in a gas tight three-electrode cell under nitrogen atmosphere. Unless specified otherwise, the measurements were performed using 1.0 mM solutions of the analyte in dry THF with 0.1 M [N(nBu)₄][PF₆] as the electrolyte and decamethylferrocene (Fc*) as the internal standard. Differential pulse voltammetry measurements were performed with 50 mV pulse amplitudes and 4 mV data intervals. All potentials listed herein were determined by cyclic voltammetry at 100 mVs⁻¹ scan rates and referenced to a saturated calomel electrode (SCE) in decamethylferrocene 0/+ to 0.102 V (THF).⁴⁶ Bulk electrolysis were conducted on two compartment cells separated by a glass frit, Pt mesh working electrode, Pt mesh counter electrode and Ag/Ag⁺ (in MeCN) non aqueous reference electrode

Synthesis of compound 5. A 10 mL scintillation vial with a stir bar was charged with [3]·HCl (99 mg, 0.29 mmol), NaN(Si(CH₃)₂)₂ (58 mg mg, 0.31 mmol) and 4 mL of dry THF. The resulting mixture was stirred at 25°C for 45 min, which resulted in a pale-yellow solution with white suspension. The heterogeneous mixture was filtered through a plug of celite into a clean 20 mL scintillation vial containing compound 1 (78 mg, 0.145 mmol) in 2 mL THF. An immediate color change was noticed from black to black green solution. The resulting mixture was stirred at 25°C for 2 h. The black green solution was filtered through a plug of celite and the volatiles were removed under vacuum. The vacuum dried black green residue was dissolved in a minimum amount of dichloromethane and titrated with 15 mL of hexanes resulting with a black green precipitate, which was further washed with hexane and the volatiles were removed under vacuum. 110 mg, 90% yield; ¹H-NMR (300 MHz, CDCl₃): δ 7.31-7.28 (m, 8H), 7.22-7.18 (m, 12H), 6.96 (s, 4H), 6.84 (s, 2H), 2.36 (s, 6H), 2.00 (s, 12H). ¹³C-NMR (75 MHz, CDCl₃): δ 182.18, 143.19, 139.13, 136.82, 135.80, 129.65, 129.31, 127.69, 127.00, 126.77, 21.41, 18.60. HRMS (ESI) for [C₄₉H₄₄FeN₂S₄]⁺[M]⁺ Calcd. 844.1737 Found. 844.1732. Anal. Calcd. for: C₄₉H₄₄FeN₂S₄: C, 69.57, H, 5.36, N, 3.31; Found: C, 69.50, H, 5.60, N, 3.46.

Synthesis of compound 6. A 10 mL scintillation vial with stir bar was charged with [4]·HCl (100 mg, 0.29 mmol), NaN(Si(CH₃)₂)₂ (58 mg mg, 0.31 mmol) and 4 mL of dry THF. The resulting mixture was stirred at 25 °C for 45 mins, which resulted in a yellow solution with white suspension. The heterogeneous mixture was filtered through a plug of celite into a clean 20 mL scintillation vial containing compound 1 (78 mg, 0.145 mmol) in 2 mL THF. An immediate color change was noticed from black to black green solution. The resulting mixture was stirred at 25 °C for 2 h. The black colored solution was filtered through a pad of celite and dried. The dried black residue was dissolved in minimum amount of dichloromethane solvent and the compound was precipitated out using hexane 15 mL resulting in a black green precipitate, which was further washed with hexane and dried under vacuum for 24 h. 103 mg, 84% yield; ¹H-NMR (300 MHz, CDCl₃): δ 7.28-7.26 (m, 8H), 7.23-7.18 (m, 12H), 6.95 (s, 4H), 3.72 (s, 4H), 2.34 (s, 6H), 2.26 (s, 12H). ¹³C-NMR (75 MHz, CDCl₃): δ 181.3, 143.06, 138.20, 137.26, 136.39, 129.62, 127.65, 127.02, 51.97, 21.35, 18.75. HRMS (ESI) for [C₄₉H₄₆FeN₂S₄]⁺[M]⁺ Calcd. 846.1893 Found. 846.1888. Anal. Calcd. for: C₄₉H₄₆FeN₂S₄: C, 69.48, H, 5.47, N, 3.31; Found: C, 68.75, H, 5.41, N, 3.22.

Synthesis of compound 7. A 10 mL scintillation vial with stir bar was charged with [3]·HCl (99 mg, 0.29 mmol), NaN(Si(CH₃)₂)₂ (58 mg mg, 0.31 mmol) and 4 mL of dry THF. The resulting mixture was stirred at 25 °C for 45 mins, which resulted in a yellow solution with white suspension. The heterogeneous mixture was filtered through a plug of celite into a clean 20 mL scintillation vial containing compound 2 (96 mg, 0.145 mmol) in 2 mL THF. An immediate generation of homogenous black solution occurred due to the breaking of dimer and adduct formation. The resulting mixture was stirred at 25 °C for 2 h. The black colored solution was filtered through a pad of celite and dried. The dried dark grey residue was dissolved in a minimum amount of dichloromethane solvent and the compound was precipitated out using 15 mL diethyl ether resulting in a dark grey precipitate, which was further washed with diethyl ether and dried under vacuum for 24 h. 129 mg, 92% yield; ¹H-NMR (300 MHz, CDCl₃): δ 7.27-7.24 (m, 8H), 6.94 (s, 4H), 6.82 (s, 2H), 6.76-6.73 (m, 8H), 3.82 (s, 12 H), 2.36 (s, 6H), 2.00 (s, 12H); ¹³C-NMR (75 MHz, CDCl₃): δ 181.05, 158.68, 138.94, 136.80, 136.51, 135.87, 130.73, 129.17, 126.63, 113.09, 55.33, 21.49, 18.62. HRMS (ESI) for [C₅₃H₅₂FeN₂O₄S₄]⁺[M]⁺ Calcd. 964.2159 Found. 964.2155. Anal. Calcd. for: C₅₃H₅₂FeN₂O₄S₄: C, 65.96, H, 5.43, N, 2.90; Found: C, 65.80, H, 5.46, N, 2.87.

Synthesis of compound 8. A 10 mL scintillation vial with stir bar was charged with [4]·HCl (100 mg, 0.29 mmol), NaN(Si(CH₃)₂)₂ (58 mg mg, 0.31 mmol) and 4 mL of dry THF. The resulting mixture was stirred at 25 °C for 45 mins, which resulted in a yellow solution with white suspension. The heterogeneous mixture was filtered through a plug of celite into a clean 20 mL scintillation vial containing compound 2 (96 mg, 0.145 mmol) in 2 mL THF. An immediate generation of homogenous black solution occurred due to the breaking of

dimer and adduct formation. The resulting black mixture was stirred at 25 °C for 2 h. The black colored solution was filtered through a pad of celite and dried. The dried black residue was dissolved in a minimum amount of dichloromethane solvent and the compound was precipitated out using diethyl ether 15 resulting black precipitate, which was further washed with diethyl ether and dried under vacuum for 24 h. 122 mg, 87% yield; ¹H-NMR (300 MHz, CDCl₃): δ 7.21-7.28 (m, 8H), 6.90 (s, 4H), 6.73-6.70 (m, 8H), 3.79 (s, 12H), 3.64 (s, 4 H), 2.33 (s, 6H), 2.22 (s, 12H). ¹³C-NMR (75 MHz, CDCl₃): δ 180.32, 158.70, 138.03, 137.29, 136.48, 136.39, 130.72, 129.53, 113.06, 55.33, 51.94, 21.42, 18.77. HRMS (ESI) for [C₅₃H₅₄FeN₂O₄S₄]⁺[M]⁺ Calcd. 966.2316 Found. 966.2311. Anal. Calcd. for: C₅₃H₅₄FeN₂O₄S₄: C, 65.82, H, 5.63, N, 2.90; Found: C, 64.85, H, 5.26, N, 3.06.

Computational Details. Spin unrestricted and spin restricted Density Functional Theory (DFT) calculations were carried out using the Amsterdam Density Functional (ADF) software package.⁴⁷ Triple-ζ Slater-type basis set with polarization functions (TZP) from the ADF basis-set library were employed on all atoms. Computations involving the metaGGA TPSS functional utilized the full electron basis sets. All other functionals considered in this study utilized basis functions where the core shells up to 1s for C, N, O, 2p for S, and 3p for Fe were kept frozen. The calculations were conducted constraining the geometry to no symmetry.

DFT calculations must be able to predict the correct ground-state net total spin polarization in order to obtain reliable results. To this end, we have optimized the Fe(pdt)₂(IMes) complex with different spin polarizations utilizing different functionals to confirm that our computational methodology matches the diamagnetic nature of the iron complex as confirmed by our ¹H and ¹³C spectra. Besides iron complexes with ferromagnetically coupled density between the ligands and iron metal, we also considered complexes where the iron center and the ligands are spin-polarized but with opposing spin densities, ie. antiferromagnetically coupled. This was done using the spin-flip method for broken symmetries implemented in ADF and was developed by Noodleman.⁴¹⁻⁴² The SI contains the relative energies (E_{rel}) in kcal/mol for each of the tested functionals. Unsurprisingly, the hybrid functionals predicted the S=1 spin state to be lowest in energy while the GGA functionals predicted the S = 0 state to be the energetic minimum. It was only for the Meta GGA TPSS functional that the antiferromagnetic S = 0 state (referred to as S = 0* in the text) was found to be slight lower in energy than the S = 0 by 0.3 kcal/mol. Based upon a comparison of the results obtained for the iron bis(dithiolene) complexes using the BP86 GGA and the hybrid B3LYP, and B3LYP* functionals, Jacobsen et al. concluded that BP86 is the most appropriate method to describe these systems.⁴⁰ Using the BP86 functional, we have determined that Fe(pdt)₂(IMes) has a spin state of S = 0 and is Pauli coupled. Generally speaking, the mean absolute error (MAE) for the measured distances increased as the number of unpaired electrons in the calculation increased for all of the tested functionals (see the

SI), further supporting the conclusion that the S = 0 state is the preferred spin state. A symmetrized fragment orbitals (SFO) analysis was utilized to investigate the percentage that the molecular orbitals of the constituent moieties contribute to the frontier molecular orbitals of the compound **5**. The NHC molecule, iron atom, and bis(dithiolene) ligand were chosen to be the molecular fragments for the analysis.

Acknowledgements

K.A. acknowledges support by funds from the Army Research Office (Department of Defense; W911NF-16-1-0197) and funds from The American Chemical Society Petroleum Research Fund (PRF-59893-UR7). Support by funds from Chemistry Department, Wright State University, College of Science and Mathematics for the purchase of the X-ray instrument is greatly acknowledged. The authors would like to acknowledge Dr. Gossie, Wright State University and Dr. Fratini, University of Dayton for help with low temperature single crystal X-ray diffraction measurements. S.S. acknowledges the Donors of the American Chemical Society Petroleum Research Fund (PRF-58954-UNI5) and the National Science Foundation (Award #1904825) for support of this research.

Notes and references

- (a) G. N. Schrauzer and V. Mayweg, Reaction of Diphenylacetylene with Nickel Sulfides, *J. Am. Chem. Soc.*, 1962, **84**, 3221-3221; (b) G. N. Schrauzer, V. P. Mayweg, H. W. Finck and W. Heinrich, Coordination Compounds with Delocalized Ground States. Bisdithiodiketone Complexes of Iron and Cobalt1, *J. Am. Chem. Soc.*, 1966, **88**, 4604-4609.
- (a) R. Eisenberg and H. B. Gray, Noninnocence in Metal Complexes: A Dithiolene Dawn, *Inorg. Chem.*, 2011, **50**, 9741-9751; (b) S. Sproules and K. Wieghardt, Dithiolene radicals: Sulfur K-edge X-ray absorption spectroscopy and Harry's intuition, *Coord. Chem. Rev.*, 2011, **255**, 837-860; (c) E. I. Stiefel, *Dithiolene Chemistry: Progress in Inorganic Chemistry* Wiley, New York, 2004.
- (a) A. P. Abbott, P. R. Jenkins and N. S. Khan, Novel complexes with new electro-optic properties, *J. Chem. Soc., Chem. Commun.*, 1994, 1935-1936; (b) H. Lv, T. P. A. Ruberu, V. E. Fleischauer, W. W. Brennessel, M. L. Neidig and R. Eisenberg, Catalytic Light-Driven Generation of Hydrogen from Water by Iron Dithiolene Complexes, *J. Am. Chem. Soc.*, 2016, **138**, 11654-11663; (c) K. L. Marshall, G. Painter, K. Lotito, A. G. Noto and P. Chang, Transition Metal Dithiolene Near-IR Dyes and Their Applications in Liquid Crystal Devices, *Molecular Crystals and Liquid Crystals*, 2006, **454**, 47/[449]-479/[481]; (d) U. T. Mueller-Westerhoff, B. Vance and D. Ihl Yoon, The synthesis of dithiolene dyes with strong near-IR absorption, *Tetrahedron*, 1991, **47**, 909-932; (e) T. Nakajima, Y. Yabushita and I. Tabushi, Amino acid synthesis through biogenetic-type CO₂ fixation, *Nature*, 1975, **256**, 60; (f) N. Robertson and L. Cronin, Metal bis-1,2-dithiolene complexes in conducting or magnetic crystalline assemblies, *Coord. Chem. Rev.*, 2002, **227**, 93-

- 127; (g) K. Wang and E. I. Stiefel, Toward Separation and Purification of Olefins Using Dithiolene Complexes: An Electrochemical Approach, *Science*, 2001, **291**, 106; (h) H. Liu, X. Li, C. Shi, D. Wang, L. Chen, Y. He and J. Zhao, First-principles prediction of two-dimensional metal bis(dithiolene) complexes as promising gas sensors, *PCCP*, 2018, **20**, 16939-16948.
4. W. R. McNamara, Z. Han, P. J. Alperin, W. W. Brennessel, P. L. Holland and R. Eisenberg, A Cobalt–Dithiolene Complex for the Photocatalytic and Electrocatalytic Reduction of Protons, *J. Am. Chem. Soc.*, 2011, **133**, 15368-15371.
5. (a) L. Dang, M. F. Shibl, X. Yang, A. Alak, D. J. Harrison, U. Fekl, E. N. Brothers and M. B. Hall, The Mechanism of Alkene Addition to a Nickel Bis(dithiolene) Complex: The Role of the Reduced Metal Complex, *J. Am. Chem. Soc.*, 2012, **134**, 4481-4484; (b) D. J. Harrison, N. Nguyen, A. J. Lough and U. Fekl, New Insight into Reactions of Ni(S₂C₂(CF₃)₂)₂ with Simple Alkenes: Alkene Adduct versus Dihydrodithiin Product Selectivity Is Controlled by [Ni(S₂C₂(CF₃)₂)₂]⁻ Anion, *J. Am. Chem. Soc.*, 2006, **128**, 11026-11027.
6. A. Zarkadoulas, E. Koutsouri and C. A. Mitsopoulou, A perspective on solar energy conversion and water photosplitting by dithiolene complexes, *Coord. Chem. Rev.*, 2012, **256**, 2424-2434.
7. (a) A. L. Balch, Cleavage reactions of cobalt and iron dithiolate compounds. Five-coordinate complexes, *Inorg. Chem.*, 1967, **6**, 2158-2162; (b) R. Yu, K. Arumugam, A. Manepalli, Y. Tran, R. Schmehl, H. Jacobsen and J. P. Donahue, Reversible, Electrochemically Controlled Binding of Phosphine to Iron and Cobalt Bis(dithiolene) Complexes, *Inorg. Chem.*, 2007, **46**, 5131-5133; (c) T. Selby-Karney, D. A. Grossie, K. Arumugam, E. Wright and P. Chandrasekaran, Structural and spectroscopic characterization of five coordinate iron and cobalt bis(dithiolene)-trimethylphosphine complexes, *J. Mol. Struct.*, 2017, **1141**, 477-483.
8. (a) J. A. McCleverty, N. M. Atherton, N. G. Connelly and C. J. Winscom, Transition-metal dithiolenes. Part VII. Five- and six-co-ordinate Lewis-base complexes of cobalt and iron bisdithiolenes, *Journal of the Chemical Society A: Inorganic, Physical, Theoretical*, 1969, 2242-2257; (b) J. A. McCleverty and D. G. Orchard, Transition metal dithiolene complexes. Part XVIII. Adducts of cobalt and iron bis-1,2-dithiolenes containing ethyldiphenylphosphine and bis(1,2-diphenylphosphino)ethane, *Journal of the Chemical Society A: Inorganic, Physical, Theoretical*, 1971, 626-631; (c) J. A. McCleverty and B. Ratcliff, Transition-metal dithiolene complexes. Part XIV. Five-co-ordinate phosphine and phosphite adducts of cobalt and iron bis(diaryl-1,2-dithiolenes), *Journal of the Chemical Society A: Inorganic, Physical, Theoretical*, 1970, 1631-1637; (d) A. K. Patra, E. Bill, E. Bothe, K. Chlopek, F. Neese, T. Weyhermüller, K. Stobie, M. D. Ward, J. A. McCleverty and K. Wieghardt, Electronic Structure of Mononuclear Bis(1,2-diaryl-1,2-ethylenedithiolato)iron Complexes Containing a Fifth Cyanide or Phosphite Ligand: A Combined Experimental and Computational Study, *Inorg. Chem.*, 2006, **45**, 7877-7890.
9. (a) J. Locke, J. A. McCleverty, E. J. Wharton and C. J. Winscom, Iron and cobalt nitrosyl 1,2-dithiolenes, *Chemical Communications (London)*, 1966, 677-678; (b) J. A. McCleverty, N. M. Atherton, J. Locke, E. J. Wharton and C. J. Winscom, Transition metal-dithiolene complexes. III. Nitrosyl complexes of iron and cobalt, *J. Am. Chem. Soc.*, 1967, **89**, 6082-6092; (c) J. A. McCleverty and B. Ratcliff, Transition-metal dithiolene complexes. Part XIII. Nitric oxide adducts of iron bis(diaryl-1,2-dithiolenes), *Journal of the Chemical Society A: Inorganic, Physical, Theoretical*, 1970, 1627-1631; (d) P. Surawatanawong, S. Sproules, F. Neese and K. Wieghardt, Electronic Structures and Spectroscopy of the Electron Transfer Series [Fe(NO)L₂]^z (z = 1⁺, 0, 1⁻, 2⁻, 3⁻; L = Dithiolene), *Inorg. Chem.*, 2011, **50**, 12064-12074.
10. K. Ray, E. Bill, T. Weyhermüller and K. Wieghardt, Redox-Noninnocence of the S,S'-Coordinated Ligands in Bis(benzene-1,2-dithiolato)iron Complexes, *J. Am. Chem. Soc.*, 2005, **127**, 5641-5654.
11. S. Friedle, D. V. Partyka, M. V. Bennett and R. H. Holm, Synthesis of metal dithiolene complexes by Si–S bond cleavage of a bis(silanylthio)alkene, *Inorg. Chim. Acta*, 2006, **359**, 1427-1434.
12. A. J. Arduengo, J. R. Goerlich and W. J. Marshall, A stable diaminocarbene, *J. Am. Chem. Soc.*, 1995, **117**, 11027-11028.
13. (a) D. M. Flanigan, F. Romanov-Michailidis, N. A. White and T. Rovis, Organocatalytic Reactions Enabled by N-Heterocyclic Carbenes, *Chem. Rev.*, 2015, **115**, 9307-9387; (b) M. N. Hopkinson, C. Richter, M. Schedler and F. Glorius, An overview of N-heterocyclic carbenes, *Nature*, 2014, **510**, 485; (c) R. S. Menon, A. T. Biju and V. Nair, Recent advances in N-heterocyclic carbene (NHC)-catalysed benzoin reactions, *Beilstein Journal of Organic Chemistry*, 2016, **12**, 444-461; (d) E. Peris, Smart N-Heterocyclic Carbene Ligands in Catalysis, *Chem. Rev.*, 2018, **118**, 9988-10031; (e) R. Singh and S. P. Nolan, N-Heterocyclic carbenes: Advances in transition metal and organic catalysis, *Annual Reports Section "B" (Organic Chemistry)*, 2006, **102**, 168-196. A. Obanda, K. Valerius, J. T. Mague, S. Sproules and J. P. Donahue, Group 10 Metal Dithiolene Bis(isonitrile) Complexes: Synthesis, Structures, Properties, and Reactivity, *Organometallics*, 2020, **39**, 2854-2870.
14. (a) T. Weskamp, F. J. Kohl, W. Hieringer, D. Gleich and W. A. Herrmann, Highly Active Ruthenium Catalysts for Olefin Metathesis: The Synergy of N-Heterocyclic Carbenes and Coordinatively Labile Ligands, *Angew. Chem. Int. Ed.*, 1999, **38**, 2416-2419; (b) S. B. Garber, J. S. Kingsbury, B. L. Gray and A. H. Hoveyda, Efficient and Recyclable Monomeric and Dendritic Ru-Based Metathesis Catalysts, *J. Am. Chem. Soc.*, 2000, **122**, 8168-8179; (c) J. Huang, E. D. Stevens, S. P. Nolan and J. L. Petersen, Olefin Metathesis-Active Ruthenium Complexes Bearing a Nucleophilic Carbene Ligand, *J. Am. Chem. Soc.*, 1999, **121**, 2674-2678; (d) M. Scholl, S. Ding, C. W. Lee and R. H. Grubbs, Synthesis and Activity of a New Generation of Ruthenium-Based Olefin Metathesis Catalysts Coordinated with 1,3-Dimesityl-4,5-dihydroimidazol-2-ylidene Ligands, *Org. Lett.*, 1999, **1**, 953-956; (e) M. Scholl, T. M. Trnka, J. P. Morgan and R. H. Grubbs, Increased ring closing metathesis activity of ruthenium-based olefin metathesis catalysts coordinated with imidazolin-2-ylidene ligands, *Tetrahedron Lett.*, 1999, **40**, 2247-2250.

15. (a) S. Díez-González, N. Marion and S. P. Nolan, N-Heterocyclic Carbenes in Late Transition Metal Catalysis, *Chem. Rev.*, 2009, **109**, 3612-3676; (b) G. C. Fortman and S. P. Nolan, N-Heterocyclic carbene (NHC) ligands and palladium in homogeneous cross-coupling catalysis: a perfect union, *Chem. Soc. Rev.*, 2011, **40**, 5151-5169; (c) E. A. B. Kantchev, C. J. O'Brien and M. G. Organ, Palladium Complexes of N-Heterocyclic Carbenes as Catalysts for Cross-Coupling Reactions—A Synthetic Chemist's Perspective, *Angew. Chem. Int. Ed.*, 2007, **46**, 2768-2813; (d) N. Marion and S. P. Nolan, Well-Defined N-Heterocyclic Carbenes–Palladium(II) Precatalysts for Cross-Coupling Reactions, *Acc. Chem. Res.*, 2008, **41**, 1440-1449.
16. (a) M. Albrecht, J. R. Miecznikowski, A. Samuel, J. W. Faller and R. H. Crabtree, Chelated Iridium(III) Bis-carbene Complexes as Air-Stable Catalysts for Transfer Hydrogenation, *Organometallics*, 2002, **21**, 3596-3604; (b) A. C. Hillier, H. M. Lee, E. D. Stevens and S. P. Nolan, Cationic Iridium Complexes Bearing Imidazol-2-ylidene Ligands as Transfer Hydrogenation Catalysts, *Organometallics*, 2001, **20**, 4246-4252; (c) D. Wang and D. Astruc, The Golden Age of Transfer Hydrogenation, *Chem. Rev.*, 2015, **115**, 6621-6686.
17. (a) S. Burling, B. M. Paine, D. Nama, V. S. Brown, M. F. Mahon, T. J. Prior, P. S. Pregosin, M. K. Whittlesey and J. M. J. Williams, CH Activation Reactions of Ruthenium N-Heterocyclic Carbene Complexes: Application in a Catalytic Tandem Reaction Involving CC Bond Formation from Alcohols, *J. Am. Chem. Soc.*, 2007, **129**, 1987-1995; (b) S. Gaillard, C. S. J. Cazin and S. P. Nolan, N-Heterocyclic Carbene Gold(I) and Copper(I) Complexes in C–H Bond Activation, *Acc. Chem. Res.*, 2012, **45**, 778-787; (c) Y. Wang, D. Wei, Y. Wang, W. Zhang and M. Tang, N-Heterocyclic Carbene (NHC)-Catalyzed sp³ β-C–H Activation of Saturated Carbonyl Compounds: Mechanism, Role of NHC, and Origin of Stereoselectivity, *ACS Catalysis*, 2016, **6**, 279-289.
18. (a) F. Cisnetti and A. Gautier, Metal/N-Heterocyclic Carbene Complexes: Opportunities for the Development of Anticancer Metallo-drugs, *Angew. Chem. Int. Ed.*, 2013, **52**, 11976-11978; (b) W. Liu and R. Gust, Update on metal N-heterocyclic carbene complexes as potential anti-tumor metallo-drugs, *Coord. Chem. Rev.*, 2016, **329**, 191-213; (c) L. Oehninger, R. Rubbiani and I. Ott, N-Heterocyclic carbene metal complexes in medicinal chemistry, *Dalton Transactions*, 2013, **42**, 3269-3284; (d) M.-L. Teyssot, A.-S. Jarrousse, M. Manin, A. Chevy, S. Roche, F. Norre, C. Beaudoin, L. Morel, D. Boyer, R. Mahiou and A. Gautier, Metal-NHC complexes: a survey of anti-cancer properties, *Dalton Transactions*, 2009, 6894-6902; (e) T. Zou, C.-N. Lok, P.-K. Wan, Z.-F. Zhang, S.-K. Fung and C.-M. Che, Anticancer metal-N-heterocyclic carbene complexes of gold, platinum and palladium, *Curr. Opin. Chem. Biol.*, 2018, **43**, 30-36; (f) J. F. Arambula, R. McCall, K. J. Sidoran, D. Magda, N. A. Mitchell, C. W. Bielawski, V. M. Lynch, J. L. Sessler and K. Arumugam, Targeting antioxidant pathways with ferrocenylated N-heterocyclic carbene supported gold(i) complexes in A549 lung cancer cells, *Chemical Science*, 2016, **7**, 1245-1256; (g) P. J. Barnard, M. V. Baker, S. J. Berners-Price and D. A. Day, Mitochondrial permeability transition induced by dinuclear gold(I)–carbene complexes: potential new antimetastatic antitumour agents, *J. Inorg. Biochem.*, 2004, **98**, 1642-1647; (h) R. McCall, M. Miles, P. Lascuna, B. Burney, Z. Patel, K. J. Sidoran, V. Sittaramane, J. Kocerha, D. A. Grossie, J. L. Sessler, K. Arumugam and J. F. Arambula, Dual targeting of the cancer antioxidant network with 1,4-naphthoquinone fused Gold(I) N-heterocyclic carbene complexes, *Chemical Science*, 2017, **8**, 5918-5929; (i) J. Selvakumar, M. H. Miles, D. A. Grossie and K. Arumugam, Synthesis and molecular structure of biologically significant bis(1,3-dimesityl-4,5-naphthoquinoimidazol-2-ylidene)gold(I) complexes with chloride and dichloridoaurate counter-ions, *Acta Crystallographica Section C*, 2019, **75**, 462-468.
19. (a) P.-C. Chiang and J. W. Bode, On the Role of CO₂ in NHC-Catalyzed Oxidation of Aldehydes, *Org. Lett.*, 2011, **13**, 2422-2425; (b) W. Jeong, E. J. Shin, D. A. Culkin, J. L. Hedrick and R. M. Waymouth, Zwitterionic Polymerization: A Kinetic Strategy for the Controlled Synthesis of Cyclic Poly(lactide), *J. Am. Chem. Soc.*, 2009, **131**, 4884-4891; (c) V. Nair, S. Vellalath, M. Poonoth and E. Suresh, N-Heterocyclic Carbene-Catalyzed Reaction of Chalcones and Enals via Homo-enolate: an Efficient Synthesis of 1,3,4-Trisubstituted Cyclopentenones, *J. Am. Chem. Soc.*, 2006, **128**, 8736-8737; (d) S. S. Sohn, E. L. Rosen and J. W. Bode, N-Heterocyclic Carbene-Catalyzed Generation of Homo-enolates: γ-Butyrolactones by Direct Annulations of Enals and Aldehydes, *J. Am. Chem. Soc.*, 2004, **126**, 14370-14371.
20. D. Enders and J. Han, Synthesis of enantiopure triazolium salts from pyroglutamic acid and their evaluation in the benzoin condensation, *Tetrahedron: Asymmetry*, 2008, **19**, 1367-1371.
21. C. S. P. McErlean and A. C. Willis, Application of an Intramolecular Stetter Reaction to Access trans,syn,trans-Fused Pyrans, *Synlett*, 2009, 233-236.
22. H. U. Vora and T. Rovis, *Aldrichimica Acta* 2011, **44**, 3.
23. C. Burstein and F. Glorius, Organocatalyzed Conjugate Umpolung of α,β-Unsaturated Aldehydes for the Synthesis of γ-Butyrolactones, *Angew. Chem. Int. Ed.*, 2004, **43**, 6205-6208.
24. J. Mahatthananchai, P. Zheng and J. W. Bode, α,β-Unsaturated Acyl Azoliums from N-Heterocyclic Carbene Catalyzed Reactions: Observation and Mechanistic Investigation, *Angew. Chem. Int. Ed.*, 2011, **50**, 1673-1677.
25. (a) H. A. Brown, Y. A. Chang and R. M. Waymouth, Zwitterionic Polymerization to Generate High Molecular Weight Cyclic Poly(Carbosiloxane)s, *J. Am. Chem. Soc.*, 2013, **135**, 18738-18741; (b) H. A. Brown and R. M. Waymouth, Zwitterionic Ring-Opening Polymerization for the Synthesis of High Molecular Weight Cyclic Polymers, *Acc. Chem. Res.*, 2013, **46**, 2585-2596; (c) D. A. Culkin, W. Jeong, S. Csihony, E. D. Gomez, N. P. Balsara, J. L. Hedrick and R. M. Waymouth, Zwitterionic Polymerization of Lactide to Cyclic Poly(Lactide) by Using N-Heterocyclic Carbene Organocatalysts, *Angew. Chem. Int. Ed.*, 2007, **46**, 2627-2630.
26. M. Y. Lui, A. K. L. Yuen, A. F. Masters and T. Maschmeyer, Masked N-Heterocyclic Carbene-Catalyzed Alkylation of Phenols with Organic Carbonates, *ChemSusChem*, 2016, **9**, 2312-2316.

27. (a) A. P. Blum, T. Ritter and R. H. Grubbs, Synthesis of N-Heterocyclic Carbene-Containing Metal Complexes from 2-(Pentafluorophenyl)imidazolidines, *Organometallics*, 2007, **26**, 2122-2124; (b) G. W. Nyce, S. Cshony, R. M. Waymouth and J. L. Hedrick, A General and Versatile Approach to Thermally Generated N-Heterocyclic Carbenes, *Chemistry – A European Journal*, 2004, **10**, 4073-4079.
28. B. C. Norris, D. G. Sheppard, G. Henkelman and C. W. Bielawski, Kinetic and Thermodynamic Evaluation of the Reversible N-Heterocyclic Carbene–Isothiocyanate Coupling Reaction: Applications in Latent Catalysis, *The Journal of Organic Chemistry*, 2011, **76**, 301-304.
29. D. J. Cardin, B. Cetinkaya, E. Cetinkaya and M. F. Lappert, Carbene complexes. Part I. Electron-rich olefins as a source of carbene complexes of platinum(II) and palladium(II); and some experiments with (CF₃)₂CN₂, *J. Chem. Soc., Dalton Trans.*, 1973, 514-522.
30. T. M. Trnka, J. P. Morgan, M. S. Sanford, T. E. Wilhelm, M. Scholl, T.-L. Choi, S. Ding, M. W. Day and R. H. Grubbs, Synthesis and Activity of Ruthenium Alkylidene Complexes Coordinated with Phosphine and N-Heterocyclic Carbene Ligands, *J. Am. Chem. Soc.*, 2003, **125**, 2546-2558. S. Naumann, M. Buchmeiser, Liberation of N-heterocyclic carbenes (NHCs) from thermally labile progenitors: protected NHCs as versatile tools in organo- and polymerization catalysis, *Catal. Sci. Technol.*, 2014, **4**, 2466.
31. (a) E. K. Bullough, M. A. Little and C. E. Williams, *Organomet.*, 2013, **32**, 570; (b) B. Gorodetsky, T. Ramnial, N. R. Branda and J. A. C. Clyburne, Electrochemical reduction of an imidazolium cation: a convenient preparation of imidazol-2-ylidenes and their observation in an ionic liquid, *Chem. Commun.*, 2004, 1972-1973; (c) G. D. Robillard, C. H. Devillers, D. Kunz, H. E. Cattetey, E. Digard and J. Andrieu, *J. Org. Lett.*, 2013, **15**, 4410.
32. (a) J. P. Canal, T. Ramnial, D. A. Dickie and J. A. C. Clyburne, From the reactivity of N-heterocyclic carbenes to new chemistry in ionic liquids, *Chem. Commun.*, 2006, 1809-1818; (b) M. Feroci, I. Chiarotto, M. Orsini and A. Inesi, Electrogenerated NHC as an organocatalyst in the Staudinger reaction, *Chem. Commun.*, 2010, **46**, 4121-4123; (c) M. Feroci, I. Chiarotto, M. Orsini, G. Sotgiu and A. Inesi, Reactivity of Electrogenerated N-Heterocyclic Carbenes in Room-Temperature Ionic Liquids. Cyclization to 2-Azetidinone Ring via C-3/C-4 Bond Formation, *Adv. Synth. Catal.*, 2008, **350**, 1355-1359.
33. F. E. Hahn and M. C. Jahnke, Heterocyclic Carbenes: Synthesis and Coordination Chemistry, *Angew. Chem. Int. Ed.*, 2008, **47**, 3122-3172.
34. K. M. Kuhn and R. H. Grubbs, A Facile Preparation of Imidazolium Chlorides, *Org. Lett.*, 2008, **10**, 2075-2077.
35. (a) S. Jiang, T. Zhang, X. Zhang, G. Zhang and B. Li, Nitrogen heterocyclic carbene containing pentacoordinate iron dicarbonyl as a [Fe]-hydrogenase active site model, *Dalton Transactions*, 2015, **44**, 16708-16712; (b) K. Riener, S. Haslinger, A. Raba, M. P. Högerl, M. Cokoja, W. A. Herrmann and F. E. Kühn, Chemistry of Iron N-Heterocyclic Carbene Complexes: Syntheses, Structures, Reactivities, and Catalytic Applications, *Chem. Rev.*, 2014, **114**, 5215-5272.
36. (a) T. Hashimoto, S. Urban, R. Hoshino, Y. Ohki, K. Tatsumi and F. Glorius, Synthesis of Bis(N-heterocyclic carbene) Complexes of Iron(II) and Their Application in Hydrosilylation and Transfer Hydrogenation, *Organometallics*, 2012, **31**, 4474-4479; (b) B. Li, T. Liu, C. V. Popescu, A. Bilko and M. Y. Darensbourg, Synthesis and Mössbauer Characterization of Octahedral Iron(II) Carbonyl Complexes Fe₂(CO)₃L and Fe₂(CO)₂L₂: Developing Models of the [Fe]-H₂ase Active Site, *Inorg. Chem.*, 2009, **48**, 11283-11289.
37. K. Arumugam, M. Selvachandran, A. Obanda, M. C. Shaw, P. Chandrasekaran, S. L. Caston Good, J. T. Mague, S. Sproules and J. P. Donahue, Redox-Active Metallodithiolene Groups Separated by Insulating Tetraphosphinobenzene Spacers, *Inorg. Chem.*, 2018, **57**, 4023-4038.
38. A. W. Addison, T. N. Rao, J. Reedijk, J. van Rijn and G. C. Verschoor, Synthesis, structure, and spectroscopic properties of copper(II) compounds containing nitrogen-sulphur donor ligands; the crystal and molecular structure of aqua[1,7-bis(N-methylbenzimidazol-2'-yl)-2,6-dithiaheptane]copper(II) perchlorate, *J. Chem. Soc., Dalton Trans.*, 1984, 1349-1356.
39. The distortion from an ideal square pyramidal geometry can be computed into an equation to analyze structural fluctuations for compounds 5-8. The following formula is used to compute τ , $\tau = (\beta - \alpha)/60$. For an ideal square pyramidal geometry, $\alpha = \beta = 180^\circ$ (α is defined as $\angle S1-Fe-S3$; β is defined as $\angle S2-Fe-S4$) and the apex position is occupied by CNHC atom. Subsequently, for an ideal trigonal bipyramidal geometry, $\alpha = 120^\circ$ (α is defined as $\angle S1-Fe-S3$ or $\angle S1-Fe-CNHC$ or $\angle S3-Fe-CNHC$) and $\beta = 180^\circ$ (β is defined as $\angle S2-Fe-S4$). For an ideal square pyramidal geometry, $\tau = 0$, while for an ideal trigonal bipyramidal $\tau = 1$. By computing τ , the degree of deviation from an ideal square pyramidal geometry can be deduced for compounds for 5-8.
40. The manuscript is currently under preparation that details the extent of distortion for a variety of iron bis(dithiolene) complexes bearing an NHC ligand. Preliminary screening reveals both steric and electronic effects contribute to the distortion exerted by an NHC unit.
41. H. Jacobsen and J. P. Donahue, Computational Study of Iron Bis(dithiolene) Complexes: Redox Non-Innocent Ligands and Antiferromagnetic Coupling, *Inorg. Chem.*, 2008, **47**, 10037-10045.
42. (a) L. Noodleman, Valence bond description of antiferromagnetic coupling in transition metal dimers, *The Journal of Chemical Physics*, 1981, **74**, 5737-5743; (b) L. Noodleman and E. J. Baerends, Electronic structure, magnetic properties, ESR, and optical spectra for 2-iron ferredoxin models by LCAO-X.alpha. valence bond theory, *J. Am. Chem. Soc.*, 1984, **106**, 2316-2327.
43. (a) A. D. Becke, Density-functional exchange-energy approximation with correct asymptotic behavior, *Physical Review A*, 1988, **38**, 3098-3100; (b) M. Ernzerhof and G. E. Scuseria, Assessment of the Perdew–Burke–Ernzerhof exchange–correlation functional, *The Journal of Chemical Physics*, 1999, **110**, 5029-5036; (c) S. Grimme, Accurate description of van der Waals complexes by density functional theory including empirical corrections, *J. Comput. Chem.*, 2004, **25**, 1463-1473; (d) J. P. Perdew,

- Density-functional approximation for the correlation energy of the inhomogeneous electron gas, *Physical Review B*, 1986, **33**, 8822-8824; (e) J. P. Perdew, Erratum: Density-functional approximation for the correlation energy of the inhomogeneous electron gas, *Physical Review B*, 1986, **34**, 7406-7406; (f) J. P. Perdew, K. Burke and M. Ernzerhof, Generalized Gradient Approximation Made Simple, *Phys. Rev. Lett.*, 1996, **77**, 3865-3868; (g) J. P. Perdew, A. Ruzsinszky, G. I. Csonka, O. A. Vydrov, G. E. Scuseria, L. A. Constantin, X. Zhou and K. Burke, Restoring the Density-Gradient Expansion for Exchange in Solids and Surfaces, *Phys. Rev. Lett.*, 2008, **100**, 136406; (h) M. Reiher, O. Salomon and B. Artur Hess, Reparameterization of hybrid functionals based on energy differences of states of different multiplicity, *Theor. Chem. Acc.*, 2001, **107**, 48-55; (i) V. N. Staroverov, G. E. Scuseria, J. Tao and J. P. Perdew, Comparative assessment of a new nonempirical density functional: Molecules and hydrogen-bonded complexes, *The Journal of Chemical Physics*, 2003, **119**, 12129-12137; (j) P. J. Stephens, F. J. Devlin, C. F. Chabalowski and M. J. Frisch, Ab Initio Calculation of Vibrational Absorption and Circular Dichroism Spectra Using Density Functional Force Fields, *The Journal of Physical Chemistry*, 1994, **98**, 11623-11627; (k) J. Tao, J. P. Perdew, V. N. Staroverov and G. E. Scuseria, Climbing the Density Functional Ladder: Nonempirical Meta-Generalized Gradient Approximation Designed for Molecules and Solids, *Phys. Rev. Lett.*, 2003, **91**, 146401; (l) Y. Zhang and W. Yang, Comment on "Generalized Gradient Approximation Made Simple", *Phys. Rev. Lett.*, 1998, **80**, 890-890.
44. M. M. D. Wilde and M. Gravel, Bis(amino)cyclopropenylidenes as Organocatalysts for Acyl Anion and Extended Umpolung Reactions, *Angew. Chem. Int. Ed.*, 2013, **52**, 12651-12654.
45. W. L. F. Armarego and C. L. L. Chai, *Purification of laboratory chemicals*, Amsterdam: Butterworth-Heinemann, 2003.
46. J. R. Aranzaes, M.-C. Daniel and D. Astruc, Metallocenes as references for the determination of redox potentials by cyclic voltammetry — Permethylated iron and cobalt sandwich complexes, inhibition by polyamine dendrimers, and the role of hydroxy-containing ferrocenes, *Can. J. Chem.*, 2006, **84**, 288-299.
47. (a) C. F. Guerra, J. G. Snijders, G. t. Velde and E. J. Baerends, Towards an order-N DFT method, *Theor. Chem. Acc.*, **99**, 391-403; (b) G. te Velde, F. M. Bickelhaupt, E. J. Baerends, C. Fonseca Guerra, S. J. A. van Gisbergen, J. G. Snijders and T. Ziegler, Chemistry with ADF, *J. Comput. Chem.*, 2001, **22**, 931-967; (c) E. J. Z. Baerends, T.; Atkins, A.J.; Autschbach, J.; Baseggio, O.; Bashford, D.; Bérces, A. Bickelhaupt, F.M.; Bo, C.; Boerrigter, P.M.; Cavallo, L.; Daul, C.; Chong, D.P.; Chulhai, D.V.; Deng, L.; Dickson, R.M.; Dieterich, J.M.; Ellis, D.E.; van Faassen, M.; Fan, L.; Fischer, T.H.; Fonseca Guerra, C.; Franchini, M.; Ghysels, A.; Giammona, A.; van Gisbergen, S.J.A.; Goez, A.; Götz, A.W.; Groeneveld, J.A.; Gritsenko, O.V.; Grüning, M.; Gusarov, S.; Harris, F.E.; van den Hoek, P.; Hu, Z.; Jacob, C.R.; Jacobsen, H.; Jensen, L.; Joubert, L.; Kaminski, J.W.; van Kessel, G.; König, C.; Kootstra, F.; Kovalenko, A.; Krykunov, M.V.; van Lenthe, E.; McCormack, D.A.; Michalak, A.; Mitoraj, M.; Morton, S.M.; Neugebauer, J.; Nicu, V.P.; Noodleman, L.; Osinga, V.P.; Patchkovskii, S.; Pavanello, M.; Peeples, C.A.; Philippsen, P.H.T.; Post, D.; Pye, C.C.; Ramanantoanina, H.; Ramos, P.; Ravenek, W.; Rodríguez, J.I.; Ros, P.; Rüger, R.; Schipper, P.R.T.; Schlüns, D.; van Schoot, H.; Schreckenbach, G.; Seldenthuis, J.S.; Seth, M.; Snijders, J.G.; Solà, M.; Stener, M.; Swart, M.; Swerhone, D.; Tognetti, V.; te Velde, G.; Vernooijs, P.; Versluis, L.; Visscher, L.; Visser, O.; Wang, F.; Wesolowski, T.A.; van Wezenbeek, E.M.; Wiesenekker, G.; Wolff, S.K.; Woo, T.K.; Yakovlev, A.L. ADF2017, SCM, Theoretical Chemistry, Vrije Universiteit, Amsterdam, The Netherlands, <http://www.scm.com>.