


Cite this: *RSC Adv.*, 2022, 12, 3027

Lipase-catalyzed acylation of levoglucosan in continuous flow: antibacterial and biosurfactant studies†

Marcelo A. do Nascimento,^{ae} Juan P. C. Vargas,^b José G. A. Rodrigues,^c Raquel A. C. Leão,^a Patricia H. B. de Moura,^d Ivana C. R. Leal,^d Jonathan Bassut,^{id e} Rodrigo O. M. A. de Souza,^{id a} Robert Wojcieszak^e and Ivaldo Itabaiana Jr ^{id *ef}

Studies involving the transformation of lignocellulosic biomass into high value-added chemical products have been intensively conducted in recent years. Its matrix is mainly composed of cellulose, hemicellulose and lignin, being, therefore, an abundant and renewable source for obtaining several platform molecules, with levoglucosan (LG) standing out. This anhydrous carbohydrate can be acylated to obtain carbohydrate fatty acid esters (CFAEs). Here, these compounds were obtained *via* enzymatic acylation of LG, commercially obtained (Start BioScience®), with different acyl donors in continuous flow. Through the experimental design using a model reaction, it was possible to optimize the reaction conditions, temperature and residence time, obtaining a maximum conversion at 61 °C and 77 min. In addition, there was a productivity gain of up to 100 times in all comparisons made with the batch system. Finally, CFAEs were applied in tests of interfacial tension and biological activity. For a mixture of 4- and 2-*O*-lauryl-1,6-anhydroglucopyranose (MONLAU), the minimum interfacial tension (IFT^{min}) obtained was 96 mN m⁻¹ and the critical micelle concentration (CMC) was 50 mM. Similar values were obtained for a mixture of 4- and 2-*O*-palmitoyl-1,6-anhydroglucopyranose (MONPAL), not yet reported in the literature, of 88 mN m⁻¹ in 50 mM. For a mixture of 4- and 2-*O*-esteryl-1,6-anhydroglucopyranose (MONEST) and 4- and 2-*O*-oleoyl-1,6-anhydroglucopyranose (MONOLE), CMC was higher than 60 mM and IFT^{min} of 141 mN m⁻¹ and 102 mN m⁻¹, respectively. Promising data were obtained for minimal inhibitory concentration (MIC) and minimal bactericidal concentration (MBC) of MONLAU against *Staphylococcus aureus* strains at 0.25 mM.

Received 5th November 2021
Accepted 12th January 2022

DOI: 10.1039/d1ra08111j

rsc.li/rsc-advances

Introduction

Carbohydrates, also called sugars, are the most abundant organic substances in the biological world that make up more than 50% of the dry weight of the earth's biomass, being

essential for life.^{1,2} These compounds can be partially acylated *via* chemo or enzymatic reactions in batch or continuous flow conditions with different acyl donors to access a variety of mono- or oligosaccharide esters which may have valuable properties with wide applications in the food, pharmaceutical, detergent, agricultural, fine chemical and personal care industries.^{3,4}

In this scenario, levoglucosan (1,6-anhydro-β-D-glucopyranose) (LG) might be a promising chemical platform for the synthesis of carbohydrate fatty acid esters, denoted as CFAEs. LG is a major product of cellulose pyrolysis and it can be converted to different high added-value chemicals such as glucose, 5-hydroxymethylfurfural, furfural, sorbitol, levoglucosenone and, as highlighted in this work, CFAEs (Fig. 1).⁵

The structure–activity relationship of CFAEs depends on the degree of substitution of the hydroxyl groups, in addition to the size and amount of unsaturation in the alkyl chains,^{3,6} which can generate compounds with a hydrophobic–lipophilic balance (HLB) of great industrial interest as emulsifying agents, stabilizers in conventional systems and also promising biological activities as already mentioned.^{7–9} It is important to

^aBiocatalysis and Organic Synthesis Group, Chemistry Institute, Federal University of Rio de Janeiro, CEP 21941-909, Brazil

^bNanotechnology Engineering Program, COPPE, Universidade Federal do Rio de Janeiro, UFRJ, Rio de Janeiro, Brazil

^cInstitute of Chemistry, Federal University of Rio de Janeiro, University City, 21941-909, Rio de Janeiro, Brazil

^dLaboratory of Natural Products and Biological Assays, Department of Natural Products and Food, Pharmacy Faculty, Federal University of Rio de Janeiro, 21941-902, Rio de Janeiro, Brazil

^eUniv. Lille, CNRS, Centrale Lille, Univ. Artois, UMR 8181 – UCCS – Unité de Catalyse et Chimie du Solide, F-59000, Lille, France

^fLaboratory of Technological Biochemistry and Biocatalysis, Department of Biochemical Engineering, School of Chemistry, Federal University of Rio de Janeiro, 21941-909, Rio de Janeiro, Brazil. E-mail: ivaldo@eq.ufrj.br

† Electronic supplementary information (ESI) available. See DOI: 10.1039/d1ra08111j






Fig. 1 General representation for highly valued products obtained by chemical and biochemical transformation of LG.

highlight that these compounds are also biodegradable and non-toxic.¹⁰

A challenging task in the chemical synthesis of CFAEs is the selective acylation of a single secondary hydroxyl group due to the presence of multiple functional groups.¹¹ Thus, several protection and deprotection steps are necessary to achieve the selective incorporation of a single ester function, consuming a large amount of energy and/or reagents contrary to the principles defined by “Green Chemistry” that requires the development of sustainable technologies with decreased environmental impact and lesser possibility of side products.¹²

Accordingly, an alternative is enzymatic processes, where lipases (triacylglycerol hydrolases, EC 3.1.1.3) are the most suitable class of enzymes for esterification and transesterification reactions. In this context, Pulido and Gotor (1994)¹³ observed the regioselective monoacylation of the secondary hydroxyl, 4-*O*-, of carbohydrates with ester oximes in 1,4-dioxane. Galletti *et al.* (2007)¹⁴ proposed the acetylation of levoglucosan in the presence of several acyl donors using acetonitrile and ionic liquids as solvent. The esterification occurred mainly in the 4-*O*-hydroxyl. The same was observed by Nascimento *et al.* (2019)¹¹ for long-chain acyl donor. Holmström and Pedersen (2020)¹⁵ obtained high degrees of regioselectivity in the acetylation of functionalized glycosides.

As shown in several studies,^{11,13–17} high yields and regioselectivity can be obtained under milder operating conditions alongside high purity of the final product.¹⁸ However, a recurrent barrier persists,¹⁹ *i.e.*, instability of the enzyme at high temperatures, inhibition by solvent and/or reagents and products, or under turbulent flow conditions, making application difficult for application on an industrial scale. Nevertheless, studies on enzymatic immobilization have been widely disseminated by several research group that report better robustness and stability of these biocatalysts, what are suitable for these applications.^{12,19,20}

Here, we used Lipase B from *Candida antarctica* (CALB) immobilized in epoxy resin,²¹ denominated **CalB_Epoxy**, in the synthesis of CFAEs *via* LG esterification/transesterification reaction with different acyl donors. We have also shown that CFAEs which were synthesized in batch¹¹ can be readily scalable to a continuous flow system where temperature and residence

time conditions have been optimized with the aid of the design of experiments (DoE) statistical approach.

Finally, the obtained CFAEs for a mixture of 4 and 2: *O*-lauryl-1,6-anhydroglucopyranose (MONLAU), *O*-palmitoyl-1,6-anhydroglucopyranose (MONPAL), *O*-estearyl-1,6-anhydroglucopyranose (MONEST) and *O*-oleoyl-1,6-anhydroglucopyranose (MONOLE), respectively, were applied in biological and surface activity tests, where promising results were obtained.

Results and discussion


One of the biggest obstacles in continuous flow chemistry is the handling of solids that can form during the chemical reaction or in the piping that precedes the entry of reagents into the reactor, usually leading to clogging of the channels. Furthermore, the low solubility of the starting materials can make flow-through synthesis a challenging task.^{22,23} Thus, and in order to obtain the best reaction conditions, the reagents were prepared at a concentration of 40 and 100 mM, respectively, and submitted to solubility tests at different temperatures (see ESI – 1.4 – Table S3†).

According to the results obtained, it is possible to observe the insolubility of levoglucosan in acetonitrile at temperatures less than or equal to 50 °C. For the acyl donors: lauric acid, vinyl palmitate and vinyl oleate were soluble in all conditions analyzed, whereas laurate and vinyl stearate were soluble at temperatures greater than or equal to 50 °C. In addition, for the molar concentrations analyzed, the solubility profile remained constant, varying only with the temperature change. Therefore, due to the low solubility of levoglucosan at temperature below 50 °C and for the experimental conditions available in the laboratory, it was necessary to heat the starting material before pumping it into the fixed bed reactor.

Optimization using DoE in a model reaction

The model reaction used was the esterification of levoglucosan with lauric acid (Scheme 1). In the application of the experimental planning, the input variables used were temperature (X_1) and residence time (X_2). And as response variable (Y) the conversion (%). In the CCD, 11 experiments were performed at random and the matrix of experiments with the obtained conversion (%) responses are shown in Table S1 (see ESI – 1.2†). After the execution of all experiments, a second-order equation was obtained:

$$Y = b_0 + b_1T + b_2t + b_1T^2 + b_1t^2 + b_1Tt \quad (1)$$



Scheme 1 Esterification reaction of levoglucosan with lauric acid. * Reactor volume packed with N435: 7.854 mL.



where, Y = conversion response (%); b_i = coefficients of the generated model for T (temperature) and t_r (residence time); the model equation coefficients were obtained using eqn (2) (ref. 24–26) below:

$$b = (X'X)^{-1}(X'Y) \quad (2)$$

From the generated model it was possible to build the response surface and the level curve (Fig. 2a and b).

Based on response surface and levels curve analysis (Fig. 2a and b) can verify that at temperatures below 54 °C and short residence times (10–25 min) the conversion remained below 20%, increasing linearly with the increase of these two parameters, until reaching a maximum conversion (55%) at 61 °C and 77 min. It was also observed that for temperatures above 69 °C and residence times less than 30 min, the conversion tends to remain below 22%, indicating that there is dependence between these two factors. Besides, there is an ideal working range between 60–62 °C and 70–80 min for maximizing conversion.

In the esterification reaction using carboxylic acid, one of the products formed is water, a strong nucleophile in deacylation of the acyl enzyme, shifting the balance in the opposite direction, *i.e.*, hydrolysis of the product (this is a thermodynamically controlled synthesis)^{27,28} and, consequently, decreasing the reaction yield. To shift the balance towards the products and increase yield, it is common to use in excess of one of the reagents or remove one of the products, usually water.²⁹

Alternatively, it is possible to obtain the products desired by changing the leaving group of the acylating agent, avoiding, for example, formation of water in the reaction medium and, furthermore, improve the solubility of the starting materials. Thus, after optimizing the temperature and residence time, the best conditions obtained were applied in the model reaction by varying the acyl donor output group (Fig. 3).

Better results were obtained using vinyl laurate as acylating agent, with conversion 74%, selectivity **3I-a** of 75% and **3III-a** of 25%. The formation of 3-*O*-lauryl-1,6-anhydro-glucopyranose (selectivity **3II-a**) was not observed. According to Boissière-Junot *et al.*³⁰ the orientation of the hydroxyls (axial or equatorial) or neighboring substituent governs the regioselectivity of acetylation catalyzed by lipase. When the C3-OH is axial a high regioselectivity is observed for the adjacent axial C4- or C2-OH.³⁰ Moreover, studies carried out by Uriarte *et al.*³¹ suggest that in




Fig. 3 Acylation of levoglucosan with different acyl donors in 61 °C and residence time 77 min using **N435** as biocatalyst. Selectivity **3I-a**: 4-*O*-lauryl-1,6-anhydroglucopyranose; selectivity **3II-a**: 3-*O*-lauryl-1,6-anhydroglucopyranose and selectivity **3III-a**: 2-*O*-lauryl-1,6-anhydroglucopyranose. Analyses were performed in a gas chromatograph equipped with a mass spectrometry detector (GC-MS).

chair conformation, interactions by intramolecular hydrogen bonds that occur between C3-OH and oxygen in C6, both in the axial position, stabilize the molecule, decreasing its reactivity as a nucleophile. As a result a preference for **3I-a** or **3III-a** must be observed.³⁰

The low conversion and loss of selectivity obtained using ethyl laurate 40% may be related to its partial solubility under applied experimental conditions. Contrary to that, the reaction in the presence of completely soluble lauric acid showed a conversion 55%, 19% lower compared to the vinyl donor, however, better than the values reported by Galletti *et al.*¹⁴ 40% conversion using lauric acid for the batch reaction for about five days.

Another point to be highlighted is the vinyl alcohol formed that tautomerizes to acetaldehyde irreversibly, a more stable compound that also contributes to shifting the chemical balance in the direct direction of the reaction, thereby increasing the yield of the desired ester.³²

The results suggest the great potential of vinyl acyl donors for the synthesis of levoglucosan esters in continuous flow, not only for their solubility, but also for the best conversions and selectivities obtained.

Synthesis of long-chain CFAEs

Finally, after screening, the promising results obtained in the acylation of levoglucosan with vinyl donor motivated us to evaluate a complementary in flow approach regarding the regioselectivity presented by lipase *Candida antarctica* B immobilized in epoxy support by our research group, called **CalB_Epoxy**, applied to obtain CFAEs of levoglucosan in the batch condition.¹¹

The transesterification reaction was carried out in the presence of a series of vinyl esters containing alkyl chains of variable length (C12–C18), as well as in the presence of biocatalyst **N435** used as reference. The results are shown in Table 1.

As a result, the formation of by-products using **N435** was not observed. In contrast, there was the formation of by-products in the presence of **CalB_Epoxy**, respectively (Table 1).




Fig. 2 (a) Response surface and (b) level curve of the conversion in the model reaction of esterification of levoglucosan with lauric acid relating temperature and time.



Regarding regioselectivity, **N435** showed similar values between 23% for **3IIa-d**, and 77% for **3Ia-d** for all acyl donors used, *i.e.*, there was a preference for the hydroxyl linked to C4 for the acylation of levoglucosan. However, there was the formation of by-products using **CalB_Epoxy**, reducing the selectivity for the desired products (Table 1 – entries 5–8). In contrast to observed by Do Nascimento *et al.*¹¹ for the batch reaction, with **N435** being more selective to produce lauric esters (99% at 50 °C) while **CalB_Epoxy** was more selective to produce oleic esters (83% at 55 °C) of LG, respectively.

Compared to the batch system,¹¹ the productivity values obtained for the continuous flow reaction were higher in all samples, even with the loss of selectivity for CFAEs using **CalB_Epoxy** (Table 2). Calculations obtained for specific activity showed that **CalB_Epoxy** presented 41.33 U mg⁻¹ of protein, a value approximately 2.5 times higher than that obtained for **N435** of 16.53 U mg⁻¹ of protein. These results suggest that the high specific activity for **CalB_Epoxy** may have contributed to significant gains for productivity, although the selectivity values for CFAEs are lower than for **N435**.

For a better understanding of the results, it is important to highlight the differences between the **N435** and **CalB_Epoxy** biocatalysts. According to Ortiz *et al.*³³ **N435** is a CALB immobilized by hydrophobic adsorption *via* interfacial activation on a Lewatit VP OC 1600 resin used as a support. The same protocol was applied by Do Nascimento *et al.*²¹ in the immobilization of CALB, but in epoxy acrylate resin ECR 8205 (Purolite®), denominated **CalB_Epoxy**. Therefore, the difference between the two biocatalysts lies in the support used for their immobilization and, consequently, in the way these enzymes

attach themselves to the solid material, which reflects on their specific chemical, biochemical, mechanical and kinetic properties.³³


As mentioned, in both biocatalysts the lipase was adsorbed on the hydrophobic surface of the support *via* an interfacial activation protocol.²¹ This method of immobilization has some advantages, for example, they can almost completely retain their activity.³⁴ Moreover, it is less sensitive to variation in experimental conditions because there is no conformational equilibrium to be changed.^{35,36} However, as it is a reversible immobilization and based on hydrophobic interactions, enzymes can detach from the support and go to the reaction medium, resulting in a loss of enzymatic charge with a subsequent decrease in its activity.

For biocatalysts **N435** and **CalB_Epoxy**, leaching of the enzyme was not observed after the passage of pure solvent through the fixed bed reactor. However, the reagents, as well as the products formed during the reaction can lead to the desorption of the lipase molecules into to reaction medium as well as the polymeric components of the support, contaminating the product, as reported by Hirata *et al.*³⁷ where free fatty acids and di or mono-glycerides, which have recognized detergent properties, favored enzyme desorption. Furthermore, bimolecular aggregates may have been formed in the immobilization and/or leaching during the reaction with subsequent interaction between two open forms of the lipase.³⁸ When this occurs, the effect can be quite negative because these dimers will be immobilized along with the monomeric enzyme molecules. This phenomenon may have occurred with **CalB_Epoxy**, changing its properties.^{38,39}

Evaluation of interfacial tension between oil/water in different concentrations of CFAEs

Table 3 shows the theoretical values of HLB calculated for CFAs using the Becher equation (eqn (1)).⁴⁰ These values between 4.42 and 5.91 and the low solubility in water, especially for compounds from 2–4, suggest an application as a wetting agent or as oil-soluble hydrophobic (w/o) emulsifiers.⁴¹

Table 1 Regioselectivity in the enzymatic acetylation of levoglucosan in CH₃CN with vinyl aliphatic esters in continuous flow. Analyses were performed in a GC-MS^a



Entry	Immobilized biocat.	Acyl donor	Regioselectivity (%)			
			3I	3II	3III	By-products
1	N435	a	75	—	25	—
2		b	76	—	24	—
3		c	73	—	27	—
4		d	75	—	25	—
5	CalB_Epoxy	a	33	—	10	57
6		b	50	—	15	35
7		c	43	—	17	40
8		d	45	—	14	41

^a **N435** = Novozym 435 (*Candida antarctica* lipase B), immobilized on macroporous acrylic type ion exchange resin; **CalB_Epoxy** = *Candida antarctica* lipase B immobilized on epoxy support by our research group. ^b 2 equivalent, a: vinyl laurate, b: vinyl palmitate, c: vinyl stearate and d: vinyl oleate.

Table 2 Comparative between the values of conversion and productivity for batch and continuous flow systems

Entry	Immobilized enz. ^a	Acyl donor ^b	Conversion (%)	Productivity ^c	
				Batch	Continuous flow
1	N435	a	74	0.030	3.660
2		b	76	0.034	4.370
3		c	57	0.034	3.511
4		d	85	0.035	5.211
5	CalB_Epoxy	a	59	0.003	0.511
6		b	99	0.010	1.500
7		c	99	0.009	1.481
8		d	97	0.014	1.420

^a **N435** = Novozym 435 (*Candida antarctica* lipase B), immobilized on macroporous acrylic type ion exchange resin; **CalB_Epoxy** = *Candida antarctica* lipase B immobilized on epoxy support by our research group. ^b 2 equivalent, a: vinyl laurate, b: vinyl palmitate, c: vinyl stearate and d: vinyl oleate. ^c Productivity = mg of product. h⁻¹ S_A⁻¹ of biocatalyst.



In this context, behavioral trend values of CFAEs were measured through experimental studies by the pendant drop method,⁴² of the minimum interfacial tension (IFT^{\min}) in function of the concentration of CFAEs in oil-water (Fig. 4).

The IFT^{\min} is one of the key parameters that govern the compatibility between the components of the blend.⁴³ It is results from the difference in energy between molecules at a fluid interface when compared to their bulk counterparts.⁴⁴ Compounds with properties such as surfactants tend to distribute themselves at interfaces between phases with different degrees of polarity, forming a molecular film that reduces interfacial and surface tension.

In this context, a drop in the voltage value at the water–oil interface was observed with the increase of the molar concentration of CFAEs in the oil and it remained practically constant after 50 mM, thus revealing a state in which the surfaces of the tested samples are saturated with the CFAEs molecules. From this specific concentration, called critical micelle concentration (CMC), thermodynamically stable molecular aggregates begin to form. The structure of the surfactant and the experimental condition define how these molecules are submerged within the fluid.⁴⁵

Also, it is possible to note that compounds 1-2 (MONLAU and MONPAL) showed a similar behavior, with a considerable decrease in IFT^{\min} below CMC, in the range of 248 mN m⁻¹ and

86 mN m⁻¹, and with non-significant variations for high concentrations of CFAEs, above the CMC. Thus, it is suggested a greater affinity of these molecules with the aqueous medium, compared to compounds 3 and 4, so that these compounds were able to spread, and also reduce the tension at the interface with the aqueous medium more effective, at 50 mM. In contrast, MONEST and MONOLE the CMC was greater than 60 mM and IFT^{\min} in the range of 137-105 mN m⁻¹.

Preliminary evaluation of the active properties of the MONLAU product were reported by Galletti *et al.*,¹⁴ which obtained an HLB value of 5.9 and 45 mN m⁻¹ at 0.019 ng mm². Furthermore, Kootery *et al.*⁴⁶ suggest that monoacylated levoglucosan esters can be considered as representative renewable surfactants.

These results corroborate those obtained in this work of the 92 mN m⁻¹ at 53 mM for MONLAU. Moreover, similar results were obtained for MONPAL e MONOLE, without description in the literature, with 88 mN m⁻¹ at 53 mM and 102.48 mN m⁻¹ at 60 mM, which indicates lipophilic water in oil emulsifier.

Antibacterial assays

The results referring the minimal inhibitory concentration and minimal bactericidal concentration, of esterified compounds against *S. aureus* strains, are shown in Table 4. Among the tested substances MONEST and levoglucosan were not able to inhibit the growth of the selected strains at concentration ≤ 1 mM. Concerning the esterified products, MONLAU and MONOLE showed higher activity, highlighting MONLAU which presented MIC values lower than that of the lauric acid substrate. MONLAU was able to inhibit the growth of all MSSA and MRSA strains at 0.25 mM. Since, MRSA presents a multi-drug-resistant pattern for variable antimicrobial classes, such macrolides, fluoroquinolones, aminoglycosides, tetracyclines, and lincosamides,⁴⁷ this is a promising result, once the World Health Organization (WHO) considers methicillin-resistant and vancomycin intermediate, and resistant, *S. aureus* to be a high priority pathogen.

Due to this promising result, bactericidal and bacteriostatic modes of action of MONLAU were evaluated, and the MBC assay indicated the bactericidal effect of MONLAU against all strains (MIC = MBC), including for methicillin-resistant *Staphylococcus aureus*. This was the first report on the antibacterial activity of levoglucosan fatty acid esters. MONLAU, against *S. aureus*, showed higher activities compared to other CFAEs reported in literature, such sucrose monolaurate (MIC = MBC = 0.4 mM),⁴⁸




Fig. 4 Relationship between the concentration of CFAs and interfacial tension.

Table 3 Interface active properties of CFAs

Entry	Compound ^a	Oil solubility (mg mL ⁻¹)	HLB value	IFT^{\min} (mN m ⁻¹)	CMC (mM)
1	MONLAU	>100	5.91	92	53
2	MONEST	>100	4.80	88	50
3	MONPAL	>100	4.42	137	64
4	MONOLE	>100	4.61	105	62

^a Mixture of 4- and 2-: *O*-lauryl-1,6-anhydroglucopyranose (MONLAU), *O*-palmitoyl-1,6-anhydroglucopyranose (MONPAL), *O*-esteryl-1,6-anhydroglucopyranose (MONEST) and *O*-oleoyl-1,6-anhydroglucopyranose (MONOLE).

Table 4 Minimal inhibitory concentration (MIC) and minimal bactericidal concentration (MBC) of products and substrates against *Staphylococcus aureus* strains

Compound	ATCC 25923 MSSA	ATCC 29213 MSSA	ATCC 33591 MRSA	517 MRSA
MONLAU	0.25 mM 0.25 mM ^a	0.25 mM 0.25 mM ^a	0.25 mM 0.25 mM ^a	0.25 mM 0.25 mM ^a
MONEST	>1.0 mM	>1.0 mM	>1.0 mM	>1.0 mM
MONPAL	>1.0 mM	>1.0 mM	1.0 mM	>1.0 mM
MONOLE	1.0 mM	0.5 mM	0.25 mM	1.0 mM
Lauric acid	0.5 mM	0.5 mM	0.5 mM	0.5 mM
Levogluconan	>1.0 mM	>1.0 mM	>1.0 mM	>1.0 mM
Methicillin	0.5 µg mL ⁻¹	0.25 µg mL ⁻¹	4.0 µg mL ⁻¹	4.0 µg mL ⁻¹

^a Minimal bactericidal concentration (MBC). MSSA: methicillin-sensitive *Staphylococcus aureus*. MRSA: methicillin-resistant *Staphylococcus aureus*.

methyl α -D-glucopyranoside monolaurate (MIC = 188 µg mL⁻¹)⁴⁹ and sucrose monocaprato (MIC > 400 µg mL⁻¹).⁵⁰

Experimental

Materials

N435 (immobilized CALB on Lewatit VP OC 1600 resin), Lipase B from *Candida antarctica* (CALB, soluble form) was purchased from Novozymes®, 1,6-anhydroglucopyranose (levoglucosan) was purchased from Start BioScience®. Epoxy resin Purolite® ECR8205F was purchased from Purolite International Limited. Aliphatic carboxylic acids, vinyl esters, NO-bis (trimethylsilyl) trifluoroacetamide (BSTFA) were purchased from Sigma-Aldrich. All other reagents used were of analytical grade.

Lipase-catalyzed acylation of levoglucosan in continuous flow

Experimental setup. In order to run the experiment as described, it is recommended to have the following equipment: reactor (Omnifit BenchMark® Microbore boron silicate glass column) of 15 mm in internal diameter and 100 mm in length and variable bed height, PTFE tubing (0.5 mm internal diameter), tubing fittings, one syringe pumps (Asia, Syrris), or equivalent, one stirrer-hotplates to initial solution (see ESI – 1.1 – Fig. S1†).

In a flask, the solution containing the starting materials was prepared and immersed in a shaking bath. Syrris® syringe pumps were used to transport the solution containing the reagents into the fixed bed reactor at different flow rates, according to the previously defined residence time. Solubility tests were performed (see ESI – 1.4 – Table S3†).

Design of experiments

In order to find the optimal experimental conditions of the variables (temperature and residence time) in the synthesis of levoglucosan esters in continuous flow, the experimental design approach was used. The DoE main objective was to increase the conversion in the esterification reaction of levoglucosan with lauric acid, according to the Scheme 1.

It is important to point out that the significant experimental variables that interfere in the system were selected according to the screening performed for the model reaction in a batch system.¹¹ After the selection of the significant variables, the optimization step was performed using the Central Composite

Design (CCD). At this stage, the previously selected variables were studied with five levels, which allowed establishing the best working conditions in the synthesis of levoglucosan esters in continuous flow.⁶

The CCD makes it possible to carry out the response surface methodology because the variables used are worked on at five levels. These levels ($-\alpha$, -1 , 0 , $+1$, $+\alpha$), are coded by convention and correspond to the five levels studied for the two variables (X_1 and X_2), where ($-\alpha$ is the lowest level, $+\alpha$ is the top level and 0 is the center point). Table 5 shows the coded levels and the experimental values studied for the two variables temperature and residence time in the CCD.

These levels have been defined to ensure that all areas of the parameter space can be explored, regardless of the range of factors (see ESI – 1.2 – Table S1†).

A stock solution containing levoglucosan and lauric acid (1 : 1, 40 mM) in acetonitrile was prepared and added to a 50 mL flask and kept under agitation at 300 rpm at 55 °C. Then, was pumped into the fixed bed reactor packed with **N435**, 2.0 g, to a total volume of 7.854 mL. Flow rates of 87–524 µL min⁻¹ was used and, in addition, different temperatures were evaluated, ranging from 50 to 70 °C. It is known that in a continuous flow, the reaction time is determined by the ratio between the reactor volume and the flow used in the system. A triplicate was performed for the best experimental condition found.

After optimizing the temperature and residence time, the influence of the acyl donor on the LG esterification or transesterification reaction was analyzed.

Transesterification of levoglucosan

After the screening step where the influence of temperature, residence time and acyl donor were investigated in the levoglucosan acetylation reaction, the best results were applied for the synthesis of levoglucosan esters containing alkyl chains of

Table 5 Levels used in central composite planning for two variables

Variable	Level				
	Code	(−1414)	(−)	0	(+) (+1414)
(X_1) Temperature (°C)	T	50	53	60	67 70
(X_2) Residence time (min)	tr	15	26	52.5	79 90



variable length, with up to 18 saturated carbons and unsaturated, biocatalyzed by **N435** and **CalB_Epoxy**, CALB immobilized in epoxy resin by our research group,²¹ according to the Scheme 2.

The reaction progress was monitored by TLC. After stopping the reaction, CFAEs were separated by flash chromatography (eluent: ethyl acetate/hexane 1 : 1—Developer: sulfuric acid/ethanol). To determine the selectivity of the lipases in the position of the hydroxyl groups, the final product, obtained as a mixture of 4-*O'* and 2-*O'* levoglucosan monoesters (MONLAU, MONPAL, MONEST and MONOLE) were silylated with BSTFA and analyzed by gas chromatography with a mass spectrometry detector (GC-MS) (see ESI – 2.1 – Fig. S3 and S4†).

The elucidation of CFAEs has been extensively carried out by NMR: ¹H, ¹³C, HMBC, HSQC, ¹H, ¹H-COSY (“Correlation Spectroscopy”) and NOEdiff (NOE “difference spectroscopy”) spectra, FTIR spectrum ν (cm⁻¹, KBr), HRMS (High-Resolution Mass Spectrometry) by our research group and the results have been published by Do Nascimento *et al.*¹¹ for further details, see ESI 2.4 – Fig. S5–S8 and S10–S13.†

Determination of productivity

In batch

$$\text{Productivity (P)} : \frac{V \times C_r \times f_r \left(\frac{\text{PM}_{\text{PROD}}}{\text{PM}_{\text{REAG}}} \right) \text{conv} \times \text{sel}}{100 \times S_A \times T}$$

In flow

$$\text{Productivity (P)} : \frac{F \times C_r \times f_r \left(\frac{\text{PM}_{\text{PROD}}}{\text{PM}_{\text{REAG}}} \right) \text{conv} \times \text{sel}}{100 \times S_A}$$

where, V is the reaction volume (mL), F is the flow rate (mL min⁻¹), T is the reaction time, C_r is the concentration (mg mL⁻¹), f_r is the purity LG (%), PM_{PROD} and PM_{REAG} are molecular weight of product and reagent (mg mol⁻¹), respectively, conv is the conversion (%), sel is the selectivity for CFAEs (%), S_A is the specific activity.

Determination of surface activity

The HLB value has been estimated by the ¹H NMR spectrum through the integration of the proton signals from the lipophilic and hydrophilic parts of the molecule. These values represent an empirical numerical correlation of the emulsifying

and solubilizing properties of different surface-active agents.⁴¹ The term I_{gph} denotes the integration amplitude of the hydrophilic groups and I_{tot} refers to the total integration amplitude of the protons in the molecule. The ratio H describes the relative hydrophilic part of the molecule and can be used to estimate HLB value applying the equation described of Rabaron *et al.*⁵¹ (eqn (3)).

$$H = I_{\text{gph}}/I_{\text{tot}}$$

$$\text{HLB} = 60H/(H + 2) \quad (3)$$

The HLB scale ranges from 0 to 20. In the range of 3.5 to 6.0, surfactants are more suitable for use in W/O emulsions. Surfactants with HLB values in the 8 to 18 range are most commonly used in O/W emulsions (Becher, 1988).⁴⁰


For the interface tension (IFT) measurements, a two-step process was used. First, compounds were dissolved in olive oil and pendant drop of organic solutions were formed in aqueous environments. To achieve this, the CFAs solutions were loaded into a syringe connected to a U-shape needle. Then, the needle was placed at the bottom of a quartz cell containing distilled water (dispersion medium), and by moving the plunger downward, a drop of volume 15 μ L was formed at the tip of the needle. The so-formed drop is illuminated by means of a light source, and a high-resolution CCD camera was used to capture the drop profile (see ESI – 2.2 - Tables S4–S8†). The determination of the interfacial tension, was calculated according to the Young–Laplace equation defined for the pendant drop (PD) method. A Goniometer OCA25 (DataPhysics Instruments, Germany) was used at room temperature and 1 atm pressure, and the droplet profiles were captured every 1 s for a total period of 20 s. CMC values were determined by plotting the interfacial tension *vs.* the concentration (in mM).

Antibacterial assays

For the antibacterial activity assessment *Staphylococcus aureus* species were used, one clinical and three strains from American Type Culture Collection (ATCC). All information about the bacterial species, correspondent codes, isolation source, and susceptibility profile is described in Table S9 (see ESI – 2.3†). The clinical strain was isolated from patients of Clementino Fraga Filho University Hospital (HUCFF), UFRJ, RJ, Brazil.⁵²

Minimal inhibitory concentration (MIC)

The MIC of the compounds (MONLAU, MONPAL, MONOLE, MONEST, lauric acid and levoglucosan) was determined by the broth microdilution technique in 96-microtiter plates,⁵³ using a bacterial solution equivalent to a 0.5 McFarland scale (10⁸ CFU mL⁻¹) subsequently diluted 1 : 10 (10⁷ CFU mL⁻¹). The samples were prepared at final concentrations of 1.0, 0.5, 0.250, 0.125, 0.0625 and 0.03125 mM. The bacterial growth was evaluated using resazurin. After 24 h, 20 μ L of aqueous resazurin solution (0.85%) was added in each well, and then the microplate was incubated again for 4 h at 37 °C.



Scheme 2 Esterification reaction of levoglucosan with different vinyl esters. * Reactor volume: 7.854 mL (2.0 g of immobilized enzyme). (a) vinyl laurate, (b) vinyl palmitate, (c) vinyl stearate and (d) vinyl oleate.



Minimal bactericidal concentration (MBC)

The MBC assay of MONLAU was determined according to Isenberg.⁵⁴ To perform this assay three concentrations of active samples were used at concentrations 0.250 mM (MIC), 0.5 (2×-MIC), 1.0 mM (4×-MIC).

Conclusions

The high potential of using immobilized enzymes in continuous processes is widely recognized. In this work, it was possible to maximize the synthesis of CFAEs in continuous flow through the DoE where a significant gain in productivity and regioselectivity for C4-OH compared to the batch process was observed. In addition, CFAEs were applied in surface activity tests through the analysis of minimum interfacial tension and in biological tests through the evaluation of antibacterial activity in species of *Staphylococcus aureus*, the clinical strain was isolated from patients of Clementino Fraga Filho University Hospital (UFRJ-RJ-Brazil) and three strains of American Type Culture Collection (ATCC).

A preliminary analysis of MONLAU reported by Galletti *et al.* (2007)¹⁴ suggests good properties as a surfactant at low concentrations of the ester. Similar results were found for MONPAL, not yet reported in the literature, at different concentrations of CFAEs. Finally, analyses of the biological activity of CFAEs showed that MONLAU showed promising results in tests of antibacterial activity compared to other CFAEs reported in the literature.

Conflicts of interest

There are no conflicts to declare.

Acknowledgements

Authors thank Fundação Carlos Chagas Filho de Amparo à Pesquisa do Estado do Rio de Janeiro for Jovem Cientista do Nosso Estado Grant Number: E-203.267/2017, Conselho Nacional de Desenvolvimento Científico e Tecnológico (BR) for Universal Grant Number 429974/2018-3, Centrale Lille, the CNRS, and Lille University as well as the Centrale Initiative Foundation, for their financial contribution. LIA CNRS France-Brazil “Energy & Environment” and Métropole Européenne de Lille (MEL) for “Cat-BioInnov” project are also acknowledged. Authors also thank Professor Dr Kátia Regina Netto dos Santos, coordinator of the Hospital Infection Laboratory at the UFRJ, Department of Medical Microbiology, Institute of Microbiology and Surfaces and Thin Films and Protective Coatings Laboratory-Metalmat/UFRJ.

References

- 1 C. C. Wang, J. C. Lee, S. Y. Luo, S. S. Kulkarni, Y. W. Huang, C. C. Lee, K. L. Chang and S. C. Hung, *Nature*, 2007, **446**, 896–899.
- 2 R. V. Stick, *Carbohydrates: The Sweet Molecules of Life*, 2001.
- 3 N. Park and M. K. Walsh, *LWT*, 2021, **147**, 111664.
- 4 M. C. P. Gonçalves, J. P. Romanelli, J. R. Guimarães, A. C. Vieira, B. P. de Azevedo and P. W. Tardioli, *Crit. Rev. Biotechnol.*, 2021, **41**, 865–878.
- 5 I. Itabaiana Junior, M. Avelar Do Nascimento, R. O. M. A. De Souza, A. Dufour and R. Wojcieszak, *Green Chem.*, 2020, **22**, 5859–5880.
- 6 H. M. El-laithy, O. Shoukry and L. G. Mahran, *Eur. J. Pharm. Biopharm.*, 2011, **77**, 43–55.
- 7 S. W. Chang and J. F. Shaw, *Nat. Biotechnol.*, 2009, **26**, 109–116.
- 8 N. Petkova, *Biointerface Res. Appl. Chem.*, 2021, **11**, 12055–12067.
- 9 J. Andrés, M. Vargas, J. Orduña, M. Bernardo, G. Metzker, E. Gomes, M. Boscolo, S. Paulo, E. Sciences and S. Jose, *Carbohydr. Res.*, 2020, **489**, 107957.
- 10 Y. Zheng, M. Zheng, Z. Ma, B. Xin, R. Guo and X. Xu, *Sugar Fatty Acid Esters*, AOCS Press, 2012.
- 11 M. Avelar do Nascimento, L. Ester Gotardo, E. Miguez Bastos, F. C. L. Almeida, R. A. C. Leão, R. O. M. A. de Souza, R. Wojcieszak and I. Itabaiana, *Sustainability*, 2019, **11**, 6044.
- 12 V. S. Gamayurova, M. E. Zinov'eva, K. L. Shnaider and G. A. Davletshina, *Catal. Ind.*, 2021, **13**, 58–72.
- 13 R. Pulido and V. Gotor, *Carbohydr. Res.*, 1994, **252**, 55–68.
- 14 P. Galletti, F. Moretti, C. Samorì and E. Tagliavini, *Green Chem.*, 2007, **9**, 987–991.
- 15 T. Holmstrøm and C. M. Pedersen, *Eur. J. Org. Chem.*, 2020, **2020**, 4612–4615.
- 16 V. Gotor, *Org. Process Res. Dev.*, 2002, **6**, 420–426.
- 17 E. Busto, V. Gotor-Fernández and V. Gotor, *Chem. Soc. Rev.*, 2010, **39**, 4504–4523.
- 18 R. T. Otto, U. T. Bornscheuer, C. Syldatk and R. D. Schmid, *J. Biotechnol.*, 1998, **64**, 231–237.
- 19 J. Chapman, A. Ismail and C. Dinu, *Catalysts*, 2018, **8**, 238.
- 20 J. J. Bassi, L. M. Todero, F. A. P. Lage, G. I. Khedy, J. D. Ducas, A. P. Custódio, M. A. Pinto and A. A. Mendes, *Int. J. Biol. Macromol.*, 2016, **92**, 900–909.
- 21 M. A. Do Nascimento, L. E. Gotardo, R. A. C. Leão, A. M. De Castro, R. O. M. A. De Souza and I. Itabaiana, *ACS Omega*, 2019, **4**, 860–869.
- 22 V. R. L. J. Bloemendal, M. A. C. H. Janssen, J. C. M. Van Hest and F. P. J. T. Rutjes, *React. Chem. Eng.*, 2020, **5**, 1186–1197.
- 23 M. Guidi, P. H. Seeberger and K. Gilmore, *Chem. Soc. Rev.*, 2020, **49**, 8910–8932.
- 24 B. Barros Neto, I. S. Scarminio and R. E. Bruns, *Como Fazer Experimentos – Pesquisa e Desenvolvimento na Ciência e na Indústria*, 4th edn, 2010.
- 25 T. R. Rosa, J. G. A. Rodrigues and R. Q. de Ferreira, *Quim. Nova*, 2016, **39**, 8–14.
- 26 F. C. Vicentini, L. C. S. Figueiredo-Filho, B. C. Janegitz, A. Santiago, E. R. Pereira-Filho and O. Fatibello-Filho, *Quim. Nova*, 2011, **34**, 825–830.
- 27 W. Chulalaksananukul, J. S. Condoret, P. Delorme and R. M. Willemot, *FEBS Lett.*, 1990, **276**, 181–184.
- 28 J. S. Rhee, S. J. Kwon and J. J. Han, in *Enzymes in Nonaqueous Solvents*, Humana Press, New Jersey, 2003, vol. 15, pp. 135–150.



- 29 U. T. Bornscheuer and R. J. Kazlauskas, in *Hydrolases in Organic Synthesis*, 2006, pp. 141–183.
- 30 N. Boissière-Junot, C. Tellier and C. Rabiller, *J. Carbohydr. Chem.*, 1998, **17**, 99–115.
- 31 I. Uriarte, P. Écija, R. Lozada-Garcia, P. Çarçabal and E. J. Cocinero, *ChemPhysChem*, 2018, **19**, 766–773.
- 32 G. D. Yadav and A. H. Trivedi, *Enzyme Microb. Technol.*, 2003, **32**, 783–789.
- 33 C. Ortiz, M. L. Ferreira, O. Barbosa, J. C. S. dos Santos, R. C. Rodrigues, Á. Berenguer-Murcia, L. E. Briand and R. Fernandez-Lafuente, *Catal. Sci. Technol.*, 2019, **9**, 2380–2420.
- 34 E. A. Manoel, J. C. S. dos Santos, D. M. G. Freire, N. Rueda and R. Fernandez-Lafuente, *Enzyme Microb. Technol.*, 2015, **71**, 53–57.
- 35 A. Bastida, P. Sabuquillo, P. Armisen, R. Fernández-Lafuente, J. Hugué and J. M. Guisán, *Biotechnol. Bioeng.*, 1998, **58**, 486–493.
- 36 R. Fernandez-Lafuente, P. Armisen, P. Sabuquillo, G. Fernández-Lorente and J. M. Guisán, *Chem. Phys. Lipids*, 1998, **93**, 185–197.
- 37 D. B. Hirata, T. L. Albuquerque, N. Rueda, J. J. Virgen-Ortiz, V. G. Tacias-Pascacio and R. Fernandez-Lafuente, *J. Mol. Catal. B: Enzym.*, 2016, **133**, 117–123.
- 38 J. M. Palomo, M. Fuentes, G. Fernández-Lorente, C. Mateo, J. M. Guisán and R. Fernández-Lafuente, *Biomacromolecules*, 2003, **4**, 1–6.
- 39 J. M. Palomo, C. Ortiz, M. Fuentes, G. Fernandez-Lorente, J. M. Guisán and R. Fernandez-Lafuente, *J. Chromatogr. A*, 2004, **1038**, 267–273.
- 40 P. Becher, *Encyclopedia of Emulsion Technology: Basic Theory, Measurement, Applications*, New York, Marcel Dek., 1988, vol. 3.
- 41 J. T. Davies, *Proc. Int. Congr. Surf. Act.*, 2nd, 1957, 426–438.
- 42 E. Y. Arashiro and N. R. Demarquette, *Mater. Res.*, 1999, **2**, 23–32.
- 43 S. Wu, *Polym. Eng. Sci.*, 1987, **27**, 335–343.
- 44 J. D. Berry, M. J. Neeson, R. R. Dagastine, D. Y. C. Chan and R. F. Tabor, *J. Colloid Interface Sci.*, 2015, **454**, 226–237.
- 45 G. Könnicker, J. Regelman, S. Belanger, K. Gamon and R. Sedlak, *Ecotoxicol. Environ. Saf.*, 2011, **74**, 1445–1460.
- 46 K. Parambath Kootery, H. Jiang, S. Kolusheva, T. P. Vinod, M. Ritenberg, L. Zeiri, R. Volinsky, D. Malferrari, P. Galletti, E. Tagliavini and R. Jelinek, *ACS Appl. Mater. Interfaces*, 2014, **6**, 8613–8620.
- 47 A. M. Alghammal, H. F. Hetta, A. Elkelish, D. H. H. Alkhalifah, W. N. Hozzein, G. E. S. Batiha, N. El Nahhas and M. A. Mabrok, *Infect. Drug Resist.*, 2020, **13**, 3255–3265.
- 48 K. M. Park, S. J. Lee, H. Yu, J. Y. Park, H. S. Jung, K. Kim, C. J. Lee and P. S. Chang, *Food Sci. Biotechnol.*, 2018, **27**, 401–409.
- 49 X. Zhang, W. Wei, X. Cao and F. Feng, *J. Sci. Food Agric.*, 2015, **95**, 1631–1637.
- 50 J. D. Monk, A. K. Hathcox and L. R. Beuchat, *Food Microbiol.*, 1996, **13**, 213–225.
- 51 A. Rabaron, G. Cavé, F. Puisieux and M. Seiller, *Int. J. Pharm.*, 1993, **99**, 29–36.
- 52 I. C. R. Leal, K. R. N. Dos Santos, I. I. Júnior, O. A. C. Antunes, A. Porzel, L. Wessjohann and R. M. H. Kuster, *Planta Med.*, 2010, **76**, 47–52.
- 53 CLSI, *Performance Standards for Antimicrobial Susceptibility Testing*, Twenty-Second Informational Supplement Clinical and Laboratory Standards Institute, 2015, vol. 32.
- 54 H. D. Isenberg, *J. Antimicrob. Chemother.*, 1988, **22**, 73–86.

