Chemical Science



ISSN 2041-6539



Chemical Science



EDGE ARTICLE

View Article Online
View Journal | View Issue



Cite this: Chem. Sci., 2016, 7, 5635

[3- N_2 -o- $C_2B_{10}H_{11}$][BF₄]: a useful synthon for multiple cage boron functionalizations of o-carborane†

Da Zhao and Zuowei Xie*

Received 9th April 2016 Accepted 7th June 2016

DOI: 10.1039/c6sc01566b

www.rsc.org/chemicalscience

A simple and efficient method for selective cage B(3) multiple functionalization of o-carborane is described. Reaction of $[3-N_2-o-C_2B_{10}H_{11}][BF_4]$ with various kinds of nucleophiles gave a very broad spectrum of cage B(3)-substituted o-carborane derivatives, $3-X-o-C_2B_{10}H_{11}$ (X = OH, SCN, NH₂, NO₂, N₃, CF₃, PO(C₆H₅)₂, etc). This reaction may serve as another efficient [^{18}F]-radiolabeling method of carborane clusters for positron emission tomography applications.

Introduction

Carboranes, 3-dimensional relatives of benzenes, are a class of boron hydride clusters in which one or more BH vertices are replaced by CH units.1 Carboranes and organic molecules display different electronic, physical, chemical and geometrical properties, which highlights the feasibility or necessity to produce hybrid molecules incorporating both of these two types of fragments.2a,b Indeed, functional carboranes are now finding a broad range of applications encompassing organic synthesis, polymers, catalysis, metal-organic frameworks, electronic devices and more.2c-r As a result, considerable attention has been directed towards the functionalization of carborane molecules.3 In contrast to the relatively well-studied methods for cage carbon functionalization of carboranes, 1,4 selective cage boron functionalization of carboranes still represents a challenging task and developing new methodologies for selective boron derivatization is eagerly desired.^{5,6}

Diazonium compounds $(R-N_2^+X^-)$ constitute an important group of intermediates that have found wide applications in organic synthesis. Many prominent named reactions associated with aryl diazonium salts have been developed since their first discovery in 1858. In sharp contrast, diazonium derivatives of carboranes are little known. It has been documented that o-carboranyl diazonium salts are non-isolable, can only be prepared in situ and undergo substitution reactions with the reaction solvent, usually inorganic acids, in the presence of copper salts. Recently, we have reported the

synthesis of a stable and isolable o-carboranyl diazonium salt, $[3-N_2-o-C_2B_{10}H_{11}][BF_4]$.^{11a} It serves as an ideal precursor for the generation of 1,3-dehydro-o-carborane¹¹ and boron-centered carboranyl radicals.¹²

On the other hand, it has been reported that B(9)-carboranyl iodonium salt can react with nucleophiles.13 Very recently, a similar approach for the functionalization of closo-borates via nucleophilic substitution reactions of the corresponding iodonium zwitterions has been developed.14 However, in these cases, only limited nucleophiles are tolerated and the chemoselectivity of the reaction is highly dependent on the nature of the nucleophiles or the reaction conditions. 13c,14 As the most widely investigated among the carborane family, general and versatile methods for selected cage boron functionalization of o-carboranes still remain very limited.⁵ Previously, our group has reported that utilizing 3-diazonium-o-carborane tetrafluoroborate as the starting material, selective B(3)-arylation of o-carborane can be achieved via the aromatic ene reaction of 1,3-dehydro-o-carborane or a visible-light mediated B-C(sp²) coupling of a carboranyl boron-centered radical. However, the substrate scope is only limited to arenes.11a,12

Considering that dinitrogen is an excellent leaving group, carboranyl diazonium salt may easily undergo a substitution reaction in the presence of a nucleophile. Moreover, compared to aryl diazonium salts, carboranyl diazonium salt may exhibit higher reactivity due to the electron deficient nature of the boron atom and lack of conjugation between the carborane cage and the diazonium group. Herein, we report a proof-of-concept study demonstrating that carboranyl diazonium salt can serve as a powerful synthon for selective cage boron functionalization of *o*-carboranes (Scheme 1).

Results and discussion

3-Diazonium-o-carborane tetrafluoroborate ([3-N₂-o-C₂B₁₀H₁₁] [BF₄]; 1) was prepared in 77% isolated yield, by treatment of

Department of Chemistry and State Key Laboratory of Synthetic Chemistry, The Chinese University of Hong Kong, Shatin, New Territories, Hong Kong, China. E-mail: zxie@cuhk.edu.hk


† Electronic supplementary information (ESI) available: Experimental details, complete characterization data, and X-ray data in CIF format for 4 and 6. CCDC 1473011 and 1473012. For ESI and crystallographic data in CIF or other electronic format see DOI: 10.1039/c6sc01566b

Functionalization of arene via aryl diazonium salt:

$$N_2BF_4$$
 [Cu], X^- or heat $X = F$, Cl, Br, I, CN, etc

Previous work: Limited examples or poor chemoselectivity

This work: Broad scope of nucleophiles & simple, highly efficient



Scheme 1 Functionalization of arene and *o*-carborane *via* diazonium salt.


3-amino-o-carborane with 1.5 equivalents of nitrosonium tetrafluoroborate.^{11a} It is noted that the stability of 1 is dependent upon the counterions used and BF₄ $^-$ offers the highest thermal stability of the salt among the anions examined, such as PF₆ $^-$ and Cl $^-$. A 1.0 g batch of carboranyl diazonium salt 1 stored at -5 °C showed no signs of decomposition over four months.

With this stable precursor in hand, we found that precursor 1 reacted rapidly with various nucleophiles (2) in acetonitrile, providing the corresponding B(3)-substituted *o*-carboranes in good to excellent yields (Table 1). Treatment of 1 with 1 equivalent of strong (charged) nucleophiles, such as halide ions, gave the corresponding halogenated carboranes in excellent yields in <5 min (Table 1, entry 1).

A large variety of nucleophiles, including inorganic salts, water, alcohols, acids, organometallic reagents, ketones, nitriles and phosphine oxides are compatible with this reaction, resulting in the formation of B–C, B–N, B–P, B–O, B–S and B–X (X = F, Cl, Br, I) bonds. More importantly, various functional groups that were previously unable to be introduced into the carborane unit can now be installed in a very simple and efficient manner. For instance, common functional groups can be easily installed on the o-carborane cage boron position using simple inorganic salts in 5 min, affording the corresponding B(3)-functionalized o-carboranes 3–14 (Table 1, entries 1–12). Reaction of precursor 1 with Grignard reagents or lithium amides also gave the B(3)-substituted o-carboranes in moderate to good yields (Table 1, entries 13 and 14).

Weak nucleophiles also work well in this reaction. For instance, in the presence of 10 equivalents of alcohols or water, B(3)-oxygenated carboranes 17 were produced in 81–98% yield (Table 1, entry 15). However, no desired product was observed for *tert*-butyl alcohol, probably due to the steric hindrance imposed by the *tert*-butyl group. Instead, 3-F-o-carborane 3a, generated *via* decomposition of precursor 1, was the only isolated product. Compared to other neutral nucleophiles, the reaction of nitriles is slower even at elevated temperature (Table 1, entry 17).¹⁵

Table 1 Reaction of nucleophiles with precursor 1^a



 a Reaction conditions: precursor **1** (0.1 mmol) was treated with nucleophile **2** (0.1 mmol for inorganic salt and phosphine oxide; **1.0** mmol for alcohol, acid and ketone; **0.4** mmol for Grignard reagent and lithium amide; nitriles were utilized as solvent) in CH₃CN solution for 5 min; yields of isolated products are given. b Deboronation occurred during purification. c CuCF₃ was prepared in situ from TMSCF₃, CuSCN and Cs₂CO₃ in acetonitrile. 16 d -78 °C, THF, **15** min. e The only product was 3-F-C₂B₁₀H₁₁. f 50 °C, 6 h.

The reactivity of precursor 1 towards nucleophiles containing P=O and S=O double bonds was also examined. For example, the reaction of dimethyl sulfoxide furnished compound 20 after hydrolysis (Table 1, entry 18). Although ³¹P and ¹¹B NMR spectra indicated high conversions, reactions with phosphine oxide nucleophiles resulted in lower yields due to the deboronation of the product during the purification process (Table 1, entry 19). ¹⁷ Notably, this metal-free approach provides a rare example of B-carboranyl phosphines. ¹⁸ The rich chemistry of the carboranyl diazonium salt towards various nucleophiles suggests that it can

Edge Article

serve as a very promising synthon for selective cage boron functionalization of *o*-carboranes. It is noteworthy that the reaction also works well when performed on a 0.5 mmol scale.¹⁷


All new compounds were fully characterized by ¹H, ¹³C, and ¹¹B NMR spectroscopy as well as HRMS spectrometry.

The molecular structures of compounds **4** and **6** were further confirmed by single-crystal X-ray analyses.¹⁷

Interestingly, precursor 1 did not react with anhydrous ether (Scheme 2, eqn (1)); however, it reacted rapidly with wet ethereal solvents. For instance, upon treatment with wet diethyl ether, compound 13, resulting from the C–O bond cleavage of ether, was isolated in 95% yield (Scheme 2, eqn (2)). When treated with anhydrous THF, polymerization occurred, leading to gel formation, which suggests the intermediacy of cationic species (Scheme 2, eqn (3)).¹⁷ When *tert*-butyl methyl ether was examined under the same reaction conditions, compound 17a, bearing a methoxy substituent at B(3) position, was formed quantitatively (Scheme 2, eqn (4)), which may shed some light on the reaction mechanism (*vide infra*).

The nucleophilic reaction of the carboranyl diazonium salt was expected to proceed through an S_N1 type of mechanism (Scheme 3).9,19 Although precursor 1 is stable in solution, it can undergo nucleophile-induced heterolytic B-N bond cleavage, producing a boronium intermediate A.14 Similar to the reaction of the dinitrogen derivatives of closo-borates, the rate-determining step is the B-N bond cleavage.9 The resultant reactive boronium intermediate can be trapped by various nucleophiles. For instance, when charged nucleophiles such as inorganic salts were employed as nucleophiles, the corresponding substituted compounds 3-14 were formed in very high yields within 5 min. If the nucleophiles are strong bases, the addition products 15-17 might also be produced via the intermediacy of 1,3-dehydro-o-carborane intermediates.11 Addition of neutral nucleophiles to the boronium intermediate A, alcohols for example, generates an oxonium ion B, which is further deprotonated by the BF₄⁻ anion to afford 17. For weakly nucleophilic ethers, such as tert-butyl methyl ether, no reaction occurs under anhydrous conditions. However, in the presence of a catalytic amount of water, the oxygenated products,

Scheme 2 Reaction of precursor 1 with ethers.



Scheme 3 Proposed reaction pathways.

such as **17a**, were produced within 5 min. This reaction probably proceeds through a sequence of C–H bond cleavage/isobutylene elimination in intermediate C, which is generated by the nucleophilic addition of the ether to the naked boron vertex of intermediate **A**.²⁰ The role of the catalytic amount of water is to facilitate the isobutylene elimination that was detected by GC-MS analyses. The formation of HBF₄ was also confirmed by ¹¹B and ¹⁹F NMR spectra.¹⁷

The present strategy provides a straightforward and practical access to cage boron functionalized o-carboranes. It has been documented that 18 F-labelled ($t_{1/2} = 109.8 \text{ min}$) carboranes are promising radiotracers in Positron Emission Tomography (PET). Previously, 18 [F]-fluorination of o-carborane was achieved by nucleophilic substitution of a B(9)-carboranyl iodonium bromide. 21 However, the overall synthesis time of 20 min limits its possible application, probably due to the low reactivity of the carboranyl iodonium bromide. As a proof of concept, we opted to improve the efficiency of the fluorination process by using precursor 1 as the starting material. Under similar reaction conditions to those reported in the literature, 17 the fluorinated product 3a was formed quantitatively within 1 min and it can be easily purified (eqn (5)).

Conclusions

A practical method for selective cage boron functionalization of *o*-carborane has been developed. By utilizing B-carboranyl diazonium salt as a synthon, a large class of *o*-carborane derivatives bearing previously inaccessible functional groups can now be efficiently prepared, which may find applications in materials sciences.

This work demonstrates that B-carboranyl diazonium salt can serve not only as a source of boron-centered radicals¹² or 1,3-dehydro-*o*-carborane,¹¹ but also as a source of boronium

cations in the presence of nucleophiles. 9.20 These intermediates serve different purposes and are complementary to each other, building up a useful toolbox for cage boron functionalization of *o*-carboranes.

Compared to aryl diazonium salts, the exceptionally high reactivity of B-carboranyl diazonium salt may be due to the lack of conjugation between the carborane cage and the diazonium group. Such a method may find useful applications in the efficient and fast synthesis of ¹⁸F-labelled *o*-carborane derivatives for medical applications.²¹

Acknowledgements

Chemical Science

The work described in this paper was supported by grants from the Research Grants Council of the Hong Kong Special Administration Region (Project No. CUHK7/CRF/12G and 14304115) and NSFC/RGC Joint Research Scheme (Project No. N_CUHK442/14).

Notes and references

- 1 (a) R. N. Grimes, *Carboranes*, Academic Press, 2nd edn, New York, 2011; (b) N. S. Hosmane, *Boron Science: New Technologies and Applications*, CRC Press, Boca Raton, 2012; (c) J. Zhang and Z. Xie, *Acc. Chem. Res.*, 2014, 47, 1623–1633.
- 2 For selected reviews and examples on applications of ocarboranes, see: (a) R. N. Grimes, Dalton Trans., 2015, 44, 5939-5956; (b) Z. Xie, Acc. Chem. Res., 2003, 36, 1-9; (c) S. Mukherjee and P. Thilagar, Chem. Commun., 2016, 52, 1070-1093; (d) X. Li, H. Yan and Q. Zhao, Chem.-Eur. J., 2016, 22, 1888–1898; (e) C. E. Housecroft, J. Organomet. Chem., 2015, 798, 218-228; (f) M. F. Hawthorne and A. Pushechnikov, Pure Appl. Chem., 2012, 84, 2279–2288; (g) M. Scholz and E. Hey-Hawkins, Chem. Rev., 2011, 111, 7035-7062; (h) C. A. Reed, Acc. Chem. Res., 2010, 43, 121-128; (i) A. F. Armstrong and J. F. Valliant, Dalton Trans., 2007, 4240-4251; (j) J. F. Valliant, K. J. Guenther, A. S. King, P. Morel, P. Schaffer, O. O. Sogbein and K. A. Stephenson, Coord. Chem. Rev., 2002, 232, 173-230; (k) Z. Xie, Coord. Chem. Rev., 2002, 231, 23-46; (l) J. J. Schwartz, M. A. Mendoza, N. Wattanatorn, Y. Zhao, V. T. Nguyen, A. M. Spokoyny, C. A. Mirkin, T. Baše and P. S. Weiss, J. Am. Chem. Soc., 2016, 138, 5957-5967; (m) K. O. Kirlikovali, J. C. Axtell, A. Gonzalez, A. C. Phung, S. I. Khan and A. M. Spokoyny, Chem. Sci., 2016, DOI: 10.1039/c6sc01146b; (n) B. J. Eleazer, M. D. Smith and D. V. Peryshkov, Organometallics, 2016, 35, 106-112; (o) Y. O. Wong, M. D. Smith and D. V. Peryshkov, Chem.-Eur. J., 2016, 22, 6764-6767; (p) J. Estrada, C. A. Lugo, S. G. McArthur and V. Lavallo, Chem. Commun., 2016, 52, 1824–1826; (q) S. G. McArthur, L. Geng and V. Lavallo, Inorg. Chem. Front., 2015, 2, 1101-1104; (r) J. Estrada, D. H. Woen, F. S. Tham, G. M. Miyake and V. Lavallo, Inorg. Chem., 2015, 54, 5142-5144.
- 3 For selected examples, see: (*a*) G. R. Kracke, M. R. VanGordon, Y. V. Sevryugina, P. J. Kueffer, K. Kabytaev, S. S. Jalisatgi and M. F. Hawthorne,

- ChemMedChem, 2015, 10, 62-67; (b) D. Zhao, J. Zhang and Z. Xie, J. Am. Chem. Soc., 2015, 137, 13938-13942; (c) D. Zhao, J. Zhang and Z. Xie, J. Am. Chem. Soc., 2015, 137, 9423-9428; (d) D. Zhao, J. Zhang and Z. Xie, Angew. Chem., Int. Ed., 2014, 53, 12902-12906; (e) K. R. Wee, Y. J. Cho, J. K. Song and S. O. Kang, Angew. Chem., Int. Ed., 2013, 52, 9682-9685; (f) D. Brusselle, P. Bauduin, L. Girard, A. Zaulet, C. Viñas, F. Teixidor, I. Ly and O. Diat, Angew. Chem., Int. Ed., 2013, 52, 12114-12118; (g) K. R. Wee, Y. J. Cho, S. Jeong, S. Kwon, J. D. Lee, I. H. Suh and S. O. Kang, J. Am. Chem. Soc., 2012, 134, 17982-17990; (h) A. M. Cioran, A. D. Musteti, F. Teixidor, Ž. Krpetić, I. A. Prior, Q. He, C. J. Kiely, M. Brust and C. Viñas, J. Am. Chem. Soc., 2012, 134, 212-221; (i) K. R. Wee, W. S. Han, D. W. Cho, S. Kwon, C. Pac and S. O. Kang, Angew. Chem., Int. Ed., 2012, 51, 2677-2680; (j) M. Koshino, T. Tanaka, N. Solin, K. Suenaga, H. Isobe and E. Nakamura, Science, 2007, 316, 853.
- 4 (a) D. Zhao and Z. Xie, *Coord. Chem. Rev.*, 2016, **314**, 14–33; (b) Z. Oiu and Z. Xie, *Dalton Trans.*, 2014, **43**, 4925–4934.
- 5 For reviews on the functionalization of cage borons, see: (*a*)
 D. Olid, R. Nuñez, C. Viñas and F. Teixidor, *Chem. Soc. Rev.*,
 2013, 42, 3318–3336; (*b*) Z. Qiu, *Tetrahedron Lett.*, 2015, 56,
 963–971; (*c*) Z. Xie, *Sci. China: Chem.*, 2014, 57, 1061–1063.
- 6 For recent examples on catalytic B-H functionalization of $C_2B_{10}H_{12}$, $CB_{11}H_{12}^-$, $C_2B_9H_{11}^{2-}$ and boranes, see: (a) Y. Quan and Z. Xie, Angew. Chem., Int. Ed., 2016, 55, 1295-1298; (b) H. Lyu, Y. Quan and Z. Xie, Angew. Chem., Int. Ed., 2015, 54, 10623-10626; (c) K. Cao, Y. Huang, J. Yang and J. Wu, Chem. Commun., 2015, 51, 7257-7260; (d) Y. Quan and Z. Xie, J. Am. Chem. Soc., 2015, 137, 3502-3505; (e) Z. Qiu, Y. Quan and Z. Xie, J. Am. Chem. Soc., 2013, 135, 12192–12195; (f) H. C. Johnson, C. L. McMullin, S. D. Pike, S. A. Macgregor and A. S. Weller, Angew. Chem., Int. Ed., 2013, 52, 9776-9780; (g) D. Liu, L. Dang, Y. Sun, H.-S. Chan, Z. Lin and Z. Xie, J. Am. Chem. Soc., 2008, 130, 16103-16110; (h) E. Molinos, S. K. Brayshaw, G. Kociok-Köhn and A. S. Weller, Organometallics, 2007, 26, 2370-2382; (i) I. Rojo, F. Teixidor, R. Kivekäs, R. Sillanpää and C. Viñas, J. Am. Chem. Soc., 2003, 125, 14720-14721.
- 7 For selected reviews on aryl diazonium compounds, see: (a) F. Mo, G. Dong, Y. Zhang and J. Wang, *Org. Biomol. Chem.*, 2013, **11**, 1582–1593; (b) D. P. Hari and B. Konig, *Angew. Chem.*, *Int. Ed.*, 2013, **52**, 4734–4743; (c) S. Mahouche-Chergui, S. Gam-Derouich, C. Mangeney and M. M. Chehimi, *Chem. Soc. Rev.*, 2011, **40**, 4143–4166; (d) C. Galli, *Chem. Rev.*, 1988, **88**, 765–792; (e) M. R. Heinrich, *Chem.-Eur. J.*, 2009, **15**, 821–833.
- 8 For selected examples, see: (a) J. P. Griess, Ann. Chem., 1858, 106, 123–125; (b) T. Sandmeyer, Ber. Dtsch. Chem. Ges., 1884, 17, 1633–1635; (c) R. Pschorr, Ber. Dtsch. Chem. Ges., 1896, 29, 496–501; (d) M. Gomberg and W. E. Bachmann, J. Am. Chem. Soc., 1924, 46, 2339–2343; (e) G. Balz and G. Schiemann, Ber. Dtsch. Chem. Ges. B, 1927, 60, 1186–1190; (f) H. Meerwein, E. Buchner and K. van Emsterk, J. Prakt. Chem., 1939, 152, 237–266; (g) M. P. Doyle, B. Siegfried and J. F. Dellaria Jr, J. Org. Chem., 1977, 42,

Edge Article

2426-2431; (h) K. Kikukawa and T. Matsuda, Chem. Lett., 1977, 159-162.

- 9 (a) B. Ringstrand, P. Kaszyński, V. G. Young and Z. Janoušek, *Inorg. Chem.*, 2010, 49, 1166–1179; (b) B. Ringstrand, P. Kaszyński and V. G. Young Jr, *Inorg. Chem.*, 2011, 50, 2654–2660.
- 10 (a) L. I. Zakharkin and V. N. Kalinin, *Izv. Akad. Nauk SSSR*, *Ser. Khim.*, 1967, 2585–2586; (b) Y. Sevryugina, R. Julius and M. F. Hawthorne, *Inorg. Chem.*, 2010, 49, 10627–10634; (c) R. A. Kasar, G. M. Knudsen and S. B. Kahl, *Inorg. Chem.*, 1999, 38, 2936–2940; (d) L. I. Zakharkin, V. N. Kalinin and V. V. Gedymin, *J. Organomet. Chem.*, 1969, 16, 371–379.
- (a) D. Zhao, J. Zhang and Z. Xie, *Angew. Chem., Int. Ed.*, 2014,
 8488–8491; (b) D. Zhao, J. Zhang and Z. Xie, *Chem.-Eur. J.*, 2015, 21, 10334–10337.
- 12 D. Zhao and Z. Xie, *Angew. Chem., Int. Ed.*, 2016, 55, 3166-3170.
- 13 For selected examples, see: (a) V. V. Grushin, Acc. Chem. Res.,
 1992, 25, 529-536; (b) V. V. Grushin, I. I. Demkina and
 T. P. Tolstaya, Inorg. Chem., 1991, 30, 4860-4863; (c)
 V. V. Grushin and T. M. Shcherbina, J. Organomet. Chem.,
 1985, 292, 105-117.
- 14 P. Kaszyński and B. Ringstrand, Angew. Chem., Int. Ed., 2015, 54, 6576–6581.
- 15 B. Ringstrand and P. Kaszynski, *Acc. Chem. Res.*, 2013, **46**, 214–225, and references therein.
- 16 G. Danoun, B. Bayarmagnai, M. F. Grunberg and L. J. Gooßen, *Angew. Chem., Int. Ed.*, 2013, **52**, 7972–7975.

- 17 For experimental details, safety precautions, complete characterization data, and X-ray data, see the ESI.†
- selected examples on B-P bond formation of o-carboranes, see: (a) K. Z. Kabytaev, A. V. Safronov, Y. V. Sevryugina, C. L. Barnes, S. S. Jalisatgi and M. F. Hawthorne, *Inorg. Chem.*, 2015, 54, 4143-4150; (b) M. Spokoyny, C. D. Lewis, G. Teverovskiy and S. L. Buchwald, Organometallics, 2012, 31, 8478-8481; (c) Bregadze, I. D. Kosenko, I. A. Lobanova, V. Z. A. Starikova, I. A. Godovikov and I. B. Sivaev, Organometallics, 2010, 29, 5366-5372; (d) R. Bernard, D. Cornu, D. Luneau, D. Naoufal, J.-P. Scharff and P. Miele, J. Organomet. Chem., 2005, 690, 2745-2749; (e) C. Viñas, R. Núñez, F. Teixidor, R. Sillanpää and R. Kivekäs, Organometallics, 1999, 18, 4712-4717; (f) S. A. Jasper, R. B. Jones, J. Mattern, J. C. Huffman and L. J. Todd, Inorg. Chem., 1994, 33, 5620-5624; (g) F. Teixidor, J. Casabo, A. M. Romerosa, C. Viñas, J. Rius and C. Miravitlles, J. Am. Chem. Soc., 1991, 113, 9895-9896.
- 19 A. Jankowiak, A. Baliński, J. E. Harvey, K. Mason, A. Januszko, P. Kaszyński, V. G. Young and A. Persoons, J. Mater. Chem. C, 2013, 1, 1144–1159.
- 20 I. Zharov, Z. Havlas, A. M. Orendt, D. H. Barich, D. M. Grant, M. G. Fete and J. Michl, *J. Am. Chem. Soc.*, 2006, **128**, 6089–6100.
- 21 K. B. Gona, V. Gómez-Vallejo, D. Padro and J. Llop, *Chem. Commun.*, 2013, 49, 11491–11493.