

PAPER

View Article Online
View Journal | View IssueCite this: *Dalton Trans.*, 2023, **52**,
5085Received 9th February 2023,
Accepted 22nd March 2023

DOI: 10.1039/d3dt00421j

rsc.li/dalton

Synthesis of imidazolium-based pentacoordinated organofluorosilicate and germanate salts†

Jan Gnidovec,^{id a,b} Evelin Gruden,^{id a} Melita Tramšek,^{id a} Jernej Iskra,^{id c}
Jaroslav Kvičala^{id d} and Gašper Tavčar^{id *a,b}

The syntheses of a new triphenyldifluorogermanate and various pentacoordinated organofluorosilicates are presented. The fluorogermane and fluorosilane compounds were obtained from the corresponding chlorosilanes and chlorogermane by halogen substitution with KF. Subsequent reaction with the imidazolium-based fluoride reagent [IPrH][F] (1,3-bis(2,6-diisopropylphenyl)imidazolium fluoride) led to the formation of [IPrH][Ph₃SiF₂] (**1**), [IPrH][Ph₂SiF₃] (**3**), [IPrH][Et₂SiF₃] (**4**), [IPrH][PhSiF₄] (**5**), [IPrH][EtSiF₄] (**6**) and [IPrH][Ph₃GeF₂] (**7**). All the products obtained were characterised by NMR, Raman spectroscopy and X-ray diffraction. The results were supported by DFT calculations of the structurally optimised compounds.

Introduction

Pentacoordinated organofluorosilicates ([R_{5-n}SiF_n][−]; n = 1–4) are silicon-based compounds containing a combination of organic and fluoride substituents. They initially attracted great interest due to their excellent properties as fluoride ion donors.^{1,2} Their advantage is their improved solubility compared to readily available alkali metal fluorides and their increased stability compared to the widely used tetraalkylammonium fluorides.³ Notable examples are the [Me₃SiF₂][−] and [Ph₃SiF₂][−] anions which are an attractive choice for nucleophilic fluorination.^{2,4} Since the 1980s, when they were first structurally characterised, there have been numerous reports of unique pentacoordinated organofluorosilicate structures (Fig. 1).^{2,5–12}

Organofluorosilicate structures adopt a trigonal-bipyramidal configuration with the fluorine generally in the axial position.² Only in one case has fluorine been observed in equatorial position, when the sterically demanding nature of a bulky bidentate ligand forced the fluorine atom into the equatorial plane.¹⁰ These pentacoordinated compounds contain 1 to 4

fluorine atoms, as shown in Fig. 1. The remaining positions are occupied by various organic substituents. Most of them are simple alkyl or aryl groups, although it has been shown that silicon can form heterocyclic structures with bidentate ligands.^{8–10,13} In certain cases, organofluorosilicates can even form the backbone of a coordination polymer.¹⁴ Due to the Lewis acidic properties of silanes, polydentate silicon-based ligand acceptors have been developed, leading to various hypervalent polysilicon species.^{9,15} In this way, the properties of fluorosilanes can be tuned to increase their efficiency in complexing fluoride anion, making them a useful fluoride sensor.¹² Zwitterionic pentacoordinated organofluorosilicates are an example of neutral organofluorosilicates in which the




Fig. 1 Examples of previously structurally characterised organofluorosilicates.

^aDepartment of Inorganic Chemistry and Technology, Jožef Stefan Institute, Jamova 39, 1000 Ljubljana, Slovenia. E-mail: gasper.tavcar@ijs.si

^bJožef Stefan International Postgraduate School, Jamova 39, 1000 Ljubljana, Slovenia

^cDepartment of Chemistry and Biochemistry, University of Ljubljana, Faculty of Chemistry and Chemical Technology, Večna pot 113, 1000 Ljubljana, Slovenia

^dDepartment of Organic Chemistry, University of Chemistry and Technology, Prague, Technická 5, 166 28 Prague 6, Czech Republic

† Electronic supplementary information (ESI) available: NMR, Raman and HRMS spectra of isolated compounds, crystal structure data and computational results. CCDC 2240916–2240923. For ESI and crystallographic data in CIF or other electronic format see DOI: <https://doi.org/10.1039/d3dt00421j>

silicon is negatively charged while an adjacent group contains a positive charge.¹¹

Organofluorogermanates, on the other hand, are extremely rare. To our knowledge, there are only a few reports of structurally characterised organofluorogermanates $[\text{C}_{23}\text{H}_{17}\text{O}][(\text{CF}_3\text{CF}_2)_3\text{GeF}_2]$ and $[(\text{CH}_3)_4\text{N}][(\text{CF}_3)_3\text{GeF}_2]$.^{16,17} Interestingly, all of them contain strongly electron-withdrawing groups. Like their silicon relatives, fluorogermanates prefer trigonal-bipyramidal configuration with fluorine atoms in the axial positions.


In 2016 the imidazolium-based fluorination reagent $[\text{IPrH}][\text{F}]$ was developed in our research group.¹⁸ It has the ability to act as a free fluoride reagent and its reactivity has already been used for the synthesis of discrete $[\text{GeF}_5]^-$, $[\text{SiF}_5]^-$, $[\text{VOF}_4]^-$, $[\text{NbF}_6]^-$, $[\text{TaF}_6]^-$, $[\text{Me}_3\text{AlF}]^-$ and $[(n\text{-Bu})_3\text{AlF}]^-$ anions.^{19–22} The preparation and structural characterisation of these anions was also possible due to the stabilising ability of the $[\text{IPrH}]^+$ cation. The steric bulkiness of the cation and its interactions with the anions allow the stabilisation of discrete anions, that would otherwise form polynuclear species, by hindering their interactions with neighbouring anions. A good example of this is the stabilisation of discrete $[\text{GeF}_5]^-$ anions in the structure of $[\text{IPrH}][\text{GeF}_5]$, which tends to form octahedrally coordinated $[\text{GeF}_6]^{2-}$ units in the presence of sterically less demanding counterions.¹⁹ So far, the reactivity of $[\text{IPrH}][\text{F}]$ has been tested on inorganic compounds known for their strong Lewis acidity^{20,22} as well as on SiF_4 and GeF_4 .¹⁹

In this work, we were interested in reactivity of $[\text{IPrH}][\text{F}]$ with even weaker Lewis acidic species. The formation and stability of organofluorosilicates depends strongly on the organic substituents attached to the silicon centre. Strong electron-withdrawing groups help in the formation of pentacoordinated anionic species. It is therefore not surprising that there is a distinct lack of organofluorosilicates with electron-donating groups. To this end, we have attempted to synthesise organofluorosilicates with electron donating ethyl groups. Due to the extremely small number of organofluorogermanate compounds, we also decided to test the reactivity towards an organofluorogermane.


Results and discussion

In this work we prepared 5 organofluorosilicate salts $[\text{IPrH}][\text{R}_{4-n}\text{SiF}_{n+1}]$ ($n = 1-3$, $\text{R} = \text{Ph}$, Et) (1–6) and 1 phenylfluorogermanate $[\text{IPrH}][\text{Ph}_3\text{GeF}_2]$ (7) starting from the corresponding organochlorosilanes or chlorogermane using KF and the imidazolium-based fluoride reagent $[\text{IPrH}][\text{F}]$. The products were prepared according to the synthetic procedures shown in Schemes 1 and 2.

In the first step, we converted organochlorosilanes $[\text{R}_{4-n}\text{SiCl}_n]$ ($n = 1-3$, $\text{R} = \text{Ph}$, Et) and triphenylchlorogermane $[\text{Ph}_3\text{GeCl}]$ to the corresponding organofluorosilanes $[\text{R}_{4-n}\text{SiF}_n]$ ($n = 1-3$, $\text{R} = \text{Ph}$, Et) and triphenylfluorogermane $[\text{Ph}_3\text{GeF}]$ by using excess amounts of at least 5 equivalents of KF per chlorine atom of the starting compound. Excess amounts were used



Scheme 1 The general synthetic procedure for preparation of organofluorosilicates $[\text{IPrH}][\text{R}_{4-n}\text{SiF}_{n+1}]$ ($n = 1-3$, $\text{R} = \text{Ph}$, Et) (1–6).



Scheme 2 The synthetic procedure for preparation of $[\text{IPrH}][\text{Ph}_3\text{GeF}_2]$ (7).

to ensure complete exchange of the halides. After completion of the reaction, the reaction mixture was filtered through PTFE filters to remove KCl and the remaining KF. In the second step, the filtrate was added to the equimolar solution of $[\text{IPrH}][\text{F}]$ and stirred overnight. The crystals of the products were isolated by slowly evaporating the solvent in a closed system under static vacuum, the driving force being a temperature difference between 2 parts of the vessel. The crystals of compounds $[\text{IPrH}][\text{Ph}_3\text{SiF}_2]$ (1), $[\text{IPrH}][\text{Ph}_2\text{SiF}_3]$ (3), $[\text{IPrH}][\text{Et}_2\text{SiF}_3]$ (4), $[\text{IPrH}][\text{PhSiF}_4]$ (5), $[\text{IPrH}][\text{EtSiF}_4]$ (6) and $[\text{IPrH}][\text{Ph}_3\text{GeF}_2]$ (7) were obtained in satisfactory quality and were used for characterisation and further analysis. The ethylfluorosilicates proved to be significantly less stable than their phenyl counterparts. Under reduced dynamic pressure, the ethylfluorosilicates decomposed back into the reactants and the volatile fluorosilane was removed from the reaction mixture. As a result, only $[\text{IPrH}][\text{F}]$ remained in the vessel, which was confirmed by NMR spectroscopy and X-ray diffraction.

In the next sections, the structural properties of triorganodifluorosilicates, diorganotrifluorosilicates and organotetrafluorosilicates are going to be presented simultaneously for the ethyl and phenyl counterparts. The properties of the triphenyldifluorogermanate are presented separately.

Triorganodifluorosilicates $[\text{IPrH}][\text{R}_3\text{SiF}_2]$ ($\text{R} = \text{Ph}$, Et)

$[\text{IPrH}][\text{Ph}_3\text{SiF}_2]$ (1) was prepared in quantitative yield according to the general procedure given above. Many attempts were made to synthesise $[\text{IPrH}][\text{Et}_3\text{SiF}_2]$ (2), but all were unsuccessful.






Fig. 2 The crystal structure of $[\text{IPrH}][\text{Ph}_3\text{SiF}_2]$ (**1**). The ellipsoids are drawn at 50% probability. All hydrogen atoms are omitted except for those on the imidazolium ring. Selected bond lengths (Å): Si(1)–F(1) 1.743(1), Si(1)–F(2) 1.746(1), Si(1)–C(1) 1.898(2), Si(1)–C(7) 1.906(2) and Si(1)–C(13) 1.907(2).

ful. Unfortunately, we were not even able to observe the formation of **2** in solution. By following the progress of the reaction between $[\text{Et}_3\text{SiF}]$ and $[\text{IPrH}][\text{F}]$ in MeCN in an NMR tube, we found that the silane $[\text{Et}_3\text{SiF}]$ did not react and only the reactants were visible in the reaction mixture. Our results indicate that **2** cannot be prepared under these conditions (Fig. S4 and S5 in the ESI†). Compound **1** crystallises in the monoclinic space group $P2_1/c$ and its asymmetric unit is shown in Fig. 2. It contains an imidazolium cation $[\text{IPrH}]^+$ and a triphenyldifluoro-silicate anion $[\text{Ph}_3\text{SiF}_2]^-$, which adopts a trigonal bipyramidal geometry with the fluorine atoms occupying the axial positions. The two Si–F bond distances in the anion are very similar. The Si(1)–F(1) has a bond distance of 1.743(1) Å, while Si(1)–F(2) has a distance of 1.746(1) Å. This difference is quite small compared to other crystal structures of this anion where this difference in Si–F bond length is more pronounced (1.734(1) and 1.721(1) Å; 1.753(2) and 1.718(2) Å; 1.732(1) and 1.726(1) Å).^{5,23,24} This could be due to the packing or the similarly strong hydrogen bonds that both fluorine atoms form with imidazolium cations. The $[\text{Ph}_3\text{SiF}_2]^-$ anion interacts with two adjacent cations to form 2 hydrogen bonds, each with one fluorine atom. The first hydrogen bond forms between F(1) and the hydrogen atom at the C2 position C(22)–H(22)⋯F(1) with an H⋯F distance of 2.0841(9) Å, while the second hydrogen bond forms between F(2) and the hydrogen atom on the backbone of the symmetrically generated unit C(24)–H(24)⋯F(2) with an H⋯F distance of 2.0383(9) Å. All significant interactions between cations and anions of the structurally characterised compounds are described with bond lengths and bond angles and are listed in Table S4 in the ESI.† Significant interactions were defined according to the calculations for $[\text{IPrH}][\text{GeF}_5]$ and $[\text{IPrH}][\text{SiF}_5]$.¹⁹

Diorganotrifluorosilicates $[\text{IPrH}][\text{R}_2\text{SiF}_3]$ (R = Ph, Et)

$[\text{IPrH}][\text{Ph}_2\text{SiF}_3]$ (**3**) and $[\text{IPrH}][\text{Et}_2\text{SiF}_3]$ (**4**) were prepared in quantitative yield according to the general procedure listed above. Compound **3** crystallises in the monoclinic space group $P2_1/c$, while **4** crystallises in the form of a solvate with MeCN (**4**·MeCN) in the monoclinic space group $P2_1/n$. The crystal




Fig. 3 The crystal structure of $[\text{IPrH}][\text{Ph}_2\text{SiF}_3]$ (**3**). The asymmetric unit contains two ion pairs of **3**. For clarity only one ion pair is shown. The ellipsoids are drawn at 50% probability. All hydrogen atoms are omitted except for those on the imidazolium ring. Selected bond lengths (Å): Si(1)–F(1) 1.645(1), Si(1)–F(2) 1.691(1), Si(1)–F(3) 1.711(1), Si(1)–C(1) 1.897(2), Si(1)–C(7) 1.901(2).




Fig. 4 The crystal structure of $[\text{IPrH}][\text{Et}_2\text{SiF}_3]$ (**4**). The ellipsoids are drawn at 50% probability. All hydrogen atoms are omitted except for those on the imidazolium ring. Selected bond lengths (Å): Si(1)–F(1) 1.733(1), Si(1)–F(2) 1.648(1), Si(1)–F(3) 1.729(1), Si(1)–C(1) 1.875(2), Si(1)–C(3) 1.885(2).

structures of **3** and **4** are shown in Fig. 3 and 4, respectively. In the crystal structure of **3**, the asymmetric unit is composed of two ion pairs. For clarity, only one is shown in Fig. 3.

The asymmetric unit of **4**·MeCN contains one molecule of MeCN. For clarity only the structure of **4** is shown in Fig. 4. The complete asymmetric units of **3** and **4**·MeCN are shown in Fig. S23 and S24 in the ESI.† The geometry of both structurally characterised $[\text{Et}_2\text{SiF}_3]^-$ and $[\text{Ph}_2\text{SiF}_3]^-$ anions is trigonal bipyramidal, with 2 fluorine atoms occupying the axial positions and the remaining fluorine atom located in the equatorial plane. In both structures, the equatorially bonded fluorine atoms have shorter Si–F bonds compared to the axially positioned ones. On average, the Si–F bond lengths in $[\text{Et}_2\text{SiF}_3]^-$ (1.703 Å) are longer than in $[\text{Ph}_2\text{SiF}_3]^-$ (1.682 Å). In both trifluorosilicates, the cation and the anion interact with each other and form 3 interactions. In compound **3**, two hydrogen bonds are formed between the hydrogen atom at C2 position and the two nearest fluorine atoms (one axial and one equatorial). The hydrogen bond lengths of the fluorine atoms in the axial position (anion 1: F(3)⋯H(32) 2.233(1) Å and anion 2: F(6)⋯H(62) 2.226(1) Å) are similar to those in the equatorial



plane (anion 1: F(1)⋯H(32) 2.298(1) Å and anion 2: F(4)⋯H(62) 2.343(1) Å). Furthermore, in compound **3**, the fluorine atom in equatorial position interacts with hydrogen atoms on one of the adjacent phenyl rings (anion 1: F(1)⋯H(69) 2.368(1) Å and anion 2: F(4)⋯H(52) 2.424(1) Å).

In **4**, 3 hydrogen bonds are formed, one between the hydrogen atom at the C2 position and the nearest fluorine atom in the axial position (F(1)⋯H(12) 2.058(1) Å), while the other two are formed between the remaining two fluorine atoms and the hydrogen atoms on the backbone of the symmetrically generated imidazolium ring (F(2)⋯H(15)^I 2.307(1) Å and F(3)⋯H(14) 2.507(1) Å).

Organotetrafluorosilicates [IPrH][RSiF₄] (R = Ph, Et)

[IPrH][PhSiF₄] (**5**) and [IPrH][EtSiF₄] (**6**) were prepared according to the general procedure listed above. To ensure quantitative conversion of chlorosilane to fluorosilane, the reaction time for halide exchange was doubled compared to other silanes. Compound **5** crystallises in the triclinic space group *P* $\bar{1}$, while **6** crystallises in the monoclinic space group *I*2*a*. The crystal structures of **5** and **6** are shown in Fig. 5 and 6, respectively.




Fig. 5 The crystal structure of [IPrH][PhSiF₄] (**5**). The asymmetric unit contains two ion pairs of **5**. For clarity only domain A of ion pair 1 is shown. The ellipsoids are drawn at 50% probability. All hydrogen atoms are omitted except for those on the imidazolium ring. Selected bond lengths (Å): Si(1A)–F(1A) 1.654(2), Si(1A)–F(2A) 1.718(2), Si(1A)–F(3A) 1.584(3), Si(1A)–F(4A) 1.660(2), Si(1A)–C(1A) 1.901(5).




Fig. 6 The crystal structure of [IPrH][EtSiF₄] (**6**). The asymmetric unit contains two ion pairs of **6**. For clarity only ion pair 1 is shown. The ellipsoids are drawn at 50% probability. All hydrogen atoms are omitted except for those on the imidazolium ring. Selected bond lengths (Å): Si(1)–F(1) 1.701(1), Si(1)–F(2) 1.673(1), Si(1)–F(3) 1.618(1), Si(1)–F(4) 1.614(1), Si(1)–C(1) 1.862(3).

The asymmetric units of the two compounds consist of two ion pairs. For clarity, only one of each is presented. The complete asymmetric units of **5** and **6** are shown in Fig. S25 and S27 in the ESI.† Both crystal structures feature a discrete [IPrH]⁺ cation and a discrete [PhSiF₄][−] or [EtSiF₄][−] anion, which have a trigonal bipyramidal geometry with 2 fluorine atoms in axial and 2 fluorine atoms in equatorial positions. In addition, the positions of the fluorine atoms in **5** are disordered. Two preferred orientations were modelled and refined, with domain A occupancy of 70% (anion 1) and 56% (anion 2) and domain B occupancy of 30% (anion 1) and 44% (anion 2). The disorder of the fluorine atoms is the result of two possible positions of the fluorine atoms in the anion due to the rotation of the fluorine atoms around the C(1A)–Si(1A) axis. The atoms in domain B are rotated by 90° compared to those in domain A. Due to high disorder in **5**, the bond lengths and angles are not discussed further. However, they are given in Table S3 in the ESI.†

Compound **6** has longer Si–F bonds at axial positions (anion 1: Si(1)–F(1) 1.701(1) Å and Si(1)–F(2) 1.672(1) Å, anion 2: Si(2)–F(5) 1.698(1) Å, Si(2)–F(6) 1.670(1) Å) than in equatorial position (anion 1: Si(1)–F(3) 1.618(1) Å and Si(1)–F(4) 1.614(1), anion 2: Si(2)–F(7) 1.621(1) Å and Si(2)–F(8) 1.617(1) Å). Similar behaviour was observed in the structures of **3** and **4** and is consistent with the literature data for [MeSiF₄][−].²⁵ Moreover, the angles between the two axially positioned fluorine atoms (anion 1: F(1)–Si(1)–F(2) 174.87(7)°, anion 2: F(5)–Si(2)–F(6) 174.56(7)°) are smaller than the ideal angle of 180°. This distortion can be attributed to the influence of the organic substituent on the remaining fluorine atoms.

In **5** the [PhSiF₄][−] interacts with 3 imidazolium cations and forms several interactions. Due to high disorder, the interactions are not discussed in detail. All relevant interactions are provided in Table S4 in the ESI.† In the structure we have noticed the interactions between hydrogen atom at C2 position of the imidazolium ring and the closest axial and equatorial fluorine atoms. The other two fluorine atoms interact with the hydrogen atoms on the backbone of the imidazolium ring of the symmetrically generated unit.

In compound **6**, the [EtSiF₄][−] anion interacts with 3 imidazolium cations and forms 5 relevant interactions. In each anion of the asymmetric unit, 2 fluorine atoms (one in axial and one in equatorial position) are oriented towards the hydrogen atom at C2 position of the cation and form 2 interactions (anion 1: F(1)⋯H(22) 2.050(1) Å and F(4)⋯H(22) 2.379(1) Å, anion 2: F(5)⋯H(52) 2.105(1) Å and F(8)⋯H(52) 2.317(1) Å). The other 2 fluorine atoms interact with hydrogen atoms on the backbone of the neighbouring imidazolium ring (anion 1: F(2)⋯H(55) 2.162(1) Å and F(3)⋯H(54) 2.280(1) Å, anion 2: F(6)⋯H(24) 2.200(1) Å and F(7)⋯H(25) 2.229(1) Å). There is also an interaction between the fluorine atom in the axial position and the hydrogen atoms of the neighbouring phenyl rings of the symmetrically generated unit (anion 1: F(2)⋯H(72) 2.460(1) Å, anion 2: F(6)⋯H(28) 2.478(1) Å).



Table 2 Experimental and calculated bond distances of [IPrH][Ph_{4-n}SiF_{n+1}] (*n* = 1–4)

Compound	<i>d</i> (Si–C)/Å		<i>d</i> (Si–F)/Å	
	Experimental	Calculated	Experimental	Calculated
[IPrH][Ph ₃ SiF ₂] (1)	1.907(2)	1.92	1.746(1) ^a	1.85 ^{a,b}
	1.906(2)	1.92	1.743(1) ^{a,b}	1.71 ^a
	1.898(2)	1.91		
[IPrH][Ph ₂ SiF ₃] (3)	1.901(2)	1.898(2)	1.711(1) ^{a,b}	1.710(1) ^{a,b}
	1.897(2)	1.897(2)	1.691(1) ^a	1.689(1) ^a
		1.90	1.645(1) ^b	1.646(1) ^b
[IPrH][PhSiF ₄] (5)			1.718(2) ^{b,c}	1.751(3) ^{b,c}
			1.660(2) ^{a,c}	1.656(3) ^{a,c}
			1.654(4) ^{a,b,c}	1.625(4) ^{a,b,c}
			1.584(3) ^c	1.574(5) ^c
			1.62(1) ^{a,b,c}	
			1.644(4) ^{a,c}	
[IPrH][SiF ₅] ^d			1.59(1) ^{b,c}	
			1.594(4) ^c	
			1.585(4) ^c	

^a Axial position of fluorine atoms. ^b Si–F distances, where fluorine atoms form the strongest interactions with the hydrogen atom at C2 position. ^c Si–C and Si–F distances for domain A, where the anions are disordered. ^d Values reported in ref. 19.

where R = Ph, Et. It seems that the addition of fluorine atoms in the structures increases the stability of silicate and germanate anions, and thus favours the formation of more fluorinated products.

Conclusions

In this work we present a comprehensive study on the reactivity of organofluorosilanes and an organofluorogermane with the imidazolium-based naked fluoride reagent [IPrH][F]. It possesses a sterically demanding cation that has been shown to be able to stabilise discrete anions. For this reason, [IPrH][F] seemed to be a good choice for the preparation of related organofluorosilicates and germanates.

Starting from commercially available organochlorosilanes, triphenylchlorogermane and potassium fluoride, a series of organofluorosilanes and Ph₃GeF were prepared, which were further reacted with [IPrH][F] to form a triphenyldifluorogermanate [IPrH][Ph₃GeF₂] (7) and organofluorosilicates [IPrH][R_{4-n}SiF_{n+1}] (*n* = 1–3, R = Ph, Et) (1–6), with the exception of [IPrH][Et₃SiF₂] (2). The reactions described are simple and selective without the need for purification of the reaction mixtures. Where possible, the newly prepared compounds were characterised by X-ray structural analysis, NMR and Raman spectroscopy. Here we present the synthesis of ethyl-substituted organofluorosilicates (4, 6), which were previously unknown. Despite their weaker Lewis acidity, which is the result of its electron donating substituents, they were able to form stable organofluorosilicate salts. Unfortunately, compound 2 did not form, even in solution. An interesting discovery was also compound 7, which is an extremely rare example of an organofluorogermanate salt and the first structurally characterised one with aromatic substituents. In this work we have shown that [IPrH][F] is a suitable reagent for the formation of new organofluorogermanate and organofluorosili-

cate species. The experimentally obtained results are in agreement with the DFT calculations. The results show that in the [IPrH][R_{4-n}SiF_n] (*n* = 1–3, R = Ph, Et) series, the stability of the products increases with increasing number of fluorine atoms. Calculations for the germanium series have yielded similar results.

Experimental

General methods

All manipulations were performed under an argon atmosphere using Schlenk and glovebox techniques. To avoid possible side reactions of fluoride with glassware, homemade FEP (tetrafluoroethylene–hexafluoropropylene) reaction vessels with PTFE (polytetrafluoroethylene) valves were used. Before use, the FEP reaction vessels were dried under vacuum conditions at room temperature. MeCN was purified using Vigor solvent purification system. MeCN-d₃ was dried over activated 3 Å molecular sieves. Molecular sieves were activated by heating at 150 °C *in vacuo* overnight. [IPrH][F] was synthesised according to the published procedure.¹⁸ KF was dried overnight *in vacuo* at 150 °C and stored in a glovebox before use. All other commercially available reagents were used as received.

NMR spectroscopy

NMR samples were prepared in a glovebox under an inert atmosphere. To prevent the compounds from coming into contact with glass, the samples were measured using homemade FEP inlays for NMR tubes. The spectra were recorded at the Slovenian NMR Centre (National Institute of Chemistry) using a Bruker AVANCE NEO 600 or 400 MHz NMR spectrometer. Spectra were recorded at 298 or 233 K. The chemical shifts of ¹H and ¹³C were referenced to the residual signals of the deuterated solvent and are given relative to tetramethyl-



- 20 Ž. Zupanek, M. Tramšek, A. Kokalj and G. Tavčar, *Inorg. Chem.*, 2018, **57**, 13866–13879.
- 21 Ž. Zupanek, M. Tramšek, A. Kokalj and G. Tavčar, *J. Fluor. Chem.*, 2019, **227**, 109373.
- 22 E. Gruden, M. Tramšek and G. Tavčar, *Organometallics*, 2022, **41**, 41–51.
- 23 P. D. Prince, M. J. Bearpark, G. S. McGrady and J. W. Steed, *Dalton Trans.*, 2007, 271–282.
- 24 A. F. Hill, J. S. Ward and Y. Xiong, *Organometallics*, 2015, **34**, 5057–5064.
- 25 C. Bolli, J. Gellhaar, C. Jenne, M. Keßler, H. Scherer, H. Seeger and R. Uzun, *Dalton Trans.*, 2014, **43**, 4326–4334.
- 26 F. Klanberg and E. L. Muetterties, *Inorg. Chem.*, 1968, **7**, 155–160.
- 27 R. K. Harris, E. D. Becker, S. M. De Cabral Menezes, P. Granger, R. E. Hoffman and K. W. Zilm, *Magn. Reson. Chem.*, 2008, **46**, 582–598.
- 28 CrysAlisPro, Version 1.171.37.33 (release 27-03-2014 CrysAlis171.NET), Agilent Technologies, (compiled Mar 27 2014, 17:12:48).
- 29 R. C. Clark and J. S. Reid, *Acta Crystallogr., Sect. A: Found. Crystallogr.*, 1995, **51**, 887–897.
- 30 G. M. Sheldrick, *Acta Crystallogr., Sect. A: Found. Adv.*, 2015, **71**, 3–8.
- 31 O. V. Dolomanov, L. J. Bourhis, R. J. Gildea, J. A. K. Howard and H. Puschmann, *J. Appl. Crystallogr.*, 2009, **42**, 339–341.
- 32 Diamond – Crystal and Molecular Structure Visualization, <https://www.crystalimpact.com/diamond/>, accessed 13 January 2023.
- 33 M. J. Frisch, G. W. Trucks, H. B. Schlegel, G. E. Scuseria, M. A. Robb, J. R. Cheeseman, G. Scalmani, V. Barone, G. A. Petersson, H. Nakatsuji, X. Li, M. Caricato, A. Marenich, J. Bloino, B. G. Janesko, R. Gomperts, B. Mennucci, H. P. Hratchian, J. V. Ortiz, A. F. Izmaylov, J. L. Sonnenberg, D. Williams-Young, F. Ding, F. Lipparini, F. Egidi, J. Goings, B. Peng, A. Petrone, T. Henderson, D. Ranasinghe, V. G. Zakrzewski, J. Gao, N. Rega, G. Zheng, W. Liang, M. Hada, M. Ehara, K. Toyota, R. Fukuda, J. Hasegawa, M. Ishida, T. Nakajima, Y. Honda, O. Kitao, H. Nakai, T. Vreven, K. Throssell, J. A. Montgomery, Jr., J. E. Peralta, F. Ogliaro, M. Bearpark, J. J. Heyd, E. Brothers, K. N. Kudin, V. N. Staroverov, T. Keith, R. Kobayashi, J. Normand, K. Raghavachari, A. Rendell, J. C. Burant, S. S. Iyengar, J. Tomasi, M. Cossi, J. M. Millam, M. Klene, C. Adamo, R. Cammi, J. W. Ochterski, R. L. Martin, K. Morokuma, O. Farkas, J. B. Foresman and D. J. Fox, *Gaussian 16 (Version B01)*, Gaussian Inc., Wallingford, CT, 2016.
- 34 J. P. Perdew, K. Burke and M. Ernzerhof, *Phys. Rev. Lett.*, 1996, **77**, 3865.
- 35 S. Grimme, J. Antony, S. Ehrlich and H. Krieg, *J. Chem. Phys.*, 2010, **132**, 154104.
- 36 S. Grimme, S. Ehrlich and L. Goerigk, *J. Comput. Chem.*, 2011, **32**, 1456–1465.
- 37 F. Weigend and R. Ahlrichs, *Phys. Chem. Chem. Phys.*, 2005, **7**, 3297–3305.
- 38 F. Weigend, *Phys. Chem. Chem. Phys.*, 2006, **8**, 1057–1065.

