

Cite this: *Chem. Sci.*, 2024, **15**, 3610

All publication charges for this article have been paid for by the Royal Society of Chemistry


Received 1st December 2023  
Accepted 24th January 2024DOI: 10.1039/d3sc06436k  
[rsc.li/chemical-science](http://rsc.li/chemical-science)

Inherent chirality was proposed by Böhmer<sup>1,2</sup> in 1994 to describe the handedness of calix[4]arene<sup>3–5</sup> derivatives devoid of any symmetry elements except for a  $C_1$  rotation axis. As a typical example, introduction of four different substituents into either upper or lower rims of a conventional cone-configured calix[4]arene can lead to a pair of isomers which are non-superimposable mirror images of each other. The inherent chirality has now been widely used<sup>6</sup> by macrocyclic and supramolecular chemists as a specific term to differentiate curved chiral macrocycles different from well-known central, axial, planar and helical molecules.

Despite unique stereochemical structures, tantalizing properties and potential applications in chemistry, materials and life science, inherently chiral compounds have remained largely underexplored.<sup>6,7</sup> Research progress has been very slow since the concept of inherent chirality was introduced nearly thirty years ago. One of the main obstacles impeding and delaying the development of the field is the difficulty in accessing necessary optically active inherently chiral materials. Indeed, most reported methods to obtain enantiopure inherently chiral macrocycles, for instance, rely on the tedious and inefficient resolution<sup>8</sup> of racemic samples by means of analytical HPLC using expensive columns coated with various chiral stationary phases and

multistep asymmetric synthesis employing chiral auxiliary groups.<sup>9</sup> The only attempts to synthesize inherently chiral calixarenes through lipase-catalyzed enantioselective acetylation of tris(*O*-2-hydroxyethylated)calixarenes<sup>10</sup> and to construct tetraazacalix[4]arene from intramolecular C–N bond formation under the catalysis of  $Pd(dba)_2/(R)$ -SEGPHOS<sup>11</sup> suffer from very low chemical yield and enantioselectivity, respectively.

The breakthroughs in the catalytic enantioselective synthesis of inherently chiral compounds have not been reported until



Scheme 1 Previous and current studies on catalytic enantioselective synthesis of inherently chiral calixarene derivatives.

<sup>a</sup>College of Chemistry, Beijing Normal University, Beijing 100875, China. E-mail: [ycheng@bnu.edu.cn](mailto:ycheng@bnu.edu.cn)

<sup>b</sup>Laboratory of Molecular Recognition and Function, Institute of Chemistry, Chinese Academy of Sciences, Beijing 100190, China

<sup>c</sup>MOE Key Laboratory of Bioorganic Phosphorous Chemistry and Chemical Biology, Department of Chemistry, Tsinghua University, Beijing 100084, China. E-mail: [tongshuo@mail.tsinghua.edu.cn](mailto:tongshuo@mail.tsinghua.edu.cn); [wangmx@mail.tsinghua.edu.cn](mailto:wangmx@mail.tsinghua.edu.cn)

† Electronic supplementary information (ESI) available. CCDC 2303849, 2303851, 2303853, 2304018, 2304020 and 2304021. For ESI and crystallographic data in CIF or other electronic format see DOI: <https://doi.org/10.1039/d3sc06436k>




very recently<sup>12–14</sup> (Scheme 1). In 2020, we<sup>12</sup> showed that linear precursors **A** undergo transition metal catalyzed enantioselective macrocyclization reaction to afford ABCD-type chiral tetraazacalix[4]aromatics **B** with an ee of >99%. It represents the extraordinary example of excellent transfer and expression of chirality from catalyst Pd/(*R,S*<sub>P</sub>)-JOSIPHOS to inherently chiral macrocycles in the step of palladium-catalyzed C–N cross coupling reaction. Catalytic enantioselective desymmetrization reactions of calix[4]arene derivatives featuring C–Br/C–H bond coupling were developed successfully last year.<sup>13,14</sup> In the presence of PdBr<sub>2</sub> and chiral BINOL-derived phosphoramidite as a catalyst, calix[4]arene substrates **C** are transformed into 9H-fluorene-embedded inherently chiral calixarenes **D** via a trans-annular C–H arylation in which an unusual enantioselective 1,5-palladium migration step is probably involved.<sup>13</sup> Interestingly, the palladium-catalyzed reaction of the same reactants **C** using a chiral bifunctional phosphine-carboxylate ligand proceeds through an alternative C–H arylation pathway to deliver fluorenone-bearing inherent chiral calix[4]arene products **E**.<sup>14</sup> The catalyst-steering divergent reactions of readily available starting materials provide attractive methods to generate structurally diversified inherently chiral calixarenes.

Almost all inherently chiral calix[4](het)arenes reported to date are composed of differently substituted *m*-(het)arylene units or varied heteroaromatic rings.<sup>6–10,12–14</sup> For instance, enantiomers **D** and **E** contain fluorene and fluorenone segments, respectively, in addition to *m*-phenylene rings.<sup>13,14</sup> As an exciting example, the nitrogen-bridged ABCD-type calix[4]arenes **B** are assembled from benzene, pyridine, pyrimidine and triazine rings with the same methylamino linkages.<sup>12</sup> Heterocalix[*n*]aromatics, or heteroatom-bridged calix(*n*)(het)arenes, are an important class of synthetic macrocyclic host compounds in (supra)molecular chemistry.<sup>15–19</sup> They are easily prepared and functionalized by a number of methods. Fine self-tuning of the macrocyclic conformation and cavity owing to the interplay between heteroatom linkages and their adjacent (hetero)aromatic rings endows heterocalix[*n*]aromatics with powerful and versatile properties<sup>15–20</sup> including selective molecular recognition and self-assembly, fabrication of sophisticated (supra)molecular architectures and functional materials, and formation and stabilization of high-valent aryl-copper species.<sup>21</sup> In contrast to conventional calixarenes, heterocalixaromatics enjoy a very rich diversity of molecular structures which results from the various combinations of heteroatoms and aromatic components. One of the most unique and advantageous structural features is the generation of inherent chirality by merely varying heteroatoms in the linking positions of three-dimensional calix[4](het)arene skeletons consisting of the same aromatic moieties.<sup>11,22</sup> In other words, calix[4](het)arenes lose their molecular symmetries simply due to the variation of connection atoms in between aromatic units. We report herein a distinct type of inherently chiral N<sub>3</sub>,O-calix[2]arene[2]triazine in which symmetrically aligned aromatic rings are bridged by one oxygen and three nitrogen atoms. We show that these unprecedented chiral macrocycles **G** were synthesized with excellent enantioselectivity from the chiral phosphoric acid-catalyzed

intramolecular nucleophilic aromatic substitution (S<sub>N</sub>Ar) reaction of reactants **F** (Scheme 1). Further development of the molecular diversity of inherently chiral N<sub>3</sub>,O-calix[2]arene[2]triazines is demonstrated by post-macrocyclization functionalization reactions on the NH bridge without decrease of enantiopurity. To the best of our knowledge, inherently chiral calix[4](het)arenes obtained from the variation of bridging atoms are extremely rare, and successful synthesis of highly enantiopure isomers has not yet been achieved under asymmetric catalysis.<sup>11,22</sup> There is also no precedent that chiral phosphoric acid acts as an organic catalyst to promote the intramolecular S<sub>N</sub>Ar reaction inducing significantly the inherent chirality of macrocyclic molecules.

We commenced our study with *de novo* synthesis of the racemic form of inherently chiral N<sub>3</sub>,O-calix[2]arene[2]triazines by means of a fragment coupling strategy<sup>15</sup> (Scheme 2). In the presence of diisopropylethylamine (DIPEA) or K<sub>2</sub>CO<sub>3</sub> as an acid scavenger, 3-aminophenol and its derivatives **1a–e** underwent two directional nucleophilic aromatic substitution reactions with 1,3-dichlorotriazine derivatives **2a–f** at room temperature or refluxing in tetrahydrofuran (THF) to afford compounds **3a–j** in 52–80% yields (Scheme S3†). Further treatment of **3a–j** with 1,3-phenylenediamine derivatives **4a–c** at ambient temperature under basic conditions in acetone or THF led to the formation of intermediates **5a–l** in good to excellent yields (Scheme S5† summarizes chemical yields of each individual compound **5a–l**). It should be addressed that the S<sub>N</sub>Ar reaction of **3a–j** with **4a–c** took place preferentially and selectively on the aryloxy-substituted chlorotriazine ring rather than the arylamino-substituted one because the former is more electron-deficient than the latter (Scheme S5†). Macrocyclization of linear precursors **5a–l** appeared not trivial. Following a conventional protocol to promote the intramolecular S<sub>N</sub>Ar reaction of **5a** in



Scheme 2 Synthesis of racemic N<sub>3</sub>,O-calix[2]arene[2]triazines 6.



refluxing acetonitrile using  $\text{Cs}_2\text{CO}_3$  as a base, desired macrocyclization product **6a** was obtained in a yield not exceeding 15%. Employment of other inorganic and organic bases, and different solvents did not improve the formation of **6a** at all (entries 1–8, Table S1†). Pleasingly, the intramolecular  $\text{S}_{\text{N}}\text{Ar}$  reaction of **5a** proceeded efficiently under acidic conditions. For example, in the presence of 0.5 to 1 equivalent of  $\text{HCl}$ ,  $\text{CH}_3\text{CO}_2\text{H}$ , *p*-TsOH and  $\text{CF}_3\text{CO}_2\text{H}$  (TFA), compound **5a** underwent cyclization reaction to yield **6a** as the sole heteracalixaromatic product in chemical yields increasing from 66% to 68%, 71% and 73% (entries 9–13, Table S1†). The target product **6a** was obtained in 76% yield when 2 equivalents of TFA were applied (entry 14, Table S1†). The TFA-mediated intramolecular  $\text{S}_{\text{N}}\text{Ar}$  reaction was very general, and all intermediates **5** tested were converted into the corresponding macrocycles **6** in the yields ranging from 69% to 92% (Scheme 2).

The structures of products were determined by mass spectrometry, NMR and X-ray crystallography. As revealed unambiguously by single crystal X-ray diffraction, the resulting  $\text{N}_3\text{O}$ -calix[2]arene[2]triazines such as **6a**, **6d** and **6j** adopt the 1,3-alternate conformation in the solid state (Fig. 1). Two face-to-face paralleled benzene rings are at perpendicular positions relative to the plane defined by four bridging heteroatoms while two distal triazine rings are inclined to the plane yielding a V-shaped cavity which is occupied by the benzyl group attached to the phenol ring. Both nitrogen and oxygen atoms in linking positions form conjugations with their adjacent triazine rings. We were also delighted to find that all racemic samples of compounds **6a–l** were resolved into enantiomers by means of HPLC using columns coated with chiral stationary phases (ESI†). It suggested that enantiomers of 1,3-alternate conformers of inherently chiral  $\text{N}_3\text{O}$ -calix[2]arene[2]triazines **6** are stable and do not racemize, rendering them attractive targets of asymmetric synthesis.

Given the facile formation of macrocycles **6** under acidic conditions, we set to investigate the asymmetric reaction of **5a** as a model reaction using chiral phosphoric acids<sup>23,24</sup> and *N*-triflyl phosphoramide<sup>24</sup> **Cat-1** to **Cat-18** as organic catalysts (Fig. 2). After first round screening both for activity and enantioselectivity (Table S2†), **Cat-16**, which is derived from (*S*)-6,6-(9-anthryl)-2,2',3,3'-tetrahydro-1,1'-spirobi[indene]-7,7'-diol, stood as a promising catalyst (entry 16, Table S2†). Further extensive examination of reaction conditions showed that the **Cat-16** catalyzed reaction at 0 °C for 24 h in  $\text{CCl}_4$  gave product




Fig. 2 Chiral phosphoric acid catalysts selected to catalyze the macrocyclization of **5a**.

$S_{\text{ic}}$ -**6a** in 23% yield with 86% ee (entry 11, Table S3†). Chemical yields increased when the reaction was extended from 36 to 72 h (entries 12 and 13, Table S2†). Decreased enantioselectivity was observed with the increase of reaction temperatures (entries 15–17, Table S3†). Other solvents including xylenes, toluene, benzene, acetone and acetonitrile led to either low conversion or diminished enantioselectivity (entries 1–10, Table S3†). It should also be pointed out that a catalyst loading of 30 mol% is necessary as chemical yield dropped with the decrease of catalyst loading to 20 mol% (entry 18, Table S3†). Change of concentration of the substrate did not affect the enantioselectivity though the conversion was slightly affected (entries 19 and 20, Table S3†). To improve enantioselectivity, additives such as molecular sieves (3–5 Å),  $\text{Ag}_2\text{CO}_3$ ,  $\text{AgOAc}$  and  $\text{K}_2\text{CO}_3$  were tested in order to remove  $\text{HCl}$  released from the  $\text{S}_{\text{N}}\text{Ar}$  reaction in question, on the basis that  $\text{HCl}$  can promote non-enantioselective macrocyclization of **5a** (*vide supra*). In all cases, additives appeared beneficial to increase conversion while attaining the same level of enantioselectivity (entries 21–30, Table S3†). For example, addition of  $\text{K}_2\text{CO}_3$  (0.6 equiv.) to the **Cat-16** (30 mol%) catalyzed reaction system led to the formation of product  $S_{\text{ic}}$ -**6a** (87% ee) in a yield of 38% (entry 29, Table S3†). Employment of equimolar  $\text{K}_2\text{CO}_3$  led to a diminished yield (entry 30, Table S3†). Finally, and gratifyingly, excellent enantioselective catalysis was achieved at 0 °C in a mixture of  $\text{CCl}_4$  and cyclohexane when  $\text{K}_2\text{CO}_3$  (0.6 equiv.) was used as an additive, affording highly enantiopure  $S_{\text{ic}}$ -**6a** in 36% yield and 93% ee (entry 40, Table S3†). Switch of the solvent system back to 1,2-dichloroethane resulted in a drastic decrease of the ee value of the product to 41% (entry 42, Table S3†).

Under the optimized conditions, the scope of the chiral phosphoric acid **Cat-16**-catalyzed synthesis of inherently chiral  $\text{O}_3\text{N}$ -calix[4]aromatics **6a–l** from the cyclization of **5a–l** was surveyed. As illustrated in Scheme 3, reactants which bear an alkoxy group such as methoxy (**5a**), ethoxy (**5b**) and isopropoxy (**5c**) on the triazine moiety were transformed into the corresponding  $S_{\text{ic}}$ -(+)-**6a–c** with an ee of 91.4–94.2%. A slight decrease of the ee value was observed when a larger benzyloxy (**5d**) is attached on triazine. The enantioselectivity of product  $S_{\text{ic}}$ -**6d** was nevertheless improved readily to 98.4% ee by recrystallization. Good enantioselectivity was attained when an electron-donating ( $S_{\text{ic}}$ -**6f**) or electron-withdrawing ( $S_{\text{ic}}$ -**6e**) group is present on the benzene ring of the aryloxy group at the lower rim. Functional groups such as ester (**5g**) were tolerated. More remarkably, irrespective of the nature of the substituent on the benzene ring





Fig. 1 X-ray crystal and molecular structures of (±)-**6a** (A), (±)-**6d** (B) and (±)-**6j** (C) with top (top) and side (bottom) views.





**Scheme 3** Catalytic enantioselective synthesis of inherently chiral N<sub>3</sub>,O-calix[2]arene[2]triazines 6a–l and the X-ray molecular structure of Sic-6b. <sup>a</sup>Reaction was performed in a mixture of CCl<sub>4</sub> and cyclohexane (v/v = 13/7). <sup>b</sup>Reaction was carried out in pure CCl<sub>4</sub>. <sup>c</sup>The ee value was obtained after recrystallization. <sup>d</sup>Enantiomer of Cat-16 was used as the chiral catalyst.

embedded in the linear structure, macrocyclization of substrates 5g–j proceeded with good to excellent enantiocontrol, furnishing products *Sic*-(+)-6g–j with high enantiopurity. It should be noted that the marginal difference of enantioselection was observed when alkoxy was replaced with methyl (5l) on the triazine ring. Noticeably, however, substitution of alkoxy by an aryl group (5k) on the triazine subunit led to a considerable decrease of the enantioselectivity of asymmetric catalysis. The distinct substituent effect between aryl, alkyl and alkoxy on enantioselectivity implied the role of triazine played in catalysis


(*vide infra*). As expected, macrocyclization of 5a and 5b applying the enantiomer of Cat-16 gave rise to *Ric*-(–)-6a and *Ric*-(–)-6b (Scheme 3).

The absolute configuration of inherently chiral products was assigned based on the single crystal X-ray molecular structure of *Sic*-6b,<sup>25</sup> which is the predominant enantiomer from the Cat-16 catalyzed reaction of 5b (Scheme 3, Fig. S4 and Table S7†), and the comparison of circular dichroism spectra between chiral products (Fig. S7–S18†). Based on the outcomes of asymmetric induction and substituent effects aforementioned, we proposed a catalytic reaction process in which chiral phosphoric acid Cat-16 forms a pair of hydrogen bonds with the secondary amino-s-triazine moiety of 5 (Fig. 3, left). The catalytic mode was supported by theoretical calculations. According to DFT calculations (Fig. S28†), two optimized pre-organized complexes between the chiral catalyst Cat-16 and substrate 5a were obtained. The complex (Fig. 3, right) leading to *Sic*-(+)-6a has a lower binding energy by 5 kcal mol<sup>–1</sup> in comparison to the complex leading to the enantiomer (Fig. S28†). Revealed by the computed structure (Fig. 3, right), the spiro phosphoric acid forms a pocket in which phosphoric acid binds secondary amino-s-triazine nicely through a pair of hydrogen bonds. One of the anthracene substituents of the catalyst forms evidently non-covalent π–π interactions with the benzene while the other forms C–H–π interaction with *t*-butyl of the substrate. There is also a weak interaction between phosphoric acid and the terminal amino group (*d*<sub>O–N</sub> = 3.885 Å) that brings *Sic*Ar reaction sites in close proximity to each other (*d*<sub>C–N</sub> = 2.801 Å). It is therefore the favourable multiple non-covalent bond interactions between the chiral catalyst and substrate that lead to enantioselective macrocyclization through the intramolecular *S<sub>N</sub>Ar* reaction (Fig. 3). To verify the computed mode of asymmetric catalysis, substrate 5m was designed and synthesized (ESI†). Without the N–H binding site, the Cat-16 catalysed reaction of 5m under identical conditions gave the corresponding product *Sic*-8a merely in 7% yield and 42% ee (Scheme 4). It may be worth noting that activation of amino-substituted heteroaromatic rings using a chiral phosphoric acid to facilitate the *S<sub>N</sub>Ar* reaction would open a new route to construct axial chiral bi(hetero)aryls.

The secondary amino moiety in the bridging position and the ester on the benzene ring would provide useful handles for derivatization of the acquired products. To demonstrate the synthetic value of the method, and also to develop the diversity of inherently chiral N<sub>3</sub>,O-calix[2]arene[2]triazines, functionalization reactions on the bridging amino group were conducted




**Fig. 3** Proposed (left) and computed (right) catalytic mode for enantioselective macrocyclization.

Scheme 4 Control reaction of **5m** under identical catalytic conditions.

(Scheme 5). Thus, assisted by NaH, enantioenriched compounds *S<sub>ic</sub>-6a* (90.0–92.0% ee) and *S<sub>ic</sub>-6g* (96.8% ee) underwent *N*-alkylation with methyl iodide (**7a**), allylic bromide (**7b**), propargyl bromide (**7c**) and 4-bromobenzyl bromide (**7d**) efficiently at 0 °C in DMF. The corresponding *N*-functionalized products *S<sub>ic</sub>-8a–h* (90.4–96.7% ee)<sup>25</sup> were isolated almost quantitatively. *N*-Arylation product *S<sub>ic</sub>-10* (89.9% ee) was also prepared straightforwardly from the reaction between *S<sub>ic</sub>-6a* (90.0% ee) and 9,9'-(6-chloro-1,3,5-triazine-2,4-diyl)bis(9H-carbazole) **9** in refluxing acetone when Cs<sub>2</sub>CO<sub>3</sub> was present. The conceivable versatile reactivities of unsaturated carbon–carbon bonds, C<sub>aryl</sub>–Br bonds introduced and of the ester group installed on the benzene ring would now render compounds *S<sub>ic</sub>-8a–h* a unique and invaluable platform for further fabrication of a wide variety of unprecedented chiral chemical entities.

The inherent chirality of heteracalix[4](het)arenes was found very stable and no racemization took place after heating *S<sub>ic</sub>(+)-6a* at 150 °C for 48 h in *o*-dichlorobenzene (Scheme S8†). The stability of inherent chirality was attributable to the presence of 2-(benzyloxy)-5-(*tert*-butyl)-1,3-phenylene in which bulky substituents on both upper and lower rims inhibit the inversion of the macrocyclic ring. It is especially worth addressing that all resulting N<sub>3</sub>,O-calix[2]arene[2]triazines are distinct inherently chiral calix[4](het)arene species. In contrast to well-documented inherently chiral calix[4](het)arenes which are composed of different (hetero)aromatic subrings and the same bridging units, the inherent chirality of all N<sub>3</sub>,O-calix[2]arene[2]triazines in question stems from non-superimposable mirror imaged macrocyclic scaffolds generated by the introduction of different heteroatom linkages. Compound *S<sub>ic</sub>(+)-8a*<sup>25</sup> represents, for instance, an extraordinary example as one oxygen atom difference in the bridging position causes desymmetrization of the



<sup>[a]</sup> *S<sub>ic</sub>-6a* (92.0% ee) was used. <sup>[b]</sup> *S<sub>ic</sub>-6a* (90.0% ee) was used. <sup>[c]</sup> *S<sub>ic</sub>(+)-6g* (96.8% ee) was used.

Scheme 5 Synthetic applications.

molecular structure. The outcomes demonstrated an alternative approach to the design and synthesis of inherently chiral heteracalix[4](het)arenes simply by taking advantage of variable heteroatom linkages.

## Conclusions

In conclusion, we have shown the synthesis of highly enantio-pure inherently chiral N<sub>3</sub>,O-calix[2]arene[2]triazines from enantioselective macrocyclization of linear precursors by means of chiral phosphoric acid-catalyzed intramolecular nucleophilic aromatic substitution reaction. We have also demonstrated efficient functionalization of the resulting macrocycles based on the reactions of bridging the secondary amino moiety. The synthetic method developed and the inherent chirality induced by heteroatom-dictated desymmetrization would open a new opportunity to design and construct a wide variety of diverse chiral macrocycles of unique structural features and potential applications. Study on the functionalization and molecular recognition of inherently chiral heteracalixaromatics<sup>26</sup> is being actively pursued in this laboratory and the results will be reported in due course.

## Data availability

All experimental and computational data are available in ESI.†

## Author contributions

Ying Cheng, Shuo Tong and Mei-Xiang Wang conceived the project and supervised the study. Xing-Chi Li conducted experiments and Xu-Dong Wang performed computational calculations. All authors contributed to discussions and wrote the manuscript.

## Conflicts of interest

There are no conflicts to declare.

## Acknowledgements

We thank the National Natural Science Foundation of China (21920102001, 22050005, 21821001, and 22371161) and Tsinghua University Dushi Program for financial support.

## Notes and references

- V. Böhmer, D. Kraft and M. Tabatabai, *J. Inclusion Phenom. Mol. Recognit. Chem.*, 1994, **19**, 17.
- A. Dalla Cort, L. Mandolini, C. Pasquini and L. Schiaffino, *New J. Chem.*, 2004, **28**, 1198.
- C. D. Gutsche, *Calixarenes: An Introduction*, Royal Society of Chemistry, Cambridge, U.K, 2nd edn, 1989.
- Calixarenes 2001*, ed. Z. Asfari, V. Böhmer, J. Harrowfield, J. Vicens and M. Saadioui, Kluwer Academic, The Netherlands, 2001.



5 *Calixarenes and Beyond*, ed. P. Neri, J. L. Sessler and M.-X. Wang, Springer, Switzerland, 2016.

6 A. Szumna, *Chem. Soc. Rev.*, 2010, **39**, 4274, and references cited therein.

7 (a) G. E. Arnott, *Chem.-Eur. J.*, 2018, **24**, 1744, and references cited therein; (b) M. Tang and X. Yang, *Eur. J. Org. Chem.*, 2023, **26**, e202300738.

8 For selected examples, see: (a) V. Böhmer, L. Merkel and U. Kunz, *J. Chem. Soc., Chem. Commun.*, 1987, 896; (b) L. Zetta, A. Wolff, W. Vogt, K.-L. Platt and V. Böhmer, *Tetrahedron*, 1991, **47**, 1911; (c) S.-Y. Li, Y.-W. Xu, J.-M. Liu and C.-Y. Su, *Int. J. Mol. Sci.*, 2011, **12**, 429; (d) J.-T. Li, L.-X. Wang, D.-X. Wang, L. Zhao and M.-X. Wang, *J. Org. Chem.*, 2014, **79**, 2178; (e) S. J. Nemat, H. Jedrzejewska, A. Prescimone, A. Szumna and K. Tiefenbacher, *Org. Lett.*, 2020, **22**, 5506; (f) D. Sokolova, G. M. Piccina and K. Tiefenbacher, *Angew. Chem., Int. Ed.*, 2022, **61**, e202203384; (g) P. Lhotak, *Org. Biomol. Chem.*, 2022, **20**, 7377; (h) S.-Y. Guo, Z.-A. Zhang, S. Tong, Q.-H. Guo, R. Hua and M.-X. Wang, *Chem. Sci.*, 2023, **14**, 8393; (i) W. Zhu, Y. Cheng, S. Tong and M.-X. Wang, *Org. Lett.*, 2023, **25**, 5105.

9 For selected examples, see: (a) P. C. B. Page, H. Heaney and E. P. Sampler, *J. Am. Chem. Soc.*, 1991, **121**, 6751; (b) K. Iwamoto, H. Shimizu, K. Araki and S. Shinkai, *J. Am. Chem. Soc.*, 1993, **115**, 3997; (c) Y.-D. Cao, J. Luo, Q.-Y. Zheng, C.-F. Chen, M.-X. Wang and Z.-T. Huang, *J. Org. Chem.*, 2004, **69**, 206; (d) Z.-X. Xu, C. Zhang, Z.-T. Huang and C.-F. Chen, *Org. Lett.*, 2008, **10**, 477; (e) W.-Z. Zhang, K. Yang, S.-Z. Li, H. Ma, J. Luo, K.-P. Wang and W.-S. Chung, *Eur. J. Org. Chem.*, 2015, 765; (f) M. Tlustý, H. Dvoráková, J. Cejka, M. Kohout and P. Lhotak, *New J. Chem.*, 2020, **44**, 6490; (g) L. Hodson, K. J. Visagie, M.-P. Smith, L. Loots, D. Kuter, T. M. Snayer and G. E. Arnott, *Chem. Commun.*, 2021, **57**, 11045; (h) F.-M. Yang, J. Luo, Y.-C. Chen and W.-S. Chung, *Synthesis*, 2023, **55**, 2786.

10 J. K. Browne, M. A. McKervey, M. Pitarch, J. A. Russell and J. S. Millership, *Tetrahedron Lett.*, 1998, **39**, 1787.

11 K. Ishibashi, H. Tsue, H. Takahashi and R. Tamura, *Tetrahedron: Asymmetry*, 2009, **20**, 375.

12 S. Tong, J.-T. Li, D.-D. Liang, Y.-E. Zhang, Q.-Y. Feng, X. Zhang, J. Zhu and M.-X. Wang, *J. Am. Chem. Soc.*, 2020, **142**, 14432.

13 X. Zhang, S. Tong, J. Zhu and M.-X. Wang, *Chem. Sci.*, 2023, **14**, 827.

14 Y.-Z. Zhang, M.-M. Xu, X.-G. Si, J.-L. Hou and Q. Cai, *J. Am. Chem. Soc.*, 2022, **144**, 22858.

15 (a) M.-X. Wang, *Chem. Commun.*, 2008, 4541; (b) M.-X. Wang, *Acc. Chem. Res.*, 2012, **45**, 182.

16 W. Maes and W. Dehaen, *Chem. Soc. Rev.*, 2008, **37**, 2393.

17 H. Tuse, K. Ishibashi and R. Tamura, *Top. Heterocycl. Chem.*, 2008, **17**, 73.

18 D.-X. Wang and M.-X. Wang, *Azacalixaromatics in Calixarenes and Beyond*, ed. P. Neri, J. L. Sessler and M.-X. Wang, Springer, 2016, ch. 14, pp. 363–397.

19 R. Hudson and J. L. Katz, *Oxacalixarenes in Calixarenes and Beyond*, ed. P. Neri, J. L. Sessler and M.-X. Wang, Springer, 2016, ch. 15, pp. 399–420.

20 D.-X. Wang and M.-X. Wang, *Acc. Chem. Res.*, 2020, **53**, 1364.

21 Q. Zhang, S. Tong and M.-X. Wang, *Acc. Chem. Res.*, 2022, **55**, 2796.

22 M.-X. Wang and H.-B. Yang, *J. Am. Chem. Soc.*, 2004, **126**, 15412.

23 (a) D. Uraguchi and M. Terada, *J. Am. Chem. Soc.*, 2004, **126**, 5356; (b) T. Akiyama, J. Itoh, K. Yokota and K. Fuchibe, *Angew. Chem.*, 2004, **116**, 1592; *Angew. Chem., Int. Ed.*, 2004, **43**, 1566; (c) M. Terada, *Chem. Commun.*, 2008, 4097; (d) T. Akiyama, J. Itoh and K. Fuchibe, *Adv. Synth. Catal.*, 2006, **348**, 999.

24 (a) D. Nakashima and H. Yamamoto, *J. Am. Chem. Soc.*, 2006, **128**, 9626; (b) M. Rueping, W. Jeawsuwan, A. P. Antonchick and B. J. Nachtsheim, *Angew. Chem.*, 2007, **119**, 2143; *Angew. Chem., Int. Ed.*, 2007, **46**, 2097; (c) M. Rueping, B. J. Nachtsheim, S. A. Moreth and M. Bolte, *Angew. Chem.*, 2008, **120**, 603; *Angew. Chem., Int. Ed.*, 2008, **47**, 593; (d) D. Parmar, E. Sugiono, S. Raja and M. Rueping, *Chem. Rev.*, 2014, **114**, 9047.

25 Deposition Numbers 2303849 (racemic **6a**), 2303851 (racemic **6d**), 2303853 (racemic **6j**), 2304018 (racemic **8d**), 2304020 (enantiopure **S<sub>ic</sub>-6b**) and 234021 (enantiomer **S<sub>ic</sub>-8a**) contain the supplementary crystallographic data for this paper.†

26 Our preliminary study showed that *S<sub>ic</sub>-(+)-6a* and *R<sub>ic</sub>-(-)-6a* are able to form a 1:1 complex with *S*-naproxen in  $\text{CDCl}_3$  with the association constant being  $374 \text{ M}^{-1}$  and  $389 \text{ M}^{-1}$ , respectively.

