RSC Advances



PAPER

View Article Online
View Journal | View Issue



Cite this: RSC Adv., 2017, 7, 20548

Rh(III)-catalyzed sequential C-H activation and annulation: access to N-fused heterocycles from arylazoles and α -diazocarbonyl compounds†

Jiyang Jie,‡^a Haoyi Li,‡^a Songxiao Wu,^a Qingyu Chai,^a Haining Wang*^b and Xiaobo Yang ^b*^a

Received 20th March 2017 Accepted 29th March 2017

DOI: 10.1039/c7ra03262e

rsc.li/rsc-advances


A mild and simple protocol has been developed for the synthesis of N-fused heterocycles from arylazoles and α -diazocarbonyl compounds *via* Rh(III)-catalyzed sequential C-H activation and annulation. Three kinds of N-fused heterocycles derived from arylpyrazoles, arylbenzimidazoles and aryl 1,2,4-triazole were successfully obtained. The reactivity of the other four arylazole analogues was also investigated.

Introduction

N-fused heterocycles are important due to their widespread applications in the pharmaceutical industry and advanced materials.^{1,2} Synthesis of various N-fused heterocycles has thus attracted much attention, particularly for drug development.³ Most syntheses have been completed *via* traditional approaches under harsh reaction conditions like high temperature and microwave irradiation.4 The past decades have seen the development of numerous protocols via transition-metal-catalyzed coupling for this purpose,5 among which Cu- or Pd-catalyzed coupling reactions6 and transition-metal-catalyzed C-H activation7 represent two attractive strategies in forming carboncarbon and carbon-heteroatom bonds. For instance, Fu et al. reported several elegant examples of copper-catalyzed cascade reactions to access N-fused heterocycles from aryl halides and active methylene compounds in 2011 and 2012 (Scheme 1a). 6a,b,d Later, Macgregor and co-workers reported a more direct Rh- or Ru-catalyzed synthesis of pyrazoloisoquinolines using C-H activation of 3-phenylpyrazoles with aryl- and alkylalkynes⁷, avoiding the use of aryl halides; however, an excess amount of oxidant was needed (Scheme 1b). Therefore, the development of a more efficient and greener synthetic approach to access Nfused heterocycles via C-H activation remains highly desirable.

Carbene migratory insertion is a well established method to directly functionalize C–H bonds, with α -diazocarbonyl compounds predominantly used in the directing group (DG)-

assisted Rh(III)-, Co(III)- and Ru(II)-catalyzed C-H activation.9 In 2012, Yu and co-workers first developed chelation-assisted Rh(III)-catalyzed intermolecular cross-coupling of diazomalonates with arene C-H bonds.10 Recently, Li et al. developed a new Ru(II)-catalyzed intermolecular coupling of diazo substrates with arenes to access indoles,11 and the Glorius group reported the first example of Co(III)-catalyzed C-H bond activation of imines with diazo compounds for the synthesis of isoquinolin-3-ones.12 Directing groups previously used in this type of transformation included, among others, imines, 13 oximes,14 amines,15 hydrazines,16 azides,17 azobenzenes18 and amides, 19 delivering the corresponding isoquinolines, benzothiazine, lactams, isoquinolones, indoles and indolines. In view of the importance of N-fused heterocycles and our continued interest in Rh(III)-catalyzed C-H activation reactions and the construction of N-fused heterocycles,20 we envision using arylazoles and diazo compounds as the substrates in a Rh(III)-



Scheme 1 Synthesis of N-fused heterocycles from arylazoles.

[&]quot;Institute of Catalysis for Energy and Environment, College of Chemistry & Chemical Engineering, Shenyang Normal University, Shenyang, Liaoning, 110034, P. R. China. E-mail: bxy1223@gmail.com

^bDepartment of Chemistry, FQRNT Centre for Green Chemistry and Catalysis, McGill University, 801 Sherbrooke St. W., Montreal, Quebec H3A 0B8, Canada. E-mail: haining.wang2@mail.mcgill.ca

[†] Electronic supplementary information (ESI) available. See DOI: 10.1039/c7ra03262e

[‡] These two authors contributed equally to this work.

catalyzed sequential C-H activation and annulation that would provide valuable N-fused heterocycles (Scheme 1c).

Results and discussion

To verify the assumption mentioned above, 3,5-diphenyl-1Hpyrazole (1a) and ethyl 2-diazo-3-oxobutanoate (2a) were selected as coupling partners (Table 1). We studied the reaction between 1a and 2a in the presence of [Cp*RhCl₂]₂ (5 mol%) and AgSbF₆ (20 mol%) at room temperature under Ar atmosphere in THF (2 mL) for 12 h. Fortunately, the desired product 3aa was obtained in 37% yield (Table 1, entry 1). Encouraged by this result, three other solvents were tested, and the results revealed that THF was the best (Table 1, compare entries 2-4). Then, considering the conversion of the substrate, the reaction time was prolonged to 24 h, and we were pleased to see that the yield of 3aa directly increased to 92% (Table 1, entry 5). Furthermore, to investigate the catalytic activity of other transition-metal catalysts, the performances of four other metal catalyst systems were compared. No desired product 3aa was obtained using [Cp*(IrCl₂)]₂/AgNTf₂ or Ru(PPh₃)₃Cl₂/AgSbF₆ as the catalyst system (Table 1, entries 7 and 8). Cp*Co(CO)I₂/AgSbF₆ was less effective; the desired product 3aa was obtained in 24% yield (Table 1, entry 6). By changing the silver salt to AgNTf₂, the yield of 3aa declined to 75%. In addition, no conversion of the substrate was observed without Ag salt or Rh salt (Table 1, entries 10, 11 and 12).

Having established the feasibility of Rh(III)-catalyzed sequential C-H activation and annulation to deliver N-fused

Table 1 Rh(\parallel)-catalyzed sequential C-H activation and annulation of 3,5-diphenyl-1*H*-pyrazole (**1a**) with ethyl 2-diazo-3-oxobutanoate (**2a**): optimization of conditions^a

Entry	Cat. (mol%)	Reaction time	Solvent	Yield ^b (%)
1	[Cp*RhCl ₂] ₂ (5)/AgSbF ₆ (20)	12 h	THF	37
2	[Cp*RhCl ₂] ₂ (5)/AgSbF ₆ (20)	12 h	DCE	Trace
3	[Cp*RhCl ₂] ₂ (5)/AgSbF ₆ (20)	12 h	MeOH	Trace
4	[Cp*RhCl ₂] ₂ (5)/AgSbF ₆ (20)	12 h	EtOH	9
5	[Cp*RhCl ₂] ₂ (5)/AgSbF ₆ (20)	24 h	THF	92
6	Cp*Co(CO)I ₂ (5)/AgSbF ₆ (20)	24 h	THF	24
7	[Cp*(IrCl ₂)] ₂ (5)/AgNTf ₂ (20)	24 h	THF	Trace
8	Ru(PPh ₃) ₃ Cl ₂ (5)/AgSbF ₆ (20)	24 h	THF	0
9	[Cp*RhCl ₂] ₂ (5)/AgNTf ₂ (20)	24 h	THF	75
10	None	24 h	THF	0
11	$[Cp*RhCl_2]_2$ (5)	24 h	THF	0
12	AgSbF ₆ (20)	24 h	THF	0

 $[^]a$ Reaction conditions: 1a (0.2 mmol), 2a (0.25 mmol), catalyst, and solvent (2 mL) were stirred at room temperature under Ar. b Isolated yield.

heterocycles, the reactions of various 5-phenyl-1H-pyrazoles with α -diazocarbonyl compounds were explored to examine the scope and generality of the present process. As depicted in Table 2, at first, 3,5-diphenyl-1H-pyrazole (1a) was kept as the representative reaction partner. A variety of diazoketoesters participated well in this transformation, affording the corresponding products in 84–99% yields (3aa–3ae). α -Diazo- β -diketo

Table 2 Rh(III)-catalyzed sequential C-H activation and annulation of substituted 3,5-diphenyl-1H-pyrazole with α -diazocarbonyl compounds a,b

 R^1 =H, R^3 =CO₂Me, **3cb**, 85%^c R^1 =Me, R^3 =CO₂Me, **3db**, 99%^c

3nb, 25%^c; 3nb', 74%^c

3ob. 99%

 $[^]a$ Reaction conditions: 1 (0.2 mmol), 2 (0.25 mmol), $[Cp*RhCl_2]_2$ (5 mol%)/AgSbF₆ (20 mol%), and THF (2 mL) were stirred at room temperature (ca. 25 °C) under Ar for 24 h. b Isolated yield. c The reaction temperature is 60 °C. d The reaction time is 48 h. e The reaction temperature is 40 °C.

substrates 2f-h also showed good efficiency in the cyclization to give the desired products in good yields. Among them, unsymmetrical diketone 2h performed well; only one regioisomer of 3ah was obtained in 72% yield. Then we found that the reaction of 3-phenyl-5-(p-tolyl)-1H-pyrazole (1b) with methyl 2-diazo-3-oxobutanoate (2b) afforded a mixture of regioisomeric products 3bb and 3bb' in 97% total yield with a ratio of 1:1 based on NMR analysis of the crude mixture. For 5-methyl-3-phenyl-1*H*-pyrazole, substituents at the *ortho* or *para* position including methyl, methoxy, trifluoromethyl, iodo, fluoro, chloro or bromo showed less reactivity; by raising the reaction temperature to 60 °C, all reacted well with diazoketoester (2b) and symmetrical diketone 2i, which was prepared from cyclohexane-1,3-dione, providing 3cb-3lb and 3ci in good vields. Interestingly, a methyl substituent at the *meta* position (1m) underwent the transformation with good regioselectivity to give a single regioisomer in 77% yield (3mb). However, the reaction of the methoxy substituent at the *meta* position (1n) gave separable regioisomeric products 3nb and 3nb' in 25% and 74% yield, respectively. Additionally, 3-methyl-5-(thiophen-3-yl)-1H-pyrazole (10) also presented good reactivity in this reaction, giving the desired product 3ob in 99% yield. Pleasingly, a gramscale reaction was also conducted under standard conditions, and 3aa was obtained in 95% yield.

Table 3 Rh(III)-catalyzed sequential C-H activation and annulation of other arylazoles with α -diazocarbonyl compounds a,b

Next, other arylazoles were examined to broaden the scope of this protocol (Table 3). Unfortunately, only the coupling of 2phenylbenzimidazole (1p) with 2-diazo-3-oxobutanoate (2a) was successful under the optimal reaction conditions above, affording the coupling product 4a in 95% yield. The structure of 4a was confirmed by NMR and HRMS (see ESI†). To our delight, by raising the reaction temperature to 100 °C, the cyclization product 3pa was isolated in 63% yield. This indicates the difference in N-nucleophilicity among indole, pyrazole and imidazole, which is consistent with previous reports. 6e,7m,n At the same reaction temperature, other α-diazocarbonyl compounds also reacted with 1p, giving 3pa-3ph in moderate yields. Chloro and trifluoromethyl substituents of 2-phenylbenzimidazole at the para position were readily converted into the desired products in good yields (3qc, 3ra, 3rb). Notably, this tandem reaction was also compatible with 5-phenyl-1H-1,2,4-triazole to form 3sa in 51% yield. However, for other arylazoles, including aryl imidazole, aryl dihydroimidazole, aryl 1,2,3-triazole and aryl tetrazole, no desired product was observed under our conditions (3tb-3wb).

To better understand this chemistry, we used Density Functional Theory (DFT) calculations to study the activation energy, specifically with regard to the C–H activation step.²¹ The C–H bond activation is facilitated by the basic pyrazole group in the substrate S1 *via* TS1 with an activation energy of 26.2 kcal mol⁻¹. However, the activation energy increases to 33.2 kcal mol⁻¹ when we use imidazole S2 as the substrate (ESI Fig. 1†). These results are consistent with the experimental observations.

Considering the previous reports⁹⁻²¹ on chelation-assisted Rh(III)-catalyzed intermolecular cross-coupling of diazo compounds and the formation of coupling product **4a** described above, we believe that this protocol likely involves sequential C–H activation and intramolecular cyclization; thus a plausible mechanism is proposed (Scheme 2). To begin with, anion exchange occurs between [Cp*RhCl₂]₂ and AgSbF₆, followed by the formation of a cationic Rh(III) species. Then the arylazole **1** undergoes directed C–H cleavage with Cp*Rh(III) to

Scheme 2 Proposed reaction pathway.

^a Reaction conditions: 1 (0.2 mmol), $[Cp*RhCl_2]_2$ (5 mol%)/AgSbF₆ (20 mol%), and THF (2 mL) were stirred at 100 °C under Ar for 24 h. ^b Isolated yield. ^c The reaction temperature is 60 °C.

Paper **RSC Advances**

give a five-membered rhodacyclic intermediate I. Next, the diazo compound 2 coordinates with it to form the intermediate II. Subsequently, migratory insertion of the carbene into the Rh-C bond gives the intermediate III, which undergoes protodemetalation to afford the alkylated product IV, releasing the Rh(III) catalyst for a new catalytic cycle. Finally, an enol intermediate is generated in situ by the tautomerization of intermediate IV, and gives the final product 3 via nucleophilic cyclization.

Conclusions

To sum up, we have developed a facile and practical method for the synthesis of valuable N-fused heterocycles via Rh(III)-catalyzed sequential C-H activation and annulation, using arylazoles and α-diazocarbonyl compounds as substrates. In this protocol, pyrazoles, benzimidazoles and one 1,2,4-triazole were obtained, and most of the pyrazoles were synthesized smoothly and efficiently under mild conditions. Additionally, a comparison of the reactivity for other arylazoles was conducted. This provides a new and alternative method for the construction of diverse and potentially bioactive N-fused heterocycles for drug candidate screening.

Acknowledgements

This work was supported by the National Natural Science Foundation of China (21402126), Natural Science Foundation of Liaoning Province (20141090), Program for Excellent Talents in University of Liaoning Province (LJQ2014121), and Doctoral Fund of Shenyang Normal University.

Notes and references

- 1 (a) E. Vitaku, D. T. Smith and J. T. Njardarson, J. Med. Chem., 2014, 57, 10257; (b) L. M. De Coen, T. S. A. Heugebaert, D. García and C. V. Stevens, Chem. Rev., 2016, 116, 80.
- 2 (a) T. Itoh, Chem. Rev., 2012, 112, 4541; (b) N. A. Romero and D. A. Nicewicz, Chem. Rev., 2016, 116, 10075.
- 3 (a) N. R. Candeias, L. C. Branco, P. M. P. Gois, C. A. M. Afonso and A. F. Trindade, Chem. Rev., 2009, 109, 2703; (b) N. M. Evdokimov, S. Van slambrouck, P. Heffeter, L. Tu, B. Le Calvé, D. Lamoral-Theys, C. J. Hooten, P. Y. Uglinskii, S. Rogelj, R. Kiss, W. F. A. Steelant, W. Berger, J. J. Yang, C. G. Bologa, A. Kornienko and I. V. Magedov, J. Med. Chem., 2011, 54, 2012; (c) A. T. Baviskar, C. Madaan, R. Preet, P. Mohapatra, V. Jain, A. Agarwal, S. K. Guchhait, C. N. Kundu, U. C. Banerjee and P. V. Bharatam, J. Med. Chem., 2011, 54, 5013; (d) W. Wei, J. Wen, D. Yang, J. Du, J. You and H. Wang, Green Chem., 2014, 16, 2988.
- 4 (a) G. Dyker, W. Stirner and G. Henkel, Eur. J. Org. Chem., 2000, 1433; (b) N. Okamoto, K. Sakurai, M. Ishikura, K. Takeda and R. Yanada, Tetrahedron Lett., 2009, 50, 4167.
- 5 (a) I. Nakamura and Y. Yamamoto, Chem. Rev., 2004, 104, 2127; (b) N. T. Patil and Y. Yamamoto, Chem. Rev., 2008, 108, 3395; (c) M. Álvarez-Corral, M. Muñoz-Dorado and I. Rodríguez-García, Chem. Rev., 2008, 108, 3174; (d)

- I. V. Seregin, A. W. Schammel and V. Gevorgyan, Tetrahedron, 2008, 64, 6876; (e) X.-F. Wu, H. Neumann and M. Beller, Chem. Rev., 2013, 113, 1; (f) C. Chen, G. Shang, J. Zhou, Y. Yu, B. Li and J. Peng, Org. Lett., 2014, 16, 1872; (g) X. Fan, M. Yan, Y. Wang and X. Zhang, J. Org. Chem., 2015, 80, 10536.
- 6 (a) J. Lu and H. Fu, J. Org. Chem., 2011, 76, 4600; (b) R. Xie, H. Fu and Y. Ling, Chem. Commun., 2011, 47, 8976; (c) L. Yan, D. Zhao, J. Lan, Y. Cheng, Q. Guo, X. Li, N. Wu and J. You, Org. Biomol. Chem., 2013, 11, 7966; (d) T. Liu and H. Fu, Synthesis, 2012, 44, 2805; (e) J. Peng, G. Shang, C. Chen, Z. Miao and B. Li, J. Org. Chem., 2013, 78, 1242; (f) L.-R. Wen, X.-J. Jin, X.-D. Niu and M. Li, J. Org. Chem., 2015, 80, 90; (g) P. Ruiz-Castillo and S. L. Buchwald, Chem. Rev., 2016, 116, 12564.
- 7 (a) G. Meng, H.-Y. Niu, G.-R. Qu, J. S. Fossey, J.-P. Li and H.-M. Guo, Chem. Commun., 2012, 48, 9601; (b) S. J. Hwang, S. H. Cho and S. Chang, J. Am. Chem. Soc., 2008, 130, 16158; (c) X. Li and M. Zhao, J. Org. Chem., 2011, **76**, 8530; (d) G. Zhao, C. Chen, Y. Yue, Y. Yu and J. Peng, J. Org. Chem., 2015, 80, 2827; (e) N. Umeda, H. Tsurugi, T. Satoh and M. Miura, Angew. Chem., Int. Ed., 2008, 47, 4019; (f) A. G. Algarra, W. B. Cross, D. L. Davies, Q. Khamker, S. A. Macgregor, C. L. McMullin and K. Singh, J. Org. Chem., 2014, 79, 1954; (g) X. Li and M. Zhao, J. Org. Chem., 2011, 76, 8530; (h) R. Wang and J. R. Falck, J. Organomet. Chem., 2014, 759, 33; (i) S. Karthik, J. Ajantha, C. M. Nagaraja, S. Easwaramoorthi and T. Gandhi, Org. Biomol. Chem., 2016, 14, 10255; (j) H. Ikemoto, T. Yoshino, K. Sakata, S. Matsunaga and M. Kanai, J. Am. Chem. Soc., 2014, 136, 5424; (k) M. Choi, J. Park, N. K. Mishra, S.-Y. Lee, J. H. Kim, K. M. Jeong, J. Lee, Y. H. Jung and I. S. Kim, Tetrahedron Lett., 2015, 56, 4678; (l) B. Li, B. Zhang, X. Zhang and X. Fan, Chem. Commun., 2017, 53, 1297; (m) S.-S. Li, Y.-Q. Xia, F.-Z. Hu, C.-F. Liu, F. Su and L. Dong, Chem.-Asian J., 2016, 11, 3165; (n) I. E. Iagafarova, D. V. Vorobyeva, A. S. Peregudov and S. N. Osipov, Eur. J. Org. Chem., 2015, 4950.
- 8 (a) H. M. L. Davies and J. R. Manning, *Nature*, 2008, **451**, 417; (b) Y. Xia, Y. Zhang and J. Wang, ACS Catal., 2013, 3, 2586.
- 9 A. Ford, H. Miel, A. Ring, C. N. Slattery, A. R. Maguire and M. A. McKervey, Chem. Rev., 2015, 115, 9981.
- 10 W.-W. Chan, S.-F. Lo, Z. Zhou and W.-Y. Yu, J. Am. Chem. Soc., 2012, 134, 13565.
- 11 Y. Li, Z. Qi, H. Wang, X. Yang and X. Li, Angew. Chem., Int. Ed., 2016, 55, 11877.
- 12 D. Zhao, J. H. Kim, L. Stegemann, C. A. Strassert and F. Glorius, Angew. Chem., Int. Ed., 2015, 54, 4508.
- 13 (a) S.-G. Lim, J. H. Lee, C. W. Moon, J.-B. Hong and C.-H. Jun, Org. Lett., 2003, 5, 2759; (b) Z.-M. Sun, S.-P. Chen and P. Zhao, Chem.-Eur. J., 2010, 16, 2619; (c) P. Zhao, F. Wang, K. Han and X. Li, Org. Lett., 2012, 14, 3400; (d) Z. Qi, S. Yu and X. Li, Org. Lett., 2016, 18, 700; (e) H. Wang, L. Li, S. Yu, Y. Li and X. Li, Org. Lett., 2016, 18, 2914; (f) M. Choi, J. Park, N. K. Mishra, S.-Y. Lee, J. H. Kim, K. M. Jeong, J. Lee, Y. H. Jung and I. S. Kim, Tetrahedron Lett., 2015, 56, 4678; (g) J. H. Kim, S. Greßies and

RSC Advances Paper

- F. Glorius, Angew. Chem., Int. Ed., 2016, 55, 5577; (h) J. Wang, S. Zha, K. Chen, F. Zhang and J. Zhu, Org. Biomol. Chem., 2016, 14, 4848; (i) X. G. Li, M. Sun, Q. Jin, K. Liu and P. N. Liu, J. Org. Chem., 2016, 81, 3901; (j) Y. Cheng and C. Bolm, Angew. Chem., Int. Ed., 2015, 54, 12349.
- 14 (a) K. Parthasarathy and C.-H. Cheng, J. Org. Chem., 2009, 74, 9359; (b) P. C. Too, S. H. Chua, S. H. Wong and S. Chiba, J. Org. Chem., 2011, 76, 6159; (c) T. K. Hyster and T. Rovis, Chem. Commun., 2011, 47, 11846; (d) Z. Shi, D. C. Koester, M. Boultadakis-Arapinis and F. Glorius, J. Am. Chem. Soc., 2013, 135, 12204; (e) B. Li, P. Jiao, H. Zhong and J. Huang, Synlett, 2013, 24, 2431; (f) R. S. Phatake, P. Patel and C. V. Ramana, Org. Lett., 2016, 18, 292.
- (a) N. K. Mishra, M. Choi, H. Jo, Y. Oh, S. Sharma, S. H. Han,
 T. Jeong, S. Han, S.-Y. Lee and I. S. Kim, *Chem. Commun.*,
 2015, 51, 17229; (b) H. Chu, P. Xue, J.-T. Yu and J. Cheng,
 J. Org. Chem., 2016, 81, 8009.
- 16 (a) J. Wang, M. Wang, K. Chen, S. Zha, C. Song and J. Zhu, Org. Lett., 2016, 18, 1178; (b) C. Song, C. Yang, F. Zhang, J. Wang and J. Zhu, Org. Lett., 2016, 18, 4510.

- 17 S. Gupta, J. Han, Y. Kim, S. W. Lee, Y. H. Rhee and J. Park, J. Org. Chem., 2014, 79, 9094.
- 18 J.-Y. Son, S. Kim, W. H. Jeon and P. H. Lee, *Org. Lett.*, 2015, 17, 2518.
- 19 (a) G. Song, D. Chen, C.-L. Pan, R. H. Crabtree and X. Li, J. Org. Chem., 2010, 75, 7487; (b) M. Presset, D. Oehlrich, F. Rombouts and G. A. Molander, Org. Lett., 2013, 15, 1528; (c) J. Shi, J. Zhou, Y. Yan, J. Jia, X. Liu, H. Song, H. E. Xu and W. Yi, Chem. Commun., 2015, 51, 668; (d) L. Shi, K. Yu and B. Wang, Chem. Commun., 2015, 51, 17277; (e) Y. Wu, P. Sun, K. Zhang, T. Yang, H. Yao and A. Lin, J. Org. Chem., 2016, 81, 2166.
- 20 (a) X. Yang, Y. Jin, H. Liu, Y. Jiang and H. Fu, RSC Adv., 2012,
 2, 11061; (b) X. Yang, Y. Luo, Y. Jin, H. Liu, Y. Jiang and H. Fu, RSC Adv., 2012,
 2, 8258; (c) X. Yang, J. Jie, H. Li and M. Piao, RSC Adv., 2016,
 6, 57371; (d) J. Jie, H. Li, M. Piao and X. Yang, Heterocycles, 2016,
 92, 1215; (e) H. Li, J. Jie, S. Wu, X. Yang and H. Xu, Org. Chem. Front., 2017,
 4, 250.
- 21 S. Qu and C. J. Cramer, J. Org. Chem., 2017, 82, 1195.