

**Finding a Soft Spot for Vanadium: A P-Bound OCP Ligand**

Journal:	<i>ChemComm</i>
Manuscript ID	CC-COM-02-2019-001500.R1
Article Type:	Communication

SCHOLARONE™
Manuscripts

Finding a Soft Spot for Vanadium: A P-Bound OCP Ligand

Lauren N. Grant,^a J. Krzystek,^b Balazs Pinter,^c Joshua Telser,^d Hansjörg Grützmacher,^e and Daniel J. Mindiola*^a

Received 00th January 20xx,
Accepted 00th January 20xx

DOI: 10.1039/x0xx00000x


www.rsc.org/

Transmetallation studies with the phosphoethynolate ion [OCP]⁻, have largely resulted in coordination according to classical Lewis Acid-Base theory. That is, for harder early transition metal ions, O-bound coordination has been observed, whereas in the case of a softer late transition metal ions, P-bound coordination predominates. Herein, we report the use of a V(III) complex, namely [(nacnac)VCl(OAr)] (1) (nacnac⁻ = [ArNC(CH₃)₂CH; Ar = 2,6-ⁱPr₂C₆H₃], to transmetallate [OCP]⁻ and bind via the P-atom as [(nacnac)V(OAr)(PCO)] (2), the first example of a 3d early transition metal that binds [OCP]⁻ via the P-atom. Full characterization of this molecule including HFEPN spectroscopy, SQUID measurements, and theoretical studies are presented.

The ligand class formulated as [XCE]⁻, where X = chalcogen (O, S, Se) and E = pnictogen (N, P, As), has long featured the ubiquitous, ambidentate [OCN]⁻ and [SCN]⁻ ligands.¹ A facile and bulk synthesis of another [XCE]⁻ ligand, the phosphoethynolate ion, [OCP]⁻,¹ has lately been developed, opening a gateway to new chemistry given the softer and more polarizable nature of phosphorous in comparison with the aforementioned nitrogen congener.¹ In recent years [OCP]⁻ has

demonstrated impressive reactivity ranging from organic transformations to new p-, d-, and f-block complexes, which also demonstrate its ambidentate ability. Thus, in “harder” and electropositive metal ions, O-bound coordination of [OCP]⁻ is expected to be dominant,^{3,4i-o} whereas with highly electron-rich W(0) and low-valent Ge, coordination is via the P-atom.^{4a-f} However, the reactivity of [OCP]⁻ with early transition metals remains limited.²⁻⁴ As steps in this direction, we recently reported the divergent reactivity of sodium phosphoethynolate, [Na(OCP)(1,4-dioxane)_{2.5}], with early transition metals.^{2,3} Specifically, reaction with Ti(III) and Sc(III) halide precursors afforded respectively a decarbonylated species having a [Ti₂P₂] core, and reductive coupling of [OCP]⁻, to yield a diphosphoethynolate ion, [OCPPCO]⁴⁻, stabilized by two Sc(III) ions (Scheme 1).^{2,3} Based on Hard/Soft Acid Base (HSAB) theory we inquired if we could achieve the uncommon phosphaketenyl binding mode^{4q} (P-bound) to an early-transition metal ion. Although one anticipates that making the metal electron-rich should promote P-binding to the now softer metal site, reductive decarbonylation readily prevents identification of such putative species.^{4h} Herein, we show how a vanadium(III) precursor can allow us to isolate and fully characterize a high-spin P-bound vanadium(III)-PCO complex that is resistant to decarbonylation. In addition to structural characterization and DFT analysis, we report high-frequency and -field EPR (HFEPN) spectral data and solid-state magnetization data in accord with such species being a high-spin d² complex (S = 1 ground state).

Following the procedure used for the previously reported Ti and Sc complexes, [(nacnac)VCl(OAr)]⁵ (1) (nacnac⁻ = [ArNC(CH₃)₂CH; Ar = 2,6-ⁱPr₂C₆H₃] was treated with 1 equiv. of [Na(OCP)(diox)_{2.5}]^{1a} (diox = 1,4-dioxane) as a slurry in toluene at room temperature, and the mixture was allowed to stir for 12 h.



Scheme 1. Dinuclear, early-transition metal complexes prepared from trivalent precursors followed by reductive

[a,*] Prof. Dr. D. J. Mindiola, Department of Chemistry, University of Pennsylvania, Philadelphia, PA 19104 (USA). E-mail: mindiola@sas.upenn.edu

[b] Dr. J. Krzystek, National High Magnetic Field Laboratory, Florida State University, Tallahassee, FL 32310 (USA)

[c] Prof. Dr. B. Pinter, Department of Chemistry, Universidad Técnica Federico Santa María, Valparaíso, 2390123 (Chile)

[d] Prof. Dr. J. Telser, Department of Biological, Physical and Health Sciences, Roosevelt University, Chicago, IL 60605 (USA)

[e] Prof. Dr. H. Grützmacher, Department of Chemistry and Applied Biosciences, ETH Zürich, Vladimir-Prelog Weg 1, Hönggerberg, 8093, Zürich, (Switzerland)


† For funding, we thank the University of Pennsylvania and the Chemical Sciences, Geosciences, and Biosciences Division, Office of Basic Energy Sciences, Office of Science, U.S. DOE (DEFG02-07ER15893 to D.J.M., synthesis and characterization). L.N.G. thanks NSF GRFP for funding. Part of this work was performed at NHMFL which is supported by the NSF award DMR-1644779 and by the State of FL. We thank Riccardo Suter for providing [Na(OCP)(dioxane)_{2.5}], Dr. Andrew Ozarowski (NHMFL) for his EPR simulation and fit program SPIN and Prof. Eric Schelter and Ekaterina Lapsheva for collecting SQUID data and for insightful discussions.

Supporting information for this article is available on the WWW under <http://www.angewandte.org>

Electronic Supplementary Information (ESI) available: [details of any supplementary information available should be included here]. See DOI: 10.1039/x0xx00000x

decarbonylation (Ti, left) or reductive coupling (Sc, right) of the diphosphaethynolate ion $[\text{OCP}]^-$. Ar represents 2,6- $i\text{Pr}_2\text{C}_6\text{H}_3$.

Examination of the crude reaction mixture revealed formation of a new species displaying broadened resonances in the ^1H NMR spectrum in the 0.5 to 9 ppm range, whereas the paramagnetic resonances attributable to the starting material **1** were no longer present (see Fig. S1 (ESI)).⁶ Although $[\text{Na}(\text{OCP})(\text{diox})_{2.5}]$ is not soluble in toluene, it is critical that coordinating solvents such as THF be avoided in the protocol since their use results in rapid decomposition of the product. To characterize the identity of the green material formed in this reaction, a concentrated toluene solution stored for 16 hours at -35°C afforded green crystals of $[(\text{nacnac})\text{V}(\text{OAr})(\text{PCO})]$ (**2**) (Scheme 2) in 89% yield for which V–P, as opposed to V–O connectivity was unambiguously confirmed by single crystal X-ray diffraction (XRD) studies (Figs. 1 and S3 (ESI)).⁶



Scheme 2. Synthesis of complex **2** from **1** (which is prepared from $[(\text{nacnac})\text{VCl}_2]$ by reaction with NaOAr).⁵ For all instances, Ar represents 2,6- $i\text{Pr}_2\text{C}_6\text{H}_3$. Three possible canonical forms for **2** are represented, of which **2c** is unlikely based on a combination of metrical parameters and theoretical studies.

In addition to XRD confirming the atypical binding mode of $[\text{OCP}]^-$ to an early-transition metal center, the coordination geometry of the phosphaeethynolate ligand reveals an unusual angle due to significant pyramidalization at the phosphorus atom. In contrast to $[(\text{nacnac})\text{Sc}(\text{OAr})(\text{OCP})(\text{THF})]$, where much like the $[\text{Na}(\text{OCP})(\text{diox})_{2.5}]$ precursor the $[\text{OCP}]^-$ ion binds through the O-atom in a near-linear fashion with the metal center ($\angle\text{Sc}-\text{O}-\text{C}$ of $170.89(2)^\circ$) with no other exceptional metrical features, the binding mode of $[\text{OCP}]^-$ in **2** is essentially perpendicular to the V–P vector, with a $\angle\text{V}-\text{P}-\text{C}$ of $85.93(5)^\circ$. This resembles the binding mode of $[\text{OCP}]^-$ with late transition metals such as Re (92.6°) and Au (86.2°).^{4a-c} However, linearity within the $[\text{OCP}]^-$ moiety is still maintained in **2** with a P1–C42–O2 angle of $176.31(1)^\circ$. A strong $\nu(\text{C}-\text{O})$ of 1876 cm^{-1} was seen in the IR spectrum (Fig. S2 (ESI)),⁶ which is of higher energy with respect to the other few known P-bound $[\text{OCP}]^-$ moieties involving late transition metals (Co: 1851 cm^{-1} , Re: 1885 cm^{-1}).^{4a,b} Notably, the long V–P bond distance of $2.4502(4)\text{ \AA}$ significantly diverges from other P-bound examples of $[\text{OCP}]^-$ to 3d metals such as (Co–P: $2.263(1)\text{ \AA}$ and Cu–P: $2.2244(5)\text{ \AA}$).^{4b} It is also significantly longer than known vanadium phosphinidene complexes (V–P: $2.14\text{--}2.19\text{ \AA}$).⁷⁻⁹ Perhaps the most notable metrical parameter in the solid-state structure of **2** is the P–C bond length of $1.6360(18)\text{ \AA}$, which falls more in the range of a single or double bond, but is clearly longer than that of the precursor $[\text{Na}(\text{OCP})(\text{dme})_2]_2$, $1.575(3)\text{ \AA}$ and $1.589(3)\text{ \AA}$.^{1a} The C–

O bond length of $1.174(2)\text{ \AA}$ is commonly observed for systems possessing a double and triple bond, and is slightly shorter than the aforementioned precursor, $1.203(4)\text{ \AA}$ and $1.213(3)\text{ \AA}$.^{1a} In accordance with these observations, Scheme 2 presents the most reasonable canonical structures to be considered. As shown in Scheme 2, structures **2a** and **2b**, are each quite reasonable based on the P–C and C–O distances and will be discussed further in light of computational studies. The metallophosphaketene resonance **2c** is the least likely, on basis of metrical parameters, chiefly the longer V–P distance, observed in the solid-state structure. Further characterization of **2** by an Evans NMR spectroscopic method confirmed the $S = 1$ nature of **2** in toluene solution at room temperature, i.e., a high-spin d^2 complex with a $\mu_{\text{eff}} = 2.47$. The solid-state magnetic behavior of **2** was also probed using DC susceptibility. As shown in Fig. S4 (ESI),⁶ **2** clearly has an $S = 1$ ground state with a sharp decrease in χT below $\sim 20\text{ K}$ due to the effect of zero-field splitting (zfs). The phenomenon of zfs is key to the desired behavior of d-block single ion magnets, which property is not the goal here. However, the zfs itself, chiefly the magnitude and sign of the axial zfs parameter, D , provides valuable information on electronic structure, such as the nature of the frontier MOs and the accessibility of electronic excited states.¹⁰ It was not possible via fitting to distinguish between a positive and a negative value for D , which is a common phenomenon.¹⁰⁻¹³ As a result, the possibility of rhombic zfs (E parameter) was not explored. Our fit suggested a magnitude of D on the order of 4 cm^{-1} , exhibiting temperature independent paramagnetism (TIP), which is qualitatively the consequence of the contribution of thermally populated spin triplet excited states not accounted for by the simple $S = 1$ spin Hamiltonian (sH) employed.^{11,12}




Fig. 1. Molecular structure of complex **2**, with nacnac $i\text{Pr}$ groups and all hydrogen atoms omitted for clarity (left); expansion of the PCO^- ligand coordinated to the $[(\text{nacnac})\text{V}(\text{OAr})]^+$ scaffold showing only the first coordination sphere of the latter (right). All thermal ellipsoids are shown at the 50% probability level. And bond lengths are reported in angstroms (\AA) and angles in degrees ($^\circ$).

To further address the electronic structure of complex **2**, HFEPR spectra were recorded on a ground polycrystalline sample, analogous to that used for the DC susceptibility measurements. The spectra produced by **2** could be recognized as originating from a triplet ($S = 1$) state with moderate zfs. The number and intensities of turning points could indeed be readily simulated. An HFEPR spectrum at a relatively low frequency (113 GHz , Fig. S5)⁶ provided the best representation of a triplet powder pattern,¹⁴ but the frequency was too low to unambiguously determine the absolute sign of D .¹⁴ For this reason, a spectrum at higher frequency (257 GHz , Fig 2) was

needed. The relative amplitudes of the experimental turning points belonging to the transitions originating from the $|S, M_S\rangle = |1, 0\rangle$ spin sublevel are higher than those originating from one of the $|1, \pm 1\rangle$ sublevels, which clearly shows the positive sign of D . To determine the full set of $S = 1$ sH parameters, particularly the g values and rhombic parameter E , a complete field-frequency dependence of the HFEP transitions was recorded.¹⁵ Such a dependence is shown in Fig 3, from which the sH parameters can be extracted with high precision and accuracy as: $D = +2.60(1) \text{ cm}^{-1}$, $|E| = 0.346(5) \text{ cm}^{-1}$, $g_x = 1.962(5)$, $g_y = 1.938(5)$, $g_z = 1.953(5)$.

The sH parameters obtained for **2** can be placed into the context of those for related species inasmuch as possible. We have previously employed HFEP to extract the zfs of a V(III) complex related to **2**, [(nacnac)V(N₃)(Ntol₂)] (tol = 4-MeC₆H₄) (**3**), which gave $D = -1.65 \text{ cm}^{-1}$, $|E| = 0.20 \text{ cm}^{-1}$.¹⁶ This previously studied complex shares the nacnac ligand with **2** and is likewise four-coordinate. That the D values for two such [(nacnac)V(III)(X)(Y)] complexes ($X = \text{OAr}$, $Y = \text{PCO}$ in **2**; $X = \text{N}_3^-$, $Y = \text{NR}_2$ in **3**) are of opposite sign indicates the difficulty in generalizing the correlation between zfs and coordination sphere of a metal ion. An ideally tetrahedral d² complex should exhibit no zfs, and homoleptic four-coordinate V(III) complexes such as [V(ArⁿCl)₄]⁻ (ArⁿCl = C₆Cl₅, 2,4,6-trichlorophenyl, or 2,6-dichlorophenyl) indeed exhibit smaller magnitude zfs ($0.45 \leq |D| \leq 0.59 \text{ cm}^{-1}$) than that observed here for such heteroleptic complexes,¹⁷ which is expected based on ligand-field theory for pseudo-tetrahedral d² complexes. Clearly, a greater database on zfs in four-coordinate V(III) complexes is needed to further evaluate the magnitude and absolute sign of the zfs in **2**. Moreover, the unusual [PCO]⁻ coordination means that **2** may even be distinct amongst mononuclear [(nacnac)V(III)(X)(Y)] complexes.




Fig. 2. An HFEP spectrum of **2** recorded at 10 K and 257 GHz (black trace). A V^{IV} impurity signal at $g = 1.98$ ($\sim 9.3 \text{ T}$) has been removed. Coloured traces are simulations using sH parameters: $S = 1$, $|D| = 2.62 \text{ cm}^{-1}$, $|E| = 0.36 \text{ cm}^{-1}$, $g_x = 1.96$, $g_y = 1.94$, $g_z = 1.95$. The particular turning points are labelled by canonical direction (x, y, z) of the magnetic field with respect to the \mathbf{D} tensor and have subscripts 1 for those that belong to the transition originating from the $|S, M_S\rangle = |1, 0\rangle$ spin sublevel while those with subscripts 2 originate from one of the $|1, \pm 1\rangle$ sublevels.




Fig. 3. A 2D field vs. frequency (energy) plot of HFEP spectral turning points in **2**. Squares are experimental data; curves were simulated using sH parameters as in the text. Red curves: turning points with magnetic field parallel to the x -axis of the zfs tensor; blue curves: $B_0 \parallel y$; black curves: $B_0 \parallel z$. The two vertical dashed lines indicate the frequencies at which the spectra shown in Figures S7 and 2, respectively, were recorded.

To elucidate the electronic structure of **2** in detail and reveal the most likely canonical resonance of the [OCP]⁻ ligand in **2**, we turned to DFT (B3LYP/def2-TZVP(-f)).⁶ These calculations on non-truncated models revealed an equilibrium structure for **2** with metrics in good agreement with the X-ray structural parameters. For example, the quasi-perpendicular $\angle \text{V-P-C}$ angle (85.0°) and distances V-P (2.46 \AA), P-C (1.66 \AA) and C-O (1.17 \AA) are well reproduced by these DFT calculations, substantiating the method of calculation. To characterize the bending and bonding of the V-PCO motif in **2** we used quasi-restricted orbitals (QROs). QROs offer a reliable and intuitive description of the unrestricted wavefunctions (with different α and β orbital subsets) through a conceptual picture resembling that of the restricted open-shell solution. The two singly occupied QRO's of **2**, d_{z^2} and $d_{x^2-y^2}$, (Fig 4; see also Fig. S6 (ESI))⁶ unmistakably reveal metal-centered unpaired electrons without notable delocalization to any of the ligands, i.e., a d² V(III) center, conforming to a spin density of 1.96 calculated for the vanadium of **2** using the Löwdin method, and consistent with the HFEP results. Close inspection of the orbital entitled $\pi_{\text{PCO}+d_{xz}}$ in Fig. 4 unveils the vanadium-PCO σ bond, which is the main V-PCO interaction, as originating from the overlap of one of the π orbitals of the [OCP]⁻ fragment and a σ -type vanadium d orbital, d_{xz} , in the quasi-tetrahedral arrangement of **2**.

The spatial characteristics of this PCO π -bond account for most of the striking structural features observed, such as the elongated P-C and significantly shortened C-O distances in the [OCP]⁻ ligand. These spatial features indicate that this π_{PCO} orbital has bonding character along P-C and antibonding character along C-O, which, under normal circumstances like in the free ligand, lead to a short P-C and a long C-O bond (i.e. to P=C-O⁻ formally). In **2** however, electron donation takes place from this π_{PCO} orbital to the vanadium, witnessed by the $\pi_{\text{PCO}+d_{xz}}$ molecular orbital in Figure 4, from which partial depopulation of π_{PCO} results in diminished attraction and

diminished repulsion along the P–C and C–O axes. The latter effects lead to the weakening of the P–C and strengthening of the C–O bonds relative to that of free $[\text{OCP}]^-$, consistent with the structural features in **2**. Also, due to the polarization of this PCO π -bond towards the phosphorous atom, the $\angle\text{V–P–C}$ deviates from angles typical of side-on or κ^2 coordination and towards a perpendicular κ^1 -type binding at P. As a result, this MO analysis implies that the formal electronic structure of $[\text{OCP}]^-$ changes from $\text{P}\equiv\text{C–O}^-$ (in free ligand) to $^-\text{P}=\text{C}=\text{O}$ upon binding to V(III) and, concomitantly, that the Lewis structure of **2** is dominated by resonance **2a** (Scheme 2). The $[\text{OCP}]^-$ structural motif is further supported by the Mayer bond indices of 1.75 and 2.06 calculated for P=C and C=O, respectively, in **2**.

In summary, we report the first example of the $[\text{OCP}]^-$ reagent binding to a highly electropositive 3d early transition metal, vanadium(III), via the P-atom, and the unexpected $\angle\text{V–P–C}$ of $85.93(5)^\circ$. Our perception is that the HSAB concept's "soft-soft" argument, which can be translated into the MO framework as the electron-rich V(III) having a better orbital energy match with phosphorous orbitals than with oxygen orbitals, explains the site-selective P-binding of $[\text{OCP}]^-$ to vanadium(III). On the other hand, the almost perpendicular binding and elongated/shortened P–C/C–O bonds of $[\text{OCP}]^-$ can be best rationalized, as detailed above, through the characteristics of the π_{PCO} orbital that forms the σ interaction with the vanadium center (Fig 4).




Fig. 4. Relevant QROs of triplet **2**. The upper two QROs represent the V-centered unpaired electrons, while HOMO-4 depicts the V–P σ interaction. Isodensity values are ± 0.04 .

Conflicts of interest

There are no conflicts to declare.

Notes and references

† CCDC 1890542 contains the supplementary crystallographic data for **2**. These data can be obtained free of charge from The Cambridge Crystallographic Data Centre via www.ccdc.cam.ac.uk/data_request/cif.

- (a) F. F. Puschmann, D. Stein, D. Heift, C. Hendriksen, Z. A. Gal, H.-F. Grützmacher and H. Grützmacher, *Angew. Chem. Int. Ed.*, 2011, **50**, 8420; (b) A. R. Jupp and J. M. Goicoechea, *Angew. Chem. Int. Ed.*, 2013, **52**, 10064; (c) J. M. Goicoechea and H. Grützmacher, *Angew. Chem. Int. Ed.*, 2018, **57**, 16968;

- F. Tambornino, A. Hinz, R. Köppe and J. M. Goicoechea, *Angew. Chem. Int. Ed.*, 2018, **57**, 8230.
- L. N. Grant, B. Pinter, B. C. Manor, H. Grützmacher and D. J. Mindiola, *Angew. Chem. Int. Ed.*, 2018, **57**, 1049.
- L. N. Grant, B. Pinter, B. C. Manor, R. Suter, H. Grützmacher and D. J. Mindiola, *Chem. Eur. J.*, 2017, **23**, 6272.
- (a) S. Alidori, D. Heift, G. Santiso-Quinones, Z. Benkő, H. Grützmacher, M. Caporali, L. Gonsalvi, A. Rossin and M. Peruzzini, *Chem. Eur. J.*, 2012, **18**, 14805; (b) L. Liu, D. A. Ruiz, F. Dahcheh, G. Bertrand, R. Suter, A. M. Tondreau and H. Grützmacher, *Chem. Sci.*, 2016, **7**, 2335; (c) A. R. Jupp, M. B. Geeson, J. E. McGrady and J. M. Goicoechea, *Eur. J. Inorg. Chem.*, 2016, **2016**, 639; (d) S. Yao, Y. Xiong, T. Szilvási, H. Grützmacher and M. Driess, *Angew. Chem. Int. Ed.*, 2016, **55**, 4781; (e) Y. Wu, L. Liu, J. Su, J. Zhu, Z. Ji and Y. Zhao, *Organometallics*, 2016, **35**, 1593; (f) Y. Xiong, S. Yao, T. Szilvási, E. Ballester-Martínez, H. Grützmacher and M. Driess, *Angew. Chem. Int. Ed.*, 2017, **56**, 4333; (g) T. P. Robinson, M. J. Cowley, D. Scheschkewitz and J. M. Goicoechea, *Angew. Chem. Int. Ed.*, 2014, **54**, 683; (h) M. Joost, W. J. Transue and C. C. Cummins, *Chem. Commun.*, 2017, **53**, 10731; (i) G.-L. Hou, B. Chen, W. J. Transue, Z. Yang, H. Grützmacher, M. Driess, C. C. Cummins, W. T. Borden and X.-B. Wang, *J. Am. Chem. Soc.*, 2017, **139**, 8922; (j) R. J. Gilliard, Jr., R. Suter, E. Schrader, Z. Benkő, A. L. Rheingold, H. Grützmacher and J. D. Protasiewicz, *Chem. Commun.*, 2017, **53**, 12325; (k) D. Heift, Z. Benkő and H. Grützmacher, *Dalton Trans.*, 2014, **43**, 5920; (l) C. Camp, N. Settineri, J. Lefèvre, A. R. Jupp, J. M. Goicoechea, L. Maron and J. Arnold, *Chem. Sci.*, 2015, **6**, 6379; (m) C. J. Hoerger, F. W. Heinemann, E. Louyriac, L. Maron, H. Grützmacher and K. Meyer, *Organometallics*, 2017, **36**, 4351; (n) R. J. Gilliard, D. Heift, Z. Benkő, J. M. Keiser, A. L. Rheingold, H. Grützmacher and J. D. Protasiewicz, *Dalton Trans.*, 2018, **47**, 666; (o) S. Bestgen, Q. Chen, N. H. Reesa and J. M. Goicoechea, *Dalton Trans.*, 2018, **47**, 13016; (p) Y. Mei, D.-J. Wu, J. E. Borger and Hansjörg Grützmacher, *Angew. Chem. Int. Ed.*, 2018, **57**, 5512; (q) D. W. N. Wilson, A. Hinz and J. M. Goicoechea, *Angew. Chem. Int. Ed.*, 2018, **57**, 2188.
- B. L. Tran, B. Pinter, A. J. Nichols, C.-H. Chen, F. T. Konopka, R. Thompson, J. Krzystek, A. Ozarowski, J. Telser, M.-H. Baik, K. Meyer and D. J. Mindiola, *J. Am. Chem. Soc.*, 2012, **134**, 13035.
- See Supporting Information.
- U. J. Kilgore, H. Fan, M. Pink, E. Urnezis, J. D. Protasiewicz and D. J. Mindiola, *Chem. Commun.*, 2009, 4521.
- F. Basuli, B. C. Bailey, J. C. Huffman, M.-H. Baik and D. J. Mindiola, *J. Am. Chem. Soc.*, 2004, **126**, 1924.
- T. W. Graham, K. A. Udachin, M. Z. Zgierski and A. J. Carty, *Organometallics*, 2011, **30**, 1382.
- B. Sieklucka and D. Pinkowicz, *Molecular Magnetic Materials: Concepts and Applications*, 2017, 1.
- R. Boča, *Coord. Chem. Rev.*, 2004, **248**, 757.
- R. Boča, *Magnetic Parameters and Magnetic Functions in Mononuclear Complexes Beyond the Spin-Hamiltonian Formalism*. In: Mingos, D. M. P. (ed) *Magnetic Functions Beyond the Spin-Hamiltonian. Structure and Bonding*, 2005, **117**, 1. Springer, Berlin, Heidelberg.
- J. Krzystek and J. Telser, *Dalton Trans.*, 2016, **45**, 16751.
- J. Telser, *eMagRes*, 2017, **6**, 207.
- J. Krzystek, A. Ozarowski and J. Telser, *Coord. Chem. Revs.*, 2006, **250**, 2308.
- B. L. Tran, J. Krzystek, A. Ozarowski, C.-H. Chen, M. Pink, J. A. Karty, J. Telser, K. Meyer and D. J. Mindiola, *Eur. J. Inorg. Chem.*, 2013, **2013**, 3916.
- P. J. Alonso, J. Forniés, M. A. GarcíaMonforte, A. Martín and B. Menjón, *Chem. Eur. J.*, 2005, **11**, 4713.
- W. Mowat, A. J. Shortland, N. J. Hill and G. Wilkinson, *J. Chem. Soc., Dalton Trans.*, 1973, 770.

