

ChemComm

Chemical Communications

rsc.li/chemcomm



ISSN 1359-7345



COMMUNICATION

Matthew D. Lloyd *et al.*

A novel colorimetric assay for α -methylacyl-CoA racemase 1A (AMACR; P504S) utilizing the elimination of 2,4-dinitrophenolate



Cite this: *Chem. Commun.*, 2017, 53, 5087

Received 18th January 2017,
Accepted 17th March 2017

DOI: 10.1039/c7cc00476a

rsc.li/chemcomm

A novel colorimetric assay for α -methylacyl-CoA racemase 1A (AMACR; P504S) utilizing the elimination of 2,4-dinitrophenolate†

Maksims Yevglevskis,^a Guat L. Lee,^a Amit Nathubhai,^a Yoana D. Petrova,^a Tony D. James,^{ib} Michael D. Threadgill,^a Timothy J. Woodman^a and Matthew D. Lloyd^{ib}*^a


α -Methylacyl-CoA racemase (AMACR; P504S) regulates branched-chain fatty acid degradation, activates Ibuprofen and is a recognised cancer drug target. A novel, facile colorimetric assay was developed based on elimination of 2,4-dinitrophenolate. The assay was used to test 5 known inhibitors, determining IC₅₀ and K_i values, reversibility and characterizing irreversible inhibition.

α -Methylacyl-CoA racemase (AMACR;‡ P504S; E.C. 5.1.99.4) performs a key role in branched-chain fatty acid β -oxidation and the pharmacological activation of Ibuprofen and related drugs.^{1,2} AMACR catalyses its reaction by a deprotonation/reprotonation mechanism,^{3–5} in which either epimer of a 2-methylacyl-CoA or 2-arylpropanoyl-CoA substrate is converted to a near 1:1 mixture of epimers.^{3,5} Conversion of *R*-2-methylacyl-CoAs to their *S*-epimers enables degradation by β -oxidation.^{1,6} *R*-Ibuprofen and most related *R*-2-APA drugs¹ are pharmacologically activated by conversion to their corresponding acyl-CoA esters, before epimerization by AMACR. Hydrolysis of the epimeric products gives a mixture of *R*- and *S*-2-APA drugs, with the *R*-product recycled by the same pathway. Thus, inactive *R*-2-APA drugs are converted into their *S*-enantiomers, which are potent inhibitors of cyclooxygenase.⁷

Concentrations of AMACR are increased in all prostate cancers,^{8,9} and in several other cancers.^{10–13} Increased AMACR catalytic activity has also been reported in prostate cancer.^{14,15} Reducing the cellular AMACR 1A^{15–17} protein using siRNA or shRNA reduces proliferation of prostate cancer cell lines *via* a pathway which is synergistic with androgen withdrawal.¹⁵ Some advanced prostate cancer cell lines revert from androgen-independent to androgen-dependent status¹⁷ upon AMACR knock-down. Consequently, AMACR has attracted considerable interest as a novel prostate cancer drug target^{16,18–20} and biomarker.^{1,21}

Despite this interest, few inhibitors of AMACR have been reported. The majority of inhibitors are rationally designed acyl-CoA derivatives,^{18–20,22} which do not comply with Lipinski's guidelines²³ and need to be delivered as carboxylate prodrugs. Drug delivery can be limited by *in vivo* conversion of the prodrug to the acyl-CoA drug, although preliminary *in vivo* studies²⁴ appear promising. *N*-Dodecyl-*N*-methylcarbamoyl-CoA **1**, a transition-state analogue, is the most potent inhibitor reported to date.¹⁹ Several non-specific protein-modifying agents also inhibit AMACR.¹⁶ No structure–activity relationships have been reported, probably due to the difficulties in assaying enzyme activity.

AMACR catalyses the conversion of either *R*- or *S*-2-methylacyl-CoA esters into a near 1:1 mixture of epimers^{3,5} (Scheme 1A). Assaying AMACR activity is challenging due to the reversibility of the reaction and the difficulties in differentiating the epimeric products since the stereochemical centre undergoing a change in configuration is remote from the stereochemical centres in the CoA moiety. Several endpoint assays have been reported and used for inhibitor testing, such as those using chiral acyl-CoA substrates^{18,19} or their derivatives.²⁵ Notably, Wilson *et al.*¹⁶ reported screening



Scheme 1 Reactions catalysed by AMACR. (A) Conversion of *R*- or *S*-2-methylacyl-CoA into an epimeric mixture, as illustrated by ibuprofenoyl-CoA **2**. Assays commonly use an acyl-CoA substrate with α -³H which is 'washed out',¹⁶ or unlabelled substrate in the presence of ²H₂O followed by ¹H NMR analysis of the 'washed in' product.^{3,5} (B) The elimination of HF from (2*R*,3*R*)-3-fluoro-2-methyldecanoyl-CoA **3** to give unsaturated product **4**.

^a Drug & Target Development, Department of Pharmacy & Pharmacology, University of Bath, Claverton Down, Bath BA2 7AY, UK. E-mail: M.D.Lloyd@bath.ac.uk;

Fax: +44-1225-386114; Tel: +44-1225-386786

^b Department of Chemistry, University of Bath, Claverton Down, Bath BA2 7AY, UK

† Electronic supplementary information (ESI) available: Experimental details including synthesis of compounds, kinetic plots, inhibitor testing. See DOI: 10.1039/c7cc00476a




of a library of 5000 compounds using a modified radiolabelled assay, and identified several non-specific protein modifying agents that were potent inhibitors of AMACR. Their assay was used for kinetic analysis of AMACR, but it is an endpoint assay measuring substrate conversion at a single time-point and extensive manipulation of samples is required. In addition the assay is likely to be complicated by the presence of a kinetic isotope effect. A continuous circular dichroism (CD) assay for the closely related *Mycobacterium tuberculosis* enzyme (MCR), which follows the conversion of either *R*- or *S*-ibuprofenoyl-CoA 2 to a near racemic mixture,²⁶ has been used for limited testing of inhibitors.²² This assay is not subject to a kinetic isotope effect, but is low-throughput as only one sample can be analysed at the same time.

It has also been discovered that AMACR catalyses the elimination of fluoride from substrates such as 3-fluoro-2-methyldecanoyl-CoA (Scheme 1B), probably *via* an E1cb mechanism.²⁷ This reaction has the advantage of being irreversible, and the methyl peaks of substrate and product do not overlap in the ¹H spectrum allowing a simple route to determine the extent of conversion. This reaction has been used for the preliminary characterization of known inhibitors.²⁸

The elimination of fluoride from 3-fluoro-2-methyldecanoyl-CoA led us to consider whether a colorimetric leaving group might eliminate from a suitable substrate. The pK_a of HF (3.2)²⁹ and 2,4-dinitrophenol (3.93)³⁰ are similar, suggesting the conjugate bases will have similar leaving-group ability. Significantly, 2,4-dinitrophenolate 5 is a well-characterized chromophore ($\epsilon_{354} = 15\,300\text{ M}^{-1}\text{ cm}^{-1}$).³¹ AMACR is known to accept substrates with a wide variety of side-chain structures,¹ and hence the use of a substrate containing a 2,4-dinitrophenoxy- moiety as a chromogenic leaving group was investigated.

Racemic 3-(2,4-dinitrophenoxy)-2-methylpropanoyl-CoA 6 was synthesized (Scheme 2) by reaction of Sanger's reagent 7 with 2-methylpropan-1,3-diol 8 under basic conditions. Treatment of alcohol 9 with CrO₃ and H₂SO₄ resulted in the racemic acid 10. This was coupled with CoA-SH^{5,27,28} to give the desired substrate 6.


Incubation of 6 with active AMACR resulted in an elimination reaction (Scheme 3A). The sample containing active enzyme possessed an intense yellow colour, which was absent from negative controls (Scheme 3B). ¹H NMR analysis confirmed that 2,4-dinitrophenolate 5 and the predicted unsaturated product 11 were formed (Scheme 3C). Incubation of substrate 6 with active



Scheme 3 Elimination of 2,4-dinitrophenolate 5 from 3-(2,4-dinitrophenoxy)-2-methylpropanoyl-CoA 6 by AMACR. (A) Reaction catalysed by enzyme; (B) NMR tubes showing samples containing active enzyme (yellow) and heat-inactivated enzyme (clear); (C) ¹H NMR spectra (500 MHz) of samples in B showing position of methyl groups of substrate 6 (red circles) and product 11 (green circles); (D) time course of reaction followed by monitoring absorbance of 2,4-dinitrophenolate 5 at 354 nm, showing negative control (heat-inactivated enzyme; red) and active enzyme (green). Assays comprised 40 μM 6 and recombinant AMACR (9.25 μg) in 100 μL NaH₂PO₄-NaOH, pH 7.4. Data are means \pm SDM ($n = 3$).

enzyme in a microtitre plate led to a rapid increase in absorbance at 354 nm, which was not observed in controls containing heat-inactivated enzyme (Scheme 3D). This reaction occurred in an enzyme concentration-dependent manner (ESI,† Fig. S1 and S2).

Assay conditions were optimized (ESI,† Fig. S3–S5) by investigating the effect of various additives on enzymatic activity. Addition of 2-mercaptoethanol, dithiothreitol or BSA had no significant effect on activity. Addition of 0.1% (v/v) Triton X-100 to the assay mixture resulted in a modest reduction of activity. This is probably due to the formation of micelles, as the amounts used are above the reported critical micellar concentrations for Triton X-100 (0.19 and 1.25 mM).³² In contrast, addition of 1.5% (w/v) *N*-lauroyl-sarcosine reduced activity to levels observed in negative controls. The enzyme proved to be tolerant of DMSO (ESI,† Fig. S5), up to at least 8% (v/v) ($\sim 1.12\text{ M}$). Michaelis-Menten kinetic behaviour was observed under the optimized assay conditions (ESI,† Fig. S6) and the following kinetic parameters were determined: $K_m = 58\ \mu\text{M}$; $V_{\text{max}} = 112\ \text{nmol min}^{-1}\ \text{mg}^{-1}$; $k_{\text{cat}} = 0.088\ \text{s}^{-1}$; $k_{\text{cat}}/K_m = 1517\ \text{M}^{-1}\ \text{s}^{-1}$. Optimised conditions for the assay were $\sim 8\ \mu\text{g}$ enzyme in 200 μL NaH₂PO₄-NaOH aq., pH 7.4 containing 40 μM 6 and up to 8% (v/v) DMSO at 30 °C, monitoring at 354 nm. Typical rates for active enzyme and negative controls



Scheme 2 Synthesis of substrate 6. Reagents & conditions: (i) Na metal, 83%; (ii) CrO₃, conc. H₂SO₄, acetone, 67%; (iii) carbonyldiimidazole, CH₂Cl₂; (iv) CoA-SH (Li⁺)₃, 0.1 M NaHCO_{3(aq)}/THF (1:1).






Fig. 1 Inhibitors used in assay. (A) Chemical structures and determined IC_{50} values from dose-response curves; (B) example assay in 96-well plate. The pink colour results from the presence of the inhibitor Rose Bengal **14** and the yellow colour from the 2,4-dinitrophenolate **5** produced by the enzyme from substrate **6**; (C) dose-response curve for Rose Bengal **14**.

(means \pm SD, $n = 3$) are 33 ± 1.9 and 0.72 ± 0.24 nmol min⁻¹ mg⁻¹. The limit of detection is ~ 1.4 nmol min⁻¹ mg⁻¹. The dynamic range is ~ 1.4 –255 nmol min⁻¹ mg⁻¹. The sensitivity threshold is ~ 0.5 –1.0 nmol min⁻¹ mg⁻¹. The Z' value³³ for the assay was 0.906 (see ESI,† for further details on how these values were calculated). The presence of other enzymes such as branched-chain acyl-CoA oxidases (ACOXs) could potentially generate a false positive signal, but this can be avoided by using *R-6* as ACOXs are known to be specific for *S*-2-methylacyl-CoA substrates.⁶

This new colorimetric assay for AMACR was validated using a series of known inhibitors for which bench-mark values are available (Fig. 1). *N*-Dodecyl-*N*-methylcarbamoyl-CoA **1** and ibuprofenoyl-CoA **2** were chosen as representative acyl-CoA inhibitors. Ebselen **12**, Ebselen Oxide **13** and Rose Bengal **14** were chosen as examples of the non-specific protein modifying agents reported by Wilson *et al.*¹⁶ Inhibitor potency was initially assessed using dose-response curves to determine IC_{50} values (see Fig. 1B and C for example).

N-Dodecyl-*N*-methylcarbamoyl-CoA **1** had a very low IC_{50} value of 0.4 nM determined by this assay, confirming its very high potency. This compares with the previous value of 98 nM, determined with an endpoint assay.¹⁹ Ibuprofenoyl-CoA **2** had a modest potency with an IC_{50} value of 554 nM. For both inhibitors activity was restored upon rapid dilution of the inhibitor, showing reversible inhibition. Michaelis-Menten kinetic analysis of compounds **1** and **2** showed that they were both competitive inhibitors of AMACR, with K_i values of 0.65 and 60 nM, respectively (Fig. 2 and ESI,† Fig. S7 and S8). It is noteworthy that both compounds appear to be much more potent when assayed using this method compared with other methods. K_i values of 98 nM and 56 μ M were previously reported in the literature for *N*-dodecyl-*N*-methylcarbamoyl-CoA **1**¹⁹ and racemic ibuprofenoyl-CoA **2**³⁴ (with native rat enzyme), respectively. The reasons for the difference in apparent potency are not clear, but it probably




Fig. 2 Analysis of inhibitor **1**. (A) Reversibility experiment, showing enzymatic activity is restored upon dilution of concentrated enzyme with 13.5 nM **1** to 0.27 nM **1**; (B) kinetic analysis to determine K_i value for **1**.

results from differences in assay conditions including micelle formation by acyl-CoA substrates and inhibitors.

Analysis of the non-specific protein modification agents **12**, **13** and **14**, identified by Wilson *et al.*¹⁶ as AMACR inhibitors, confirmed their potency. Ebselen **12** gave an IC_{50} value of 133 nM, compared to the previously reported value¹⁶ of 2.789 μ M. Rapid dilution of Ebselen **12** resulted in part restoration of activity in our assay (ESI,† Fig. S7), consistent with irreversible inhibition. This is consistent with the complex behaviour observed by Wilson *et al.*¹⁶

Similarly, Ebselen Oxide **13** was a potent inhibitor with an IC_{50} value of 774 nM compared to the previously reported value¹⁶ of 795 nM. Rapid dilution of **13** did not result in full restoration of activity, suggesting (covalent) irreversible inhibition. Further investigation showed that non-saturating time-dependent inhibition of AMACR occurred, with a second-order rate constant of 116 ± 8 M⁻¹ s⁻¹ (Fig. 3 and ESI,† Fig. S10), consistent with **13** being a non-specific inhibitor operating by a one-step mechanism. Compound **14** was a reversible inhibitor ($IC_{50} = 3259$ nM), compared to 10 000 nM.¹⁶ Prolonged incubation caused photolytic protein degradation (ESI,† Fig. S9).

AMACR has attracted much attention as both a novel drug target and cancer biomarker since the first reports of its involvement in prostate cancer in 2003.¹⁵ Exploitation of this discovery has been very limited due to the absence of a suitable assay. Our novel colorimetric assay provides a versatile platform for detailed characterization of inhibitors and determination of




Fig. 3 Inactivation of AMACR by Ebselen Oxide **13**. (A) Time-dependent inactivation of recombinant AMACR by Ebselen Oxide **13**. Concentrations of Ebselen Oxide **13** (inset) are given for the pre-incubation phase before dilution of the enzyme with substrate **6** in the assay; (B) dependence of inhibition on Ebselen Oxide **13** concentration. First order rate constants (s⁻¹) derived from **A** was plotted vs. Ebselen Oxide **13** concentration in the pre-incubation phase. Data are mean \pm standard error.



structure–activity relationships. The assay also allows for inter-rogation of the complex biology of AMACR and its role in lipid metabolism and cancer. Similar colorimetric assays could potentially be adapted for use with other enzymes catalysing reactions *via* enolate intermediates, including other racemases,³⁵ acyl-CoA oxidases^{36,37} and other enzymes,^{38,39} several of which are of academic, medicinal, or biotechnological interest.

This work was funded by Prostate Cancer UK (S10-03 and PG14-009), a University of Bath Overseas Research Studentship, and a Biochemical Society Summer Vacation studentship (2015). The authors are members of the Cancer Research @ Bath (CR@B) network.

Notes and references

‡ Abbreviations used: AMACR, α -methylacyl-CoA racemase splice variant 1A; 2-APA, 2-arylpropanoic acid (profen); BSA, bovine serum albumin; CoA, coenzyme A; MCR, 2-methylacyl-CoA racemase from *M. tuberculosis*.

- M. D. Lloyd, M. Yevglevskis, G. L. Lee, P. J. Wood, M. D. Threadgill and T. J. Woodman, *Prog. Lipid Res.*, 2013, **52**, 220–230.
- C. Reichel, R. Brugger, H. Bang, G. Geisslinger and K. Brune, *Mol. Pharmacol.*, 1997, **51**, 576–582.
- D. J. Darley, D. S. Butler, S. J. Prideaux, T. W. Thornton, A. D. Wilson, T. J. Woodman, M. D. Threadgill and M. D. Lloyd, *Org. Biomol. Chem.*, 2009, **7**, 543–552.
- P. Bhaumik, W. Schmitz, A. Hassinen, J. K. Hiltunen, E. Conzelmann and R. K. Wierenga, *J. Mol. Biol.*, 2007, **367**, 1145–1161.
- T. J. Woodman, P. J. Wood, A. S. Thompson, T. J. Hutchings, G. R. Steel, P. Jiao, M. D. Threadgill and M. D. Lloyd, *Chem. Commun.*, 2011, **47**, 7332–7334.
- P. P. VanVeldhoven, K. Croes, S. Asselberghs, P. Herdewijn and G. P. Mannaerts, *FEBS Lett.*, 1996, **388**, 80–84.
- J. A. Mitchell, P. Akarasereenont, C. Thiernemann, R. J. Flower and J. R. Vane, *Proc. Natl. Acad. Sci. U. S. A.*, 1993, **90**, 11693–11697.
- J. Luo, S. Zha, W. R. Gage, T. A. Dunn, J. L. Hicks, C. J. Bennett, C. N. Ewing, E. A. Platz, S. Ferdinandusse, R. J. Wanders, J. M. Trent, W. B. Isaacs and A. M. De Marzo, *Cancer Res.*, 2002, **62**, 2220–2226.
- Z. Jiang, B. A. Woda, K. L. Rock, Y. D. Xu, L. Savas, A. Khan, G. Pihan, F. Cai, J. S. Babcook, P. Rathanaswami, S. G. Reed, J. C. Xu and G. R. Fanger, *Am. J. Surg. Path.*, 2001, **25**, 1397–1404.
- Z. Jiang, G. R. Fanger, B. F. Banner, B. A. Woda, P. Algate, K. Dresser, J. C. Xu, S. G. Reed, K. L. Rock and P. G. Chu, *Cancer Detect. Prev.*, 2003, **27**, 422–426.
- A. K. Witkiewicz, S. Varambally, R. L. Shen, R. Mehra, M. S. Sabel, D. Ghosh, A. M. Chinnaiyan, M. A. Rubin and C. G. Kleer, *Cancer Epidemiol., Biomarkers Prev.*, 2005, **14**, 1418–1423.
- C.-F. Li, F.-M. Fang, J. Lan, J.-W. Wang, H.-j. Kung, L.-T. Chen, T.-J. Chen, S.-H. Li, Y.-H. Wang, H.-C. Tai, S.-C. Yu and H.-Y. Huang, *Clin. Cancer Res.*, 2014, **20**, 6141–6152.
- Z. Jiang, G. R. Fanger, B. A. Woda, B. F. Banner, P. Algate, K. Dresser, J. C. Xu and P. G. G. Chu, *Hum. Pathol.*, 2003, **34**, 792–796.
- C. Kumar-Sinha, R. B. Shah, B. Laxman, S. A. Tomlins, J. Harwood, W. Schmitz, E. Conzelmann, M. G. Sanda, J. T. Wei, M. A. Rubin and A. M. Chinnaiyan, *Am. J. Pathol.*, 2004, **164**, 787–793.
- S. Zha, S. Ferdinandusse, S. Denis, R. J. Wanders, C. M. Ewing, J. Luo, A. M. De Marzo and W. B. Isaacs, *Cancer Res.*, 2003, **63**, 7365–7376.
- B. A. P. Wilson, H. Wang, B. A. Nacev, R. C. Mease, J. O. Liu, M. G. Pomper and W. B. Isaacs, *Mol. Cancer Ther.*, 2011, **10**, 825–838.
- K. Takahara, H. Azuma, T. Sakamoto, S. Kiyama, T. Inamoto, N. Ibuki, T. Nishida, H. Nomi, T. Ubai, N. Segawa and Y. Katsuoka, *Anticancer Res.*, 2009, **29**, 2497–2505.
- A. J. Carnell, I. Hale, S. Denis, R. J. A. Wanders, W. B. Isaacs, B. A. Wilson and S. Ferdinandusse, *J. Med. Chem.*, 2007, **50**, 2700–2707.
- A. J. Carnell, R. Kirk, M. Smith, S. McKenna, L.-Y. Lian and R. Gibson, *ChemMedChem*, 2013, **8**, 1643–1647.
- A. Morgenroth, E. A. Urusova, C. Dinger, E. Al-Momani, T. Kull, B. G. Glattig, H. Frauendorf, O. Jahn, F. M. Mottaghy, S. N. Reske and B. D. Zlatopolskiy, *Chem. – Eur. J.*, 2011, **17**, 10144–10150.
- P.-Y. Lin, K.-L. Cheng, J. D. McGuffin-Cawley, F.-S. Shieu, A. C. Samia, S. Gupta, M. Cooney, C. L. Thompson and C. C. Liu, *Biosensors*, 2012, **2**, 377–387.
- M. Pal, M. Khanal, R. Marko, S. Thirumalairajan and S. L. Bearne, *Chem. Commun.*, 2016, **52**, 2740–2743.
- C. A. Lipinski, F. Lombardo, B. W. Dominy and P. J. Feeney, *Adv. Drug Delivery Rev.*, 2001, **46**, 3–26.
- C. Festuccia, G. L. Gravina, A. Mancini, P. Muzi, E. Di Cesare, R. Kirk, M. Smith, S. Hughes, R. Gibson, L.-Y. Lian, E. Ricevuto and A. J. Carnell, *Anti-Cancer Agents Med. Chem.*, 2014, **14**, 1031–1041.
- W. R. Shieh and C. S. Chen, *J. Biol. Chem.*, 1993, **268**, 3487–3493.
- D. Ouazia and S. L. Bearne, *Anal. Biochem.*, 2010, **398**, 45–51.
- M. Yevglevskis, G. L. Lee, M. D. Threadgill, T. J. Woodman and M. D. Lloyd, *Chem. Commun.*, 2014, **50**, 14164–14166.
- M. Yevglevskis, G. L. Lee, J. Sun, S. Zhou, X. Sun, G. Kociok-Köhn, T. D. James, T. J. Woodman and M. D. Lloyd, *Org. Biomol. Chem.*, 2016, **14**, 612–622.
- D. C. Harris, *Quantitative Chemical Analysis*, W. H. Freeman, New York, 8th edn, 2010.
- M. Y. Moridani, A. Siraki and P. J. O'Brien, *Chem.-Biol. Interact.*, 2003, **145**, 213–223.
- S. Chafaa, J. Meullemeestre, M. J. Schwing, F. Vierling, V. Bohmer and W. Vogt, *Helv. Chim. Acta*, 1993, **76**, 1425–1434.
- D. Y. Yu, F. Huang and H. Xu, *Anal. Methods*, 2012, **4**, 47–49.
- J. H. Zhang, T. D. Y. Chung and K. R. Oldenburg, *J. Biomol. Screening*, 1999, **4**, 67–73.
- W. Schmitz, R. Fingerhut and E. Conzelmann, *Eur. J. Biochem.*, 1994, **222**, 313–323.
- M. E. Tanner, *Acc. Chem. Res.*, 2002, **35**, 237–246.
- A. R. Dewanti and B. Mitra, *Biochemistry*, 2003, **42**, 12893–12901.
- X. J. Liu, G. S. Deng, X. S. Chu, N. Li, L. Wu and D. Li, *Bioorg. Med. Chem. Lett.*, 2007, **17**, 3187–3190.
- R. B. Hamed, E. T. Batchelar, I. J. Clifton and C. J. Schofield, *Cell. Mol. Life Sci.*, 2008, **65**, 2507–2527.
- A. K. Casey, M. A. Hicks, J. L. Johnson, P. C. Babbitt and P. A. Frantom, *Biochemistry*, 2014, **53**, 2915–2925.

