

Cite this: *Chem. Sci.*, 2018, 9, 2552

A chemiluminescent probe for cellular peroxynitrite using a self-immolative oxidative decarbonylation reaction†

Jian Cao,^{ab} Weiwei An,^{ab} Audrey G. Reeves^a and Alexander R. Lippert^{*,abc}

Peroxyntirite (ONOO⁻) is a highly reactive oxygen species which has been recognized as an endogenous mediator of physiological activities like the immune response as well as a damaging agent of oxidative stress under pathological conditions. While its biological importance is becoming clearer, many of the details of its production and mechanism of action remain elusive due to the lack of available selective and sensitive detection methods. Herein, we report the development, characterization, and biological applications of a reaction-based chemiluminescent probe for ONOO⁻ detection, termed as PNCL. PNCL reacts with ONOO⁻ via an isatin moiety through an oxidative decarbonylation reaction to initiate light emission that can be observed instantly with high selectivity against other reactive sulphur, oxygen, and nitrogen species. Detailed studies were performed to study the reaction between isatin and ONOO⁻, which confirm selectivity for ONOO⁻ over NO₂[•]. PNCL has been applied for ONOO⁻ detection in aqueous solution and live cells. Moreover, PNCL can be employed to detect cellular ONOO⁻ generated in macrophages stimulated to mount an immune response with lipopolysaccharide (LPS). The sensitivity granted by chemiluminescent detection together with the specificity of the oxidative decarbonylation reaction provides a useful tool to explore ONOO⁻ chemistry and biology.

Received 28th November 2017

Accepted 31st January 2018

DOI: 10.1039/c7sc05087a

rsc.li/chemical-science

Introduction

Peroxyntirite (ONOO⁻) is a highly reactive oxygen species that can be formed biologically from a diffusion-controlled reaction between superoxide (O₂^{•-}) and nitric oxide (NO[•]).¹ In biological systems, ONOO⁻ has long been known as a deleterious species due to its oxidative damage to lipids, proteins, and nucleic acids.² Abnormal regulation of ONOO⁻ in living systems is associated with diseases such as ischemia-reperfusion,³ diabetes,⁴ cardiac dysfunction,⁵ inflammatory conditions,⁶ auto-immune, and neurodegenerative diseases.^{7,8} However, ONOO⁻ has been proposed to act as a physiological mediator in certain contexts, for example in immune system function⁹ and ischemic preconditioning.¹⁰ Macrophages produce nitric oxide and superoxide to form ONOO⁻ to attack invading pathogens.¹¹ Macrophage cells recognize lipopolysaccharide (LPS), a component of the bacterial cell wall, which induces high expression of inducible nitric oxide synthase (iNOS).¹² The effect of LPS is

amplified by release of IFN-β, providing a mechanism for robust local production of NO[•] and ONOO⁻. Although important biological functions of ONOO⁻ are still emerging, there remains a lack of efficient and selective methods to monitor its production in real time to study its effects in living systems. Therefore, the development of precise approaches for detecting ONOO⁻ is crucial to provide a more detailed understanding of its complex biological effects.

Much of the difficulty of studying the biological roles of ONOO⁻ stems from its complex chemistry. ONOO⁻ coexists with its protonated form ONOOH (pK_a 6.8) under physiological pH, and can directly undergo two-electron reactions with thiols,¹³ CO₂,¹⁴ and carbonyls.¹⁵ The H⁺ and CO₂ adducts, ONOOH and ONOOCO₂⁻, rapidly decompose via homolytic cleavage reactions to form secondary radicals such as hydroxyl radical (HO[•]), carbonate radical (CO₃^{•-}), and nitrogen dioxide (NO₂[•]), which mediate potent one-electron chemistry, including tyrosine nitration.¹⁶ ONOO⁻ can also form from reaction between O₂ and NO[•] produced by photolysis of Angeli's salt under alkaline conditions.¹⁷ Under physiological conditions, Angeli's salt has been widely used as a nitroxyl (HNO) donor for biological studies. Reaction of HNO with O₂ also proceeds at neutral pH to form a potent oxidative species and represents an important decomposition pathway of HNO.^{18,19} While there are similarities and differences between ONOO⁻ and the autooxidation product of HNO formed at neutral pH, the details of this

^aDepartment of Chemistry, Southern Methodist University, Dallas, TX 75275-0314, USA. E-mail: alippert@smu.edu

^bCenter for Drug Discovery, Design, and Delivery (CD4), Southern Methodist University, Dallas, TX 75275-0314, USA

^cCenter for Global Health Impact (CGHI), Southern Methodist University, Dallas, TX 75275-0314, USA

† Electronic supplementary information (ESI) available: Additional experimental details, supplementary figures, and scanned spectra. See DOI: 10.1039/c7sc05087a



chemistry and the exact species eliciting the oxidative effect of Angeli's salt is still under debate.

The classical approach for studying biological ONOO⁻ relies on immunostaining of 3-nitrotyrosine residues,²⁰ which lacks temporal precision and is an indirect method tracking the "footprint" of ONOO⁻. A large number of reaction-based fluorescent probes have been developed in an attempt to solve this problem. These small molecule tools can react with chemical species as they are produced, enabling detection within living intact systems. Several families of reaction-based fluorescent probes for ONOO⁻ detection have been reported and applied in biological studies, such as trifluorocarbonyl-based,²¹ *N*-dearylation-based,²² and boronic ester-based²³ probes. Organoselenium-based,²⁴ and organotellurium-based²⁵ probes are advantageous because the fluorescence signal is reversible, but depend upon reaction with cellular reductants to reverse the signal.²⁶ This limited reversibility, as well as other intrinsic drawbacks of fluorescent detection methods such as photobleaching and autofluorescence highlights the need for new methods of ONOO⁻ detection. Chemiluminescence provides an attractive solution for studying biological molecules and processes due to the low background, high signal-to-noise ratio, and reversible signal of this technique.²⁷ Nanoparticle-based methods have recently been reported for chemiluminescent ONOO⁻ detection.^{28,29} These probes rely on an interesting radical-pair mechanism to generate signal from the ONOO⁻ decomposition products HO[•] and O₂^{•-}. Although these systems do display good sensitivity, complementary approaches that operate *via* direct reaction with ONOO⁻ are desirable. Chemiluminescent probes based on sterically hindered 1,2-dioxetanes offer opportunities to incorporate specific reaction-based triggers and have already been applied for detection of analytes in cellular systems and for live animal imaging.^{30–34} One fruitful design strategy uses Schaap's adamantylidene-dioxetane attached to an analyte-specific reactive handle.³⁵ A striking substituent effect for Schaap's 1,2-dioxetanes has been reported, where incorporation of an electron withdrawing group such as an acrylonitrile at the *ortho* position of the phenol results in a significant increase in the overall chemiluminescent quantum yield of 1,2-dioxetanes.³⁶ As a result, chemiluminescent emission can be observed in aqueous conditions without the addition of any polymeric enhancer solutions. We herein describe the development and application of a selective chemiluminescent probe for ONOO⁻ detection, **PNCL**, with both *in vitro* and live cell applications.

Results and discussion


Design and synthesis of PNCL

As part of a program to develop chemical probes for imaging reactive sulphur,^{29,37} oxygen,^{30,38,39} and nitrogen species, our laboratory has previously reported that isatin reacts with ONOO⁻ *via* an oxidative decarbonylation reaction to generate anthranilic acid.⁴⁰ ¹⁹F NMR probes, 5-fluoroisatin and 6-fluoroisatin, have been developed for selective detection of ONOO⁻ that utilize this newly discovered transformation. In order to expand this reaction-based detection strategy for luminogenic

detection, we adopted a self-immolative 1,6-elimination strategy for chemiluminescent ONOO⁻ detection that relies on tethering an isatin moiety to a 1,2-dioxetane chemiluminescent scaffold *via* an ether linkage (Scheme 1). The isatin serves as the reaction handle, and 1,6-elimination can be triggered after reacting with ONOO⁻ to form the anthranilic acid with the amino group positioned *para* to the benzylic ether. Luminescent emission is then initiated upon the decomposition of the 1,2-dioxetane through a chemically initiated electron exchange luminescence (CIEEL) mechanism. Synthesis of a peroxyinitrite chemiluminescent probe (**PNCL**) proceeded from commercially available 5-iodoisatin **1** by protection of the ketone group as the dimethylacetal **2** (Scheme 2). The protected 5-iodoisatin was subjected to palladium catalyzed reductive formylation with *N*-formyl saccharin as the CO source to form the aldehyde **3**,⁴¹ followed by reduction of the aldehyde with NaBH₄ and deprotection to afford benzyl alcohol **4**. The alcohol **4** was linked to the phenol **5** (ref. 36) through a Mitsunobu reaction to deliver the enol ether precursor **6**. This precursor underwent [2 + 2] cycloaddition with photogenerated singlet oxygen using Rose bengal as a sensitizer to provide the final 1,2-dioxetane **PNCL**.

Spectral response of the probes to ONOO⁻

After obtaining **PNCL**, its photophysical properties and reactivity towards ONOO⁻ were characterized. Luminescence emission spectra were collected using an F-7000 Hitachi spectrophotometer by treating 20 μM **PNCL** with 0–200 μM ONOO⁻ (Fig. 1). An emission peak centered at 525 nm was observed in the presence of ONOO⁻ with ~28-fold increase in peak luminescence intensity compared to a blank control (Fig. 1A). The reaction between **PNCL** and ONOO⁻ was analyzed by GC-MS, confirming the generation of 2-adamantanone and 2-chloro-4-acrylonitrile-3-(methoxycarbonyl)phenol (Fig. S1†). The luminescence emission slowly decayed over a course of 20 min (Fig. 1B), consistent with fast reaction kinetics of ONOO⁻ with the isatin moiety of **PNCL**,⁴⁰ followed by a slower 1,6-elimination. Interestingly, ONOO⁻ can be added to a solution of **PNCL** over several cycles with good recovery of chemiluminescent signal (Fig. S2†). This shows that the reversible chemiluminescent emission has good reproducibility and can be used monitor ONOO⁻ fluxes over time. The dose dependence of the peak chemiluminescent emission at 525 nm measured 10



Scheme 1 Spiroadamantane 1,2-dioxetane-based probe for chemiluminescent ONOO⁻ detection.





Scheme 2 Synthesis of PNCL.



Fig. 1 Response of PNCL to ONOO⁻. (A) Chemiluminescent emission spectra of 20 μM PNCL and 0 (blue trace), 5, 10, 20, 40, 80, 100, 200 μM (red trace) ONOO⁻. (B) Time scans of the chemiluminescent emission at 525 nm of 20 μM PNCL and 0 (blue trace), 5, 10, 20, 40, 80, 100, 200 μM (red trace) ONOO⁻. (C) Peak emission intensity at 525 nm of 20 μM PNCL after adding 0–200 μM ONOO⁻. (D) Time scan of the chemiluminescent emission at 525 nm of 20 μM PNCL and 200 μM SIN-1. All experiments were performed in 20 mM HEPES (pH 7.4), containing <1% DMSO.

seconds after ONOO⁻ addition showed excellent linearity, demonstrating that ONOO⁻ can be quantified using PNCL (Fig. 1C). Calibration of the response of PNCL with lower concentrations of ONOO⁻ revealed a detection limit of 6 nM (3σ) bolus ONOO⁻ (Fig. S3†), comparing favorably to previous fluorescent and chemiluminescent methods (Table S1†). The reactivity of PNCL towards SIN-1 was also evaluated (Fig. 1D), showing that PNCL can monitor the slow release of ONOO⁻ over time, as would be observed in biological systems.

The selectivity of PNCL for ONOO⁻ was tested against a panel of biologically relevant reactive sulphur, oxygen, and nitrogen species and metal cations. The response of PNCL to

200 μM ONOO⁻ was compared against other analytes in HEPES buffer (20 mM, pH 7.4), by adding 5 mM reduced glutathione (GSH), 1 mM L-cysteine, or 200 μM other reactive species. Most of the species tested displayed no significant increase in luminescence intensity over the blank control within 20 min (Fig. 2). Given the destructive reactivity of HO[•], we examined the interference of this species by treating PNCL with ONOO⁻ in the presence of the HO[•]-generating Fenton system (Fig. S4†), where only a minimal reduction in signal was observed. A small luminescence increase was observed when treating PNCL with Angeli's salt, commonly used as a nitroxyl (HNO) donor. This was consistent with previous observations using HPLC to track the reactivity of 5-fluoroisatin with Angeli's salt.³⁷

In light of studies exploring the generation of ONOO⁻ and other nitrogen oxide species from the autooxidation of NO⁻ and HNO,^{17–19} we performed a series of competition experiments with HCO₃⁻ and GSH to better understand the luminescence increase resulting from reaction of PNCL with Angeli's salt. ONOO⁻ reacts rapidly with CO₂, but slowly with GSH. HNO, on the other hand, reacts rapidly with GSH.⁴² We treated PNCL with ONOO⁻ (Fig. 3A and B) or Angeli's salt (Fig. 3C and D) in the presence of various amount of HCO₃⁻ (1 mM, 2 mM, 3 mM, 4 mM, and 5 mM). HCO₃⁻ decreased the chemiluminescence emission intensity generated by both ONOO⁻ and Angeli's salt, providing evidence that Angeli's salt can generate ONOO⁻. A similar experiment was performed by reacting PNCL (20 μM) with ONOO⁻ (Fig. 4A and B) or Angeli's salt (Fig. 4C and D) and various amounts of GSH (50 μM, 100 μM, 200 μM, and 1 mM). However, the GSH only quenched the luminescence intensity from Angeli's salt (Fig. 4B) and no significant luminescence decrease was observed from ONOO⁻ (Fig. 4A). Interestingly, substoichiometric quantities of GSH would not only reduce the total amount of luminescence, but displayed a lag time before producing a chemiluminescent signal (Fig. 4C). Angeli's salt generates HNO with a half-life of 2.3 minutes at 37 °C. The observed lag time of the chemiluminescent response is



Fig. 2 Selectivity of PNCL versus cations and reactive sulphur, oxygen, and nitrogen species. Chemiluminescent emission at 525 nm of 20 μM PNCL and 200 μM biologically relevant analytes in 20 mM HEPES (pH 7.4). Bars represent chemiluminescent emission at 525 nm at 4, 8, 12, 16, 20 min after addition of reactive species. Legend: (1) ONOO⁻, (2) Cys, (3) DEA NONOate, (4) GSH (5 mM), (5) GSNO, (6) H₂O₂, (7) Angeli's salt, (8) KO₂, (9) Na₂S, (10) Na₂S₂O₃, (11) Na₂SO₄, (12) NaNO₂, (13) HO[•], (14) OCl⁻, (15) ^tBuOOH, (16) ^tO₂, (17) Na⁺, (18) Mg²⁺, (19) K⁺, (20) Ca²⁺, (21) Zn²⁺, (22) Fe²⁺, (23) blank. Error bars are ± S.D. All experiments were performed in 20 mM HEPES or 20 mM PBS (pH 7.4), containing <1% DMSO.






Fig. 3 Inhibition of response by NaHCO_3 . (A) Time scans and (B) integrated intensity of the chemiluminescent emission of $20 \mu\text{M}$ PNCL, $200 \mu\text{M}$ ONOO^- and 0 (blue trace), 1 mM, 2 mM, 3 mM, 4 mM, and 5 mM (red trace) NaHCO_3 . (C) Time scans and (D) integrated intensity of the chemiluminescent emission of $20 \mu\text{M}$ PNCL, $200 \mu\text{M}$ Angeli's salt and 0 (blue trace), 1 mM, 2 mM, 3 mM, 4 mM, and 5 mM (red trace) NaHCO_3 . All experiments were performed in 20 mM HEPES (pH 7.4), containing <1% DMSO.

consistent with GSH scavenging the newly formed HNO more rapidly than the autooxidation reaction of HNO. Once the GSH is consumed, HNO can be oxidized to ONOO^- or another nitrogen oxide species capable of two-electron chemistry and reaction with PNCL proceeds to produce a luminescent signal.

In order to confirm this selectivity and that $\text{HCO}_3^-/\text{CO}_2$ does not scavenge HNO, we prepared a fluorescent nitroxyl (HNO) probe, herein called XF1 (Fig. 5A). HNO reacts with the diphenylphosphinobenzoyl group to form an aza-ylide that will nucleophilically attack the ester to trigger the release of the fluorophore.^{43,44} A time-dependent fluorescence turn-on was observed when treating $10 \mu\text{M}$ XF1 with $200 \mu\text{M}$ Angeli's salt (Fig. S5A and B[†]), and selectivity tests demonstrated that XF1 can distinguish HNO from other reactive sulfur, oxygen, and nitrogen species (Fig. S5C[†]). Inhibition experiments were conducted by treating XF1 with various amounts of HCO_3^- or GSH together with Angeli's salt. While no obvious inhibition of the fluorescence response was observed with the addition of HCO_3^- , the fluorescence was completely quenched with $200 \mu\text{M}$ GSH (Fig. 5B and C). These results confirm that GSH scavenges HNO generated from Angeli's salt but $\text{HCO}_3^-/\text{CO}_2$ does not scavenge HNO. Taken together, the results from Fig. 3–5 are consistent with the luminescence increase from Angeli's salt being due to the generation of ONOO^- . We do note that alternative oxidative species formed from HNO have been proposed.¹⁸ Although we cannot completely rule this out, the results here are consistent with the formation of ONOO^- from HNO and support the high specificity of PNCL for detecting




Fig. 4 Inhibition of response by glutathione. (A) Time scans and (B) integrated intensity of the chemiluminescent emission of $20 \mu\text{M}$ PNCL, $200 \mu\text{M}$ ONOO^- and 0 (blue trace), $50 \mu\text{M}$, $100 \mu\text{M}$, $200 \mu\text{M}$, and 1 mM (red trace) glutathione. (C) Time scans and (D) integrated intensity of the chemiluminescent emission of $20 \mu\text{M}$ PNCL, $200 \mu\text{M}$ Angeli's salt and 0 (blue trace), $50 \mu\text{M}$, $100 \mu\text{M}$, $200 \mu\text{M}$, and 1 mM (red trace) glutathione. All experiments were performed in 20 mM HEPES (pH 7.4), containing <1% DMSO.




Fig. 5 HNO scavenging by NaHCO_3 and GSH. (A) Design of XF1, a fluorescent probe for HNO. (B) Peak emission intensity of $10 \mu\text{M}$ XF1, $200 \mu\text{M}$ Angeli's salt and 0–5 mM NaHCO_3 . (C) Peak emission intensity of $10 \mu\text{M}$ XF1, $200 \mu\text{M}$ Angeli's salt and 0–1 mM glutathione. All experiments were performed in 20 mM HEPES (pH 7.4), containing <1% DMSO with $\lambda_{\text{ex}} = 488 \text{ nm}$.

ONOO^- . As a final mechanistic investigation, we examined the selectivity of ONOO^- versus NO_2^+ , HO^+ , and $\text{CO}_3^{\cdot-}$ (the decomposition products of ONOO^-) by treating isatin with ONOO^- in the presence of $\text{HCO}_3^-/\text{CO}_2$ or TEMPOL, a free radical scavenger (Fig. S6[†]). While $\text{HCO}_3^-/\text{CO}_2$ inhibits the reaction, TEMPOL has no effect. This suggests that isatin reacts directly with ONOO^-



and not with free radical species like NO_2^\cdot , HO^\cdot , or $\text{CO}_3^{\cdot-}$ that are formed *via* spontaneous ONOO^- decomposition.

Evaluation of PNCL for ONOO^- detection in living cells

After demonstrating that **PNCL** reacts with ONOO^- with high sensitivity and selectivity, we then proceeded to investigate the ability of **PNCL** for ONOO^- detection in living cells using a BioTek Cytation 5 plate reader. Toxicity was determined by the MTT assay, which indicated low toxicity after 24 hour treatment with **PNCL** at concentrations less than 100 μM and more than 50% viability after treating with high concentrations of 1 mM **PNCL** for 24 hours (Fig. S7†). Cellular internalization was verified by imaging the intrinsic fluorescence emission of **PNCL** (Fig. S8†). In order to evaluate the ability of **PNCL** to detect cellular delivery of ONOO^- from donor compounds, RAW 264.7 macrophages were incubated with **PNCL** for 30 minutes, washed, and then treated with 0, 200 μM , 400 μM , 800 μM , 1 mM, or 2 mM SIN-1, a commonly used ONOO^- donor that operates by simultaneously generating NO^\cdot and $\text{O}_2^{\cdot-}$. A time course of the chemiluminescent emission intensity increased to reach a maximum at 40 minutes (Fig. 6A). Both the peak and integrated luminescence intensities increased with increasing concentrations of SIN-1 (Fig. 6A and B), and a 5-fold luminescence increase was obtained with 2 mM SIN-1 compared with

a blank control. We also performed the same experiment in A549 cells (Fig. S9†). Similar results were observed and excellent linearity was achieved in living cells between 0 and 400 μM SIN-1 (Fig. S9C†). This increase in chemiluminescent emission from SIN-1 and **PNCL** was attenuated by the ONOO^- scavenger Mn(III)TMPyP (Fig. S10†), confirming that the response is due to ONOO^- .

Finally, we investigated the ability of **PNCL** to detect ONOO^- generated by macrophages stimulated to mount an immune response. Lipopolysaccharide (LPS) binds to the toll-like receptors of macrophages, activating the transcription factor $\text{NF-}\kappa\text{B}$ ultimately leading to the expression of iNOS and generation of reactive oxygen species that can produce ONOO^- .¹² Macrophages stimulated with LPS produced increased chemiluminescent signal from **PNCL** (Fig. 6C), showing a 36% increase in integrated luminescence intensity with 1000 ng mL^{-1} LPS. Treatment with the ONOO^- scavenger Mn(III)TMPyP reduces this signal back to control levels and co-incubation with the selective iNOS inhibitor 1400 W also reduces the signal from LPS and **PNCL** (Fig. S11†). These control experiments validate the ability of **PNCL** to detect cellular production of ONOO^- . Bright field images of the cells after LPS stimulation show the formation of phagosomes (Fig. 6D), a phenotypic response to LPS exposure.

Conclusions

We have designed and synthesized a reaction-based chemiluminescent probe for ONOO^- , **PNCL**. **PNCL** reacts with ONOO^- through an oxidative decarbonylation reaction to initiate light emission that can be observed instantly with high sensitivity and selectivity against other reactive sulphur, oxygen, and nitrogen species. The observed reactivity towards Angeli's salt is attributed to the generation of ONOO^- , which is supported by inhibitor experiments with HCO_3^- and GSH, providing experimental evidence of an autooxidative product of HNO. Several studies have purported that the autooxidation of HNO produces a product unique from ONOO^- , as evidenced by data showing no effect of HCO_3^- on the reaction between Angeli's salt and reactive oxygen species probe dihydrorhodamine 123.¹⁸ In our studies, however, we observe a drastic HCO_3^- dependent reduction in chemiluminescent signal from **PNCL** and Angeli's salt, consistent with ONOO^- formation from HNO under physiological conditions.¹⁹ **PNCL** was used to detect ONOO^- generated from donor compounds in multiple cell lines, as well as cellular ONOO^- generated by macrophages stimulated with LPS. Confirmation of the chemical specificity towards ONOO^- over downstream free radical products, coupled with operational compatibility in living cells establishes **PNCL** a useful and easy-to-use tool for studying ONOO^- in chemical and biological systems.

Conflicts of interest

A. R. L. discloses a financial stake in BioLum Science, LLC, a company developing asthma monitoring technology.




Fig. 6 ONOO^- detection in RAW 264.7 macrophages. (A) Time scans and (B) integrated intensity of chemiluminescence emission of RAW 264.7 macrophages incubated with 20 μM **PNCL** for 30 minutes, washed, and then treated with 0 (blue trace), 200 μM , 400 μM , 800 μM , 1 mM, and 2 mM (red trace) SIN-1. (C) Integrated chemiluminescent emission intensity of RAW 264.7 macrophages stimulated with (Cont) Vehicle control, (LPS) 1000 ng mL^{-1} LPS, or (LPS, MnTMPyP) 1000 ng mL^{-1} LPS and 25 μM Mn(III)TMPyP and then treated with 20 μM **PNCL**. (D) Brightfield images of stimulated RAW 264.7 macrophages. Error bars are S.D. from $n = 3$ –7 wells and 3 biological replicates ($n = 10$ wells from 6 biological replicates for the control experiment). Statistical significance was assessed using a two-tailed student's t -test. *** $p < 0.001$, ** $p < 0.01$.



Acknowledgements

Research reported in this publication was supported by the National Institute Of General Medical Sciences of the National Institutes of Health under award number R15GM114792. The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health. Additional support was provided by Southern Methodist University (start-up funds to A. R. L and Engaged Learning funds to A. G. R.). We acknowledge Manasa Subbarao for assistance with cell culture and Sara Lange for assistance with preliminary experiments. We thank Maciej Kukula (UT Arlington), Hamilton Lee (UT Dallas), and Prof. Jeremiah Gassen-smith (UT Dallas) for assistance with mass spectrometry. Human lung adenocarcinoma epithelial cell (A549) were purchased from ATCC and cultured in F-12K media supplemented with 10% Fetal Bovine Serum (FBS) and 1% antibiotics (penicillin/streptomycin, 100 U mL⁻¹).

Notes and references

- 1 J. S. Beckman, T. W. Beckman, J. Chen, P. A. Marshall and B. A. Freeman, *Proc. Natl. Acad. Sci. U. S. A.*, 1990, **87**, 1620.
- 2 (a) A. J. Gow, D. Duran, S. Malcolm and H. Ischiropoulos, *FEBS Lett.*, 1996, **385**, 63; (b) G. L. Squadrito and W. A. Pryor, *Free Radical Biol. Med.*, 1998, **25**, 392; (c) C. Szabó, H. Ischiropoulos and R. Radi, *Nat. Rev. Drug Discovery*, 2007, **6**, 662; (d) P. Pacher, J. S. Beckman and L. Liaudet, *Physiol. Rev.*, 2007, **87**, 315.
- 3 P. Liu, B. H. Xu, J. Quilley and P. Y. K. Wong, *Am. J. Physiol.: Cell Physiol.*, 2000, **279**, 1970.
- 4 W. Kosseljans, A. Eis, R. Sahay, D. Brockman and L. Myatt, *Am. J. Physiol.: Heart Circ. Physiol.*, 2000, **278**, 1311.
- 5 P. B. Massion, O. Feron, C. Dessy and J. L. Balligand, *Circ. Res.*, 2003, **93**, 388.
- 6 H. Wiseman and B. Halliwell, *Biochem. J.*, 1996, **313**, 17.
- 7 R. Ahmad, Z. Rasheed and H. Ahsan, *Immunopharmacol. Immunotoxicol.*, 2009, **31**, 388.
- 8 (a) J. S. Beckman, M. Carson, C. D. Smith and W. H. Koppenol, *Nature*, 1993, **364**, 584; (b) M. A. Smith, P. L. R. Harris, L. M. Sayre, J. S. Beckman and G. Perry, *J. Neurosci.*, 1997, **17**, 2653; (c) N. A. Simonian and J. T. Coyle, *Annu. Rev. Pharmacol. Toxicol.*, 1996, **36**, 83.
- 9 C. Brito, M. Naviliat, A. C. Tiscornia, F. Vuillier, G. Gualco, G. Dighiero, R. Radi and A. M. Cayota, *J. Immunol.*, 1999, **162**, 3356–3366.
- 10 S. Altug, A. T. Demiryürek, K. A. Kane and I. Kanzik, *Br. J. Pharmacol.*, 2000, **130**, 125.
- 11 M. N. Alvarez, G. Peluffo, L. Piacenza and R. Radi, *J. Biol. Chem.*, 2011, **286**, 6627.
- 12 A. T. Jacobs and L. J. Ignarro, *J. Biol. Chem.*, 2001, **276**, 47950.
- 13 W. H. Koppenol, J. J. Moreno, W. A. Pryor, H. Ischiropoulos and J. S. Beckman, *Chem. Res. Toxicol.*, 1992, **5**, 834.
- 14 S. V. Lymar and J. K. Hurst, *J. Am. Chem. Soc.*, 1995, **117**, 8867.
- 15 G. Merényi, J. Lind and S. Goldstein, *J. Am. Chem. Soc.*, 2002, **124**, 40.
- 16 G. Ferrer-Sueta and R. Radi, *ACS Chem. Biol.*, 2009, **4**, 161.
- 17 C. E. Donald, M. N. Hughes, J. M. Thompson and F. T. Bonner, *Inorg. Chem.*, 1986, **25**, 2676.
- 18 (a) K. M. Miranda, A. S. Dutton, L. A. Ridnour, C. A. Foreman, E. Ford, N. Paolucci, T. Katori, C. G. Tocchetti, D. Mancardi, D. D. Thomas, M. G. Espey, K. N. Houk, J. M. Fukuto and D. A. Wink, *J. Am. Chem. Soc.*, 2005, **127**, 722; (b) J. H. Jorolan, L. A. Buttitta, C. Cheah and K. M. Miranda, *Nitric Oxide*, 2015, **44**, 39.
- 19 R. Smulik, D. Debski, J. Zielonka, B. Michalowski, J. Adamus, A. Marcinek, B. Kalyanaraman and A. Sikora, *J. Biol. Chem.*, 2014, **289**, 35570–35581.
- 20 (a) H. Ischiropoulos, *Arch. Biochem. Biophys.*, 1998, **356**, 1; (b) R. Radi, *Proc. Natl. Acad. Sci. U. S. A.*, 2004, **101**, 4003.
- 21 (a) D. Yang, H. Wang, Z. Sun, N. Chung and J. Shen, *J. Am. Chem. Soc.*, 2006, **128**, 6004; (b) Z. Sun, H. Wang, F. Liu, Y. Chen, P. K. H. Tam and D. Yang, *Org. Lett.*, 2009, **11**, 1887.
- 22 (a) T. Peng, X. Chen, L. Gao, T. Zhang, W. Wang, J. Shen and D. Yang, *Chem. Sci.*, 2016, **7**, 5407; (b) J. Li, C. S. Lim, G. Kim, H. M. Kim and J. Yoon, *Anal. Chem.*, 2017, **89**, 8496.
- 23 X. Sun, Q. Xu, G. Kim, S. E. Flower, J. P. Lowe, J. Yoon, J. S. Fossey, X. Qian, S. D. Bull and T. D. James, *Chem. Sci.*, 2014, **5**, 3368.
- 24 F. Yu, P. Li, G. Li, G. Zhao, T. Chu and K. Han, *J. Am. Chem. Soc.*, 2011, **133**, 11030.
- 25 F. Yu, P. Li, B. Wang and K. Han, *J. Am. Chem. Soc.*, 2013, **135**, 7674.
- 26 Z. Lou, P. Li and K. Han, *Acc. Chem. Res.*, 2015, **48**, 1358–1368.
- 27 A. R. Lippert, *ACS Cent. Sci.*, 2017, **3**, 269.
- 28 W. Zhou, Y. Cao, D. Sui and C. Lu, *Anal. Chem.*, 2016, **88**, 2659–2665.
- 29 W. Zhou, S. Dong, Y. Lin and C. Lu, *Chem. Commun.*, 2017, **53**, 2122–2125.
- 30 T. S. Bailey and M. D. Pluth, *Methods Enzymol.*, 2015, **554**, 81.
- 31 L. Liu and R. P. Mason, *PLoS One*, 2010, **5**, e12024.
- 32 J. Cao, R. Lopez, J. M. Thacker, J. Y. Moon, C. Jiang, S. N. S. Morris, J. H. Bauer, P. Tao, R. P. Mason and A. R. Lippert, *Chem. Sci.*, 2015, **6**, 1979.
- 33 J. Cao, J. Campbell, L. Liu, R. P. Mason and A. R. Lippert, *Anal. Chem.*, 2016, **88**, 4995.
- 34 (a) O. Green, S. Gnaim, R. Blau, A. Eldar-Boock, R. Satchi-Fainaro and D. Shabat, *J. Am. Chem. Soc.*, 2017, **139**, 13243; (b) N. Hananya, A. E. Boock, C. R. Bauer, R. Satchi-Fainaro and D. Shabat, *J. Am. Chem. Soc.*, 2016, **138**, 13438.
- 35 A. P. Schaap, T. S. Chen, R. S. Handley, R. DeSilva and B. P. Giri, *Tetrahedron Lett.*, 1987, **28**, 1155.
- 36 O. Green, T. Eilon, N. Hananya, S. Gutkin, C. R. Bauer and D. Shabat, *ACS Cent. Sci.*, 2017, **3**, 349.
- 37 J. L. Kroll, C. A. Werchan, A. G. Reeves, K. J. Bruemmer, A. R. Lippert and T. Ritz, *Physiol. Behav.*, 2017, **179**, 99.
- 38 A. G. Reeves, M. Subbarao and A. R. Lippert, *Anal. Methods*, 2017, **9**, 3418.
- 39 M. E. Quimbar, K. M. Krenek and A. R. Lippert, *Methods*, 2016, **109**, 123.
- 40 K. J. Bruemmer, S. Merrikhihaghi, C. T. Lollar, S. N. S. Morris, J. H. Bauer and A. R. Lippert, *Chem. Commun.*, 2014, **50**, 12311.



- 41 T. Ueda, H. Konishi and K. Manabe, *Angew. Chem., Int. Ed.*, 2013, **52**, 8611.
- 42 J. M. Fukuto, M. D. Bartberger, A. S. Dutton, N. Paolucci, D. A. Wink and K. N. Houk, *Chem. Res. Toxicol.*, 2005, **18**, 790.
- 43 (a) K. Kawai, N. Ieda, K. Aizawa, T. Suzuki, N. Miyata and H. Nakagawa, *J. Am. Chem. Soc.*, 2013, **135**, 12690; (b) X. Jing, F. Yu and L. Chen, *Chem. Commun.*, 2014, **50**, 14253; (c) C. Liu, H. Wu, Z. Wang, C. Shao, B. Zhu and X. Zhang, *Chem. Commun.*, 2014, **50**, 6013; (d) K. Zheng, W. Lin, D. Cheng, H. Chen, Y. Liu and K. Liu, *Chem. Commun.*, 2015, **51**, 5754; (e) M. Ren, B. Deng, K. Zhou, J. Wang, X. Kong and W. Lin, *J. Mater. Chem. B*, 2017, **5**, 1954.
- 44 J. A. Reisz, C. N. Zink and S. B. King, *J. Am. Chem. Soc.*, 2011, **133**, 11675.

