



Cite this: *Environ. Sci.: Processes Impacts*, 2022, 24, 2032

Received 20th April 2022
Accepted 17th August 2022

DOI: 10.1039/d2em00166g

rsc.li/espi

Use and release of per- and polyfluoroalkyl substances (PFASs) in consumer food packaging in U.S. and Canada†

Laura Minet,[‡] Zhanyun Wang,^{cd} Anna Shalin,^b Thomas A. Bruton,^{§e} Arlene Blum,^{ef} Graham F. Peaslee,^g Heather Schwartz-Narbonne,^b Marta Venier,^h Heather Whitehead,^g Yan Wu^{¶h} and Miriam L. Diamond^{‡*bi}

Numerous per- and polyfluoroalkyl substances (PFASs) occur in consumer food packaging due to intentional and unintentional addition, despite increasing concern about their health and environmental hazards. We present a substance flow analysis framework to assess the flows of PFASs contained in plant fiber-based and plastic food packaging to the waste stream and environment. Each year between 2018 and 2020, an estimated 9000 (range 1100–25 000) and 940 (range 120–2600) tonnes per year of polymeric PFASs were used in 2% of food packaging in the U.S. and Canada, respectively. At least 11 tonnes per year of non-polymeric PFASs also moved through the food packaging life cycle. Approximately 6100 (range 690–13 000) and 700 (range 70–1600) tonnes per year of these PFASs were landfilled or entered composting facilities in the U.S. and Canada, respectively, with the potential to contaminate the environment. The results suggest that minimal food packaging contains intentionally added PFASs which, nonetheless, has the potential to contaminate the entire waste stream. Further, this indicates that PFASs are not needed for most food packaging. These results serve as a benchmark to judge the effectiveness of future industry and government initiatives to limit PFAS use in food packaging.

Environmental significance

Although we estimate that only ~2% of U.S. and Canadian food packaging contains intentionally added PFASs, this level of addition results in PFASs flows of 9000 (range 1100–25 000) and 940 (range 120–2600) tonnes per year. Most of the PFASs are polymeric, but we estimate a minimum of 11 tonnes per year of nonpolymeric and hence mobile and potentially bio-accumulative PFASs are contained in U.S. and Canadian food packaging. PFASs in food packaging have the potential to contaminate the waste stream (*e.g.*, recycled materials, compost, landfill leachate), and the environment from releases to air, water and soils. The estimated low usage of intentionally added PFASs suggests that alternatives are available and hence PFASs can be eliminated from this non-essential use. The prevalence of non-intentionally occurring PFASs in food packaging is more problematic. Our estimates are a benchmark for comparison, as jurisdictions deliberate PFASs reduction strategies, *e.g.*, U.S. EPA PFAS Strategic Roadmap of 2021, and as other jurisdictions may follow the 7 U.S. states that have adopted legislative bans on PFAS use in food contact materials.

1 Introduction

Per- and polyfluoroalkyl substances (PFASs) are a class of more than 4700 non-polymeric (*e.g.*, perfluoroalkyl acids) and polymeric (*e.g.*, fluoropolymers, perfluoropolyethers, and side-chain fluorinated polymers) substances.¹ PFASs have been detected in paper- and plastic-based consumer food packaging, *e.g.*, popcorn microwave bags,² dessert/bread wrappers,³ Tex Mex food packaging³ and “eco-friendly” plant fiber-based food packaging.⁴ Previous studies have revealed the presence of perfluoroalkylcarboxylic acids (PFCAs), perfluoroalkanesulfonic acids (PFASAs), fluorotelomer alcohols (FTOHs) and phosphate esters (PAPs and diPAPs) in food packaging,^{3,5–10} with recent evidence of the use of polymers such as side-chain fluorinated polymers.^{11,12} A recent European-wide survey has confirmed the widespread use of PFASs in disposable food packaging and tableware.¹³

The strong chemical and biological stability of the per-fluorinated carbon moieties^{1,14} leads to the high persistence and accumulation of many PFASs in the environment,^{15,16} biota^{17,18}

^aDepartment of Civil Engineering, University of Victoria, Canada

^bDepartment of Earth Sciences, University of Toronto, Canada. E-mail: miriam.diamond@utoronto.ca

^cEmpa – Swiss Federal Laboratories for Materials Science and Technology, Technology and Society Laboratory, Switzerland

^dInstitute of Environmental Engineering, ETH-Zürich, Switzerland

^eGreen Science Policy Institute, USA

^fDepartment of Chemistry, University of California, Berkeley, USA

^gDepartment of Physics, University of Notre Dame, USA

^hO'Neill School of Public and Environmental Affairs, Indiana University, USA

ⁱSchool of the Environment, University of Toronto, Canada

† Electronic supplementary information (ESI) available. See <https://doi.org/10.1039/d2em00166g>

‡ This research was conducted while the first author was at the University of Toronto, Canada.

§ Current address: California Department of Toxic Substances Control 1001 I St., Sacramento, CA, 95814, USA

¶ Key Laboratory of Geographic Information Science (Ministry of Education), School of Geographic Sciences, East China Normal University, Shanghai 200241, China



and humans,^{19,20} with the potential for serious environmental and human health harm.^{21,22} Some PFASs present in food packaging can be transferred to food or air through food contact and heating processes, leading to human exposure through ingestion and inhalation.^{23,24} Disposal of packaging waste can lead to the environmental release of PFASs *via* landfill leachate,^{25,26} compost application²⁷ and biosolids, and effluent discharges from wastewater treatment plants (resulting from inputs due to human exposure).^{28–30}

The United States (U.S.), through the Food and Drug Administration (FDA), has a list of PFASs authorized for use in food packaging;³¹ however, information is lacking on the actual usage of these PFASs in food packaging production, or whether manufacturers are using non-authorized PFASs. Canadian regulatory oversight of PFASs use in food packaging is more complicated due to the lack of a public list of authorized PFASs (ESI, Section S1 and Table S1†).³² However, as a signatory to the Stockholm Convention, Canada restricted the use, sale and importation of PFOS and PFOS-containing products in 2008 with exemptions, and in 2012 restrictions were applied to PFOA, its salts and precursors, namely long-chain or LC-PFCAs, again with specific exemptions.^{33–35} A further tightening of these regulations with reduced exemptions was recently introduced to the Canadian Parliament.³⁶

The U.S. Environmental Protection Agency (EPA) initiated the PFOA (perfluorooctanoic acid) 2010/2015 Stewardship program in 2006, including PFOA, its long-chain homologues and their precursors. In 2020, a voluntary agreement was struck between the U.S. FDA and manufacturers of grease-proofing agents used in paper and paperboard food packaging, to phase out the use of PFASs derived from 6 : 2 FTOH by 2024.³⁷ This agreement was prompted by new evidence suggesting significant potential human health risks arising from chronic dietary exposure.³⁸ Further, a recent report from the Organisation for Economic Co-operation and Development (OECD) has argued for more regulation of these compounds in food packaging.³⁹

Since the 2006 agreement, PFASs manufacturers in North America and Europe largely replaced long-chain PFASs with, for example, shorter-chain homologues and other PFASs such as perfluoropolyethers. Short-chain replacements are as persistent as long-chain PFASs, and even more difficult to remove from water treatment systems.⁴⁰ The other major replacement, polymeric PFASs, notably side-chain fluorinated polymers, also present environmental and human health concerns because they contain non-polymeric PFASs impurities and can degrade into non-polymeric, persistent and mobile PFASs.^{11,14,28,41,42}

Little is known about the fate of PFASs used in food packaging, such as how much is released into the environment and from where these releases occur. Substance flow analysis (SFA) is a useful tool for understanding the major inputs and outputs of a system in order to prioritise action,⁴³ including, for example, PFASs in carpets.⁴⁴ To date, the stock and flows of PFASs that transfer PFASs used in food packaging to various waste streams have not been estimated. Our goals were to develop a general SFA framework for PFASs in food packaging and, as a case study, apply it to the U.S. and Canada for a year

representative of 2018 to 2020. This case study illustrates the potential of PFASs used in food packaging to contaminate the waste stream and environment, and it provides a benchmark against which to judge the effectiveness of jurisdictional initiatives to limit the use of PFASs in food packaging.

2 Methods

Substance flow analysis framework

The SFA framework includes the three life stages of food packaging: production, use and end-of-life (Fig. 1 and ESI Section S2†). Five transient and permanent sinks of PFASs were identified: food packaging storage, human storage, landfill storage, destruction by incineration, and environmental storage. Eqn (1) describes the variation of PFAS mass (M_{PFAS}) in food packaging over time.

$$\begin{aligned} \frac{dM_{\text{PFAS}}}{dt} &= \text{inflow}_{\text{PFAS}} - \text{outflow}_{\text{PFAS}} \\ &= F_{\text{FP}} + F_{4,1} - (F_1 + F_{\text{production}} + F_{\text{storage}}) \end{aligned} \quad (1)$$

where F_{FP} is the flow of PFASs produced for food packaging, $F_{4,1}$ is the flow from discarded food packaging that is recycled back to food packaging, F_1 is the flow from the sales of food packaging, $F_{\text{production}}$ is the flow released to the environment (*i.e.*, air and surface water) during the production of food packaging, and F_{storage} is the outflow in stored food packaging.

Fig. 1 describes the flows of PFASs present in consumer food packaging. The framework includes releases during PFASs production as a potential source to the environment,¹ as identified by $F_{\text{PFAS prod}}$ in Fig. 1. We also note that municipal compost originates from post-consumer use of food packaging that is sent to municipally run composting facilities and biosolids originate from wastewater treatment plants. Additionally, we have included the land application of biosolids for agricultural purposes through the flow as $F_{\text{bio. to env}}$.

Based on their PFAS content and origin (described in ESI Section S1†), food packaging can be divided into three categories. Category 1 (“functional” category) corresponds to food packaging with PFASs intentionally added, including plant fiber-based packaging with sufficient PFAS(s) to confer water- and grease-repellency, such as popcorn bags⁴⁵ and plant fiber-based (marketed as “compostable”) bowls,³ as well as plastic packaging with fluoropolymers as extrusion aids.^{46,47} Category 2 (“non-functional” category) includes food packaging with the unintentional addition of PFASs during manufacturing (*e.g.*, from recycled materials, cross contamination, or lubricants in machinery used to produce packaging) in a quantity insufficient to provide functionality such as repellency properties.⁴⁸ Category 3 (“no PFASs” category) corresponds to food packaging free of PFASs.

SFAs were parameterised for the three categories of food packaging (see above) and two groups of PFASs: first, polymeric ones (including side-chain fluorinated polymers and perfluoropolyethers that are usually contained in paper-based food packaging,⁴⁹ and fluoropolymers that have been found in plastic-based food packaging⁵⁰), and, second, non-polymeric compounds, *e.g.*, fluorotelomer (FT) alcohols (n:2 FTOHs), FT





Fig. 1 A schematic overview of the general substance flow analysis (SFA) framework for PFASs in consumer food packaging (FP). The numbered flows are quantified in the case study. "Environment" is used to designate air and surface water. Values indicated correspond to the estimated flows of total PFASs in the U.S. (in tonnes per year).

sulfonates (n:2 FTSS), FT acrylates (n:2 FTAc), FT methacrylates (n:2 FTMAc).

Case study

We constructed two SFAs for food packaging (e.g., paperboard, plastic pouches, fast food wrapping paper, plant-fiber bowls): one for the sum of polymeric PFASs and another for the sum of non-polymeric PFASs. Many of these non-polymeric compounds can be production residuals, impurities, by-products and degradation products of polymeric PFASs, or unintentional contaminants.^{11,48}

We considered a reference year between 2018 and 2020. Only a limited number of flows could be estimated due to a lack of information. To address some of these data gaps and to recognize the uncertainty in our estimates, we provided a range of estimates approximating low, middle and high scenarios, presented as middle (low–high), for parameter values.

For the two categories of food packaging containing PFASs, *i.e.*, with functional (e.g. PFASs added to confer grease- and water-repellency) and non-functional (e.g. inadvertent PFASs that do not confer grease- and water-repellency) concentrations, we estimated the polymeric and non-polymeric PFASs content

as a middle value bounded by low and high estimates, by combining information provided in the U.S. FDA Inventory of Effective Food Contact Substance Notifications⁵¹ (ESI Table S1†) with data from the literature.

Using information provided in patents,⁵¹ we assumed the maximum concentration of polymeric PFASs in food packaging with functional PFASs addition ($C_{\text{functional}}$) to be 1.2% by weight of finished packaging (high scenario), and the middle scenario to be 0.5% (*i.e.*, approximately half the maximum value recommended for use) (Table 1 and ESI Section S3†). The minimum concentration to confer functionality such as water- and oil-repellency is 0.1% by weight of finished paper.⁵¹ For food packaging with non-functional addition of PFASs, we took 0.01% by weight of finished paper as the maximum concentration of polymeric PFASs, which is the 10% of the minimum concentration assumed for functional addition of PFASs, and 0 and 0.005% for the low and middle scenarios, respectively.

For non-polymeric PFASs in the "functional" category, we used 370, 2820 and 25 200 ng g⁻¹ for the lower, middle and high values, respectively, estimated from the list of PFASs authorized for use by the U.S. Food and Drug Administration and from Liu *et al.* (see ESI Table S1†).^{31,52} For food packaging with non-



Table 1 Input data for the substance flow analysis case study (FP = food packaging)^a

Parameter/type of FP		Low	Middle	High	Source
PFAS concentrations in FP					
Non-polymeric PFASs	$C_{\text{functional}}$	$3.7 \times 10^{-5}\%$	$2.8 \times 10^{-4}\%$	$2.5 \times 10^{-3}\%$	Literature (ESI Section S3.1)
	$C_{\text{non-functional}}$	0	$2.0 \times 10^{-5}\%$	$3.7 \times 10^{-5}\%$	Literature (ESI Section S3.1)
	$C_{\text{no PFAS}}$	0	0	0	—
Polymeric PFASs	$C_{\text{functional}}$	0.1%	0.5%	1.2%	Patent ⁵¹
	$C_{\text{non-functional}}$	0	0.005%	0.01%	Assumed
	$C_{\text{no PFAS}}$	0	0	0	—
Proportion of FP types					
$\%_{\text{functional}}$	Category “functional”	2%	2%	2%	Back calculation and assumed
$\%_{\text{non-functional}}$	Category “non-functional”	40%	50%	60%	Back calculation and assumed
$\%_{\text{no PFAS}}$	Category “no PFAS”	58%	48%	38%	Back calculation and assumed
Mass of FP in municipal solid waste					
Canada	M_{FP} (megatonnes)	5.9	7.6	8.8	Government of Canada, ⁶¹ Statistics Canada ⁶²
U.S.		57	72	83	U.S. EPA, ^{63,64} Government of Canada ⁶¹
Proportion of FP littered					
$\%_{\text{litter}}$		5.0%	15%	30%	Assumed
Proportion of FP recycled					
$\%_{\text{FP recycled}}$		10%	20%	30%	Government of Canada, ^{61,65} U.S. EPA ⁶⁴
Proportion of FP recycled to FP					
$\%_{\text{FP recycled to FP}}$		5.0%	18%	33%	U.S. EPA ⁵³
Proportion of FP intended for recycling sent to landfills or exported					
$\%_{\text{recycling to landfill}}$		5.0%	15%	30%	NY Times, ⁵⁴ CTV news, ⁵⁵ Globe and Mail ⁵⁶
Proportion of FP incinerated					
Canada	$\%_{\text{incineration}}$	3.0%	5.0%	8.0%	Statistics Canada, ⁵⁷ Solid Waste & Recycling ⁵⁸
U.S.		4.0%	11%	17%	U.S. EPA ⁵⁹
Proportion of Canadian FP waste sent to Michigan					
$\%_{\text{Can. Waste to Mich}}$		5.0%	11%	20%	Michigan Department of Environment ⁶⁰

^a Functional refers to FP with PFASs added to confer grease- and water-repellency and non-functional refers to FP with inadvertent PFASs addition that does not confer grease- and water-repellency.

functional addition of non-polymeric PFASs, we used 370 ng g^{-1} as our upper boundary, and 0 ng g^{-1} as the minimum, with 200 ng g^{-1} as the middle value. A complete listing of data sources and explanations for these choices are provided in ESI Section S3.1.†

We note that the SFA for polymeric PFASs provides inputs to the non-polymeric PFASs SFA through the release of side chains and other processes of polymeric breakdown. However, we had insufficient data to quantify this transfer and therefore did not quantify releases from polymeric breakdown as an input to the non-polymeric PFASs SFA. All values used in the SFA are listed in Table 1 with full details in ESI Section S2.†

F_1 : flow in the sales of food packaging. The flow in the sales of food packaging (F_1) was calculated as:

$$F_1 = M_{\text{FP}} \times (C_{\text{functional}} \times \%_{\text{functional}} + C_{\text{non-functional}} \times \%_{\text{non-functional}} + C_{\text{noPFAS}} \times \%_{\text{noPFAS}}) \quad (2)$$

where M_{FP} is the mass of food packaging generated in a given year (ESI Section S2.3†); $C_{\text{functional}}$, $C_{\text{non-functional}}$, $C_{\text{no PFAS}}$ are the

concentrations in each of the three categories of food packaging, which are present in the $\%_{\text{functional}}$, $\%_{\text{non-functional}}$, and $\%_{\text{no PFAS}}$ percentages, respectively. Due to a lack of data, the percentages of food packaging types were assumed such that the estimated flows accorded with 10 000 (1080–21 600) tonnes of PFASs used in food packaging annually in the U.S., which we estimated based on data available in the literature. We used the same percentages for Canada (ESI Section S2.3 and S3.2†) (Table 1).

F_2 : flow in discarded food packaging. We assumed a short lifetime of food packaging, *i.e.*, all food packaging from food bought in a year is discarded the same year since food is typically consumed the same year it was purchased. Further, we assumed that significant loss during the lifetime of food packaging is minimal, although some losses occur.^{23,24} The flow in discarded food packaging, F_2 , is

$$F_2 = F_1 - F_{\text{intake}} \cong F_1 \quad (3)$$



Table 2 Main calculated flows of PFASs from consumer food packaging in the U.S. and Canada (in tonnes per year), along with the percentage of the total inflow (*i.e.*, F_i) they represent

		U.S.			Canada		
		Low	Middle	High	Low	Middle	High
Polymeric PFASs							
F_1	Flow in the sales of FP	1.13×10^3	9.03×10^3	2.49×10^4	1.18×10^2	9.43×10^2	2.64×10^3
F_2	Flow in discarded FP	1.13×10^3	9.03×10^3	2.49×10^4	1.18×10^2	9.43×10^2	2.64×10^3
		0%	0%	0%	0%	0%	0%
F_3	Flow in littered FP	3.39×10^2	4.51×10^2	3.74×10^3	3.55×10^1	4.71×10^1	3.96×10^2
		30%	5%	15%	30%	5%	15%
F_4	Flow in recycled FP	7.92×10^1	1.71×10^3	6.36×10^3	8.28	1.79×10^2	6.73×10^2
		7%	19%	26%	7%	19%	26%
$F_{4,1}$	Flow in recycled FP that is recycled back to FP	3.96	3.09×10^2	2.10×10^3	4.14×10^{-1}	3.22×10^1	2.22×10^2
		0%	3%	8%	0%	3%	8%
$F_{4,2}$	Flow in recycled FP that is recycled into other materials	7.13×10^1	1.15×10^3	2.35×10^3	7.45	1.20×10^2	2.49×10^2
		6%	13%	9%	6%	13%	9%
$F_{4,3}$	Flow in FP initially intended for recycling but that is ultimately landfilled or exported	3.96	2.57×10^2	1.91×10^3	4.14×10^{-1}	2.69×10^1	2.02×10^2
		0%	3%	8%	0%	3%	8%
F_5	Flow in incinerated FP	3.17×10^1	9.43×10^2	3.60×10^3	2.48	4.48×10^1	1.80×10^2
		3%	10%	14%	2%	5%	7%
F_6	Flow in landfilled and composted FP	6.81×10^2	5.92×10^3	1.12×10^4	7.20×10^1	6.72×10^2	1.39×10^3
		60%	66%	45%	61%	71%	53%
$F_{6,Michigan}$	Flow in landfilled FP sent to Michigan	—	—	—	3.60	7.39×10^1	2.78×10^2
		—	—	—	3%	8%	11%
$F_{6,Canada}$	Flow in FP landfilled or composted in Canada	—	—	—	6.84×10^1	5.98×10^2	1.11×10^3
		—	—	—	58%	63%	42%
Non-polymeric PFASs							
F_1	Flow in the sales of FP	4.19×10^{-1}	1.13×10^1	6.04×10^1	4.37×10^{-2}	1.18	6.39
F_2	Flow in discarded FP	4.19×10^{-1}	1.13×10^1	6.04×10^1	4.37×10^{-2}	1.18	6.39
		0%	0%	0%	0%	0%	0%
F_3	Flow in littered FP	1.26×10^{-1}	5.63×10^{-1}	9.06	1.31×10^{-2}	5.88×10^{-2}	9.58×10^{-1}
		30%	5%	15%	30%	5%	15%
F_4	Flow in recycled FP	2.93×10^{-2}	2.14	1.54×10^1	3.06×10^{-3}	2.23×10^{-1}	1.63
		7%	19%	26%	7%	19%	26%
$F_{4,1}$	Flow in recycled FP that is recycled back to FP	1.47×10^{-3}	3.85×10^{-1}	5.08	1.53×10^{-4}	4.02×10^{-2}	5.38×10^{-1}
		0%	3%	8%	0%	3%	8%
$F_{4,2}$	Flow in recycled FP that is recycled into other materials	2.64×10^{-2}	1.43	5.70	2.76×10^{-3}	1.50×10^{-1}	6.03×10^{-1}
		6%	13%	9%	6%	13%	9%
$F_{4,3}$	Flow in FP initially intended for recycling but that is ultimately landfilled or exported	1.47×10^{-3}	3.21×10^{-1}	4.62	1.53×10^{-4}	3.35×10^{-2}	4.89×10^{-1}
		0%	3%	8%	0%	3%	8%
F_5	Flow in incinerated FP	1.17×10^{-2}	1.18	8.72	9.19×10^{-4}	5.59×10^{-2}	4.34×10^{-1}
		3%	10%	14%	2%	5%	7%
F_6	Flow in landfilled and composted FP	2.52×10^{-1}	7.38	2.72×10^1	2.66×10^{-2}	8.38×10^{-1}	3.37
		60%	66%	45%	61%	71%	53%
$F_{6,Michigan}$	Flow in landfilled FP sent to Michigan	—	—	—	1.33×10^{-3}	9.22×10^{-2}	6.73×10^{-1}
		—	—	—	3%	8%	11%
$F_{6,Canada}$	Flow in FP landfilled or composted in Canada	—	—	—	2.53×10^{-2}	7.46×10^{-1}	2.69
		—	—	—	58%	63%	42%

where F_{intake} is the flow of PFASs from food packaging digested or inhaled by humans resulting from food heating or consumption. Since we had insufficient information to estimate the human intake flow, we assumed that this intake would be negligible compared to the amount remaining within the food packaging (ESI Section S2.4†).

F_3 : flow in littered food packaging. A small portion of food packaging waste is littered, quantified as

$$F_3 = F_2 \times \%_{\text{litter}} \quad (4)$$

where $\%_{\text{litter}}$ is the percentage of food packaging that is littered. Values for $\%_{\text{litter}}$ were assumed (ESI Section 2.5).

F_4 : flow in recycled food packaging. The flow in recycled food packaging, F_4 , is the sum of several flows which depend on the percentage of food packaging recycled ($\%_{\text{FP recycled}}$). It was divided into four parts,



$$F_4 = F_{4,1} + F_{4,2} + F_{4,3} + F_{\text{recycling}} \quad (5)$$

$$F_4 = (F_2 - F_3) \times \%_{\text{FP recycled}} \quad (6)$$

$$F_{4,1} = F_4 \times \%_{\text{FP recycled to FP}} \quad (7)$$

$$F_{4,2} \cong F_4 - F_{4,1} - F_{4,3} \quad (8)$$

$$F_{4,3} = F_4 \times \%_{\text{recycling to landfill}} \quad (9)$$

$F_{4,1}$ is the flow from recycled food packaging that is recycled back to food packaging, which depends on the percentage of food packaging recycled back to food packaging, $\%_{\text{FP recycled to FP}}$, estimated using data on the use of recycled paper in the production of new paper in the U.S.⁵³ $F_{4,2}$ is the flow from recycled food packaging that is recycled into other materials. $F_{4,3}$ is the flow from food packaging intended for recycling but that is ultimately landfilled or exported ($\%_{\text{recycling to landfill}}$), which could represent up to one-third of discarded food packaging.^{54–56} $F_{\text{recycling}}$ is the flow from the recycling process of food packaging emitted to the environment, which is assumed to be negligible due to a lack of information sufficient for quantification (ESI Section S2.6†).

F_5 : flow in incinerated food packaging. The flow of PFASs in incinerated food packaging depends on the percentage of food packaging in the waste stream that is sent to incineration ($\%_{\text{incineration}}$) (Table 1 and ESI Section 2.7†). Values for $\%_{\text{incineration}}$ were taken from U.S. and Canadian sources where waste incineration is more common in the U.S. than in Canada.^{57–59}

$$F_5 = (F_2 - F_3) \times \%_{\text{incineration}} \quad (10)$$

F_6 : flow in landfilled and composted food packaging. Food packaging that is not littered, incinerated or recycled is sent to landfill and composting facilities in the U.S., Canada, or abroad (ESI Section 2.8†). We did not have access to data distinguishing municipal garbage that is landfilled *versus* composted, and thus have combined both flows into F_6 (which separates into $F_{6\text{-compost}}$ and $F_{6\text{-landfill}}$). Some waste for landfilling from Canada (specifically Ontario) is sent to Michigan, U.S., by mutual agreement (F_6 , Michigan, $\%_{\text{Can. waste to Mich.}}$),⁶⁰ where M_{Mich} and M_{Can} represent the mass of Canadian waste sent to Michigan and Canada landfills, respectively.

$$F_6 = F_2 - F_3 - F_4 - F_5 \quad (11)$$

For Canada more specifically:

$$F_6 = F_{6,\text{Canada}} + F_{6,\text{Michigan}} \quad (12)$$

$$F_{6,\text{Michigan}} = F_6 \times \%_{\text{Can. waste to Mich.}} = F_6 \times \frac{M_{\text{Mich}}}{M_{\text{Can}}} \quad (13)$$

3 Results and discussion

The inflows of polymeric PFASs were 9000 (range 1100–25 000) and 940 (range 120–2600) tonnes per year for middle (low–high)

scenarios in the U.S. and Canada, respectively (Fig. 1 and Table 2). Of this total inflow, >99% were polymeric forms. The total estimate inflows of non-polymeric PFASs were 11 (range 0.42–60) and 1.2 (range 0.044–6.4) tonnes per year for the U.S. and Canada, respectively. Total inflows were dominated by the intentional addition of PFASs: ~80% and 35% of the polymeric and non-polymer PFASs flows, respectively, were estimated to have come from food packaging with PFASs added at levels to confer grease and water repellency (ESI Tables S9 and S10†). Conversely, 65% of the non-polymeric (environmentally mobile) inflow of 7.2 tonnes per year or ~0–18 tonnes per year combined for U.S. and Canada, was attributable to the occurrence of PFASs found in food packaging at non-functional concentrations. The total PFASs inflow was due to ~2% and 40–60% of total food packaging containing functional and non-functional addition of PFASs, respectively.

The scenario based on middle values provides the most realistic estimate of actual flows, benchmarked against an estimate of PFASs used in food packaging in the U.S. The flow of 9000 tonnes per year in the U.S. was about a third of the 21 600 tonnes per year of fluorinated substances that could be used annually for food packaging in the U.S. based on values from ~2000 and 2015 (ESI Section S3.2†).^{5,66} Note that none of the upcoming bans in food packaging by three U.S. states or the U.S. voluntary phase-out from food packaging were considered in this analysis (the ban in New York will occur after 2022).⁶⁷

All food packaging produced moved to end-of-life, according to our assumption of limited storage. For the U.S., ~70% (60–50%, low–high values) or 6100 (range 690–13 000, flows F_6 and $F_{3,2}$) tonnes of polymeric PFASs from food packaging per year went to landfill and composting facilities, ~5% (30–15%) or 450 (340–3700) tonnes per year to litter and ~10% (3–14%) or 940 (32–3600) tonnes per year to incineration. An unknown fraction of food packaging sent to landfill and/or composting was sent abroad.⁶⁸

As mentioned above, some breakdown of polymeric PFASs into non-polymeric forms is inevitable. However, degradation rates and yield, including degradation of side-chain fluorinated polymers, are highly uncertain.^{69,70} Van der Veen *et al.*⁷¹ studied releases from durable water-repellent clothes, measuring 5- to 100-fold increases of non-polymeric PFASs concentrations following weathering, probably due to the degradation and transformation of precursors used for the formation of the durable water-repellent polymers. Similar values were obtained by Schellenberger *et al.*⁷² These results suggest that even limited breakdown of the large mass of polymeric PFASs from food packaging would considerably increase the mass and flows of non-polymeric PFASs.

Releases of PFASs from food packaging to the environment occur at different life-cycle stages and from various sources. Wastewater treatment plants (WWTPs) receive PFASs from food packaging with associated releases from WWTPs to the environment through outgassing,^{73–76} biosolids applied to land,^{29,77,78} or the discharge of final effluent.^{79–82} Outgassing and leachate from landfills are substantial sources discharged to the environment. We estimated that every year at least 2.27 and 0.26 tonnes of non-polymeric PFASs are released to the U.S. and



Canadian environment, respectively, due to the limited ability of WWTP to treat PFASs contained in landfill leachate (ESI Section S2.8†). Furthermore, Lang *et al.*²⁵ estimated that the entire U.S. waste load (of which about 30% was food packaging) generates at least 563 to 638 kg per year of selected non-polymeric PFASs in landfill leachate, of which 5 : 3 fluoroelomer carboxylic acid (FTCA) was the dominant PFAS identified out of the limited PFASs analysed, noting that composition varies depending on the age of the landfill. Leachate is often directed to WWTPs that are not designed to efficiently treat PFASs.⁸³

WWTPs also receive PFASs from food packaging *via* human waste, unquantified due to a lack of data. However, several studies have highlighted the tendency of PFASs to transfer from food packaging to food,^{2,23} which leads to ingestion through food consumption.⁸⁴ The heating process of food packaging, such as popcorn microwave bags, can also lead to airborne releases of PFASs which can be inhaled.²⁴

Although unquantified, recent analyses of municipal compost from the U.S. found higher concentrations of PFAAs in composts that included food packaging than in those that did not.²⁷ Other major unquantified flows included PFASs released during the manufacturing and recycling of food packaging containing PFASs^{85–87} and from incineration.^{88,89}

4 Implications and conclusions

This analysis is the first estimate of flows of PFASs used in food packaging – a total of 9000 and 940 tonnes per year in the U.S. and Canada, respectively. Of this, over 99% was polymeric, arising from an estimated intentional addition of PFASs to only ~2% of food packaging. Despite this low rate of PFAS usage, our estimates indicate the potential to contaminate the entire waste stream, including materials intended for recycling, landfilling and litter.

In addition to illustrating the potential of PFASs used in food packaging to contaminate the environment, numerous parts of the waste stream, and new products that contain recycled paper content, these results highlight the lack of knowledge regarding the overall presence and fate of PFASs in food packaging. While substance-based use and emission estimates of PFASs species and groups have been compiled,^{42,69,90–92} it is difficult to compare the flows of PFASs in food packaging estimated here for the U.S. and Canada with PFASs used in other applications because national product-based inventories of PFAS uses and emissions have not been compiled. The one exception is of PFASs used in carpets in California, for which the flows to landfill are orders-of-magnitude lower than the flows of food packaging estimated in this study, on a per capita basis, likely due to the high recycling rate of carpets in the U.S.⁴⁴

For environmental releases, particularly glaring knowledge gaps regarding PFASs use in food packaging, which are also instructional for other PFASs uses, are releases during the production with implications for human exposure, and fate at end-of-life (including food packaging intended for recycling but that is exported to low-income countries).^{93,94} Another knowledge gap is the contribution of polymeric PFASs to non-

polymeric forms due to the presence of non-polymeric impurities, and degradation in landfills and the environment. Although total flows were dominated by polymeric PFASs, of concern are non-polymeric flows of at least ~11 and 1.2 tonnes per year in the U.S. and Canada, given the mobility and higher toxicity of these substances. These knowledge gaps stress the need for chemicals management regimes to take a comprehensive life-cycle approach that considers all stages, from manufacturing to end-of-life, and types of PFASs, when adjudicating the use of chemicals in food packaging and other applications, which is not currently the case in the U.S. or in Canada.^{31,32}

The flows of non-functionally added PFASs in food packaging (estimated at ~20% of total PFASs flows), including 7.2 and 0.75 tonnes per year in the U.S. and Canada, respectively, of non-polymeric PFASs, highlight our lack of control of PFASs entering food packaging from unintentional use. However, it is the low intentional use of PFASs in food packaging (~2% of the total food packaging) that leads to the largest releases of PFASs, and highlights hotspots/blindspots for the sound management of food packaging.^{95,96}

These results, specifically the low percentage (~2%) of food packaging with intentionally added PFASs, lend support to regulatory initiatives to eliminate the use of PFASs in food packaging in seven U.S. states,⁶⁷ the voluntary agreement brokered between the food packaging industry and the U.S. FDA to restrict which PFAS can be used in food packaging,³⁷ and initiatives among several European countries to restrict non-essential uses of PFAS.^{97–99} We question whether these initiatives would be able to control flows of PFASs from non-intentional introduction to food packaging in these jurisdictions. Canada has not yet introduced restrictions for PFASs use specific to food packaging and, as noted above, the regulatory process is neither transparent or comprehensive.^{100,101}

Given the high persistence and potential for harm of the entire class of PFASs, both polymers and non-polymers, this analysis leads to questioning the use of PFASs in food packaging in any jurisdiction.

Conflicts of interest

There are no conflicts to declare.

Acknowledgements

We thank Graham White, Anastase Rulibikiye and Abdul Afghan (Health Canada) for discussions on chemicals management, and gratefully acknowledge the assistance of Alec Gonzales, Matt Roddy, Nick Caterisano, Chase Miller and Molly LaDuca in acquiring PIGE data, and Saleena Hak for shopping. Funding was provided by Environment and Climate Change Canada, Great Lakes Protection Initiative (3000694876, GCXE21P039), Natural Sciences and Engineering Research Council of Canada (NSERC, RGPIN-2017-06654), and by charitable contributions to the Green Science Policy Institute, including from the Target Foundation.



References

- R. C. Buck, J. Franklin, U. Berger, J. M. Conder, I. T. Cousins, P. de Voogt, A. A. Jensen, K. Kannan, S. A. Mabury and S. P. van Leeuwen, Perfluoroalkyl and polyfluoroalkyl substances in the environment: Terminology, classification, and origins, *Integr. Environ. Assess. Manage.*, 2011, **7**, 513–541.
- T. H. Begley, K. White, P. Honigfort, M. L. Twaroski, R. Neches and R. A. Walker, Perfluorochemicals: Potential sources of and migration from food packaging, *Food Addit. Contam.*, 2005, **22**, 1023–1031.
- L. A. Schaidler, S. A. Balan, A. Blum, D. Q. Andrews, M. J. Strynar, M. E. Dickinson, D. M. Lunderberg, J. R. Lang and G. F. Peaslee, Fluorinated Compounds in U.S. Fast Food Packaging, *Environ. Sci. Technol. Lett.*, 2017, **4**, 105–111.
- G. Yuan, H. Peng, C. Huang and J. Hu, Ubiquitous Occurrence of Fluorotelomer Alcohols in Eco-Friendly Paper-Made Food-Contact Materials and Their Implication for Human Exposure, *Environ. Sci. Technol.*, 2016, **50**, 942–950.
- X. Trier, K. Granby and J. H. Christensen, Polyfluorinated surfactants (PFS) in paper and board coatings for food packaging, *Environ. Sci. Pollut. Res.*, 2011, **18**, 1108–1120.
- L. Schultes, G. F. Peaslee, J. D. Brockman, A. Majumdar, S. R. McGuinness, J. T. Wilkinson, O. Sandblom, R. A. Ngwenyama and J. P. Benskin, Total Fluorine Measurements in Food Packaging: How Do Current Methods Perform?, *Environ. Sci. Technol. Lett.*, 2019, **6**, 73–78.
- T. Shoeib, Y. Hassan, C. Rauert and T. Harner, Poly- and perfluoroalkyl substances (PFASs) in indoor dust and food packaging materials in Egypt: Trends in developed and developing countries, *Chemosphere*, 2016, **144**, 1573–1581.
- I. Zabaleta, E. Bizkarguenaga, D. Bilbao, N. Etxebarria, A. Prieto and O. Zuloaga, Fast and simple determination of perfluorinated compounds and their potential precursors in different packaging materials, *Talanta*, 2016, **152**, 353–363.
- I. Zabaleta, N. Negreira, E. Bizkarguenaga, A. Prieto, A. Covaci and O. Zuloaga, Screening and identification of per- and polyfluoroalkyl substances in microwave popcorn bags, *Food Chem.*, 2017, **230**, 497–506.
- B. Barhoumi, S. G. Sander and I. Tolosa, A review on per- and polyfluorinated alkyl substances (PFASs) in microplastic and food-contact materials, *Environ. Res.*, 2022, **206**, 112595.
- D. of T. S. C. (DTSC) CalEPA, *Product-Chemical Profile for Food Packaging Containing Perfluoroalkyl or Polyfluoroalkyl Substances: Discussion Draft*, 2020.
- K. L. Vorst, N. Saab, P. Silva, G. Curtzwiler and A. Stekete, Risk assessment of per- and polyfluoroalkyl substances (PFAS) in food: Symposium proceedings, *Trends Food Sci. Technol.*, 2021, **116**, 1203–1211.
- J. Straková, J. Schneider, N. Cingotti, A. Bennett, A. den Boer, K. Brabcová, S. Brosché, M. Fernandez, F. Gorre, E. Hawke, G. Jensen, N. Malval, M. Möller, S. Müller, A. Roger, F. Veillerette, B. Wahlund and J.-L. Wietor, *Throwaway Packaging, Forever Chemicals: European-wide Survey of PFAS in Disposable Food Packaging and Tableware*, 2021.
- C. F. Kwiatkowski, D. Q. Andrews, L. S. Birnbaum, T. A. Bruton, J. C. Dewitt, D. R. U. Knappe, M. V. Maffini, M. F. Miller, K. E. Pelch, A. Reade, A. Soehl, X. Trier, M. Venier, C. C. Wagner, Z. Wang and A. Blum, *Environ. Sci. Technol. Lett.*, 2020, **7**, 532–543.
- C. M. Butt, U. Berger, R. Bossi and G. T. Tomy, *Sci. Total Environ.*, 2010, **408**, 2936–2965.
- S. Chu and R. J. Letcher, Side-chain fluorinated polymer surfactants in aquatic sediment and biosolid-augmented agricultural soil from the Great Lakes basin of North America, *Sci. Total Environ.*, 2017, **607–608**, 262–270.
- M. Houde, A. O. De Silva, D. C. G. Muir and R. J. Letcher, Monitoring of perfluorinated compounds in aquatic biota: An updated review, *Environ. Sci. Technol.*, 2011, **45**, 7962–7973.
- J. P. Giesy and K. Kannan, Peer Reviewed: Perfluorochemical Surfactants in the Environment, *Environ. Sci. Technol.*, 2002, **36**, 146A–152A.
- L. S. Haug, C. Thomsen and G. Becher, Time trends and the influence of age and gender on serum concentrations of perfluorinated compounds in archived human samples, *Environ. Sci. Technol.*, 2009, **43**, 2131–2136.
- T. Zhang, Q. Wu, H. W. Sun, X. Z. Zhang, S. H. Yun and K. Kannan, Perfluorinated compounds in whole blood samples from infants, children, and adults in China, *Environ. Sci. Technol.*, 2010, **44**, 4341–4347.
- C. Lau, K. Anitole, C. Hodes, D. Lai, A. Pfahles-Hutchens and J. Seed, Perfluoroalkyl acids: A review of monitoring and toxicological finds, *Toxicol. Sci.*, 2007, **99**, 366–394.
- V. Barry, A. Winquist and K. Steenland, Perfluorooctanoic Acid (PFOA) Exposures and Incident Cancers among Adults Living Near a Chemical Plant, *Environ. Health Perspect.*, 2013, **121**, 1313–1318.
- T. H. Begley, W. Hsu, G. Noonan and G. Diachenko, Migration of fluorochemical paper additives from food-contact paper into foods and food simulants, *Food Addit. Contam., Part A*, 2008, **25**, 384–390.
- E. Sinclair, S. K. Kim, H. B. Akinleye and K. Kannan, Quantitation of gas-phase perfluoroalkyl surfactants and fluorotelomer alcohols released from nonstick cookware and microwave popcorn bags, *Environ. Sci. Technol.*, 2007, **41**, 1180–1185.
- J. R. Lang, B. McKay Allred, J. A. Field, J. W. Levis and M. A. Barlaz, National Estimate of Per-and Polyfluoroalkyl Substance (PFAS) Release to U.S. Municipal Landfill Leachate, *Environ. Sci. Technol.*, 2017, **51**, 2197–2205.
- J. P. Benskin, B. Li, M. G. Ikonomou, J. R. Grace and L. Y. Li, Per- and polyfluoroalkyl substances in landfill leachate: Patterns, time trends, and sources, *Environ. Sci. Technol.*, 2012, **46**, 11532–11540.



- 27 Y. J. Choi, R. K. Lazcano, P. Yousefi, H. Trim and L. S. Lee, Perfluoroalkyl Acid Characterization in U.S. Municipal Organic Solid Waste Composts, *Environ. Sci. Technol. Lett.*, 2019, **6**, 372–377.
- 28 A. C. Blaine, C. D. Rich, L. S. Hundal, C. Lau, M. A. Mills, K. M. Harris and C. P. Higgins, Uptake of perfluoroalkyl acids into edible crops via land applied biosolids: Field and greenhouse studies, *Environ. Sci. Technol.*, 2013, **47**, 14062–14069.
- 29 J. Yu, J. Hu, S. Tanaka and S. Fujii, Perfluorooctane sulfonate (PFOS) and perfluorooctanoic acid (PFOA) in sewage treatment plants, *Water Res.*, 2009, **43**, 2399–2408.
- 30 A. B. Lindstrom, M. J. Strynar, A. D. Delinsky, S. F. Nakayama, L. McMillan, E. L. Libelo, M. Neill and L. Thomas, Application of WWTP biosolids and resulting perfluorinated compound contamination of surface and well water in Decatur, Alabama, USA, *Environ. Sci. Technol.*, 2011, **45**, 8015–8021.
- 31 U.S. Food and Drug Administration (FDA), *Inventory of Effective Food Contact Substance (FCS) Notifications*, accessed 9 December 2020, https://www.cfsanappsexternal.fda.gov/scripts/fdcc/index.cfm?set=FCN&sort=FCN_No&order=DESC&startrow=1&type=basic&search=.
- 32 PackagingLaw.com, Canada Considering Positive List System for Food-Contact Materials, accessed 22 December 2020, <https://www.packaginglaw.com/news/canada-considering-positive-list-system-food-contact-materials>.
- 33 Stockholm Convention, *Perfluorooctanoic acid (PFOA), its salts and PFOA-related compounds*, accessed 23 November 2020, <http://chm.pops.int/Implementation/Alternatives/AlternativestoPOPs/ChemicalslistedinAnnexA/PFOA/tabid/8292/Default.aspx>.
- 34 Stockholm Convention, *The new POPs under the Stockholm Convention*, accessed 23 November 2020, <https://www.pops.int/TheConvention/ThePOPs/TheNewPOPs/tabid/2511/Default.aspx>.
- 35 United States Environmental Protection Agency, *Fact Sheet: 2010/2015 PFOA Stewardship Program*, accessed 21 September 2021, <https://www.epa.gov/assessing-and-managing-chemicals-under-tsca/fact-sheet-20102015-pfoa-stewardship-program>.
- 36 Government of Canada, *Canada Gazette, Part 1, Volume 156, Number 20: Prohibition of Certain Toxic Substances Regulations*, 2022, accessed 28 July 2022, <https://www.gazette.gc.ca/rp-pr/p1/2022/2022-05-14/html/reg2-eng.html>.
- 37 U.S. Food and Drug Administration (FDA), *Authorized Uses of PFAS in Food Contact Applications*, <https://www.fda.gov/food/chemical-contaminants-food/authorized-uses-pfas-food-contact-applications>, accessed 7 March 2022.
- 38 S. V. Kabadi, J. W. Fisher, D. R. Doerge, D. Mehta, J. Aungst and P. Rice, Characterizing biopersistence potential of the metabolite 5:3 fluorotelomer carboxylic acid after repeated oral exposure to the 6:2 fluorotelomer alcohol, *Toxicol. Appl. Pharmacol.*, 2020, **388**, 114878.
- 39 OECD, *PFASs and Alternatives in Food Packaging (Paper and Paperboard) Report on the Commercial Availability and Current Uses, 2020 Series on Risk Management No. 58*, ENV.
- 40 F. Li, J. Duan, S. Tian, H. Ji, Y. Zhu, Z. Wei and D. Zhao, Short-chain per- and polyfluoroalkyl substances in aquatic systems: Occurrence, impacts and treatment, *Chem. Eng. J.*, 2020, **380**, 122506.
- 41 A. K. Rosenmai, C. Taxvig, T. Svingen, X. Trier, B. M. A. van Vugt-Lussenburg, M. Pedersen, L. Lesné, B. Jégou and A. M. Vinggaard, Fluorinated alkyl substances and technical mixtures used in food paper-packaging exhibit endocrine-related activity in vitro, *Andrology*, 2016, **4**, 662–672.
- 42 L. Li, J. Liu, J. Hu and F. Wania, Degradation of Fluorotelomer-Based Polymers Contributes to the Global Occurrence of Fluorotelomer Alcohol and Perfluoroalkyl Carboxylates: A Combined Dynamic Substance Flow and Environmental Fate Modeling Analysis, *Environ. Sci. Technol.*, 2017, **51**, 4461–4470.
- 43 D. Laner, J. Feketitsch, H. Rechberger and J. Fellner, A Novel Approach to Characterize Data Uncertainty in Material Flow Analysis and its Application to Plastics Flows in Austria, *J. Ind. Ecol.*, 2016, **20**, 1050–1063.
- 44 J. Chen, L. Tang, W.-Q. Chen, G. F. Peaslee and D. Jiang, Flows, Stock, and Emissions of Poly- and Perfluoroalkyl Substances in California Carpet in 2000–2030 under Different Scenarios, *Environ. Sci. Technol.*, 2020, **54**, 6908–6918.
- 45 N. Seltenrich, PFAS in Food Packaging: A Hot, Greasy Exposure, *Environ. Health Perspect.*, 2020, **128**, 054002.
- 46 Arkema Kynar®, *Polymer Processing and Recycling Aids for Polyolefins and other Polymers*, accessed 7 September 2021, <https://www.extremematerials-arkema.com/en/markets-and-applications/chemical-industry-and-general-industry/fluoropolymer-processing-aids/>.
- 47 S. E. Amos, *The Application of Fluoropolymer Based Processing Aids in Polypropylene and Thermoplastic Rubber*, *TAPPI Polym. Laminations, Coat. Conf., Proc.*, 1998.
- 48 G. W. Curtzwiler, P. Silva, A. Hall, A. Ivey and K. Vorst, Significance of Perfluoroalkyl Substances (PFAS) in Food Packaging, *Integr. Environ. Assess. Manage.*, 2021, **17**, 7–12.
- 49 R. J. Letcher, S. Chu and S. A. Smyth, Side-chain fluorinated polymer surfactants in biosolids from wastewater treatment plants, *J. Hazard. Mater.*, 2020, **388**, 122044.
- 50 B. J. Henry, J. P. Carlin, J. A. Hammerschmidt, R. C. Buck, L. W. Buxton, H. Fiedler, J. Seed and O. Hernandez, A critical review of the application of polymer of low concern and regulatory criteria to fluoropolymers, *Integr. Environ. Assess. Manage.*, 2018, **14**, 316–334.
- 51 M. Eiji, T. Motohiro, K. Junsuke and M. Sho, *JP Patent*, JP2017078227A, 2014.
- 52 X. Liu, Z. Guo, E. E. Folk and N. F. Roache, Determination of fluorotelomer alcohols in selected consumer products and preliminary investigation of their fate in the indoor environment, *Chemosphere*, 2015, **129**, 81–86.
- 53 U.S. EPA, *Wastes - Resource Conservation - Common Wastes & Materials - Paper Recycling*, accessed 3 September 2020,



- <https://archive.epa.gov/wastes/conserve/materials/paper/web/html/faqs.html>.
- 54 Livia Albeck-Ripka, *Your Recycling Gets Recycled, Right? Maybe, or Maybe Not*, accessed 20 October 2020, <https://www.nytimes.com/2018/05/29/climate/recycling-landfills-plastic-papers.html>.
- 55 Christy Somos, *This is why Canadian waste keeps ending up in Asia*, accessed 20 October 2020, <https://www.ctvnews.ca/world/this-is-why-canadian-waste-keeps-ending-up-in-asia-1.4441472>.
- 56 Frances Bula, *China's tough new recycling standards leaving Canadian municipalities in a bind*, accessed 20 October 2020, <https://www.theglobeandmail.com/news/national/chinese-ban-on-foreign-recyclables-leaving-some-canadian-cities-in-the-lurch/article37536117/>.
- 57 Statistics Canada, *Human Activity and the Environment - Annual Statistics 2005 - Feature Article Solid Waste in Canada*, Ottawa, 2005.
- 58 Solid Waste & Recycling, *Incineration in Canada*, <https://www.solidwastemag.com/feature/incineration-in-canada/>, accessed 16 September 2020.
- 59 U.S. EPA, *Advancing Sustainable Materials Management*, 2019.
- 60 Michigan Department of Environment, *Report of Solid Waste Landfilled in Michigan*, 2020.
- 61 Government of Canada, *Plastics Challenge – Food Packaging*, accessed 13 August 2020, <https://www.ic.gc.ca/eic/site/101.nsf/eng/00036.html>.
- 62 Statistics Canada, *Table 38-10-0032-01 Disposal of Waste*, 2018.
- 63 U.S. EPA, *Reducing Wasted Food & Packaging: A Guide for Food Services and Restaurants*, 2015.
- 64 U.S. EPA, *Containers and Packaging: Product-Specific Data*, accessed 14 September 2020, <https://www.epa.gov/facts-and-figures-about-materials-waste-and-recycling/containers-and-packaging-product-specific-data>.
- 65 Resource Productivity & Recovery Authority (RPR), *Datacall Report 2018*, 2018.
- 66 F. Alpizar, T. Backhaus, N. Decker, I. Eilks, N. Escobar-Pemberthy, P. Fantke, K. Geiser, M. Ivanova, O. Jolliet and H.-S. Kim, UN Environment Global Chemicals Outlook II-From Legacies to Innovative Solutions: Implementing the 2030, *Agenda for Sustainable Development*, 2019.
- 67 Keller and Heckman LLP, *New York Bans PFAS in Food Packaging*, *Natl. Law Rev.*, 2020, **11**, 1.
- 68 *The Canadian Press*, *Canadian garbage on its way from Philippines to Vancouver*, accessed 8 January 2021, <https://www.cbc.ca/news/technology/canadian-garbage-from-philippines-departure-1.5156007>.
- 69 Z. Wang, I. T. Cousins, M. Scheringer, R. C. Buck and K. Hungerbühler, Global emission inventories for C4-C14 perfluoroalkyl carboxylic acid (PFCA) homologues from 1951 to 2030, part II: The remaining pieces of the puzzle, *Environ. Int.*, 2014, **69**, 166–176.
- 70 I. Liagkouridis, R. Awad, S. Schellenberger, M. M. Plassmann, I. T. Cousins and J. P. Benskin, Combined Use of Total Fluorine and Oxidative Fingerprinting for Quantitative Determination of Side-Chain Fluorinated Polymers in Textiles, *Environ. Sci. Technol. Lett.*, 2022, **9**, 30–36.
- 71 I. van der Veen, A. C. Hanning, A. Stare, P. E. G. Leonards, J. de Boer and J. M. Weiss, The effect of weathering on per- and polyfluoroalkyl substances (PFASs) from durable water repellent (DWR) clothing, *Chemosphere*, 2020, **249**, 126100.
- 72 S. Schellenberger, I. Liagkouridis, R. Awad, S. Khan, M. Plassmann, G. Peters, J. P. Benskin and I. T. Cousins, An Outdoor Aging Study to Investigate the Release of Per- and Polyfluoroalkyl Substances (PFAS) from Functional Textiles, *Environ. Sci. Technol.*, 2022, **56**, 3471–3479.
- 73 M. Shoeib, J. Schuster, C. Rauert, K. Su, S. A. Smyth and T. Harner, Emission of poly and perfluoroalkyl substances, UV-filters and siloxanes to air from wastewater treatment plants, *Environ. Pollut.*, 2016, **218**, 595–604.
- 74 I. Weinberg, A. Dreyer and R. Ebinghaus, Waste water treatment plants as sources of polyfluorinated compounds, polybrominated diphenyl ethers and musk fragrances to ambient air, *Environ. Pollut.*, 2011, **159**, 125–132.
- 75 L. Vierke, L. Ahrens, M. Shoeib, E. J. Reiner, R. Guo, W.-U. Palm, R. Ebinghaus and T. Harner, Air concentrations and particle-gas partitioning of polyfluoroalkyl compounds at a wastewater treatment plant, *Environ. Chem.*, 2011, **8**, 363.
- 76 L. Ahrens, M. Shoeib, T. Harner, S. C. Lee, R. Guo and E. J. Reiner, Wastewater treatment plant and landfills as sources of polyfluoroalkyl compounds to the atmosphere, *Environ. Sci. Technol.*, 2011, **45**, 8098–8105.
- 77 P. Guerra, M. Kim, L. Kinsman, T. Ng, M. Alaei and S. A. Smyth, Parameters affecting the formation of perfluoroalkyl acids during wastewater treatment, *J. Hazard. Mater.*, 2014, **272**, 148–154.
- 78 C. Zhang, H. Yan, F. Li, X. Hu and Q. Zhou, Sorption of short- and long-chain perfluoroalkyl surfactants on sewage sludges, *J. Hazard. Mater.*, 2013, **260**, 689–699.
- 79 L. Ahrens, S. Felizeter, R. Sturm, Z. Xie and R. Ebinghaus, Polyfluorinated compounds in waste water treatment plant effluents and surface waters along the River Elbe, Germany, *Mar. Pollut. Bull.*, 2009, **58**, 1326–1333.
- 80 L. Ahrens, Polyfluoroalkyl compounds in the aquatic environment: A review of their occurrence and fate, *J. Environ. Monit.*, 2011, **13**, 20–31.
- 81 R. Bossi, J. Strand, O. Sortkjær and M. M. Larsen, Perfluoroalkyl compounds in Danish wastewater treatment plants and aquatic environments, *Environ. Int.*, 2008, **34**, 443–450.
- 82 R. Guo, W. J. Sim, E. S. Lee, J. H. Lee and J. E. Oh, Evaluation of the fate of perfluoroalkyl compounds in wastewater treatment plants, *Water Res.*, 2010, **44**, 3476–3486.
- 83 H. O. Kwon, H. Y. Kim, Y. M. Park, K. S. Seok, J. E. Oh and S. D. Choi, Updated national emission of perfluoroalkyl substances (PFASs) from wastewater treatment plants in South Korea, *Environ. Pollut.*, 2017, **220**, 298–306.



- 84 H. P. Susmann, L. A. Schaidler, K. M. Rodgers and R. A. Rudel, Dietary Habits Related to Food Packaging and Population Exposure to PFASs, *Environ. Health Perspect.*, 2019, **127**, 107003.
- 85 Environmental Defense Fund (EDF), *Paper Mills as a Significant Source of PFAS Contamination, but Who's Watching?*, accessed 8 December 2020, <https://blogs.edf.org/health/2018/05/21/pfas-paper-mills/>.
- 86 H. A. Langberg, H. P. H. Arp, G. D. Breedveld, G. A. Slinde, Å. Høiseter, H. M. Grønning, M. Jartun, T. Rundberget, B. M. Jenssen and S. E. Hale, Paper product production identified as the main source of per- and polyfluoroalkyl substances (PFAS) in a Norwegian lake: source and historic emission tracking, *Environ. Pollut.*, 2020, 116259.
- 87 H. A. Langberg, H. P. H. Arp, G. D. Breedveld, G. A. Slinde, Å. Høiseter, H. M. Grønning, M. Jartun, T. Rundberget, B. M. Jenssen and S. E. Hale, Paper product production identified as the main source of per- and polyfluoroalkyl substances (PFAS) in a Norwegian lake: Source and historic emission tracking, *Environ. Pollut.*, 2021, **273**, 116259.
- 88 M. Schlummer, C. Sölch, T. Meisel, M. Still, L. Gruber and G. Wolz, Emission of perfluoroalkyl carboxylic acids (PFCA) from heated surfaces made of polytetrafluoroethylene (PTFE) applied in food contact materials and consumer products, *Chemosphere*, 2015, **129**, 46–53.
- 89 A. N. García, N. Viciano and R. Font, Products obtained in the fuel-rich combustion of PTFE at high temperature, *J. Anal. Appl. Pyrolysis*, 2007, **80**, 85–91.
- 90 Z. Wang, I. T. Cousins, M. Scheringer and K. Hungerbühler, Fluorinated alternatives to long-chain perfluoroalkyl carboxylic acids (PFCAs), perfluoroalkane sulfonic acids (PFSAs) and their potential precursors, *Environ. Int.*, 2013, **60**, 242–248.
- 91 Z. Wang, I. T. Cousins, M. Scheringer, R. C. Buck and K. Hungerbühler, Global emission inventories for C4-C14 perfluoroalkyl carboxylic acid (PFCA) homologues from 1951 to 2030, Part I: Production and emissions from quantifiable sources, *Environ. Int.*, 2014, **70**, 62–75.
- 92 J. Glüge, M. Scheringer, I. T. Cousins, J. C. DeWitt, G. Goldenman, D. Herzke, R. Lohmann, C. A. Ng, X. Trier and Z. Wang, An overview of the uses of per- and polyfluoroalkyl substances (PFAS), *Environ. Sci.: Processes Impacts*, 2020, **22**, 2345–2373.
- 93 Organisation for Economic Cooperation and Development (OECD), *Working towards a Global Emission Inventory of PFASs: Focus on PFCAs - Status Quo and the Way Forward*, Paris, 2015.
- 94 K. L. Law, N. Starr, T. R. Siegler, J. R. Jambeck, N. J. Mallos and G. H. Leonard, The United States' contribution of plastic waste to land and ocean, *Sci. Adv.*, 2020, **6**, 1–8.
- 95 B. Geueke, K. Groh and J. Muncke, Food packaging in the circular economy: Overview of chemical safety aspects for commonly used materials, *J. Cleaner Prod.*, 2018, **193**, 491–505.
- 96 T. Savvaides, J. P. Koelmel, Y. Zhou, E. Z. Lin, P. Stelben, J. J. Aristizabal-Henao, J. A. Bowden and K. J. Godri Pollitt, Prevalence and Implications of Per-and Polyfluoroalkyl Substances (PFAS) in Settled Dust, *Curr. Environ. Health Rep.*, 2022, 1–13.
- 97 Danish Veterinary and Food Administration, *Ban on fluorinated substances in paper and board food contact materials (FCM)*, accessed 7 March 2022, <https://www.foedevarestyrelsen.dk/english/SiteCollectionDocuments/Kemiogfoedevarekvalitet/UK-Fact-sheet-fluorinated-substances.pdf>.
- 98 SIAL Paris, *PFAS rules for 2022 set to improve eco-safety of food packaging*, accessed 7 March 2022, <https://newsroom.sialparis.com/regional-spotlight/europe/pfas-rules/>.
- 99 European Chemical Agency (ECHA), *Perfluoroalkyl chemicals (PFAS)*, accessed 7 March 2022, <https://echa.europa.eu/hot-topics/perfluoroalkyl-chemicals-pfas>.
- 100 Government of Canada, *Information Requirements For Food Packaging Submissions*, accessed 30 August 2021, <https://www.canada.ca/en/health-canada/services/food-nutrition/legislation-guidelines/guidance-documents/information-requirements-food-packaging-submissions.html>.
- 101 Government of Canada, *Substance prohibition summary for PFOA, long-chain PFCAs and related substances*, accessed 22 December 2020.

