RSC Advances



REVIEW

View Article Online



Cite this: RSC Adv., 2014, 4, 47806

Received 14th August 2014 Accepted 23rd September 2014

DOI: 10.1039/c4ra08675a

www.rsc.org/advances

Recent advances in transition-metal-catalyzed C-CN bond activations

Qiaodong Wen, Ping Lu* and Yanguang Wang*

C-C bond activation and cleavage have become more and more attractive in recent years for the great challenge of breaking C-C bonds and the great advantage of constructing new chemical bonds. Among them, the activation of inert C-CN bonds by transition metals has been widely investigated and a number of improvements have been achieved by many groups. This review mainly summarizes recent advances in this field, based on different kinds of reactions derived from C-CN bond activation, including cyanofunctionalization, cross coupling reactions and cyanation. Moreover, mechanistic studies involving transition-metal-catalyzed C-CN bond activation are also mentioned

Introduction


Nitriles are not only important in natural products, pharmaceuticals, agrochemicals, dyes, herbicides, and materials,1 but also can be transferred into other compounds with various functionalities, such as amines, amidines, amides, aldehydes, carboxylic acids, pyrazoles, triazoles, and tetrazoles,2 through C≡N transformations. Moreover, through transition metal catalyzed C-CN bond activation, nitriles can be utilized in the reactions of cyanofunctionalization, decyanative crosscoupling, and cyanation. That is why numerous methodologies related to the making as well as the breaking of C-CN bonds have been established. In contrast to the review articles considering the construction of C-CN bonds from the aspects of various transition metals and safe cyanide sources,3 reviews about the transition-metal catalyzed C-CN bond activation are seldom reported.4 Herein, we will describe the recent progress in this field based on reaction categories and mechanistic studies.

C-CN bond is relatively stronger and stable in comparison with C-X bond referring to have higher bond dissociation energy (BDE). BDEs of C-CN bonds in CH₃-CN, CH₂=CH-CN, HC≡C-CN, and Ph-CN are 121.1, 132.1, 152.1, and 132.7 kcal mol⁻¹,⁵ respectively. In comparison with these data, BDEs of C-X bonds in Ph-F, Ph-Cl, Ph-Br, and Ph-I are 127.4, 97.3, 82.7, and 66.9 kcal mol⁻¹, respectively. Thus, the selective cleavage of C-CN bonds by using transition metal catalyst is a challengeable target and of great importance in organic synthesis. Reactions involving the cleavage of C-CN bonds can be simply categorized into three types according to the utility degree of nitriles. In cyanofunctionalizations, both R and CN units are

presented in products (route a, Scheme 1). When "CN" is used as the leaving group, R-CN becomes an alternative substrate of R-X/R-OTf for cross coupling reactions which have been widely applied in making C-Si, C-P, C-B, C-H, as well as C-C bonds (route b, Scheme 1). With the disclosure of the safe cyanation reagents in the recent years, R-CN has also been developed for the operational benign cyanation reagents, such as benzyl nitrile,7 and acetonitrile.8 In these reactions, "CN" unit is successfully transferred from starting materials to target molecules (route c, Scheme 1).

2. Cyanofunctionalizations

When oxidative addition of transition metal occurred on C-CN bond, an active intermediate, acyl/alkynyl/alkenyl/alkyl metal cyanide (C-TM-CN), is formed and C-CN bond is activated and finally cleaved. The subsequent 1,2-migratory insertion of C-TM-CN species into unsaturated bond, followed by reductive elimination affords a variety of addition products and regenerates transition metal for the next catalytic cycle (Scheme 2). This transition metal catalyzed reaction is called cyano functionalization or carbocyanation. Because such reaction installs two C-C bonds simultaneously onto an unsaturated bond with great atom efficiency, this reaction is surely of great synthetic



Scheme 1 Three patterns of the utility of RCN in organic synthesis.

Department of Chemistry, Zhejiang University, Hangzhou, 310027, P. R. China. E-mail: pinglu@zju.edu.cn; orgwyg@zju.edu.cn; Fax: +86-571-87952543; Tel: +86-571-87952543

Scheme 2 General mechanism of cyanofunctionalization.

values. Up to now, outstanding results have been achieved in this area, including cyanoacylation, cyanoesterification, cyanoamidation, cyanoarenylation, cyanoalkenylation, cyanoalkynylation, and cyanoalkylation, according to the carbon types where C–CN bond is activated. Most of these cyanofunctionalizations occur on alkyne, and some of them happen on alkene and 1,2-diene. All these reactions undergo exclusively in *syn*-addition fashion with high regioselectivities when unsymmetrical triple or double bonds are applied. Moreover, these cyanofunctionalizations could be significantly promoted by adding appropriate Lewis acid.

2.1 Cyanoacylation

Early in 1994, Takaya reported palladium-catalyzed cyanoacylation of terminal acetylenes with aryloyl cyanides, leading to the formation of alkenes attached with two electronwithdrawing groups, acyl and cyano (Scheme 3).9a Under the designated palladium catalyst system, phenylacetylene (1) reacted with 4-methylbenzoyl cyanide (2a), giving the corresponding addition product (3), accompanied by trace amount of regio isomer (4) and acetylenic ketone (5). The amount of 5 was decreased when the reaction was carried out for a longer time. Thus acetylenic ketone (5) was proposed as a plausible intermediate during the reaction process. Through this intermediate, addition products (3) were formed by a sequence of Pdcatalyzed hydrocyanation and isomerization (Scheme 4). If the Pd-catalyzed hydrocyanation occurred in a different direction, Z-4 would be produced. Thus, E-4 was formed through isomerism. Although the formation of intermediate C was

Scheme 3 Pd-catalyzed cyanoacylation of phenylacetylene with 4-methylbenzoyl cyanide.

$$Ar^{1} CN Pd(0)$$

$$Ar^{2} Pd A$$

$$Ar^{1} COAr^{2}$$

$$Ar^{2} COAr^{2}$$

$$Ar^{3} COAr^{2}$$

$$Ar^{4} COAr^{2$$

Scheme 4 Proposed mechanism of Pd-catalyzed cyanoacylation.

reasonable, the route through intermediate 5 was more possible. This strategy provided a route for the regioselective preparation of electron-deficient alkenes. A detail study on the role of the mixed ligands was presented in 1996. PPh₃ was responsible for its suitable basicity and cone angle, while dppb (1,4-bis(diphenylphosphino)butane) benefitted for a higher conversion.

2.2 Cyanoesterification

In 2006, Nishihara reported palladium-catalyzed C–CN bond cleavage of cyanoformates (7) and addition to norbornenes (6), affording various norbornane derivatives (8) with two functional groups, cyano and ester (Scheme 5). ¹⁰ Benzonorbornadiene and norbornadienes were both applicable to this reaction, as well as many other cyanoformates.

In their mechanistic studies, the trans-Pd(CN)(CO₂Et)(PPh₃)₂ (A) was isolated and proved to be the key intermediate by the addition of Pd(PPh₃)₄ to ethyl cyanoformate (7a) (Scheme 6). Single crystal analysis of the Me analogue of trans-Pd(CN)(CO₂Et)(PPh₃)₂ (A) indicated that A possessed a square planar structure with two triphenylphosphines in trans positions. 10a 1,2-Migratory insertion of trans-A to the double bond of norbornene (alkoxypalladation) followed by the reductive elimination gave the cyanoesterification product (8a) in synaddition manner. Two groups (CN and COOEt) occupied the exo-positions instead of the endo-positions. That an alternative 1,2-migratory insertion of trans-A norbornene

Scheme 5 Pd-catalyzed cyanoesterification of norbornene.

В

little lower yield.

COOEt

high affinity of cyano to palladium center.10b

.CN

cyanopalladation

(cyanopalladation generating \mathbf{B}') was ruled out because of the

Meanwhile, Nakao and Hiyama reported a nickel-catalyzed C–CN bond cleavage of cyanoformates (7) and their sequential cyanoesterifications of cumulative 1,2-dienes (9), affording various 2-(1-cyanoalk-1-yl)acrylates (10), as well as their regioisomers (11) (Scheme 7). ¹¹ By reversing the molar ratio of 7a to 9a to 1.0: 1.2, 11a was formed predominantly. ¹¹ Except for cyanoformates (7), benzoyl cyanide (2b) was also suitable for this reaction under modified conditions and provided 12 with a

In their proposed mechanism, Ni catalyst inserted to $C(sp^2)$ –CN bond (Scheme 8). 1,2-Migratory insertion of C–Ni–CN (A) on 1,2-diene occurred on the less steric hindered terminal double bond with the ester group migrated to the central carbon of 1,2-diene. In this way, C was regioselectively formed through the possible complex C and C-allylnickel intermediate (C). By reductive elimination in two different types, two different products (C0 and C-C11a) were obtained. C11a was proposed to be obtained C11a was thought to be the thermodynamic product, whereas C10a

Scheme 7 Ni-catalyzed cyanoesterification/cyanoacylation on 1,2-diene.

$$E-11a + L_{n}Ni(0)$$

$$E = \frac{10a}{Ni(0)L_{n}}$$

Scheme 8 Proposed mechanism for Ni-catalyzed cyanoesterification on 1.2-diene.

was a kinetic product. This proposal was proved by the condition controlled experiments when the reaction temperature was increased to 100 $^{\circ}$ C and the amount of **10a** was significantly decreased.

In 2011, Douglas demonstrated an intramolecular Pd-catalyzed cyanoesterification, accessing to functionalized lactones (13) in moderate to excellent yields (Scheme 9).¹² The decarboxylation and decarbonylation byproducts were successfully suppressed by performing the reactions in microwave reactor at high temperature. The authors also hypothesized that the 1,2-migratory insertion was the product-determining step, rather than the step of C(sp²)–CN bond activation, because of the higher reactivity of electron-rich alkynes.

2.3 Cyanoamidation

In 2006, Takemoto reported the intramolecular cyanoamidation of alkyne and alkene through the designation of substrate structures (14, 16, and 18) (Scheme 10).¹³ Nonaromatic substrates (14) proceeded exclusively in *syn*-addition manner and provided α -alkylidene lactams (15) in moderate to good yields. In case of n=1, isomerized product 15' was obtained in 39% yield because of the 1,3-migration of the C-4 proton.

Aromatic substrate (16 and 18) provided indole derivatives 17 and 19, respectively. When cyanoamidation occurred on triple bonds, the stereoselectivity was largely depended upon the R group which was attached on the triple bonds. When R was phenyl or CH_2OTBS group, yields were moderate and the products were in exclusively *Z*-form. When R was altered to butyl, a mixture of *Z* and *E* in a ratio of 69 : 31 was isolated in 97% yield. When cyanoamidation occurred on double bonds, a quaternary carbon center was created and 3.3'-disubstituted

Scheme 9 Pd-catalyzed intramolecular cyanoesterification.

Scheme 10 Pd-catalyzed intramolecular cyanoamidations

oxindole 19 was obtained almost quantitatively, which could be used as the key intermediate in the preparation of physostigmine.

All these reactions occurred in a same manner as the previous cyanofunctionalizations. Pd(0) was oxidatively added to amido cyanides (14, 16, and 18) to form amidopalladium cyanide intermediates (A) (Scheme 11). Thus, a sequential amidopalladation occurred on triple or double bonds to generate intermediates B. Finally, amido and cyano groups were installed into both ends of triple or double bonds by reductive elimination. When two reaction sites (amido cyanide and unsaturated bonds) existed in one molecule, an intramolecular reaction happened to afford lactams successfully.

NC
$$R_2N$$
 $Pd(0)$ R_2N CN R_2N CN R_2N R_2N R_2N R_2N $Pd(II)$ R_2 R_1 R_2 R_2 R_2 R_2 R_2 R_1 R_2 R_2 R_2 R_2 R_1 R_2 R_2 R_2 R_2 R_2 R_1 R_2 R_2 R_2 R_1 R_2 R_2 R_2 R_2 R_2 R_1 R_2 R_2 R_2 R_1 R_2 R_2 R_2 R_2 R_2 R_1 R_2 R_2 R_2 R_2 R_1 R_2 R_2 R_2 R_2 R_1 R_2 R_2 R_1 R_2 R_2 R_2 R_2 R_2 R_1 R_2 R_2 R_2 R_2 R_1 R_2 R_2 R_2 R_2 R_2 R_1 R_2 R_2 R_2 R_1 R_2 R_2 R_2 R_2 R_2 R_1 R_2 R_2 R_2 R_2 R_1 R_2 R_2

Scheme 11 Proposed mechanism for Pd-catalyzed intramolecular cyanoamidations.

Scheme 12 Enantioselective synthesis of (S)-19 through Pd-catalyzed cyanoamidation.

Later on, Takemoto practiced the enantioselective synthesis of 3,3'-disubstituted oxindole (S-19) by finely tuning the catalyst, chiral ligand, additive, and solvent (Scheme 12).14 By screening these factors, a combination of Pd(dba)2, an optically active phosphoramidite, and a suitable Lewis base in decalin was optimized to be the best conditions to reach the effective enantioselective synthesis of 3,3'-disubstituted oxindole (S-19). It was also noticed that this combination was effective to a broad range of substrates and might be applied in the synthesis of more complicated molecules.

2.4 Cyanoalkenylation/cyanoalknylation/cyanoalkylation

In 2004, Nakao and Hiyama developed a Ni-catalyzed cyanoarylation of different alkynes (21) through the activation of Ar-CN bonds, installing aryl and cynao in the two ends of triple bonds and providing various β-aryl substituted alkenenitriles (Scheme 13). Lots of functional groups on the aryl nitriles (20) tolerated the reaction conditions, such as the carbonyl groups of ketone, aldehyde and ester. Cyano group also survived after reaction if equivalent molar ratio of reactants was used. The boryl also tolerated the reaction conditions which indicated

Scheme 13 Preliminary studies of Ni-catalyzed cyanoarylation on oct-4-yne.

RSC Advances Review

that a variety of products could be further derived from the boryl group through Suzuki–Miyaura couplings. Aryl nitriles bearing electron-neutral and electron-donating groups afforded the products with relatively longer time, which indicated the rate-determining step was oxidative addition of Ni(0) to C(sp²)–CN bond. When unsymmetrical alkynes were used, mixtures were obtained. However, a regioselective preparation could be realized when the steric hindrance of two groups on triple bond could be distinguished, such as methyl *versus tert*-butyl. The bulkier group and the cyano group were found to be bonded to the same carbon in the final. Unfortunately, terminal alkynes did not work under the standard reaction conditions due to its rapid trimerization or oliomerization.

They proposed a possible mechanism (Scheme 14).^{15b} Ni(0) catalyst inserted to Ar–CN bond of arenecarbonitrile (20) through oxidative addition, forming a Ni(II) complex (A). Then the Ni(II) complex (A) regioselectively added to the internal alkyne by 1,2-migratory insertion depending on the steric hindrance. A subsequently reductive elimination provided a cyanoarylation product (22) in *syn*-addition manner and regenerated the Ni(0) catalyst for the next catalytic cycle. An alternative 1,2-migratory insertion, arylnickelation instead of cyanonickelation, could not be ruled out although the repulsion between ligand and bulkier R² group was apparent.

Besides aryl nitriles, allyl cyanides (23) could be used as the substrate for the oxidative addition of Ni(0), and provided the cyanoallylation products (24) when the resulting nickel species reacted with alkynes (Scheme 15). ¹⁶ Thus, reaction between allyl

Scheme 14 Proposed mechanism for Ni-catalyzed cyanoarylation.

Scheme 15 Ni-catalyzed cyanoallylations with allyl cyanides.

cyanides (23) and oct-4-yne (21a) in the presence of 10 mol% $\rm Ni(cod)_2$ and 20 mol% $\rm P(4\text{-}CF_3\text{-}C_6H_4)_3$ provided 6-substituted (2 $\rm Z$,5 $\rm E$)-2,3-dipropylhexa-2,5-dienenitriles (24) in regio- and stereoselective way. Partial isomerism of the double bond was observed. Synthetic utility of 2,5-dienenitriles was demonstrated by selective reductions. For instance, the resulting acrylonitriles could easily be reduced into acrylaldehydes and allyl alcohols by selecting suitable reductants without contamination of the reductive products derived from alkene unit.

Addition of α -siloxyallyl cyanide (25), which derived from the reaction between acrolein and TMSCN, onto oct-4-yne (21a) under the aforementioned reaction conditions afforded (Z)-5-formyl-2,3-dipropylpent-2-enenitrile (26) in 81% yield (Scheme 16). The silyl enol ether (29) was isolated and indicated to be the key intermediate by using t-butyldimethylsilyl (TBDMS) instead of TMS. To these cyanoalkylation reactions, terminal alkynes worked well and provided 2-substituted (Z)-5-formyl-pent-2-enenitriles (30) in high regioselectivity and high stereoselective. Furthermore, the formyl group could perform the aldehyde reactions and provided α -functionalized acrolein (27) and hepta-2,6-dienenitrile (28) effectively.

A possible mechanism was proposed by authors (Scheme 17). Ni catalyst firstly added to $C(sp^3)$ –CN bond of allyl cyanides (23, 25) via oxidative addition, giving a π -allylnickel intermediate (A). 1,2-Migratory addition of π -allylnickel intermediate alkynes, and followed by reductive elimination, acrylonitriles (24, 29) were obtained. Here the products having the bulkier group at a cyano-substituted carbon were much favored in consideration of the steric hindrance, especially to cases of terminal alkynes in which 2-substituted (Z)-5-formyl-pent-2-enenitriles (30) were afforded in excellent regioselectivity.

A breakthrough work was reported in 2007. Nakao and Hiyama demonstrated a Lewis-acid assisted Ni-catalyzed cyanoarylation of alkynes with nitriles, which efficiently improved the reactivities of some electron-rich aryl cyanides (20a), comparing to the previous works (Scheme 18).^{17a} Moreover, the amount of the catalyst and the ligand was significantly decreased. By surveying the substituted group on aryl cyanides, it was indicated that selective activation of Ar–CN bond over Ar–Br/Cl was remarkable. Bromo or

Scheme 16 Ni-catalyzed cyanoallylations with α -siloxyallyl cyanide.

$$\begin{array}{c} \text{CN} & \text{NC} \\ \text{R}^{1} & \text{R}^{3} \\ \text{24, 29} & \text{L}_{n} \text{Ni}(0) \\ \text{23} & \text{CN} \\ \text{Q1} & \text{Or 25} \\ \text{NC} - \text{Ni} \\ \text{Or 25} \\ \text{R}^{1} & \text{A} & \text{R}^{1} \\ \text{C} & \text{CN} \\ \text{R}^{2} & \text{CN} \\ \text{R}^{2} & \text{CN} \\ \text{R}^{3} & \text{R}^{2} & \text{CN} \\ \text{R}^{2} & \text{CN} & \text{R}^{2} & \text{R}^{3} \\ \text{Steric hindrance: } \text{R}^{3} \geq \text{R}^{2} \\ \end{array}$$

Scheme 17 Proposed mechanism for the Ni-catalyzed cyanoallylations.

Scheme 18 LA-mediated Ni-catalyzed cyanoarylations

chloro (22b) was survived after reaction. Moreover, sterically hindered 2,6-dimethyl benzonitrile (20c) participated the reaction and the Ar-CN bond was activated if the amount of the catalyst, ligand, and Lewis acid was magnified 5 times, the reaction time was prolonged to 134 h, and the reaction temperature was raised to 100 °C. The corresponding product (22c) was obtained in 78% yield. Another interesting case was the cyanoarylation of 1-methyl-1*H*-indole-3-carbonitrile (31) over 21a. Activation of Ar-CN occurred prior to the activation of Ar-H at 2-position of indole in the presence of Lewis acid.¹⁸

In the aid of Lewis acid, cyanoalkenylation was also successfully achieved by fine tuning ligand and Lewis acid.

Thus, a series of conjugated dienenitriles were obtained in good to excellent yields (Scheme 19). Alkene configuration of *trans*-cinnamonitrile (33) was remained after reaction and the resulting dienenitrile (34) could be transferred into pyridine (35) by a sequence of reduction, 6π -electrocyclization, and aromatization. 2-Menzylidenemalononitrile (36) participated the reaction and provided dicyano-substituted 1,3-diene (37) in 81% yield. In this case, a selective activation of Ar–CN bond which appeared *trans* to phenyl group was demonstrated. Further, other cyanofunctionalizations, such as aforementioned cyanoesterification 17b and cyanoamidation 27c were all optimized in the aid of Lewis acid.

Finally, $C(sp^3)$ –CN bond in acetonitrile was activated. Reaction of acetonitrile (38) and oct-4-yne gave trisubstituted acrylonitrile (39a) in 71% yield (Scheme 20). That methyl group came from acetonitrile albeit AlMe₃ was confirmed by using CH_3CN-d_3 . Other nitriles, such as 2-(trimethylsilyl)-acetonitrile and propiononitrile, provided the corresponding addition products, but in relatively lower yields. In case of propiononitrile (40), it provided the *syn*-adduct (39b) in 24% yield only. One of the by-products was (*E*)-2-propylhex-2-enenitrile (41) (Scheme 21). Oxidative addition of Ni(0) to propiononitrile leading to the formation of ethylnickel cyanide (A), which

Scheme 19 LA-mediated Ni-catalyzed cyanoalkenylations.

Scheme 20 LA-mediated Ni-catalyzed cyanomethylation.

Scheme 21 Formation of the hydrocyanation byproducts in cyanoalkylations.

underwent β -elimination to generate highly active species, HNiCN (B). In the presence of oct-4-yne, hydrocyanation occurred and provided the by-product 41.

The dramatic effects of Lewis acids could be contributed to the coordination between R-CN and Lewis acid which accelerated the rate of oxidative addition of Ni(0) to R-CN bond (Scheme 22). Regioselectivity was ascribed to the repulsion between the bulkier R² and R when the unsymmetrical alkyne and nickel species coordinated. Ultimately, the bulkier R² group and cyano group (CN) bonded to the same carbon of alkynes. Moreover, these reactions occurred in exclusive *syn*-addition manner, the *trans*-adducts came from the isomerism of *syn*-adducts in the presence of Lewis acid or ligand, or under heat.

Short after, Nakao and Hiyama reported cyanoalkylation of alkynes with arylacetonitriles (42), also catalyzed by nickel and Lewis acid (Scheme 23). Various arylacetonitriles could undertake this reaction in good yields, including some heteroaryl derivatives. As arylacetonitriles, β -elimination resulted from the normal alkanenitriles, such as aforementioned byproduct (41), could be effectively diminished. To the symmetrical alkynes, addition occurred in an exclusive *syn*-fashion. However, to the unsymmetrical alkynes, poor regioselectivities were observed except for the cases where the steric

 $\begin{array}{c} R \\ R^1 \\ CN \\ CN \\ L \ or \ heat \\ (LA \ cat.) \\ R^1 \\ R^2 \\ CN-LA \\ R^1 \\ CN-LA \\ R^1 \\ CN-LA \\ R^1 \\ CN-LA \\ R^2 \\ CN-LA \\ R^1 \\ R^2 \\$

Scheme 22 Proposed mechanism for the LA-mediated Ni-catalyzed cyanoalkylation.

Scheme 23 LA-mediated Ni-catalyzed cyanoalkylation of alkynes with arylacetonitriles.

hindrance could be differentiated, such as methyl *versus tert*-butyl, methyl *versus* TMS. The bulkier group always linked to carbon with CN attached in the final.

To make a clear outcome of the mechanism, the authors selected optically active (S)- α -phenylpropionitrile (44) as the substrate and detected all the byproducts. It was worth noting that the recovered starting materials showed 80% ee and the product did not undertake racemization under their conditions. These facts indicated that the absolute configuration of the chiral carbon retained after these sequential steps, including the oxidative addition, 1,2-migratory insertion, and reductive elimination. The partial loss of the optical purity could be ascribed to the equilibrium between the β -elimination and its reversed 1,2-migratory insertion as illustrated in Scheme 24.

In 2010, Nakao and Hiyama reported heteroatom-directed cyanoalkylation of alkynes (Scheme 25). The heteroatoms, like nitrogen atom, played a significant role in suppressing the aforementioned unexpected β -hydride elimination via the intramolecular chelating of nitrogen with nickel center. The amino effect at the γ -position of butyronitrile (46a) was impressive. When the amino was on the β -position of propiononitrile, only trace amount of the syn-adduct was detected under the same reaction conditions. It was also noticeable that when the amino group was on δ -position of pentanenitrile, with

NC R₂ Ar LNI(0) NC Ar LA=P-ligand LA=AlMe₂Cl steric hindrance:
$$R^2 \ge R^1$$
LA-NC R_2 R_1 R_2 R_3 R_4 R_4 R_4 R_5 R_6 R_7 R_8 R_9 R_9

Scheme 24 Retention of configuration in LA-mediated Ni-catalyzed cyanoalkylation.


Scheme 25 Heteroatom-directed cyanoalkylation of alkyne.

one more methylene unit between pyrrolidine and cyano (46b), cyanoalkylation occurred, but on the γ -position of the pyrrolidyl group. This selectivity was also observed in case where 5-pyrrolidyl hexanenitrile (46c) was used. In this case, a longer reaction time was required and the hydrocyanation by-product (41) was isolated in 9%.

The above experiments indicated that a 5-membered azanickelacycle (C) was the key intermediate (Scheme 26). With one more CH_2 , a 6-membered azanickelacycle (F) (or even 7-membered in case with two more CH_2) could be converted to 5-membered azanickelacycle (C) by a sequence of β -hydrogen elimination and hydronickelation. This proposed mechanism properly explained the fact that the amino-substituted butyronitrile, valeronitrile and hexanenitrile could undertake cyanoalkylation smoothly. Similar to the above mentioned cases, the *syn*-adducts formed having a bulkier group and CN bonded to the same sp^2 -carbon when unsymmetrical alkynes were used.

Other heteroatom (O, S) in the appropriate position of alkanenitrile (48, 50) could also function as the directing heteroatom *via* its coordination to nickel center and afford the corresponding cyanoalkylation products (49, 51) (Scheme 27).

Owning to homo- and/or cross-cyclotrimerization in nickel catalyst system, it was uneasy for alkynyl cyanide (52) to undertake the catalytic insertion of Ni(0) to C(sp)-CN bond. Nevertheless, Nakao and Hiyama reported nickel/BPh₃-



Scheme 26 Proposed mechanism for the heteroatom-directed cyanoalkylation of alkyne.

Scheme 27 Examples for O, S-directed cyanoalkylation of alkyne.

catalyzed cyanoalkynylation of alkynes, accessing to conjugated enyne structures (53) (Scheme 28).²¹ With the aid of Lewis acid, the trimerization products (54) were effectively inhibited.

After investigating the substrate scope, they found that bulky silyl substituented alkynyl cyanides (55) might suppress the homo- and/or cross-cyclotrimerization of the nitriles, resulting in a lower requirement of catalyst amounts and approached excellent yields of 56 (Scheme 29). Diynyl cyanides were also suitable for this reaction. 1,2-Dienes (57) reacted with alkynyl cyanides (55) under the same conditions, giving conjugated enynes (58) in moderate to good yields. The regioselectivity was largely depending upon the substituted group attached on 1,2-diene.

The oxidative insertion of C(sp)–CN bond with Ni(0) catalyst assisted by BPh_3 was thought to be the initial step in the proposed mechanism to form key intermediate **A** (Scheme 30). Followed by the coordination with alkynes or 1,2-dienes, respectively, complexes **B** and **D** were formed. In case of the alkyne as substrate, the alkynyl group migrated from the nickel center to the less hindered carbon of the coordinated alkyne to generate **C**. In case of 1,2-diene as substrate, the alkynyl migrated from nickel center to the central carbon of 1,2-diene to generate the allylnickel species **E**. Finally, the reductive elimination of **C** or **E** took place and provided the addition products, **56** and **58** respectively, with both high regioselectivity and high stereoselectivity.

In 2011, Kurahashi and Matsubara reported an intermolecular cycloaddition of *o*-arylcarboxy benzonitriles (59) and alkynes (21) through Ni-catalyzed cleavage of C–CN and C–CO

Scheme 28 BPh₃-mediated Ni-catalyzed cyanoalknylation of alkyne.

Scheme 29 Cyanoalknylation of alkyne or 1,2-diene with silyl substituented alkynyl cyanides.

55 58
$$| N_1 | | 1$$
 56 55 $| N_2 | | 1$ 57 $| N_3 | | 1$ 58 $| N_4 | | 1$ 58 $| N_4 | | 1$ 58 $| N_4 | | 1$ 59 $| N_4 | | 1$ 50 $| N_4 | | 1$

Scheme 30 Proposed mechanism for cyanoalknylation of alkyne or 1,2-diene.

bond, affording coumarins (**60**) in good yields (Scheme 31).²² It was worth noting that aryl cyanide (**20d**) was isolated, as well as the compound (**22d**) formed from its carbocyanation with alkyne. Similarly, *o*-arylamide-substituted benzonitrile (**61**) could also undertake this reaction with oct-4-yne, which gave the quinolinone (**62**) in 83% yield.²³

In 2013, Nakao and Hiyama chose polyfluorobenzonitriles (20e, 20f) to conduct cyanoarenylation with alkynes, affording a series of corresponding polyfluoroarylcyanation products (22e, 22f) by applying Ni(cod)₂/DPEphos/BPh₃ catalytic system in the solvent of cyclopentylmethylether (CPME) (Scheme 32).24 Under their reaction conditions, C-CN bond activation was prior to C-H and C-F bond activation, so the resulting cyanoarenvlation products could be obtained smoothly without the interference with the byproducts derived from C-H and C-F bond activations. Of note, the existence of fluorine atoms on the arenes the C-CN bond activation, driving fluorobenzonitriles more reactive than non-fluorine substituted aryl nitrile which was demonstrated by competition experiment between pentafluorobenzonitrile and benzonitrile. Moreover, the dramatic effect of BPh3 and DPEphos was clearly illustrated the single crystal structure of the nickel insertion

Scheme 31 Ni-catalyzed cycloadditions.

Scheme 32 Ni-catalyzed chemoselective C-CN bond activation over C-H/C-F bond activation.

intermediate (*cis-A*) which was responsible for the subsequent 1,2-migratory insertion of Ar_F–Ni–CN on alkyne. Combination of Ni(cod)₂, DPEphos, and BPh₃ in CPME was also used for cyanoarenylation of terminal alkynes and alkenes. As an interesting example, selective activation of C–H bond albeit C–CN bond was achieved by using PCyp₃ as ligand. Thus, compound (63) was obtained from 22f in 90% yield with high chemoselectivity.

2.5 Stereoselective cyanoarenylation

In comparison with above mentioned cyanofunctionalization of alkynes, cyanoarenylation of alkenes was seldom reported. In 2008, Jacobsen reported an enantioselectively intramolecular cyanoarenylation of unactivated olefins through Ni-catalyzed C(sp²)–CN bond activation, providing indane derivatives with a quaternary chiral center in good yield (Scheme 33).²⁵ By screening the conditions and selecting a proper chiral ligand, the indane product (65) was obtained in high enantioselectivity. Combination of NiCl₂·DME and Zn was used as the Ni(0)

Scheme 33 Asymmetric intramolecular cyanoarenylation of inactivated olefin.

source, while the olefin isomerism was minimized by increasing the ratio of TangPHOS to Ni.

Meanwhile, Nakao and Hiyama also reported the intramolecular cyanoarenylation of various specially designed alkenes with Ni(cod)₂/AlMe₂Cl catalyst. A number of substrates tolerated the reaction conditions (Scheme 34).²⁶ Amino and silyl tethers provided the corresponding cyclized products 67 and 68 in 86% and 92% yields, respectively, by choosing proper ligand. Insertion of olefin into Ar–CN bond was found to be in a 5-exo-trig fashion. The new constructed ring could be enlarged to six- or seven-membered ring, as indicated by the representative compounds 69 and 70. The cyanoarenylation reaction proceeded in a *syn*-addition manner which was clearly shown by the diaster-eoselective preparation of compounds 72a and 72b (Scheme 35).

Not only the reaction could be conducted in high diastereoselectivity, but also in high enantioselectivity by using a chiral ligand, such as (R,R)-i-Pr-Foxap (Scheme 36). This asymmetric synthesis had been successfully applied in the preparation of (-)-esermethole.

Mechanism of this intramolecular cyanoarenylation was proposed based on the isolation of key intermediates 75 and 76 (Scheme 37). Structures 75 and 76 were established by their single crystal analysis. Reaction of stoichiometric 64, Ni(cod)₂, P(n-Bu)₃, and AlMe₂Cl at room temperature immediately provided η^2 -nitrile complex 75. Existence of Lewis acid facilitated the oxidative addition of Ni(0) to Ar–CN bond and afforded complex 76 after 6 h. 1,2-Migratory insertion onto intramolecular olefin provided η^2 -nitrile complex 77 via intermediate A. Finally, cyclized product was obtained via ligand exchange.

Scheme 34 Substrate diversity for the Ni-catalyzed intramolecular cyanoarenylation.

Scheme 35 Diastereoselective preparation of 72.

Scheme 36 Enantioselective preparation of 74.

Scheme 37 Proposed mechanism of the intramolecular cyanoar-enylation of olefin.

Above all, a number of cyanofunctionalizations on unsaturated alkynes, alkenes, 1,2-dienes were disclosed, including cyanoeserification, cyanoamidation, cyanoacylation, cyanoarenylation, cyanoalkenylation, cyanoalkylation, and cyanoalkynylation based on the carbon type bonded to CN group in the substrates. All of these reactions were of great synthetic value because they simultaneously installed two groups into one molecule with 100% atom efficiency. More noteworthy, reactions proceeded in *syn*-addition fashion with high stereoselectivity.

3. Cross coupling reactions

R part in nitriles (R–CN) can undertake cross coupling reactions with other components, in which CN group is regarded as a leaving group. In comparison with the typical substrates (R–X or R–OTs) for cross coupling reactions in making C–Si, C–P, C–B, C–H, as well as C–C bonds, nitrile (R–CN) provides a unique and complementary way in the preparation of functionalized compounds.

3.1 Decyanative silvlation

In 2006, Chatani reported a Rh(1)-catalyzed cleavage of C-CN bond and Si-Si bond, achieving the decyanative silylation of aryl and alkenyl nitriles (Scheme 38).^{27a} They chose [RhCl(cod)]₂ as

RSC Advances Review

Scheme 38 Rh-catalyzed Si-assisted decyanative silylation of aryl nitriles

the catalyst, and hexamethyldisilane (78) as the silvlation agent which also functioned as the auxiliary reagent in the helping of C-CN bond cleavage. In the substrate scope, dimethylphenylsilyl and benzyldimethylsilyl groups could also be introduced under the reaction conditions established for hexamethyldisilane, although with moderate yields. Steric hindrance might influence the reaction rate and reduce the yields. Heteroaryl nitriles and alkenyl nitriles were both good substrates for this transformation by selecting suitable rhodium catalyst. It was also noteworthy that the boryl group (20g) survived after reaction which opened a window for subsequently selective cross coupling to build higher π -conjugated compounds. Although fluorine on the arvl nitriles survived after the reaction, C-Cl and C-Br bond were found to be more reactive than C-CN bond.


The decyanative silylation was well extended to allyl cyanides. Thus, allyl cyanide (23a) furnished allyl silane (79) in 59% yield (Scheme 39). A relative lower reaction temperature was applied for this transformation due to the relative facile cleavage of C-CN bond and the formation of the stable π -allyl rhodium complex.28 However, when benzyl cyanide (42a) was used as the substrate, a higher reaction temperature was applied for complete conversion of benzyl cyanide. In this case, enamine formation (81) was isolated in a certain amount. The enamine byproduct (84) became dominant when 2,2-diphenylacetonitrile (82) was used.

Rh-catalyzed Si-assisted decyanative silylation of allyl/ Scheme 39 benzyl nitriles.

It was also noteworthy that even the unactivated alkyl cyanide could be used for the decyanative silvlation when triisopropyl phosphite was added (Scheme 40). Thus, 4-phenylbutanenitrile (85) provided corresponding silvlated product (86) in 48% yield with 34% recovery of the starting material. In this case, the byproduct resulting from the β-hydrogen elimination of the possible alkyl metal intermediate was not observed.

In their proposed catalytic cycle (Scheme 41), silylrhodium species (A) was generated via σ-bond metathesis between Rh-X and Si-Si bonds. Alternatively, A could be formed via an oxidative/elimination process. Migration of silvl of Rh-Si (A) to the nitrogen atom of nitrile generated η^2 -iminoacyl complex (B) as the key intermediate in this catalytic cycle.27b,c Subsequent migration of aryl/alkyl to rhodium center provided aryl/alkyl rhodium species (C) coordinated with silvl isocyanide. In the presence of disilane (75), σ-bond metathesis between C and disilane (75) gave rise to the silvlated product (R-Si) and regenerated Rh-Si (A) for completing the catalytic cycle. Meanwhile, silyl isocyanide was excluded and isomerized into thermodynamically stable silvl cyanide in the final. When benzyl cyanide (42a) was used, two possible ways leading to the formation of enamine (81) was proposed. σ-Bond metathesis between B and disilane (75) produced D and regenerated A. Intramolecular migration of silyl from imino carbon to nitrogen via aza-Brook-type rearrangement formed aminocarbene (E). Through this path, enamine was finally obtained via the carbene insertion to adjacent C-H bond. Alternatively, β-hydrogen elimination of B, and followed by hydrorhodation across C=N bond of N-silyl ketenimine (F) generated G. As a result of σ -bond metathesis between G and disilane (75), enamine (81) was afforded in a certain amount.

Scheme 40 Rh-catalyzed Si-assisted decyanative silylation of alkyl



Scheme 41 Proposed mechanism for Rh-catalyzed Si-assisted decyanative silylations.

Further investigation of decyanative silvlation revealed that 1,5-rhodium migration step might be involved after the Rhcatalyzed C-CN bond cleavage if the substrates (87) were rationally designed (Scheme 42).29 Thus a mixture of decyanative silvated isomers (88 and 89) was obtained after the silvlation occurred on aryl rhodium species (A and B) before and after 1,5migration. Product ratio was highly dependent on the acidity of Ar-H which was putatively to be activated. With anyl possessing the electron donating group ($R^2 = OMe$), the direct decyanative silylation product was dominant. In a contrast, with aryl bearing the electron withdrawing group (3,4,5-trifluorophenyl), the 1,5migrated product was isolated as a major product. 1,5-Mgiration was indicated to be irreversible by conducting a comparative experiment and the rate of 1,5-migration resulted in the product distribution. Migration distance between the metal center and the Ar-H was suitable to form the six-membered metallacyclic intermediate (or transition state), no 1,4-, 1,6- or 1,7-migration was observed. Similar 1,5-migration was observed in case when diboron was used instead of disilane in the decyanative borylation.

3.2 Decyanative phosphination

In 2011, Yang reported the phosphination of aryl nitriles with phosphate sources (90) by the Ni-catalyzed C-CN bond activation followed by C-P coupling, obtaining various phosphorus ligands (Scheme 43).30 A series of rhodium catalysts failed to catalyze this decyanative phosphination, while NiCl₂(PPh₃)₂ and NiCl₂(PCy₃)₂ were found to be effective. t-BuOK was crucial to this transformation. Aryl part of aryl nitriles could be

Scheme 42 1,5-Rhodium migration involved in decyanative silylation.

Scheme 43 Ni-catalyzed decyanative phosphination of aryl nitriles.

transferred into the corresponding aromatic diphenylphosphines (91 and 93) in good yields, except several steric hindered aryl nitriles and pyridinyl nitriles which were obtained in relatively lower yields.

In order to clarify the mechanism, a controlled experiment was conducted (Scheme 44). By using hexamethyldisilane or tetraphenyldiphosphine instead of Me₃Si-PPh₂ under the same reaction conditions, the former reaction did not occur, while the later one furnished corresponding phosphinated product in 45% yield. Thus, the possible Si-assisted C-CN bond cleavage in the presence of nickel catalyst was excluded.

A plausible mechanism was proposed (Scheme 45). Ni(II) was firstly reduced to Ni(0) (A) in the presence of base for the subsequent oxidative addition to Ar-CN bond via the complexes (B and C) between aryl nitrile and nickel. As a result, aryl nickel cyanide (D) was formed. Ligand exchange through two possible routes as proposed by the authors provided intermediate (E). Subsequent reductive elimination of E produced the phosphinated product and regenerated Ni(0) (A) for next catalytic cycle.

Decyanative borylation 3.3

In 2011, Tobisu and Chatani described the borylation of nitriles with a diboron reagent (94) through Rh(1)-catalyzed C-CN bond cleavage (Scheme 46).31a,b In the optimization studies, they found the base and phosphine ligand were quite important in improving the yield, of which 1,4-diazabicyclo[2.2.2]octane (DABCO) and Xantphos were the best choices, respectively. Various nitriles, including aryl, alkenyl and benzylic nitriles, could form the corresponding boronic esters in good yields. This method provided a new strategy for the synthesis of boronic esters, which were very useful for further derivatization.

Scheme 44 Controlled experiments.

Scheme 45 Possible mechanism for the Ni-catalyzed decyanative phosphination of aryl nitriles.

Scheme 46 Rh-catalyzed decyanative borylation of aryl cyanides.

Of significant note, C–Cl bond and specified C–B bond survived after reactions which respectively allowed the orthogonal and sequential designs for further cross coupling reactions.

The mechanistic pathway supposed that the diboron (94) played an important role in the cleavage of C–CN bond (Scheme 47). σ -Bond metathesis between Rh–Cl bond and B–B bond generated Rh–B bond and B–Cl bond which was confirmed by ¹¹B NMR analysis. This step was the same to the previous metal-silicon bond formation. Addition of Rh–B species (A) across nitrile triple bond resulted in the formation of adduct (B) which isomerized to C for the sterically favored *syn*- β -aryl elimination. As a result, R–Rh (D) was formed and borylated *via* σ -bond metathesis with diboron. Although β -carbon elimination was proposed in Chatani's report, ³¹ α a mechanism including the sequential 2,1-insertion and deinsertion could not be excluded. ³¹ α 0

Rh-Cl
$$B-B$$
 $B-C$
 $B-B$
 $B-C$
 $B-B$
 $B-C$
 $B-B$
 $B-C$
 $B-C$

Scheme 47 Rh-catalyzed decyanative borylation of nitriles.

3.4 Decyanation

Besides of the cross coupling reactions with disilane, C–CN bond could also be cleaved and transfer to C–H bond, called decyanation. Such reactions were quite useful in organic synthesis because they offered the opportunity to temporally utilize the benefits from the CN group, such as the acidity of α -H, the ortho-directing effect, and the strong electron withdrawing nature of cyano group.

In 2009, Tobisu and Chatani reported a protocol for Rhcatalyzed reductive cleavage of C–CN bond with hydrosilane as the reducing agent, converting varying nitriles to the corresponding alkanes or arenes (Scheme 48).³² They selected [RhCl(cod)]₂ as the rhodium catalyst, P(OBu)₃ as the ligand and triisopropylsilane as the reducing agent, affording decyanated products with high efficiencies. A variety of functional groups tolerated the reaction conditions, such as dimethylamino, methoxy, ester, and boronic ester. A deuterated product (98) was also gained when the corresponding deuteriosilane was used in the reaction. Besides of aryl nitriles could be decyanated, benzilic nitriles as well as normal alkyl cyanides could also be

Scheme 48 Rh-catalyzed Si-assisted decyanation of aryl nitriles.

decyanated in excellent yields without the contamination from β -hydrogen elimination.

In 2013, Maiti demonstrated nickel catalyzed decyanation of nitriles selecting tetramethyldisiloxane (TMDSO) as the hydride source (Scheme 49).³³ In the catalytic system of Ni(acac)₂, PCy₃, and AlMe₃, aryl and aliphatic cyanides reacted with TMDSO and decyanated to arenes and aliphatic hydrocarbons, respectively. A variety of functional groups on aromatic ring suffered this reaction condition, such as methoxy, ester, and carbonyl. Cleavage of Ar–SMe bond occurred prior to the cleavage of Ar–CN bond. Heterocyclic cyanides also preceded smoothly under the reaction conditions. The deuterated product was isolated when Ph₂SiD₂ was used as the reductant. The *trans*-[Ni(PCy₃)₂(CN)₂] was isolated from the reaction, but failed to catalyze the decyanation, indicating that it was the cause of catalyst deactivation.

Short after, Maiti reported nickel-catalyzed decyanation of aryl and aliphatic cyanides using hydrogen gas as the hydride source (Scheme 50). The C-CN bond was efficiently activated in assistance of AlMe₃, which was also responsible to consume the *in situ* produced HCN gas. It was worthy to note that aliphatic cyanides (103), possessing β -hydrogens, were also applicable for this decyanation without the formation of the byproduct arising from β -hydrogen elimination. A variety of functional groups tolerated the reaction conditions, such as MeO-, HO-, F-, ester, amide, ketone, as well as the tethered olefin.

Scheme 49 Ni-catalyzed LA-assisted decyanation of aryl/alkyl nitriles with silane.

Scheme 50 Ni-catalyzed LA-assisted decyanation of aryl/alkyl nitriles with hydrogen.

In 2013, Enthaler reported decyanation of aryl and alkyl cyanides catalyzed by their pre-prepared nickel complex and employed tert-butylmagnesium chloride as the hydride source (Scheme 51).³⁵ Most substituent groups could tolerant the reaction conditions except for thioether and several halogens (Br and F). In these cases, the cleavage of C–S and C-X bonds was observed. Hydrogen atom in the product came from Grignard reagent which was further confirmed by using C_2D_5MgBr . Finally, they proposed a possible mechanism, in which C–CN bond was supposed to be cleaved through a single-electron transfer (SET) step and a radical procedure (Scheme 52).

3.5 Decyanative arylation

In 2001, Miller's group firstly reported the coupling of benzonitriles and aryl Grignard reagents using Me₃P-based Ni catalysts, forming the corresponding cross-coupling biaryl products (105) (Scheme 53).³⁶ This reaction involved an C-CN bond cleavage mediated by the special Cl₂Ni(PMe₃)₂ catalyst. The

Scheme 51 Ni-catalyzed decyanation of aryl nitriles with *tert*-butyl-magnesium chloride.

Scheme 52 Proposed mechanism for Ni-catalyzed decyanation of aryl nitriles with *tert*-butylmagnesium chloride.

Scheme 53 Ni-catalyzed cross coupling between nitrile and aryl Grignard reagent.

yield was significantly promoted by the pretreatment of Grignard reagent with an alkoxide (*t*-BuOLi in most cases), which efficiently inhibited the formation of byproducts, such as 4,4'-dimethoxybiphenyl from the homocoupling, anisole from the metal-H exchange, and 4-methoxybenzophenylimine from nucleophilic addition of Grignard reagent to the cyano. By applying the combination of PhMgBr and *t*-BuOLi, a variety of aryl nitriles were coupled to phenyl, including the aryls either with electron donating group (MeO, Me₂N, Me) or electron withdrawing group (F, COOR). Moreover, the aryl nitriles could be extended to heterocyclic, such as pyridinyl and furanyl. By using the proper lithium salt, all of three cyanopyridines, 2-, 3-, and 4-cyanopyridines, could couple to phenyl magnesium chloride.

Later in 2003, Miller and Dankwardt reported the nickel catalyzed cross-coupling of aryl nitriles with the modified alkyl and alkenyl Grignard reagents, also providing corresponding styrene and alkyl arene derivatives in good yields (Scheme 54).³⁷ By the addition of either *t*-BuOLi or PhSLi to the Grignard reagents, the nucleophilic addition of Grignard to aryl nitriles could be effectively diminished. Undoubtedly, this procedure contained a nickel catalyzed activation of the C–CN bond.

In 2009, Shi reported a cross coupling of aryl nitriles with aryl/alkenyl boronic esters through Ni-catalyzed C-CN bond cleavage (Scheme 55).³⁸ With the aid of t-BuOK as a base and CuF₂ as an additive, aryl/alkenyl boronic esters coupled with

R
$$\stackrel{\square}{\longrightarrow}$$
 + R'MgZ $\stackrel{\text{NiCl}_2(\text{PMe}_3)_2}{\longrightarrow}$ R $\stackrel{\square}{\longrightarrow}$ R $\stackrel{\square}{\longrightarrow}$ R' = alkyl or alkenyl up to 80% yield Z = Ot-Bu or SPh

Scheme 54 Ni-catalyzed cross coupling between nitrile and alkyl/alkenyl Grignard reagent.

Scheme 55 Ni-catalyzed cross coupling between nitrile and boronic ester.

aryl nitriles and provided biaryls in moderate to good yields. A variety of aryl nitriles were tested, including the aryls with electron-withdrawing and electron-donating groups.

Considering the relative reactivity of C–Cl, C–CN, and C–OMe bond activations in cross-coupling reactions, the authors successfully prepared polysubstituted benzene (111) with different aryl groups by orthogonal design (Scheme 56).

In 2012, Wang developed Ni-catalyzed cross coupling of aryl nitriles and arylmanganese reagents through the cleavage of C-CN bond (Scheme 57).³⁹ The arylmanganese reagents, which had been commonly used in many cross-coupling reactions, showed high efficiency in these cross-coupling reactions. A number of aryl nitriles and arylmanganese chlorides could be used for coupling. In comparison with the Grignard reagents or boronic esters used above, this reaction had great advantages, such as mild conditions, and no requirement of base and additive.

During the process of the rhodium catalyzed decyanative silylation of aryl cyanides as previously mentioned in Scheme 41,²⁸ Ar–Rh was formed as a key intermediate in the catalytic cycle. By designation of the structure of Ar–Rh with a tethered electrophile, such as aryl chloride, C–C bond could be formed intramolecularly. Thus, substituted dibenzofurans (113) and carbazole derivatives (114) were well synthesized by the Rh-catalyzed Si-assisted C–CN bond cleavage and a sequential coupling with intramolecular Ar–Cl (Scheme 58). Besides of these, the benzylic C–CN bond could be arylated with intramolecular Ar–Cl. Thus, fluorene (116) was obtained in 60% yield,

Scheme 56 Orthogonal synthesis of triarylbenzene based on relative reactivity.

Scheme 57 Ni-catalyzed cross coupling between nitrile and arylmanganese reagent.

Scheme 58 Rh-catalyzed Si-assisted cross coupling between C-CN and Ar-Cl

accompanying with the formation of the decyanated silylation product (117) in 15% yield. This methodology used the aryl chloride instead of organometallic reagent to realize the cross coupling between C-CN and C-Cl bonds. From this point of view, it was of synthetic value because aryl chlorides were more stable and more widely existed.

3.6 Decyanative alkenylation

In 2010, Tobisu and Chatani demonstrated a Rh-catalyzed Siassisted C-CN bond cleavage of aryl nitrile (20j), followed by a Mizoroki-Heck type reaction with triethyl(vinyl)silane (118), achieving the alkenylation of aryl nitrile (119) as well as the silylated byproduct (120) (Scheme 59).40 All the resulting olefins were in trans-configuration. After screening the reaction conditions, ligand was proved to be a significant factor in affecting the selectivity of forming the desire alkenylated product over silvlated product. As a result, P(4-FC₆H₄)₃ was found to be optimal. The ratio of alkenylated product to silylated product largely depended on the substituted group on the silicon. Triethylvinylsilane provided alkenylated product and silylated product in 65% and 18% yields, while triisopropylvinylsilane afforded them in 11% and 79%, respectively. On the contrary, triisopropoxylvinylsilane afforded the alkenylated product as the sole product, no silylated product was detected. 4-Cyano-1-tosylindole coupled with triethylvinylsilane under the same reaction conditions and provided 4-vinyl substituted-1-tosylindole in 82% yield without the contamination from C-H bond activation of 2- or 3-position of indole. 3,3-Diphenylacrylonitrile, as an example of alkenyl cyanides, could also couple with triethylvinylsilane to afford dienylsilane, accordingly.

Considering the selective C-Br and C-CN bond activation by using suitable transition metal catalyst, 1,4-divinyl phenylene

Scheme 59 Rh-catalyzed decyanative Mizoroki-Heck type coupling.

Scheme 60 Orthogonal synthesis of conjugated divinylphenylene.

(122), with different substituted groups at ends could be approached (Scheme 60).

In the proposed mechanism (Scheme 61), Rh catalyst transferred to an active silylrhodium species (**A**) by exclusion of TMSCl, and then Si-assisted C–CN bond cleavage occurred and arylrhodium species (**C**) was subsequently generated via η^2 -iminoacyl complex (**B**). During this process, silyl isocyanide was formed. Finally, a Mizoroki–Heck type reaction between intermediate (**C**) and vinylsilane occurred to form the desire product via addition and elimination, accompanied by the generation of rhodium hydride species (**E**). Subsequent reaction between rhodium hydride species (**E**) and disilane regenerated silylrhodium species (**A**) for the next catalytic cycle. In some cases, silylated product was obtained from the reaction between arylrhodium species (**C**) and disilane which was described in detail in the section of the coupling with heteroatoms.

Thus, a number of σ -bonds, including C–Si, C–P, C–B, C–H, and C–C bonds, could be feasibly constructed *via* transition metal catalyzed C–CN bond cleavage and the subsequent cross coupling. Of particular note, advantageous of R–CN were obvious. First, they could be treated as a succedaneum for R–X/R–OTs in cross coupling. Second, based on the selectivity and the reactivity of C–CN, C–X, and C–O bonds, these protocols enabled orthogonal cross coupling possible for having higher conjugated system. Third, the decyanation reaction took the advantage of "CN" properties, such as α -C–H acidity, *ortho*-directing effect, and electron-withdrawing nature, and could be utilized in the synthesis of challengeable compounds.

Scheme 61 Proposed mechanism for the Rh-catalyzed decyanative Mizoroki–Heck type coupling.

4. Cyanation reactions

The "CN" part in nitriles can be used as cyanide source in cyanation reactions. In contrast to extremely toxic metal cyanides (MCN), organic nitriles are less toxic, soluble in normal organic solvents, and easy to handle with operational benign. They slowly release the cyanide anions through C–CN bond activation and are safe cyanation reagents. Meanwhile, the nature of slow release of cyanide anions during the reaction process effectively avoids the overloading of cyanide anions to transition metal center resulting from the thermodynamically stable TM–CN bond. With this respect, minimum amount of the catalyst and cyanide precursor are used in cyanations. In cyanation reactions, there are several ways to cleave the C–CN bonds, those are Si-assisted oxidative addition of TM, TM-assisted nucleophilic substitution, and TM-catalyzed α -C–H oxidation.

4.1 C-CN cleavage via Si-assisted TM migration insertion

In 2005, an iron complex, $Cp(CO)_2$ FeMe was disclosed to catalyze the C–CN bond cleavage in the photoreaction of MeCN and Et_3 SiH. Reaction afforded Et_3 SiCN in 72% yield by extrusion of CH_4 with a trace amount of Et_3 Si–Si Et_3 formation (Scheme 62).⁴¹ Here Et_3 SiCN was thought to be thermodynamically stable product, isomerized from Et_3 SiNC. This catalytic system was also applicable for the cleavage of C–CN bond of aryl nitriles.

In their proposed mechanism (Scheme 63), light triggered the exclusion of CO from $Cp(CO)_2FeMe$ and left the space for oxidative addition to Si–H bond in Et_3SiH . After reductive elimination, a key intermediate (C), with 16 valence electrons, was generated. Coordinating this species with acetonitrile in η^2 -fashion (D), followed by silyl-assisted iron migration insertion, iron isocyanide complex (E) was formed. Exclusion of silyl isocyanide (Et_3SiNC) from E regenerated A for completing the

Et₃SiH + MeCN
$$\frac{\text{Cp(CO)}_2\text{FeMe (2 mol\%)}}{\text{THF, hv, r.t., 24 h}} \rightarrow \text{Et}_3\text{SiCN + CH}_4$$

Scheme 62 Fe-catalyzed Si-assisted photocyanation of silane with acetonitrile.

Scheme 63 Proposed mechanism for the Fe-catalyzed Si-assisted photocyanation of silane with acetonitrile.

catalytic cycle. As a result, Et_3SiCN was formed after isomerism, while $Et_3Si-SiEt_3$ was formed from the reaction between key intermediate (C) and Et_3SiH *via* addition/elimination sequence.

4.2 C-CN cleavage via TM-assisted nucleophilic substitution

In 2012, Wang and Lu reported palladium-catalyzed cyanation of aryl halides (123) using benzyl cyanide (42a) as the cyano source (Scheme 64).7a The C-CN bond of benzyl cyanide was cleaved in the aid of palladium and transferred to aryl halides, leading to the formation of aryl nitriles. In the optimized conditions, Pd(OAc)2 was the best catalyst, and it was quite necessary for having K2CO3 as the base and n-Bu4NBr as the additive. A number of functional groups, such as amino, hydroxy, alkyloxy, and carbonyl, tolerated this reaction conditions and provided the desired aryl nitriles in moderate to good yields. 1,4-Palladium migration was observed in the designed substrate. Treatment of 124 under the standard reaction conditions, a mixture of 125a and 125b was obtained in 46% yield. The ratio of 125a and 125b was determined to be 1:1. 125a was the key intermediate for the preparation of Satan's medicine. Of particular note, this protocol was realized as a safety method because only a trace amount of cyanide anion was detected which, moreover, effectively inhibited the poison of the palladium catalyst resulting from the high affinity of palladium to cyanide anion.

The general Pd-catalyzed cyanation of aryl halides contained three steps, oxidative addition, ligand exchange, and reductive elimination (Scheme 65). The key obstacle for this general one was the poison of the palladium catalyst which resulted from overloading of cyanide anion to palladium center. In the cyanation with benzyl cyanide, after the oxidative addition of Pd(0) to aryl halides, **B** was formed and subsequently coordinated with benzyl cyanide to generate key intermediate (C). Thus, C–CN bond was activated and cleaved by the assistance of halide. Finally, reductive elimination afforded aryl nitriles and regenerated Pd(0) for completing the catalytic cycle.

Subsequently, Shen developed palladium catalyzed cyanation of aryl halides, introducing ethyl cyanoacetate as the cyanating reagent (Scheme 66). The combination of Pd(OAc)₂, TMEDA, DPPE, Na₂CO₃ and KI could make the best transformation from aryl halides to aryl nitriles. A broad substrate

Scheme 64 Pd-catalyzed cyanation of aryl halide with benzyl cyanide.

Scheme 65 Mechanism for the Pd-catalyzed cyanation of aryl halide with benzyl cyanide.

Scheme 66 Pd-catalyzed cyanation of aryl halide with ethyl cyanoacetate.

scope was presented, including the acetyl, hydroxy, amino, ester, and amide. 5-Bromo-1-mthylindole proceeded the reaction affording the desired 5-cyano-1-mthylindole in 52% yield without the byproduct resulting from the C-H activation of 3position of indole. The major defect for this transformation was the competitive aromatic nucleophilic substitution of enolate anion on the highly electron-deficient aryl halides. It happened spontaneously in basic environment without the participation of palladium. The authors claimed the complex of the ethyl cyanoacetate enolate and the palladium was the key intermediate in this reaction. It could form the palladium-cyano species and facilitate the cleavage of C-CN bond.

In 2012, Zhou reported Cu-catalyzed cyanation of aryl iodides using malononitrile as the cyanide source (Scheme 67).43 After screening the reaction conditions, Cu(OAc)₂, 1,10-phenanthroline, t-BuONa were all the best choices, along with KF as a beneficial additive. The intermediate, a complex of [Cu(phen)(CN)₂] (126), was formed by proceeding the reaction in the absence of aryl iodides. It was indicated that [Cu(phen)(CN)₂] could react with the aryl iodides to form the aryl nitriles in good yields with or without KF. Thus, a copper-catalyzed C-CN bond cleavage of malononitrile assisted by KF might be involved in the reaction pathway, followed by the copper-catalyzed cyanation of aryl iodides with intermediate

Later, Shen subsequently demonstrated a copper-catalyzed cyanation of 2-phenylpyridines and the 1-pyridinylindoles

Scheme 67 Cu-catalyzed KF-assisted cyanation of aryl iodides with malononitrile.

with acetonitrile (Scheme 68).8c It was hypothesized by the authors that hexamethyldisilane was employed as a Lewis acid to weaken the C-CN bond for its facile cleavage. Thus, a S_N2type nucleophilic substitution occurred on the methyl of acetonitrile and resulted in the formation of copper cyanide species for the subsequent cyanation.

4.3 C-CN cleavage via TM-catalyzed α-C-H oxidation

In 2012, Wang and Lu developed Cu-catalyzed cyanations of 2phenylpyridinederivatives (131) with benzyl cyanides as the cyanation reagents (Scheme 69).7b A variety of substrates were proper for this transformation. A number of substituted groups tolerated the reaction conditions and provided desired

Scheme 68 Cu-catalyzed cyanation of arenes with acetonitrile using O₂

Scheme 69 Cu-catalyzed cyanation of arene via aerobic C-H oxidation of benzyl cyanide.

cyanation products in excellent yields. Not only the pyridinyl group can be used as the directing group for the C-H bond activation, but pyrazolyl, pyrimidinyl also.

After further investigation, a different pathway was illustrated (Scheme 70) in comparison with previously Pd-catalyzed pathway (Scheme 65). An aerobic C–H oxidation of benzyl cyanide in the catalyst system was supposed to produce benzaldehyde cyanohydrin which was responsible for slowly releasing cyano anions for the subsequent Cu-catalyzed cyanation of arenes. It was pointed that matching two Cu-catalyzed reactions rates, those were Cu-catalyzed C–H aerobic oxidation of benzyl cyanide and heteroatom-directed *ortho-*C–H functionalization, was the key issue for approaching a successful transformation.

Afterwards, benefitted from the advantageous of benzyl cyanide as cyanation reagent and inexpensive copper salts as catalysts, cyanations of aryl halides, ^{7c} aryl boronic acids, ^{7d} and indoles ^{7e,f} were realized with nice substrate diversity and high yields (Scheme 71). Of particular note, by using the standard reaction conditions for cyanation of aryl halides, cyanation of 124 provided 125a in 50% yield without the observation of 125b.

In 2012, a copper-catalyzed oxidative cyanation of aryl halides with acetonitrile as the cyanation reagent was reported

$$\begin{array}{c} \text{Ph} & \text{CN} \\ \text{Cu(I)} \\ \text{O}_2 \\ \text{OH} & \text{PhCHO} \\ \text{OH} & \text{CN} \\ \end{array} \begin{array}{c} \text{OCN} \\ \text{O}_2 \\ \text{OH} \\ \text{CN} \\ \end{array} \begin{array}{c} \text{OCN} \\ \text{OO} \\ \text{ON} \\ \text{CU(I)} \\ \text{OO} \\ \text{ON} \\ \text{CU(I)} \\ \text{OO} \\ \text$$

Scheme 70 Mechanism for Cu-catalyzed cyanation of arene with benzyl cyanide.

123

+ Ph CN
$$\frac{\text{Cu}_2\text{O} (1.0 \text{ equiv})}{\text{DMF, air, } 130 \, ^{\circ}\text{C, } 12 \text{ h}}$$

MeO

123

$$X = 1, 92\%$$

$$X = \text{Br, } 66\%$$

Cul (1.0 equiv)
$$X = \text{Br, } 66\%$$

Cul (1.0 equiv)
$$X = \text{Br, } 66\%$$

TBHP (aq. 2.0 equiv)
$$X = \text{DMAc, air, } 130 \, ^{\circ}\text{C, } 20 \text{ h}$$
Ph 133

134, 82%

Cul (1.2 equiv)
$$X = \text{DMF, air, } 100 \, ^{\circ}\text{C, } 34 \text{ h}$$

135

136 96%

Me

Scheme 71 Selected examples for cyanation of aryl halides, aryl boronic acids, and indoles with benzyl cyanide.

Scheme 72 Cu-catalyzed cyanation of aryl halides with acetonitrile.

by Li (Scheme 72). Sa They employed Ag₂O/air as the oxidant and triphenylphosphine oxide as ligand, converting various aryl iodides and bromides to aryl nitriles in moderate to good yields. Cu(1) species was thought to be responsible for the cleavage of Ar–X to form Ar–Cu species, followed by coordination with acetonitrile. In assistance of Ag₂O/O₂, acetonitrile was oxidized and decomposed to produce formic acid and "CN" for the following cyanation. Thus, the cleavage of C–CN of acetonitrile belonged to the Cu-catalyzed oxidative cleavage (Scheme 73).

In 2013, Zhu also introduced acetonitrile as the cyanide source to the copper-mediated C2-cyanation of indoles and pyrroles without the participation of phosphine ligand (Scheme 74). With the pyrimidyl or pyridinyl on the N atom as the directing group, cyanation occurred selectively at C2-position. AgOAc and the O_2 atmosphere were used as oxidants in this reaction, also involving copper-mediated C-CN cleavage of acetonitrile via α -C-H oxidation.

Throughout the cyanation reactions by using organic nitriles as "CN" surrogates, both cleavage and formation of C-CN bond were involved. Although organic sources had significantly advantageous over traditional used metal cyanides, such as

$$Ar - CN$$

$$Ar - CU^{||}L_nOAc$$

Scheme 73 Mechanism for the Cu-catalyzed cyanation of aryl halides with acetonitrile.

Scheme 74 Cu-catalyzed cyanation of arenes with acetonitrile.

RSC Advances Review

path a
$$R-C \equiv N$$
 $\xrightarrow{M/LA}$ R $C \equiv N - \dots LA$

path b $R-C \equiv N$ $\xrightarrow{M-SiR_3'}$ \xrightarrow{N} \xrightarrow{N}

Scheme 75 Representative pathways of transition metal catalyzed C-CN cleavage.

non-toxic and soluble in common organic solvents, only few organic nitriles had been hitherto examined and developed. Further investigation in this area for easy synthesis of aryl nitriles is still highly demanded.

Conclusions

In summary, C–C σ-bond cleavage, catalyzed by transition metals with high selectivity and high efficiency, has gained lots of interests in recent years for its great usage in constructing new chemical bond. However, the high BDE of C-CN makes the C-CN bond inert and brings a huge challenge to organic chemists. Based on the aforementioned research works, it is noticed that transition metal plays a key role in activating C-CN bonds and can be categorized into four principles (Scheme 75). Most of mechanisms involve an oxidative addition of low-valent TM to C-CN bond which is greatly facilitated by adding of Lewis acid (Scheme 75, path a). In some cases, insertion of cyano into metalsilyl bond forms η^2 -iminoacyl complex which subsequently undergoes de-insertion of silyl isocyanide and results in the cleavage of C-CN bond (Scheme 75, path b). In others, transition metals function as Lewis acid which weakens C-CN bond for its sequential nucleophilic substitution (Scheme 75, path c). Besides of these, by applying the α -H acidity of nitrile, transition metal catalyzes C-H oxidation and leads to the formation of unstable cyanohydrins which releases cyanide anion by retrocyanohydrination (Scheme 75, path d). Although some functions of transition metal have been clearly shown in C-CN bond activations, some of others still remain unclear, which drives researchers devoting more endeavours in mechanistic studies.

Although above mentioned cyanofunctionalizations, crosscoupling reactions, and cyanations have well presented the utilities of nitriles in organic synthesis in recent years, the huge potentials of using the abundant, stable, and low-cost R-CN in feasibly making various C-C, C-H, as well as C-heteroatom bonds in organic synthesis still need to be explored and developed.

Acknowledgements

The authors thank the National Nature Science Foundation of China (NSFC) (21032005, 21272203).

Notes and references

- 1 (a) A. Kleemann, J. Engel, B. Kutscher and D. Reichert, Pharmaceutical Substance: Synthesis, Patents, Applications, Georg Thieme, Stuttgart, 4th edn, 2001; (b) R. C. Larock, Comprehensive Organic Transformations: A Guide to Functional Group Preparations, Wiley-VCH, Weinheim, Germany, 1989, pp. 819-995.
- 2 (a) A. J. Fatiadi, Preparation and Synthetic Applications of Cyano Compounds, ed. S. Patai and Z. Rappoport, Wiley-VCH, New York, NY, 1983; (b) Z. Rappoport, Chemistry of the Cyano Group, John Wiley & Sons, London, UK, 1970, pp. 121-312.
- 3 For reviews, see: (a) G. P. Ellis and T. M. Romney-Alexander, Chem. Rev., 1987, 87, 779-794; (b) M. Sundermeier, A. Zapf and M. Beller, Eur. J. Inorg. Chem., 2003, 3513-3526; (c) P. Anbarasan, T. Schareina and M. Beller, Chem. Soc. Rev., 2011, **40**, 5049–5067; (d) J. Kim, H. J. Kim and S. Chang, Angew. Chem., Int. Ed., 2012, 51, 11948-11959; (e) G. Yan, J. Yu and L. Zhang, Chin. J. Org. Chem., 2012, 32, 294-303; (f) Q. Wen, J. Jin, L. Zhang, Y. Luo, P. Lu and Y. Wang, Tetrahedron Lett., 2014, 55, 1271-1280.
- 4 For recent reviews, see: (a) M. Tobisub and N. Chatani, Chem. Soc. Rev., 2008, 37, 300–307; (b) C. Nájera and J. M. Sansano, Angew. Chem., Int. Ed., 2009, 48, 2452-2456; (c) X. Kou, J. Fan, X. Tong and Z. Shen, Chin. J. Org. Chem., 2013, 33, 1407–1422; (d) R. Wang and J. R. Falck, RSC Adv., 2014, 4, 1062-1066.
- 5 A. A. Zavitsas, *I. Phys. Chem. A*, 2003, **107**, 897–898.
- 6 CRC Handbook of Chemistry and Physics, ed. R. D. Lide, CRC Press, Boca Raton, FL, 82nd edn, 2001, pp. 9-74.
- 7 (a) Q. Wen, J. Jin, B. Hu, P. Lu and Y. Wang, RSC Adv., 2012, 2, 6167-6169; (b) J. Jin, Q. Wen, P. Lu and Y. Wang, Chem. Commun., 2012, 48, 9933-9935; (c) Q. Wen, J. Jin, Y. Mei, P. Lu and Y. Wang, Eur. J. Org. Chem., 2013, 4032-4036; (d) Y. Luo, Q. Wen, Z. Wu, J. Jin, P. Lu and Y. Wang, Tetrahedron, 2013, 69, 8400-8404; (e) L. Zhang, Q. Wen, J. Jin, C. Wang, P. Lu and Y. Wang, Tetrahedron, 2013, 69, 4236-4240; (f) O. Y. Yuen, P. Y. Choy, W. K. Chow, W. T. Wong and F. Y. Kwong, J. Org. Chem., 2013, 78, 3374-3378.
- 8 (a) R.-J. Song, J.-C. Wu, Y. Liu, G.-B. Deng, C.-Y. Wu, W.-T. Wei and J.-H. Li, Synlett, 2012, 23, 2491-2496; (b) C. Pan, H. Jin, P. Xu, X. Liu, Y. Cheng and C. Zhu, J. Org. Chem., 2013, 78, 9494-9498; (c) X. Kou, M. Zhao, X. Qiao, Y. Zhu, X. Tong and Z. Shen, Chem.-Eur. J., 2013, 19, 16880-16886.
- 9 (a) K. Nozaki, N. Sato and H. Takaya, J. Org. Chem., 1994, 59, 2679-2681; (b) K. Nozaki, N. Sato and H. Takaya, Bull. Chem. Soc. Jpn., 1996, 59, 1629-1637.
- 10 (a) Y. Nishihara, Y. Inoue, M. Itazaki and K. Takagi, Org. Lett., 2005, 7, 2639-2641; (b) Y. Nishihara, Y. Inoue, S. Izawa, M. Miyasaka, K. Tanemura, K. Nakajimab and K. Takagi, Tetrahedron, 2006, 62, 9872-9882.
- 11 (a) Y. Nakao, Y. Hirata and T. Hiyama, J. Am. Chem. Soc., 2006, 128, 7420-7421; (b) Y. Hirata, T. Inui, Y. Nakao and T. Hiyama, J. Am. Chem. Soc., 2009, 131, 6624-6631.

RSC Advances Review

- 12 N. R. Rondla, S. M. Levi, J. M. Ryss, R. A. V. Berg and C. J. Douglas, Org. Lett., 2011, 13, 1940-1943.
- 13 Y. Kobayashi, H. Kamisaki, R. Yanada and Y. Takemoto, Org. Lett., 2006, 8, 2711-2713.
- 14 Y. Yasui, H. Kamisaki and Y. Takemoto, Org. Lett., 2008, 10, 3303-3306.
- 15 (a) Y. Nakao, S. Od and T. Hiyama, J. Am. Chem. Soc., 2004, 126, 13904-13905; (b) Y. Nakao and T. Hiyama, Pure Appl. Chem., 2008, 80, 1097-1107.
- 16 Y. Nakao, T. Yukawa, Y. Hirata, S. Oda, J. Sato and T. Hiyama, J. Am. Chem. Soc., 2006, 128, 7116-7117.
- 17 (a) Y. Nakao, A. Yada, S. Ebata and T. Hiyama, J. Am. Chem. Soc., 2007, 129, 2428-2429; (b) Y. Hirata, A. Yada, E. Morita, Y. Nakao, T. Hiyama, M. Ohashi and S. Ogoshi, J. Am. Chem. Soc., 2010, 132, 10070-10077; (c) Y. Yasui, H. Kamisaki, T. Ishida and Y. Takemoto, Tetrahedron, 2010, 66, 1980-1989.
- 18 Y. Nakao, K. S. Kanyiva, S. Oda and T. Hiyama, J. Am. Chem. Soc., 2006, 128, 8146-8147.
- 19 A. Yada, T. Yukawa, Y. Nakao and T. Hiyama, Chem. Commun., 2009, 3931-3933.
- 20 Y. Nakao, A. Yada and T. Hiyama, J. Am. Chem. Soc., 2010, 132, 10024-10026.
- 21 Y. Nakao, Y. Hirata, M. Tanaka and T. Hiyama, Angew. Chem., Int. Ed., 2008, 47, 385-387.
- 22 K. Nakai, T. Kurahashi and S. Matsubara, J. Am. Chem. Soc., 2011, 133, 11066-11068.
- 23 For a further study on this topic, see: K. Nakai, T. Kurahashi and S. Matsubara, Org. Lett., 2013, 15, 856-859.
- 24 Y. Minami, H. Yoshiyasu, Y. Nakao and T. Hiyama, Angew. Chem., Int. Ed., 2013, 52, 883-887.
- 25 M. P. Watson and E. N. Jacobsen, J. Am. Chem. Soc., 2008, 130, 12594-12595.
- 26 Y. Nakao, S. Ebata, A. Yada, T. Hiyama, M. Ikawa and S. Ogoshi, I. Am. Chem. Soc., 2008, 130, 12874-12875.
- 27 (a) M. Tobisu, Y. Kita and N. Chatani, J. Am. Chem. Soc., 2006, 128, 8152-8153; (b) F. L. Taw, P. S. White, R. G. Bergman and M. Brookhart, J. Am. Chem. Soc., 2002, 124, 4192-4193; (c)

- F. L. Taw, A. H. Mueller, R. G. Bergman and M. Brookhart, I. Am. Chem. Soc., 2003, 125, 9808-9813.
- 28 M. Tobisu, Y. Kita, Y. Ano and N. Chatani, J. Am. Chem. Soc., 2008, 130, 15982-15989.
- 29 M. Tobisu, J. Hasegawa, Y. Kita, H. Kinutab and N. Chatani, Chem. Commun., 2012, 48, 11437-11439.
- 30 M. Sun, H.-Y. Zhang, Q. Han, K. Yang and S.-D. Yang, Chem.-Eur. J., 2011, 17, 9566-9570.
- 31 (a) M. Tobisu, H. Kinuta, Y. Kita, E. Rémond and N. Chatani, J. Am. Chem. Soc., 2012, 134, 115-118; (b) H. Kinuta, Y. Kita, E. Rémond, M. Tobisu and N. Chatani, Synthesis, 2012, 44, 2999-3002; (c) Y.-Y. Jiang, H.-Z. Yu and Y. Fu, Organometallics, 2013, 32, 926-936; (d) H. Kinuta, H. Takahashi, M. Tobisu, S. Mori and N. Chatani, Bull. Chem. Soc. Ipn., 2014, 87, 655-669.
- 32 (a) M. Tobisu, R. Nakamura, Y. Kita and N. Chatani, J. Am. Chem. Soc., 2009, 131, 3174-3175; (b) M. Tobisu, R. Nakamura, Y. Kita and N. Chatani, Bull. Korean Chem. Soc., 2010, 31, 582-587.
- 33 T. Patra, S. Agasti, Akanksha and D. Maiti, Chem. Commun., 2013, 49, 69-71.
- 34 T. Patra, S. Agasti, A. Modak and D. Maiti, Chem. Commun., 2013, 49, 8362-8364.
- 35 M. Weidauer, C. I. Someya, E. Irran and S. Enthaler, Asian J. Org. Chem., 2013, 2, 150-156.
- 36 J. A. Miller, Tetrahedron Lett., 2001, 42, 6991-6993.
- 37 J. A. Miller and J. W. Dankwardt, Tetrahedron Lett., 2003, 44, 1907-1910.
- 38 D.-G. Yu, M. Yu, B.-T. Guan, B.-J. Li, Y. Zheng, Z.-H. Wu and Z.-J. Shi, Org. Lett., 2009, 11, 3374-3377.
- 39 N. Liu and Z. Wang, Adv. Synth. Catal., 2012, 354, 1641–1645.
- 40 Y. Kita, M. Tobisu and N. Chatani, Org. Lett., 2010, 12, 1864-1867.
- 41 H. Nakazawa, K. Kamata and M. Itazaki, Chem. Commun., 2005, 4004-4006.
- 42 S. Zheng, C. Yu and Z. Shen, Org. Lett., 2012, 14, 3644-3647.
- 43 Z. Jiang, Q. Huang, S. Chen, L. Long and X. Zhou, Adv. Synth. Catal., 2012, 354, 589-592.