





Showcasing research from Professor Martin Kirk's and Professor David Shultz's laboratories, Department of Chemistry and Chemical Biology, The University of New Mexico, Albuquerque, New Mexico, USA and Department of Chemistry, North Carolina State University, Raleigh, North Carolina, USA.

Transferrable property relationships between magnetic exchange coupling and molecular conductance

Computed molecular conductances mediated by organic σ/π -bridges are compared to experimentally-determined magnetic exchange coupling parameters in a series of Donor-Bridge-Acceptor Biradical complexes at parity of the bridge fragment. For monomeric and dimeric bridges, biradical magnetic exchange coupling and computed conductance are related by a power law with an observed scale invariance. We also find that the ratio of distance dependent decay constants for bridge-mediated conductance and magnetic exchange coupling does not equal unity. The results of our observations have been cast in valence bond terms.

As featured in:



See Martin L. Kirk,
David A. Shultz *et al.*,
Chem. Sci., 2020, 11, 11425.

Cite this: *Chem. Sci.*, 2020, **11**, 11425

All publication charges for this article have been paid for by the Royal Society of Chemistry

Transferrable property relationships between magnetic exchange coupling and molecular conductance†

Martin L. Kirk,^{ab} Ranjana Dangl,^a Diana Habel-Rodriguez,^a Jing Yang,^a David A. Shultz^{*c} and Jinyuan Zhang^{†c}

Calculated conductance through $Au_n-S-Bridge-S-Au_n$ (Bridge = organic σ/π -system) constructs are compared to experimentally-determined magnetic exchange coupling parameters in a series of $Tp^{Cum,Me}ZnSQ-Bridge-NN$ complexes, where $Tp^{Cum,Me}$ = hydro-tris(3-cumenyl-1-methylpyrazolyl) borate ancillary ligand, Zn = diamagnetic zinc(II), SQ = semiquinone ($S = 1/2$), and NN = nitronyl nitroxide radical ($S = 1/2$). We find that there is a nonlinear functional relationship between the biradical magnetic exchange coupling, $J_{D \rightarrow A}$, and the computed conductance, g_{mb} . Although different bridge types (monomer vs. dimer) do not lie on the same $J_{D \rightarrow A}$ vs. g_{mb} curve, there is a scale invariance between the monomeric and dimeric bridges which shows that the two data sets are related by a proportionate scaling of $J_{D \rightarrow A}$. For exchange and conductance mediated by a given bridge fragment, we find that the ratio of distance dependent decay constants for conductance (β_g) and magnetic exchange coupling (β_J) does not equal unity, indicating that inherent differences in the tunneling energy gaps, $\Delta\epsilon$, and the bridge-bridge electronic coupling, H_{BB} , are not directly transferrable properties as they relate to exchange and conductance. The results of these observations are described in valence bond terms, with resonance structure contributions to the ground state bridge wavefunction being different for SQ-Bridge-NN and $Au_n-S-Bridge-S-Au_n$ systems.

Received 7th August 2020
Accepted 14th September 2020

DOI: 10.1039/d0sc04350h

rsc.li/chemical-science

Introduction

Electron transport in single-molecule devices is typically interrogated *via* a combination of experimental and computational probes of conductance using Metal-Bridge-Metal (M-B-M) junctions (Fig. 1A).^{1–14} To account for inherent variations in molecule-electrode binding geometries, the experimental determination of conductance (g_{mb}) typically requires hundreds to thousands of individual conductance measurements in order to construct conductance histograms, which allow researchers

to determine both conductance distributions and the most probable conductance value.¹⁵ Since conductance calculations often utilize a single M-B-M geometry and the molecular geometry in the actual junction is unknown, this can make direct comparisons between theory and experiment difficult to interpret. As a result, studies that correlate single-molecule conductance with physical observables such as electron transfer rate constants ($k_{D \rightarrow A}$)^{2,4,5,14,16} and magnetic exchange couplings




Fig. 1 (A) Metal-Bridge-Metal junctions corresponding to biradical complexes (B). (C) Bridges common to M-B-M junctions and Donor-Bridge-Acceptor complexes (B), along with ferromagnetic exchange parameters, $J_{D \rightarrow A}$ for biradical complexes (C). BCO = bicyclo[2.2.2] octane, Ph = *para*-phenylene, Th = 2,5-thiophene, Py = pyridine, PyTh = 2-pyridyl-5'-thiophene, Ph₂ = *para*-biphenylene, and Th₂ = 2,5'-bithiophene.

^aDepartment of Chemistry and Chemical Biology, The University of New Mexico, MSC03 2060, 1 University of New Mexico, Albuquerque, New Mexico 87131-0001, USA. E-mail: mkirk@unm.edu

^bCenter for High Technology Materials, The University of New Mexico, Albuquerque, New Mexico 87106, USA

^cDepartment of Chemistry, North Carolina State University, Raleigh, North Carolina 27695-8204, USA. E-mail: shultz@ncsu.edu

† Electronic supplementary information (ESI) available: Computed phenylene *I-V* plot and MPSH HOMO, synthesis, characterization, crystal structure, and magnetic data for SQ-Py-NN. CCDC 2022006 and 2022007. For ESI and crystallographic data in CIF or other electronic format see DOI: 10.1039/d0sc04350h

‡ Current address: Beijing National Laboratory for Molecular Sciences, CAS Key Laboratory of Organic Solids, Institute of Chemistry, Chinese Academy of Sciences, Beijing 100190, China.



$(J_{D \rightarrow A})^{1,17-20}$ continue to be a topic of current research. A primary impact of these efforts will be to provide key insight into orbital pathways for charge transport through molecules, which is difficult to achieve using traditional approaches. These correlations, coupled with the determination of transferable property relationships, are expected to facilitate improved molecular design concepts for molecular electronics and spintronics.

The crux of these transferable property relationships was first described by Nitzan,²¹ who derived a relationship between g_{mb} and $k_{D \rightarrow A}$ at parity of the molecular bridge. With some approximations, Nitzan was able to show that conductance could be related to the square of the D–A electronic coupling matrix element, $H_{D \rightarrow A}^2$ ($H_{D \rightarrow A}^2 = |H_{DB}H_{BA}|^2 |G_B(E)|^2$), for electron transfer (eqn (1)–(4)).²¹

$$g_{mb} \approx \frac{8e^2}{\pi^2 \Gamma_D^{(L)} \Gamma_A^{(R)} \text{FCWDS}} k_{D \rightarrow A} \quad (1)$$

$$g_{mb} = \frac{e^2}{\pi \hbar} \Gamma_{LB} \Gamma_{RB} |\overline{G_B}|^2 \quad (2)$$

$$k_{D \rightarrow A} = \frac{2\pi}{\hbar} |H_{DB}H_{BA}|^2 |G_B(E)|^2 \text{FCWDS} \quad (3)$$

$$G_B = \frac{H_{BB}^{n-1}}{\Delta \varepsilon^n} \quad (4)$$

here, the H_{ij} are pairwise electronic coupling matrix elements that connect D, B, and A, the Γ_i are broadening functions that represent the coupling between the molecule and electrodes, $\Delta \varepsilon$ is the tunneling energy gap, G_B is the bridge Green's function with the overbar indicating that G_B may differ for electron transfer (or magnetic exchange) compared to conductance, and n is the number of bridge units. These transferable property relationships indicate that any physical observable that is a function of $H_{D \rightarrow A}^2$ can be related to g_{mb} (e.g. eqn (5)).

$$J_{D \rightarrow A} \propto H_{D \rightarrow A}^2 \propto k_{D \rightarrow A} \propto g_{mb} \quad (5)$$

This leads to a corresponding relationship between these parameters and the distance decay constants (β_i) for electron transfer, conductance, and magnetic exchange as indicated in eqn (6),

$$k_{D \rightarrow A} \propto g_{mb} \propto J_{D \rightarrow A} \propto \exp(-\beta_i \times nL) \quad (6)$$

where nL is the distance spanned by the molecular bridge (e.g. $R_{DA} - R_0$).

There is both experimental and computational support for a correlation between $k_{D \rightarrow A}$, g_{mb} and $J_{D \rightarrow A}$.^{14,17,22–31} Beratan and Waldeck^{4,5} have built on Nitzan's work to show that the relationship between $k_{D \rightarrow A}$ and g_{mb} is nonlinear for alkane and peptide nucleic acid oligomer bridges, but they are related by a power law dependence. Their work supports an argument that the bridge Green's function, G_B , is not the same for electron transfer and conductance. As a result, Beratan and Waldeck^{4,5} were able to show that the distance decay constants for $k_{D \rightarrow A}$ and g_{mb} are, in general, inequivalent and a linear relationship

between $k_{D \rightarrow A}$ and g_{mb} will only occur if the individual β_i are identical.

Since the magnetic exchange interaction ($J_{D \rightarrow A}$) is proportional to $H_{D \rightarrow A}^2$ via eqn (5),^{32–36} we have initiated experiments designed to explore the correlation of $J_{D \rightarrow A}$ between donor–acceptor biradical centers that are spanned by molecular bridge units (D–B–A biradicals = SQ–Bridge–NN; SQ = $S = 1/2$ semiquinone, and NN = $S = 1/2$ nitronyl nitroxide radical, Fig. 1B and C)^{1,17–20,26,35–40} with computed molecular conductance values, g_{mb} .¹ An advantage to understanding the magnitude of g_{mb} by correlating with experimental $J_{D \rightarrow A}$ parameters is highlighted by the fact that the variable-temperature magnetic susceptibility measurements used to determine $J_{D \rightarrow A}$ are performed on solid-state samples with known geometries that have been determined by X-ray crystallography. Thus, there is a single conformation present that contributes to $J_{D \rightarrow A}$. Unlike experimental conductance measurements in a contact geometry and solution electron transfer rate measurements, the magnitude of $J_{D \rightarrow A}$ determined by solid state magnetic susceptibility measurements is not affected by structural and solvent bath induced inhomogeneities or by decoherence effects.

Relating molecular orbital pathways for $J_{D \rightarrow A}$ to orbital transport channels for g_{mb} has its own set of challenges, and these include (1) differences in the molecular fragments covalently attached to the bridge unit, (2) the molecular identity of surfactant groups, (3) variations in surface contacts, (4) the choice of electrode material and surface morphology, and (5) inhomogeneities in molecular structure (e.g., conformation) and in contact geometries. As alluded to above, an important related question was recently posited by Herrmann,¹⁴ which concerns the transferability of geometric and electronic structure contributions of the bridge moiety to $k_{D \rightarrow A}$, g_{mb} , and $J_{D \rightarrow A}$ (Fig. 2). Herein, we address these difficulties and the concept of bridge electronic structure transferability between conductance and exchange. We then use our results to explicitly determine distance decay constants (β_g) for g_{mb} from prior studies that have determined the distance dependence of $J_{D \rightarrow A}$. Our results are described in valence bond terms, with resonance structure contributions to the ground state bridge wavefunction being different for SQ–Bridge–NN and Au_n –S–Bridge–S– Au_n systems.

Bridge (B) transferability in electron transport and magnetic exchange




Fig. 2 Illustration of how bridge-mediated electronic coupling is related to conductance and magnetic exchange coupling. Top: Molecular bridge (B) connects metallic electrodes to give a Metal–Bridge–Metal device. Bottom: Molecular bridge (B) connects spin centers to facilitate magnetic exchange coupling.



This provides a path forward for understanding the power law relationship between g_{mb} and $J_{\text{D} \rightarrow \text{A}}$ in the absence of structural inhomogeneities and decoherence effects.

Experimental

General

The synthesis and characterization of **SQ-BCO-NN**, **SQ-Ph-NN**, **SQ-Th-NN**, **SQ-Ph₂-NN**, **SQ-Py-Th-NN**, and **SQ-Th₂-NN** have been published previously.^{1,17,20,26,35,39,41,42} See the ESI† for the preparation and characterization (spectroscopic data, X-ray crystal structure, and magnetic characterization) of the pyridyl-bridged biradical complex, **SQ-Py-NN**.

Resonance Raman spectroscopy

Solution (methylene chloride) resonance Raman spectra were collected in either a 180° backscattering (780 nm) or 90° scattering geometry using 407 nm excitation from a Coherent Innova 70 Kr⁺ (1 W) ion laser. The scattered radiation was passed through a longpass filter (Semrock RazorEdge) to remove Rayleigh scattered laser light and then dispersed onto a liquid nitrogen-cooled Infrared Associates CCD detector using a Princeton Acton spectrograph. The laser power at the sample was typically kept between 40 and 100 mW in order to prevent possible photo- and thermal degradation of the sample. The sample was sealed in a glass capillary tube and spun with a custom-made sample holder. All data were scan-averaged, and any individual data set with vibrational bands compromised by cosmic events were discarded.

Molecular electronic structure calculations

Thiophene and benzene calculations were performed at the density functional level of theory using the Gaussian 09W software package.⁴³ Input files were prepared using the molecular builder function in GaussView. Calculations employed the B3LYP hybrid exchange–correlation functional and a 6-31g(d',p') split valence basis set with polarizability functions was used for all atoms. Frontier molecular orbitals (MOs) were generated from fully optimized ground states.

Transport calculations

Transport calculations were performed using ATK 2016.0 v16.0,^{44–46} as detailed in our earlier work.¹ The device configuration consists of the left Au electrode, the molecular bridge (scattering region), and the right Au electrode. Both the left and right gold electrodes consist of nine layers with each layer comprised of a 3 × 3 array of Au(111) atoms, for a total of 81 gold atoms per electrode. For the electrodes, the C or z direction is periodic in the system. This is also the direction of electron transport. Initial geometry optimizations of the bridge molecules **BCO**, **Py**, **Ph**, **Th**, **Ph₂**, **Py-Th**, **Th₂** were performed using Gaussian 09 (ref. 43) with a 6-31g(d',p') basis set and the B3LYP functional. Electrode surfaces were constructed by cleaving the bulk crystal and using a copy of this to form the second electrode. The molecule was then placed on the surface of the left electrode and subsequently connected to the right electrode

through terminal sulfur atoms to create the final device geometry. Prior to creating the final device geometry, the bulk configuration was optimized using a single-zeta basis set for the gold atoms (to save computational time) and a double zeta basis set for all other atoms. The optimization was performed using ATK-DFT with a Perdew–Zunger local density approximation (LDA-PZ) exchange–correlation. After this bulk optimization, the configuration was converted into a final device geometry for all subsequent calculations. A 5 × 5 × 51 K-point sampling has been used in x, y, z-directions, respectively. The boundary conditions were Dirichlet (fixed boundary condition) in the z direction, which is the direction of transport. The boundary conditions in the x and y directions were disabled so they could not be changed but the default parameterization is periodic for the x and y directions. The NEGF formalism was used to calculate the non-equilibrium electron density of the central region of the device.

The Landauer–Büttiker formula⁴⁷ relates transmission probabilities to conductance, g_{mb} ($g_{\text{mb}} = I/V$). This formula is used to calculate the voltage dependent current, $I(V)$, across a molecular junction, which is determined by integrating the transmission function, $T(E,V)$, according to eqn (7).⁴⁷

$$I(V) = \frac{e}{h} \sum_{\sigma} \int T_{\sigma}(E) \left[f \left(\frac{E - \mu_{\text{R}}}{k_{\text{B}} T_{\text{R}}} \right) - f \left(\frac{E - \mu_{\text{L}}}{k_{\text{B}} T_{\text{L}}} \right) \right] dE \quad (7)$$

here, the Fermi–Dirac distribution functions for the right and left electrodes are given by $f(E - \mu_{\text{R}})/k_{\text{B}} T_{\text{R}}$ and $f(E - \mu_{\text{L}})/k_{\text{B}} T_{\text{L}}$, with μ_{R} and μ_{L} being the chemical potentials for the right and left electrodes, respectively, e is the charge of the electron, h is Planck's constant, eV_{bias} is the bias window equivalent to $\mu_{\text{R}} - \mu_{\text{L}}$ (where $V_{\text{bias}} = V_{\text{L}} - V_{\text{R}}$), $T_{\sigma}(E)$ is the transmission coefficient for the spin component, σ , describing the junction at an energy E and a bias voltage V_{bias} . All electron transport properties were computed using the ATK software package that includes virtual nanolab associated analysis modules. The molecular projected self-consistent Hamiltonian (MPSH) technique has been used to understand the molecular orbital origin of the resonant peaks in the transmission spectra.^{48,49}

Results and discussion

Exchange coupling in SQ–Bridge–NN complexes

Magnetic exchange coupling constants ($J_{\text{D} \rightarrow \text{A}}$) for the monomeric SQ–Bridge–NN biradical complexes (Fig. 1B and C) **SQ-BCO-NN**, **SQ-Ph-NN**, **SQ-Th-NN**,^{17,20,26,41} and for the dimeric bridge molecules **SQ-Py-Th-NN**,¹ **SQ-Ph₂-NN**,¹⁷ and **SQ-Th₂-NN**¹⁷ have previously been determined by magnetic susceptibility measurements. The dimer exchange spin Hamiltonian (eqn (8)):

$$\mathcal{H}_{\text{ex}} = -2J_{\text{D} \rightarrow \text{A}} S_{\text{SQ}} S_{\text{NN}} \quad (8)$$

has been used in the analysis of these data, including the analysis of new magnetic susceptibility data for the pyridyl-bridged complex (**SQ-Py-NN**), which yields $J_{\text{D} \rightarrow \text{A}} = +72 \text{ cm}^{-1}$ (see ESI†). Collectively, these SQ–Bridge–NN biradical systems possess a large range of $J_{\text{D} \rightarrow \text{A}}$ -values, bridge-dependent electronic structures, and both σ - and π -coupling units. Notably, all



of these molecules are *ferromagnetically* exchange coupled^{20,35,36} with $J_{D \rightarrow A}$ values that vary from a low of $J_{D \rightarrow A} = +1 \text{ cm}^{-1}$ for σ -mediated exchange in **SQ-BCO-NN**, to a markedly larger $J_{D \rightarrow A} = +220 \text{ cm}^{-1}$ for π -mediated exchange in **SQ-Th-NN**.

Exchange coupling within the context of a valence bond configuration interaction model

Given the geometric structures of these SQ-Bridge-NN biradicals, we can conveniently divide them into two distinct bridge categories: the first possesses single-ring molecular bridges, while the second contains two-ring molecular bridges where each bridge ring is connected to the other *via* a σ -bond. The electronic origins of π -mediated exchange coupling in **SQ-Ph-NN** and σ -mediated exchange in **SQ-BCO-NN** have been rigorously evaluated using a combination of spectroscopy, magnetic susceptibility, and theory.^{19,20,26,39,41,50} For π -mediated exchange, a valence-bond configuration interaction (VBCI) model has been used.^{19,20,35,36} Within the context of more traditional orbital pathways for understanding exchange coupling, the VBCI method conveniently illustrates how specific excited state configurations admix with the ground configuration to affect the magnitude of $J_{D \rightarrow A}$. The NN(SOMO), SQ(SOMO), NN-Bridge(HOMO), and the NN-Bridge(LUMO) (Fig. 3) orbitals form a convenient minimal active space to generate the ground and excited state configurations that contribute to $J_{D \rightarrow A}$ for *conjugated* SQ-Bridge-NN systems. Previous work has demonstrated the crucial role of the LUMO(B-NN) in determining the magnitude of $J_{D \rightarrow A}$.^{17,19,20,35,36,39} Resonance structures that illustrate both the spin and the charge distributions of ground state **SQ-Ph₂-NN** configurations and excited state configurations that arise from one-electron promotions within this active space are shown in Fig. 4. Here, the ground SQ-Bridge-NN configuration is represented by GC, EC1 and EC2 represent the configuration that results from an intraligand SQ(SOMO) $\rightarrow$ NN-Bridge(LUMO) electron promotion, and the NN-Bridge(HOMO) $\rightarrow$ SQ(SOMO) one-electron promotion is represented by the EC3 and EC4 resonance structures. Similar resonance structures can be




Fig. 3 Simple 4-orbital diagram showing NN-Bridge(HOMO) $\rightarrow$ SQ(SOMO) and SQ(SOMO) $\rightarrow$ NN-Bridge(LUMO) one-electron promotions that figure prominently in charge transfer configurations that contribute to magnetic exchange in SQ-Bridge-NN biradicals.




Fig. 4 Resonance structures illustrating enhanced H_{BB} , H_{DB} and H_{BA} couplings in D-B-A biradicals. The corresponding zwitterionic and biradical resonance structures for M-B-M couplings (H_{BB} , I_{LB} , I_{RB}) are higher in energy and contribute less to the conductance. All SQ-B-NN ligands have a net -1 charge. Donor and acceptor moieties are color-coded according to oxidation state: green = 9 π -electron $\text{SQ}^{1-}/7$ π -electron NN; red = 10 π -electron catecholate²⁻/8 π -electron NN^{1-} ; blue = 8 π -electron quinone⁰/6 π -electron NN^{1+} .

drawn for other SQ-Bridge-NN biradical complexes. For the *conjugated* SQ-Bridge-NN systems discussed here, the charge transfer configuration generated by the intraligand SQ(SOMO) $\rightarrow$ NN-Bridge(LUMO) one-electron promotion (EC1 and EC2) has been shown to be the dominant contributor to the ground state *ferromagnetic* exchange coupling in **SQ-Ph-NN**, with the NN-Bridge(HOMO) $\rightarrow$ SQ(SOMO) (EC3 and EC4) playing a less important role.^{17,19,20,35,36,39} The relative contributions of these two intraligand charge transfer (ILCT) configurations will be dependent on the nature of the bridge. The use of more electron donating bridge molecules (*e.g.*, thiophene) is expected to increase the relative NN-Bridge(HOMO) $\rightarrow$ SQ(SOMO) charge transfer contribution to the magnetic exchange by raising the energy of the bridge HOMO and LUMO orbitals. In addition, we have shown that bond torsions which disrupt Ph-NN π -coupling result in the ILCT transition being shifted to higher energy, with a concomitant reduction in the ILCT intensity relative to configurations that possess more planar Ph-NN conformations.⁴¹ As a result, an ILCT transition is *not* observed in **SQ-BCO-NN** and the lack of a CT feature in the visible region of the electronic absorption spectrum correlates with the weak σ -mediated exchange coupling.^{26,41} Our understanding of how the bridge HOMO and LUMO orbitals affect the magnitude of both $J_{D \rightarrow A}$ and $H_{D \rightarrow A}$ allow us to begin an assessment of how

Open Access Article. Published on 08 gollgotmännu 2020. Downloaded on 2025-08-11 14:03:50.
This article is licensed under a Creative Commons Attribution 3.0 Unported Licence.




Fig. 6 Computed current ($g_{\text{mb}} = I/V$) vs. exchange coupling ($J_{\text{D} \rightarrow \text{A}}$) for monomeric molecular bridges (red circles) and for dimeric molecular bridges (blue circles). The data indicate that current is not directly proportional to magnetic exchange coupling. The data are fit to separate empirical power law functions with identical exponents, with best fits shown as black lines. Computations were performed using a bias voltage of +2 V which approximates the intraligand charge transfer energy in SQ–Bridge–NN complexes.

the two data sets are related by a proportionate scaling of the exchange interaction, $J_{\text{D} \rightarrow \text{A}}$. As a result, we find that $J_{\text{D} \rightarrow \text{A}}$ and g_{mb} are indeed highly correlated amongst this bridge set, but the simple proportionality relationship, $J_{\text{D} \rightarrow \text{A}} \propto H_{\text{D} \rightarrow \text{A}}^2 \propto g_{\text{mb}}$, is not generally valid across a series of monomeric or dimeric bridge molecules.

Fig. 7 shows the computed current as a function of magnetic exchange coupling for thiophene and phenylene bridges (0–2 bridge units). Following the formalism of Beratan and Waldeck,⁵ these data have been fit to eqn (12),

$$g_{\text{mb}} = [g_{\text{L}=0}(J_{\text{D} \rightarrow \text{A}}^0)^{-\beta_{\text{g}}/\beta_{\text{J}}}] (J_{\text{D} \rightarrow \text{A}})^{\beta_{\text{g}}/\beta_{\text{J}}} \quad (12)$$




Fig. 7 Computed current (conductance) vs. exchange coupling for phenylene bridges (red circles) and for thiophene bridges (blue circles). Best fits to the Eq. in the text are shown as red and blue lines, respectively. Note that the value of the exponents, $m = \beta_{\text{g}}/\beta_{\text{J}}$, lead to a greater degree of linearity for the phenylene data set (see text). The black circle is located at (0,0), since $g_{\text{mb}} \rightarrow 0$ as $J_{\text{D} \rightarrow \text{A}} \rightarrow 0$. Conductance values for M–B–M systems were computed at +2 V bias using the Landauer–Büttiker formalism, which relates transmission probabilities to conductance ($g_{\text{mb}} = I/V$).

in order to understand the relative ratios of distance dependent decay constants for conductance (β_{g}) and magnetic exchange coupling (β_{J}). The best fits of eqn (12) to our g_{mb} vs. $J_{\text{D} \rightarrow \text{A}}$ data yield $\beta_{\text{g}}/\beta_{\text{J}} = 1.15$ for the phenylene series and $\beta_{\text{g}}/\beta_{\text{J}} = 1.58$ for the thiophene series (Fig. 7). Beratan and Waldeck also observed a nonlinear relationship between g_{mb} and heterogeneous electron transfer rate constants, k_{ET} .^{4,5} However, in their study they showed that $\beta_{\text{g}}/\beta_{\text{ET}} < 1$ for a series of alkanethiol and peptide nucleic acid oligomer bridges, with electron transfer decaying more rapidly with increasing bridge distance than conductance. Here, we observe the opposite correlation, with conductance falling off more rapidly with distance than the magnetic exchange coupling ($\beta_{\text{g}}/\beta_{\text{J}} > 1$), albeit using very different molecular bridges.

The relationship between g_{mb} and $J_{\text{D} \rightarrow \text{A}}$ for thiophenes is markedly more nonlinear than what we observe in the phenylene data. Eqn (13) describes how differences in these exponential

$$\beta_i = \frac{2}{L} \ln \left(\frac{\Delta \epsilon}{H_{\text{BB},i}} \right) \quad (13)$$

decay constants (β_i) directly relate to inherent differences in the tunneling energy gaps, $\Delta \epsilon$, the bridge–bridge electronic coupling, H_{BB} , and L , the length of the bridge unit.^{56,57} The tunneling energy gap ($\Delta \epsilon$) is defined as the energy required to promote an electron from the donor (anode) to the bridge LUMO, or to promote a hole from the acceptor (cathode) to the bridge HOMO, and the distance between the donor and acceptor (or the electrodes) is given by R_{DA} ($R_{\text{DA}} = R_0 + nL$). Thus, a combination of large H_{BB} and a small $\Delta \epsilon$ lead to low β -values and more shallow distance decays that promote long range magnetic superexchange coupling and conductance. Eqn (14) relates these distance decay constants to effective barrier heights, ΔE_{eff} , with m_e being the mass of the tunneling electron.^{56,57}

$$\Delta E_{\text{eff}} = \left(\frac{\hbar^2}{8m_e} \right) \beta^2 = (0.952 \text{ eV } \text{\AA}^2) \beta^2 \quad (14)$$

Beratan and Waldeck elegantly attributed their observed deviations in $\beta_{\text{g}}/\beta_{\text{ET}}$ from unity to a combination of charge transfer energy barrier differences between electron transfer in D–B–A ensembles and electron transport in M–B–M configurations, bath-induced decoherences on the bridge molecule, and bridge–bridge electronic coupling (H_{BB}).^{4,5} Their work comparing conductance values and electron transfer rate constants indicated that differences in effective barrier heights derive from differences between the electrode work function in a transport geometry and donor/acceptor redox potentials in D–B–A systems.^{4,5} With respect to β_i values, differences in the work function and redox potentials derive from different values for $\Delta \epsilon$. In general, due to the interdependence of $\Delta \epsilon$ and H_{BB} on β_i values it is difficult to separate these individual contributions to ΔE_{eff} .⁵⁸

Since decoherence effects are not important for the $J_{\text{D} \rightarrow \text{A}}$ vs. g_{mb} correlation presented here, the dominant contributors to the nonlinearity between $J_{\text{D} \rightarrow \text{A}}$ and g_{mb} must relate to differences in bridge–bridge electronic coupling (H_{BB}) and the tunneling energy



As introduced above, the resonance structures shown in Fig. 4 describe the charge and spin distributions of the ground state configuration (GC) and four excited state configurations (ECs) for the **SQ-Ph₂-NN** biradical complex. Quinoidal EC1 and EC3 possess enhanced Ph-Ph π -coupling, which provides a mechanism for increasing the magnitude of H_{BB} . Contributions of resonance structures EC1-EC4 to the ground state all lead to enhanced magnetic exchange coupling in SQ-Bridge-NN biradicals. This occurs *via* a combination of B-B, D-B, and B-A electronic couplings (H_{BB} , H_{DB} , H_{BA}), which are different than those provided by the coupling of these same bridge molecules with the electrodes ($|\overline{G}_{\text{B}}|$, I_{LB} , I_{RB}). This is partly due to the fact that only high energy zwitterionic and biradical resonance structures can be drawn for the S-Ph-Ph-S electrode linkage. In contrast, these CT configurations lead to a low-energy ILCT state at $\sim 24\,000\text{ cm}^{-1}$ in SQ-Bridge-NN biradicals, which has been shown to admix with the electronic ground state by resonance Raman spectroscopy.^{20,41} Specifically, the effect of this enhanced electronic coupling has been observed in SQ-Bridge-NN biradical compounds by optical pumping into the excited state ILCT and observing a large resonance enhancement of the phenylene quinoidal stretching

vibration, which decreases in intensity with increased SQ-Bridge and Bridge-NN bond torsions.⁴¹ For di-bridged **SQ-Ph₂-NN**, a large resonance enhancement of the 1600 cm⁻¹ quinoidal stretch is also observed when pumping into the SQ → Ph-NN ILCT band¹⁷ (Fig. 8). The nature of the charge transfer is indicated by the computed electron density difference map (EDDM) in Fig. 8, which shows net charge transfer from the SQ donor (red) to the Ph-NN acceptor fragment (blue). Thus, the nature of the charge transfer in di-bridged **SQ-Ph₂-NN** is analogous to what has been observed previously in the mono-bridged **SQ-Ph-NN** biradical,^{20,39} highlighting quinoidal resonance structure contributions to the ground state wavefunction.

Conclusions

This work provides a detailed account of the transferability of electronic coupling with regard to exchange coupling and conductance. We have developed an electronic structure description of the functional relationship between $J_{D \rightarrow A}$ and g_{mb} , which we observed to be nonlinear ($\beta_g/\beta_J \neq 1$). Plots of $J_{D \rightarrow A}$ vs. g_{mb} for monomeric and dimeric bridges lie on different curves due to a scale invariance between these bridge types, which indicates a proportionate scaling of the exchange interaction, $J_{D \rightarrow A}$. Our g_{mb} vs. $J_{D \rightarrow A}$ data yield $\beta_g/\beta_J = 1.15$ for the phenylene series and $\beta_g/\beta_J = 1.58$ for the thiophene series, and these ratios accurately replicate recently reported β -values for electron transport through oligothiophene and oligophenylene bridges.^{59,60} In addition, these β_g/β_J ratios have been used to determine conductance tunneling barrier heights for oligophenylenes ($\Delta E_{eff} = 1555$ cm⁻¹) and oligothiophenes ($\Delta E_{eff} = 941$ cm⁻¹) that span electrodes in a transport geometry. This translates to a decrease in effective barrier heights ΔE_{eff} for magnetic exchange coupling of ~25% for phenylenes and ~60% for thiophenes when compared to conductance values using these same organic bridge fragments.

Tunneling gaps and bridge-bridge electronic coupling, H_{BB} , calculated from experimental data for the phenylene- and thiophene series provide a convincing explanation for the more pronounced g_{mb} vs. $J_{D \rightarrow A}$ nonlinearity observed in the thiophene series compared to the phenylene series. Additionally, resonance Raman spectroscopy indicates that H_{BB} may be enhanced by quinoidal resonance structure contributions in SQ-Bridge-NN biradicals relative to M-B-M devices for conductance. Namely, contributions from ECN resonance structures (Fig. 4) are expected to be more pronounced for **SQ-Th_n-NN** than for **SQ-Ph_n-NN** or **Au_n-S-Bridge-S-Au_n** constructs.

Resonance enhancement of the bridge quinoidal stretching mode illustrates the key bridge distortion that is the hallmark of a large H_{BB} , and the magnitude of this excited state distortion is proportional to the degree of SQ → B-NN charge transfer, which contributes to the magnitude of the magnetic exchange interaction in SQ-Bridge-NN biradicals. From a valence bond/resonance perspective, contributing ECN resonance structures accurately reflect both the spin- and charge distribution of low-lying SQ-Bridge-NN CT states, for which the corresponding quinoidal bridge-containing excited states in M-Bridge-M

constructs lie far higher in energy. With respect to the importance of quinoidal resonance contributions, aromaticity has been shown to reduce conductance values in single molecule junctions.⁶⁵ Critically, the quinoidal resonance structure contributions in Fig. 4 result in diminished aromaticity and contribute more to exchange than to conductance, leading to $\gamma \neq 1$ and $\beta_g/\beta_J > 1$ for these molecular bridges.

Conflicts of interest

There are no conflicts to declare.

Acknowledgements

M. L. K. acknowledges NSF (CHE-1900237) for financial support. D. A. S. acknowledges financial support from NSF (CHE-1931291 and CHE-1764181). The authors thank the UNM Center for Advanced Research Computing, supported in part by the NSF, for providing high performance computing resources used in this work.

Notes and references

- 1 M. L. Kirk, D. A. Shultz, J. Y. Zhang, R. Dangi, L. Ingersol, J. Yang, N. S. Finney, R. D. Sommer and L. Wojtas, *Chem. Sci.*, 2017, **8**, 5408–5415.
- 2 Y. Tsuji, R. Movassagh, S. Datta and R. Hoffmann, *ACS Nano*, 2015, **9**, 11109–11120.
- 3 J. L. Xia, B. Capozzi, S. J. Wei, M. Strange, A. Batra, J. R. Moreno, R. J. Amir, E. Amir, G. C. Solomon, L. Venkataraman and L. M. Campos, *Nano Lett.*, 2014, **14**, 2941–2945.
- 4 R. Venkatramani, E. Wierzbinski, D. H. Waldeck and D. N. Beratan, *Faraday Discuss.*, 2014, **174**, 57–78.
- 5 E. Wierzbinski, R. Venkatramani, K. L. Davis, S. Bezer, J. Kong, Y. J. Xing, E. Borguet, C. Achim, D. N. Beratan and D. H. Waldeck, *ACS Nano*, 2013, **7**, 5391–5401.
- 6 W. B. Chen, J. R. Widawsky, H. Vazquez, S. T. Schneebeli, M. S. Hybertsen, R. Breslow and L. Venkataraman, *J. Am. Chem. Soc.*, 2011, **133**, 17160–17163.
- 7 E. D. Fung, D. Gelbwaser, J. Taylor, J. Low, J. L. Xia, I. Davydenko, L. M. Campos, S. Marder, U. Peskin and L. Venkataraman, *Nano Lett.*, 2019, **19**, 2555–2561.
- 8 W. D. Ding, M. Koepf, C. Koenigsmann, A. Batra, L. Venkataraman, C. F. A. Negre, G. W. Brudvig, R. H. Crabtree, C. A. Schmuttenmaer and V. S. Batista, *J. Chem. Theory Comput.*, 2015, **11**, 5888–5896.
- 9 P. Darancet, J. R. Widawsky, H. J. Choi, L. Venkataraman and J. B. Neaton, *Nano Lett.*, 2012, **12**, 6250–6254.
- 10 S. V. Aradhya, J. S. Meisner, M. Krikorian, S. Ahn, R. Parameswaran, M. L. Steigerwald, C. Nuckolls and L. Venkataraman, *Nano Lett.*, 2012, **12**, 1643–1647.
- 11 S. Schneebeli, M. Kamenetska, F. Foss, H. Vazquez, R. Skouta, M. Hybertsen, L. Venkataraman and R. Breslow, *Org. Lett.*, 2010, **12**, 4114–4117.
- 12 M. Kamenetska, S. Y. Quek, A. C. Whalley, M. L. Steigerwald, H. J. Choi, S. G. Louie, C. Nuckolls, M. S. Hybertsen,



- J. B. Neaton and L. Venkataraman, *J. Am. Chem. Soc.*, 2010, **132**, 6817–6821.
- 13 L. Venkataraman, *Nat. Nanotechnol.*, 2008, **3**, 187–188.
- 14 C. Herrmann, *J. Phys. Chem. A*, 2019, **123**, 10205–10223.
- 15 L. Venkataraman, J. E. Klare, I. W. Tam, C. Nuckolls, M. S. Hybertsen and M. L. Steigerwald, *Nano Lett.*, 2006, **6**, 458–462.
- 16 C. Joachim and M. A. Ratner, *Proc. Natl. Acad. Sci. U. S. A.*, 2005, **102**, 8801.
- 17 M. L. Kirk, D. A. Shultz, D. E. Stasiw, G. F. Lewis, G. B. Wang, C. L. Brannen, R. D. Sommer and P. D. Boyle, *J. Am. Chem. Soc.*, 2013, **135**, 17144–17154.
- 18 M. L. Kirk, D. A. Shultz, D. E. Stasiw, D. Habel-Rodriguez, B. Stein and P. D. Boyle, *J. Am. Chem. Soc.*, 2013, **135**, 14713–14725.
- 19 M. L. Kirk and D. A. Shultz, *Coord. Chem. Rev.*, 2013, **257**, 218–233.
- 20 M. L. Kirk, D. A. Shultz, E. C. Depperman, D. Habel-Rodriguez and R. D. Schmidt, *J. Am. Chem. Soc.*, 2012, **134**, 7812–7819.
- 21 A. Nitzan, *J. Phys. Chem. A*, 2001, **105**, 2677–2679.
- 22 A. B. Ricks, G. C. Solomon, M. T. Colvin, A. M. Scott, K. Chen, M. A. Ratner and M. R. Wasielewski, *J. Am. Chem. Soc.*, 2010, **132**, 15427–15434.
- 23 E. A. Weiss, M. J. Ahrens, L. E. Sinks, A. V. Gusev, M. A. Ratner and M. R. Wasielewski, *J. Am. Chem. Soc.*, 2004, **126**, 5577.
- 24 J. Proppe and C. Herrmann, *J. Comput. Chem.*, 2015, **36**, 201–209.
- 25 C. Herrmann and J. Elmsiz, *Chem. Commun.*, 2013, **49**, 10456–10458.
- 26 D. E. Stasiw, J. Y. Zhang, G. B. Wang, R. Dangi, B. W. Stein, D. A. Shultz, M. L. Kirk, L. Wojtas and R. D. Sommer, *J. Am. Chem. Soc.*, 2015, **137**, 9222–9225.
- 27 S. Nishizawa, J. Hasegawa and K. Matsuda, *Chem. Lett.*, 2014, **43**, 530–532.
- 28 S. Nishizawa, J. Hasegawa and K. Matsuda, *Chem. Phys. Lett.*, 2013, **555**, 187–190.
- 29 S. Nishizawa, J. Hasegawa and K. Matsuda, *J. Phys. Chem. C*, 2013, **117**, 26280–26286.
- 30 M. Shinomiya, K. Higashiguchi and K. Matsuda, *J. Org. Chem.*, 2013, **78**, 9282–9290.
- 31 Y. Tsuji, R. Hoffmann, M. Strange and G. C. Solomon, *Proc. Natl. Acad. Sci. U. S. A.*, 2016, **113**, E413–E419.
- 32 P. W. Anderson, *Phys. Rev.*, 1950, **79**, 350–356.
- 33 H. A. Kramers, *Physica*, 1934, **1**, 182–192.
- 34 P. J. Hay, J. C. Thibeault and R. Hoffmann, *J. Am. Chem. Soc.*, 1975, **97**, 4884–4899.
- 35 M. L. Kirk, D. A. Shultz, E. C. Depperman and C. L. Brannen, *J. Am. Chem. Soc.*, 2007, **129**, 1937–1943.
- 36 M. L. Kirk, D. A. Shultz and E. C. Depperman, *Polyhedron*, 2005, **24**, 2880–2884.
- 37 T. Steenbock, D. A. Shultz, M. L. Kirk and C. Herrmann, *J. Phys. Chem. A*, 2017, **121**, 216–225.
- 38 M. L. Kirk and D. A. Shultz, *Coord. Chem. Rev.*, 2013, **257**, 218–233.
- 39 M. L. Kirk, D. A. Shultz, D. Habel-Rodriguez, R. D. Schmidt and U. Sullivan, *J. Phys. Chem. B*, 2010, **114**, 14712–14716.
- 40 D. A. Shultz, K. E. Vostrikova, S. H. Bodnar, H. J. Koo, M. H. Whangbo, M. L. Kirk, E. C. Depperman and J. W. Kampf, *J. Am. Chem. Soc.*, 2003, **125**, 1607–1617.
- 41 D. A. Shultz, M. L. Kirk, J. Y. Zhang, D. E. Stasiw, G. B. Wang, J. Yang, D. Habel-Rodriguez, B. W. Stein and R. D. Sommer, *J. Am. Chem. Soc.*, 2020, **142**, 4916–4924.
- 42 M. L. Kirk and D. A. Shultz, *Coord. Chem. Rev.*, 2013, **257**, 218–233.
- 43 M. J. Frisch, G. W. Trucks, H. B. Schlegel, G. E. Scuseria, M. A. Robb, J. R. Cheeseman, G. Scalmani, V. Barone, G. A. Petersson, H. Nakatsuji, X. Li, M. Caricato, A. Marenich, J. Bloino, B. G. Janesko, R. Gomperts, B. Mennucci, H. P. Hratchian, J. V. Ortiz, A. F. Izmaylov, J. L. Sonnenberg, D. Williams-Young, F. Ding, F. Lipparini, F. Egidi, J. Goings, B. Peng, A. Petrone, T. Henderson, D. Ranasinghe, V. G. Zakrzewski, J. Gao, N. Rega, G. Zheng, W. Liang, M. Hada, M. Ehara, K. Toyota, R. Fukuda, J. Hasegawa, M. Ishida, T. Nakajima, Y. Honda, O. Kitao, H. Nakai, T. Vreven, K. Throssell, J. A. Montgomery Jr, J. E. Peralta, F. Ogliaro, M. Bearpark, J. J. Heyd, E. Brothers, K. N. Kudin, V. N. Staroverov, T. Keith, R. Kobayashi, J. Normand, K. Raghavachari, A. Rendell, J. C. Burant, S. S. Iyengar, J. Tomasi, M. Cossi, J. M. Millam, M. Klene, C. Adamo, R. Cammi, J. W. Ochterski, R. L. Martin, K. Morokuma, O. Farkas, J. B. Foresman and D. J. Fox, *Gaussian 09, Revision A.02*, Gaussian, Inc., Wallingford CT, 2016.
- 44 S. Smidstrup, T. Markussen, P. Vancraeyveld, J. Wellendorff, J. Schneider, T. Gunst, B. Verstichel, D. Stradi, P. A. Khomyakov, U. G. Vej-Hansen, M.-E. Lee, S. T. Chill, F. Rasmussen, G. Penazzi, F. Corsetti, A. Ojanpera, K. Jensen, M. L. N. Palsgaard, U. Martinez, A. Blom, M. Brandbyge and K. Stokbro, *J. Phys.: Condens. Matter*, 2019, **32**, 015901.
- 45 M. Brandbyge, J.-L. Mozos, P. Ordejon, J. Taylor and K. Stokbro, *Phys. Rev. B: Condens. Matter Mater. Phys.*, 2002, **65**, 165401.
- 46 J. M. Soler, E. Artacho, J. D. Gale, A. Garcia, J. Junquera, P. Ordejon and D. Sanchez-Portal, *J. Phys.: Condens. Matter*, 2002, **14**, 2745–2779.
- 47 M. Büttiker, Y. Imry, R. Landauer and S. Pinhas, *Phys. Rev. B: Condens. Matter Mater. Phys.*, 1985, **31**, 6207–6215.
- 48 C. Van Dyck and M. A. Ratner, *Nano Lett.*, 2015, **15**, 1577–1584.
- 49 J. Taylor, H. Guo and J. Wang, *Phys. Rev. B: Condens. Matter Mater. Phys.*, 2001, **63**, 245407.
- 50 E. C. Depperman, S. H. Bodnar, K. E. Vostrikova, D. A. Shultz and M. L. Kirk, *J. Am. Chem. Soc.*, 2001, **123**, 3133.
- 51 C. Van Dyck and M. A. Ratner, *J. Phys. Chem. C*, 2017, **121**, 3013–3024.
- 52 C. Van Dyck, V. Geskin and J. Cornil, *Adv. Funct. Mater.*, 2014, **24**, 6154–6165.
- 53 S. Priyadarshy, S. S. Skourtis, S. M. Risser and D. N. Beratan, *J. Chem. Phys.*, 1996, **104**, 9473–9481.



- 54 K. Yoshizawa, T. Tada and A. Staykov, *J. Am. Chem. Soc.*, 2008, **130**, 9406–9413.
- 55 Y. Tsuji, A. Staykov and K. Yoshizawa, *J. Am. Chem. Soc.*, 2011, **133**, 5955–5965.
- 56 H. B. Gray and J. R. Winkler, *Proc. Natl. Acad. Sci. U. S. A.*, 2005, **102**, 3534.
- 57 O. S. Wenger, *Acc. Chem. Res.*, 2011, **44**, 25–35.
- 58 O. S. Wenger, *Chem. Soc. Rev.*, 2011, **40**, 3538–3550.
- 59 D. J. Wold, R. Haag, M. A. Rampi and C. D. Frisbie, *J. Phys. Chem. B*, 2002, **106**, 2813–2816.
- 60 B. Capozzi, E. J. Dell, T. C. Berkelbach, D. R. Reichman, L. Venkataraman and L. M. Campos, *J. Am. Chem. Soc.*, 2014, **136**, 10486–10492.
- 61 L. Pauling and G. W. Wheland, *J. Chem. Phys.*, 1933, **1**, 362–374.
- 62 G. W. Wheland and L. Pauling, *J. Am. Chem. Soc.*, 1935, **57**, 2086–2095.
- 63 L. Pauling, *The Nature of the Chemical Bond*, Cornell University Press, Ithaca, NY, 3rd edn, 1960.
- 64 F. Weinhold and C. R. Landis, *Valency and bonding: a natural bond orbital donor-acceptor perspective*, Cambridge University Press, Cambridge, UK; New York, 2005.
- 65 W. B. Chen, H. X. Li, J. R. Widawsky, C. Appayee, L. Venkataraman and R. Breslow, *J. Am. Chem. Soc.*, 2014, **136**, 918–920.

