

Variable coordination of a chiral diphosphine containing an amidinium/NHC group within its backbone: μ -*P,P'*, κ^2 -*P,P'* and κ^3 -*P,C,P'* coordination modes†‡Paul D. Newman,*^a Kingsley J. Cavell^a and Benson M. Kariuki^b

Received 5th July 2012, Accepted 15th August 2012

DOI: 10.1039/c2dt31475d

A diphosphine ligand (**1**·HPF₆), which is a potential precursor to a PC_{NHC}P pincer, with a backbone containing two phenylene groups and a central bicyclic 4-aza-2-azoniabicyclo[3.2.1]oct-2-ene unit has been synthesised and coordinated to Pd(II) and Pt(II) to give *trans*-[M(κ^2 -*P,P'*-**1**·H)Cl₂]PF₆ where M = Pd (**2**) or Pt (**3a**). Single-crystal structure determinations of **2** and **3a** show the complexes to be isostructural with the diphosphine coordinated in a *trans*-spanning fashion and the amidinium unit being protonated and non-coordinated. **2** and **3a** react with CH₃I to give the dimers *trans*-[Pd₂(μ -**1**·H)₂I₄](PF₆)₂, **6**, and *trans*-[Pt₂(μ -**1**·H)₂I₄](PF₆)₂, **7**, as the major products. This bridging mode of coordination of [**1**·H]⁺ is also seen in *trans*-[Rh₂(μ -**1**·H)(1,5-COD)₂Cl₂]PF₆, **4**, and [Pt₂(μ - κ^2 -**1**·H)(dvdms)]PF₆, **5**. Upon treatment with KO^tBu complexes **2** and **3a** undergo deprotonation at the amidinium carbon to give *trans*-[M(κ^3 -*P,C,P'*-**1**)-Cl]PF₆ where M = Pd (**8**), and Pt (**9**). The related *trans*-[Rh(κ^3 -*P,C,P'*-**1**)(CO)]PF₆ (**10**) is prepared directly from **1**·HPF₆ and Rh(acac)(CO)₂; this and the palladium and platinum complexes **8** and **9** are isolated as isomeric mixtures as a consequence of a conformational isomerism. *In situ* deprotonation of **1**·HPF₆ followed by addition of Ag(CF₃SO₃) gave S_{Ag}-[Ag(κ^3 -*P,C,P'*-**1**)(CF₃SO₃)], **11**. Some preliminary studies of the reactivity of **2** and **8** in Suzuki-type reactions are reported and the Pt(0) system has been shown to be an active hydrosilylation catalyst.

Introduction

Wide angle diphosphines have been heavily exploited in homogeneous catalysis as demonstrated by the studies of van Leeuwen and others on the hydrocyanation and hydroformylation of alkenes.¹ The success of such systems, *e.g.* the Xanthos series, resides in the ability of the two phosphine donors to adopt an obtuse bite angle enforced by the rigid nature of the ligand backbone. These relatively fixed bite angles, which typically lie between 110 and 130°, assist in promoting metal-based catalysis but preclude true *trans* coordination at a square planar metal. Other diphosphines are able to coordinate to give P–M–P angles of ~180° and these are defined as *trans*-spanning ligands.^{1*e,f*,2} These are generally of two types: those with rigid backbones that strongly favour *trans*-coordination which can restrict catalysis through **2e** redox processes,^{2*a*,3} and those that are flexible and able to promote catalysis through rearrangement to active *cis*- or monodentate forms.^{2*a*,4} Depending on the nature

of the intervening ligand architecture, some *trans*-spanning diphosphines can undergo metallation at an atom in the ligand backbone to give PXP pincer complexes (where X is often carbon).⁵ Such complexes (which are not restricted to a PCP donor set) are finding increasing employment in homogeneous catalysis as they are usually robust by nature and able to withstand the high temperatures demanded by certain types of transformation notably alkane dehydrogenation.⁶ They are also proving useful in non-oxidative bond activation through metal–ligand cooperation.⁷

The range of pincer ligands containing neutral NHC carbon donors includes those with a single central carbene donor⁸ and those with two carbenes in the terminal positions⁹ while homo-leptic tricarbenes tend to be of the facial type.¹⁰ We are interested in expanded-ring NHC-based tridentates incorporating a central C_{NHC} donor and have recently reported a dipyriddy-ER-NHC (L^{Py2}) containing a bicyclic core derived from camphor.¹¹ These earlier investigations established some of the coordination chemistry of the ligand and highlighted its relative flexibility as a N,C_{NHC},N' donor with both planar and non-planar species being isolated. In [Ni(κ^3 -N,C,N'-L^{Py2})Cl]⁺ where the three donor atoms and the metal are essentially coplanar, twisting in the ligand backbone was observed giving rise to λ and δ conformational isomers.^{11*a*} However the related Pd(II) and Pt(II) complexes of L^{Py2} proved elusive. Isomerism of a different type was also observed in the [Rh(1,5-COD)(L^{Py2})]⁺ and [Ir(1,5-COD)-(L^{Py2})]⁺. The ligand occupies one axial and two equatorial sites in these trigonal bipyramidal five-coordinate complexes and the source of the isomerism here lies in the relative orientation of the

^aCardiff Catalysis Institute, School of Chemistry, Cardiff University, Cardiff, CF10 3AT, UK. E-mail: newmanp1@cardiff.ac.uk; Fax: +44 (0)29 20874030; Tel: +44 (0)29 20870464

^bSchool of Chemistry, Cardiff University, Cardiff, CF10 3AT, UK

†CCDC 891175–891178. For crystallographic data in CIF or other electronic format see DOI: 10.1039/c2dt31475d

‡We would like to acknowledge the outstanding contribution made by David J. Cole-Hamilton to chemistry in general, and to catalysis in particular. His work has been widely recognised over many years for its imagination and thoroughness in implementation.



central bicyclic structure and the pyridyl groups leading to *endo*, *endo* or *exo,exo* species.^{11b}

As coordination flexibility in multidentate ligands can have a profound effect on catalytic performance, we were interested in exploring the coordination behaviour of NHC based systems bearing donors other than pyridine groups. As part of this broader study we have opted to explore the coordination chemistry of a diphosphino-ER-NHC containing the bicyclic 6/7 membered NHC skeleton, the initial results of which are presented here.

Results and discussion

Pro-ligand synthesis

The synthesis of the pro-ligand **1**-HPF₆ is highlighted in Scheme 1 and follows an already established procedure for the related dipyrindyl analogue.^{11a} The condensation of two mols of 2-(diphenylphosphino)benzaldehyde with 1*R*,3*S*-diamino-1,2,2-trimethylcyclopentane gives the diimine as an off-white solid. Subsequent reduction with NaBH₄ produces the diamine which is readily condensed with triethylorthoformate in the presence of NH₄PF₆ to give **1**-HPF₆ as a white solid. The salt is relatively air-stable in the solid state but does slowly oxidise when in solution. The two phosphine groups in **1**-HPF₆ are different and, although the source of this distinction is remote (H *versus* CH₃ at the ϵ position), two signals are observed in the ³¹P{¹H} NMR spectrum at δ_P = −16.6 and −17.5 ppm. The ¹H NMR spectrum shows the NCHN proton at δ_H = 7.79 ppm which compares to a value of 8.14 ppm for this hydrogen in the dipyrindyl derivative L^{py2}.^{11a} The only other signals of any diagnostic value in the ¹H spectrum are those for the benzylic hydrogens which occur as three doublets of doublets between 4.84 and 4.42 ppm and a doublet at 4.28 ppm. The pro-ligand is only modestly soluble in alcohol and (although it appears to slowly oxidise in CDCl₃) can be crystallised from MeOH in air to give crystals suitable for structural analysis by single-crystal X-ray techniques. The molecular structure of the compound is shown in Fig. 1. The C–N bond lengths to the amidinium carbon in the two independent molecules of the unit cell average 1.315 Å and the C–N–C angles are 123.1° (average). These compare to values of 1.314(3) Å and 123.5(2)° and 1.307(6) Å and 123.0(4)° in the dibenzyl and dipyrindyl derivatives respectively.^{11a}

(κ^2 -P,P'-1-H) complexes

The design of the ligand is such that coordination as a bidentate through both phosphorus atoms can lead to *cis*- and *trans*-isomers although the latter may be encouraged by the lengthy and semi-rigid backbone. Initial interest was to see whether κ^2 -P,P' coordination could be achieved without activation of the amidinium group to investigate potential *cis/trans* isomerism in the bidentate systems. When one mol of **1**-HPF₆ and K₂[PdCl₄] were heated in EtOH at 70 °C a bright yellow solid precipitated from the solution. The isolated complex gives an AB pattern in the ³¹P{¹H} NMR spectrum with δ_P values of 24.7 and 22.3 ppm respectively. The large magnitude (542 Hz) of the ²J_{P-P} coupling constant suggests that the two inequivalent phosphorus donors are mutually *trans*.^{2a,b} The compound is confirmed as *trans*-[Pd(κ^2 -P,P'-1-H)Cl₂][PF₆], **2**, through determination of its molecular structure by single-crystal X-ray techniques (Fig. 2). The P–Pd–P bond angles average 173.6° for the two independent molecules in the unit cell and the average Pd–P bond length of 2.344 Å is slightly longer than the 2.322 Å and 2.316 Å observed in related *trans*-spanning diphosphine complexes of palladium.^{4a,12} The Pd–Cl bond lengths average 2.293 Å and compare with values of 2.292 and 2.312 Å for the related complexes.^{4a,12} The amidinium group has not undergone deprotonation during the preparation of **2** but it does appear pre-set for






Fig. 1 Ortep view of the molecular structure of [1-H]⁺. For clarity the PF₆[−] anion is not shown. Thermal ellipsoids are drawn at 50% probability, hydrogens are omitted for clarity. Selected bond lengths (Å) and angles (°) for [1-H]⁺: C1–N1 1.334(7), C1–N2 1.316(8), N1–C1–N2 122.7(6).



Scheme 1 Synthesis of pro-ligand **1**-HPF₆.





Scheme 2 (i) $\text{K}_2[\text{PdCl}_4]$, EtOH, Δ ; (ii) $\text{K}_2[\text{PdCl}_4]$, EtOH, Δ ; (iii) xs MeI, Δ ; (iv) KO^tBu ; (v) 0.5 $[\text{Rh}(1,5\text{-COD})\text{Cl}]_2$, MeOH, Δ ; (vi) $\text{Pt}_2(\text{dvdms})_3$; (vii) $[\text{Rh}(\text{acac})(\text{CO})_2]$, THF, Δ ; (viii) KO^tBu , $\text{Ag}(\text{OTf})$.






Fig. 2 Ortep view of the molecular structure of **2**. Thermal ellipsoids are drawn at 50% probability, hydrogens are omitted for clarity. Selected bond lengths (Å) and angles (°) for **2**: Pd1–P1 2.315(5), Pd1–P2 2.356(5), Pd1–Cl1 2.297(4), Pd1–Cl2 2.306(5), C1–N1 1.34(2) C1–N2 1.38(2), P1–Pd1–P2 173.2(2), Cl1–Pd1–Cl2 170.0(2), P1–Pd1–Cl1 85.30(16), P1–Pd1–Cl2 93.52(16), N1–C1–N2 118.3(17).

coordination upon treatment with base; this conclusion derives from the orientation of the NCN carbon which points towards the metal. This conformation is favoured by an intramolecular electrostatic interaction between the positively charged amidinium group and the chloride ligands that manifests itself as a noticeable bending of the two chlorides towards the amidinium group. Although the average Cl–Pd–Cl bond angle of 170.0° is not atypical, as similar or even greater distortions are observed in related *trans*-Pd(P₂)Cl₂ complexes,^{2a} the nature of the distortion in **2** is distinct as the flexing in these other complexes is away from the backbone of the diphosphine as dictated by steric effects.^{2a,4a}

The ¹H NMR spectrum of **2** shows the NCN hydrogen at δ_H 9.58 ppm, a shift of 1.79 ppm downfield of its position in the uncoordinated ligand. This may be a consequence of the electrostatic (possibly weak H-bonding) interaction with the Cl ligands or might be a result of its position along the *z* axis of the palladium. Downfield shifts for hydrogens orientated in relatively fixed positions over a palladium or platinum coordination plane have been noted previously.^{2i,j} All four hydrogens of the methylene groups are inequivalent as evidenced by the presence of four doublets in the ¹H NMR spectrum between 5.02 and 4.61 ppm with ²J_{H–H} coupling constants of 14.5 and 15.2 Hz. The remainder of the ¹H NMR spectrum is less informative providing little diagnostic information. The amidinium carbon is observed at δ_C = 154.5 ppm in the ¹³C{¹H} NMR spectrum of **2**. This is little shifted from its position in the spectrum for the free salt. The remainder of the ¹³C{¹H} NMR spectrum mimics closely that for **1**·HPF₆ except there is no observable coupling between the benzylic carbons and the phosphorus centre.

The complex [Pt(κ²-P,P'-**1**·H)Cl₂](PF₆) is isolated as a cream solid from the reaction of **1**·HPF₆ with K₂[PtCl₄] in EtOH. Inspection of the ³¹P{¹H} NMR spectrum of the isolated solid shows it to be a mixture of two complexes: a major species **3a** described by an AB pattern in the ³¹P{¹H} NMR spectrum and a minor species **3b** that gives a broad singlet (see below). The two complexes were separated by simple treatment with CHCl₃ which readily dissolved the major compound leaving the minor component as an insoluble solid.

The chloroform soluble complex **3a** gives an AB pattern in the ³¹P{¹H} NMR spectrum at δ_P = 20.0 (¹J_{P–Pt} = 2523 Hz) and 19.7 (¹J_{P–Pt} = 2526 Hz) ppm. The ¹H NMR spectrum is closely similar to that for *trans*-[Pd(κ²-P,P'-**1**·H)Cl₂](PF₆) with a downfield shifted NCHN resonance at 9.90 ppm. The similarity in the spectra for this complex and **2** suggests that they are isostructural which is confirmed upon inspection of the solid state structure (Fig. 3). The Pt–P bond lengths average 2.332(5) Å which are close to the value of 2.304(5) Å seen for Pt(SPANphos)Cl₂^{3b} and 2.3204(12) Å in a related complex.^{4c} The P–Pt–P angle however is significantly more obtuse at 178.9(2)° in **3a** compared to the lower values of 171.9(4)° and 174.43(5)° seen with the reported complexes.

The broad resonance seen in the ³¹P{¹H} NMR spectrum of the minor product **3b** sharpens to a singlet (δ_P = 13.5 ppm, ¹J_{P–Pt} = 2556 Hz) upon heating to 80 °C in d₆-DMSO. The ¹H NMR spectrum recorded at this temperature showed the amidinium hydrogen at 8.31 ppm which is not significantly shifted from its position in the spectrum of the uncoordinated ligand. The benzylic CH₂ hydrogens are inequivalent and appear as four separate resonances between δ_H = 5.80 and 4.77 ppm. One of the methyl groups is seen to resonate at an unusually high position (0.17 ppm) in the ¹H NMR spectrum of **3b** which might indicate a residence within the shielding region of one or more of the aromatic rings. Although the two phosphorus centres are inequivalent they clearly have the same chemical shift in the ³¹P{¹H} NMR spectrum of **3b** as only one resonance is seen. The magnitude of the ¹J_{P–Pt} coupling constant would suggest that it is not a *cis*-isomer as a larger *J* value would have been expected if the phosphines were *trans* to chloride.¹³ Thus the complex has likely *trans*-P donors but is seemingly not a monomer as the chloroform soluble compound **3a** has been shown to be this species. The most likely structure of this complex is therefore the dimer *trans*-[Pt₂(μ-**1**·H)₂(Cl)₄](PF₆)₂, although this could not be confirmed by mass spectrometry as only monomeric species were observed for the compound.

(μ-**1**·H) complexes

The 1 : 1 reaction of **1**·HPF₆ and [Rh(1,5-COD)Cl]₂ in MeOH at near reflux led to the formation of a light yellow solid. Inspection of the ¹H NMR spectrum of the isolated solid revealed the presence of coordinated COD and [**1**·H]⁺ in a 2 : 1 ratio. Other characteristic features of the ¹H NMR spectrum include the amidinium hydrogen at δ_H = 8.18 ppm and the four inequivalent benzylic methylene hydrogens which appear as four doublets between δ_H 6.5 and 5.6 ppm. The relative ratio of the [**1**·H]⁺ and COD resonances coupled with the chemical shift of the amidinium hydrogen would suggest that the isolated complex is [Rh₂(1,5-COD)₂(μ-**1**·H)Cl₂](PF₆)₂, **4**. The ³¹P{¹H} NMR spectrum consists of two doublets at δ_P 21.7 and 21.4 with ¹J_{P–Rh} being 146 Hz. Although a coordination shift is evident and both resonances show coupling to rhodium, the lack of any ²J_{P–P} coupling supports the above assignment where the amidinium diphosphine is coordinating in a bridging mode. The magnitude of the ¹J_{P–Rh} coupling is commensurate with that expected for a triarylphosphine *trans* to a C=C donor of COD as noted in several complexes of the type [Rh(1,5-COD)(PAR₃)Cl].¹⁴

The reaction of **1**·HPF₆ with Karstedt's complex Pt₂(dvdms)₃ in THF gave an air-stable cream coloured solid upon work-up.






Fig. 3 Ortep view of the molecular structure of **3a**. Thermal ellipsoids are drawn at 50% probability, hydrogens are omitted for clarity. Selected bond lengths (Å) and angles (°) for **3a**: Pt2–P3 2.329(5), Pt2–P4 2.315(4), Pt2–Cl3 2.310(5), Pt2–Cl4 2.303(4), C48–N3 1.31(2), C48–N4 1.32(2), P3–Pt2–P4 174.9(3), Cl3–Pt2–Cl4 174.73(19), P3–Pt2–Cl3 87.63(14), P3–Pt2–Cl4 94.15(14), N3–C48–N4 120.5(19).

The $^{31}\text{P}\{^1\text{H}\}$ NMR spectrum consisted of two singlets at 20.7 and 18.7 ppm each with Pt-195 satellites ($^1J_{\text{P-Pt}} = 3494$ and 3506 Hz, respectively). As for the rhodium complex **4** above, the presence of both a coordination shift and P–Pt coupling confirms coordination of the phosphine donors and the absence of any $^2J_{\text{P-P}}$ coupling indicates a μ bonding mode. Inspection of the ^1H NMR spectrum of the complex reveals the amidinium hydrogen at δ_{H} 8.18 ppm which is exactly as observed in the Rh dimer **4**. Integration of selected resonances of the diphosphine and the dvdm's reveal the ratio to be 1 : 2 and the most likely formulation of the product is $[\text{Pt}_2(\text{dvdm})_2(\mu\text{-I}\cdot\text{H})]\text{PF}_6$. This accords with similar systems reported by Lappert where comparable values of $\Delta\delta$ and $^1J_{\text{P-Pt}}$ are noted.¹⁵ This assignment is further supported upon inspection of the ^{195}Pt NMR spectrum which shows two doublets of multiplets at $\delta_{\text{Pt}} = -5553$ and -5563 ppm for the two platinum centres in the dimer. These chemical shift values again compare with related $\text{Pt}(\text{PR}_3)(\text{dvdm})_2$ complexes.^{15,16}

When **2** was reacted with excess MeI at 50 °C in CHCl_3 an orange precipitate was formed. Inspection of the $^{31}\text{P}\{^1\text{H}\}$ NMR spectrum of the product showed a singlet at $\delta_{\text{P}} = 21.1$ ppm. The ^1H NMR spectrum was similar to that for **2** albeit with notable shifts in δ_{H} for the amidinium hydrogen and the benzylic CH_2 hydrogens. The MS data showed a peak for the complex $[\text{Pd}(\text{I}\cdot\text{H})\text{I}_2]^+$ at 1063 amu confirming that the chloride ligands had been exchanged for iodide presumably through two successive oxidative addition (MeI)/reductive elimination (MeCl) cycles. The similar spectroscopic features of **2** and **6** suggest that **6** is a monomer containing *trans*-bound $[\text{I}\cdot\text{H}]^+$. The orange solid could be recrystallised from MeCN to give crystals suitable for analysis by single-crystal X-ray techniques (Fig. 4). As can be seen from the structure shown in Fig. 4 the complex has rearranged upon recrystallisation to give the $[\text{Pd}_2(\mu\text{-I}\cdot\text{H})_2\text{I}_4]\cdot\text{I}_{0.27}(\text{PF}_6)_{1.73}$ species containing two bridging $[\text{I}\cdot\text{H}]^+$ ligands. The coordination about each of the Pd centres is the expected square planar arrangement with the P-donors (and by necessity the I^- ligands) being mutually *trans*. The Pd–P bond lengths average 2.33 Å which is slightly shorter than in **2** indicating a




Fig. 4 Ortep view of the molecular structure of **6**. Thermal ellipsoids are drawn at 50% probability, hydrogens are omitted for clarity. Selected bond lengths (Å) and angles (°) for **6**: Pd1–P1 2.321(7), Pd1–P2 2.330(6), Pd1–I1 2.632(2), Pd1–I2 2.592(3), Pd2–P3 2.362(7), Pd2–P4 2.295(6), Pd2–I3 2.565(2), Pd2–I4 2.531(3), P1–Pd1–P2 166.36(18), I1–Pd1–I2 168.06(8), P1–Pd1–I1 88.31(17), P1–Pd1–I2 91.13(18), P3–Pd2–P4 167.69(17), I3–Pd2–I4 168.95(9), P3–Pd2–I3 91.30(18), P3–Pd2–I4 87.23(17).

small level of strain in the chelate. The P–Pd–P angles are unexpectedly acute at 166° compared to the 174° in **2** while the I–Pd–I angles at 169° are similar to the value of 170° seen for the Cl–Pd–Cl angle in **2**. The nature of the bending is such that the palladium atoms point inwards with regard to the cavity defined by the dimetallo macrocycle (Fig. 4). When the crystals were redissolved in CD_3CN for spectroscopic analysis, the resultant $^{31}\text{P}\{^1\text{H}\}$ and ^1H NMR spectra were complex indicating the presence of a number of species in solution. Substitution of up to four iodides in the dimer by CD_3CN is one possible source of solution complexity, however it seems likely that the $\mu\text{-I}\cdot\text{H}$ coordination mode of the diphosphine is not so robust that bidentate modes are precluded in solution.

The reaction of **3a** or **3b** (or a mixture of both) with MeI gave an orange solid **7** that showed a similar pattern to the **3a/b** mixture in its $^{31}\text{P}\{^1\text{H}\}$ NMR spectrum. Significantly however the position of the resonances had shifted ~ 10 ppm upfield commensurate with the replacement of the two chloride ligands in **3a/b** with iodide. This was confirmed by mass spectrometry where a species correlating to $[\text{Pd}(\text{I}\cdot\text{H})\text{I}_2]^+$ was observed at 1151 amu. The magnitude of the $^1J_{\text{P-Pt}}$ coupling constants were comparable to those in **3a/b** as expected for phosphorus donors situated mutually *trans*. No attempt was made to separate the mixture.

$(\kappa^3\text{-P,C,P'}\text{-I})$ complexes

In the examples discussed above the diphosphine ligand coordinates solely through the phosphorus atoms and the third potential donor (the NHC carbon) is protonated and hence non-coordinating. The solid-state structures of the platinum and palladium complexes of $\kappa^2\text{-}[\text{I}\cdot\text{H}]^+$ show this carbon to be poised for coordination upon removal of the proton so that tridentate forms should be accessible upon addition of base to **2** and **3a**. When **2** was treated with 1.2 equivalents of KO^tBu in THF the colour of the solution changed from yellow to a deep blood-red and the complex $[\text{Pd}(\kappa^3\text{-P,C,P'}\text{-I})\text{Cl}]\text{PF}_6$, **8**, was isolated as a red solid



after work-up. Inspection of the $^{31}\text{P}\{^1\text{H}\}$ NMR spectrum of **8** revealed the presence of two sets of AB doublets around 5 ppm suggesting the presence of two isomers in solution. This was confirmed upon observation of the ^1H NMR spectrum where a total of six CH_3 resonances and eight separate doublets for the benzylic methylenes were seen. Some notable features of the ^1H NMR spectrum include an unusual chemical shift for two of the methyl resonances to highfield of TMS and a large δ_{H} separation for the hydrogens of any given benzylic- CH_2 group with one resonating over 2 ppm downfield of its geminal partner. It would appear that the binding of the ligand in the κ^3 -mode has placed one of the methyl groups of each isomer in a region that experiences a local shielding effect presumably from one or more of the aromatic rings. Similarly, one geminal hydrogen of each benzylic group in either isomer resides in a deshielded zone. Isomerisation in heterodonor terdentates containing a central NHC donor has been observed by us previously in square planar and five-coordinate trigonal bipyramidal metal systems.¹¹ Conformational differences in the ligand backbone are the source of the isomerism which is dictated by the desire of the NHC donor to be canted with respect to the coordination plane as shown schematically in Fig. 5; in related square planar metal systems where the NHC does not have pendant donors the NCN link of the carbene donor is known to lie approximately perpendicular to the coordination plane.^{11a}

As noted above the $^{31}\text{P}\{^1\text{H}\}$ NMR spectrum of **8** consists of two AB patterns for the two conformers with $^2J_{\text{P-P}}$ coupling appreciably smaller at 436 and 435 Hz than observed in **2**. More surprising is the upfield shift in δ_{P} that accompanies the change from κ^2 to κ^3 binding as the reverse (downfield) trend is more often observed upon formation of smaller chelates. These more familiar downfield shifts often reflect the presence of shorter M-P bond lengths and it might be that the transition to the κ^3 mode leads to longer Pd-P bond lengths compared to those seen in **2**; unfortunately this could not be confirmed as crystals suitable for single-crystal structure determination were not forthcoming. There was no evidence of $^2J_{\text{C-P}}$ coupling to the NCN carbons in the $^{13}\text{C}\{^1\text{H}\}$ NMR spectrum of **8** where two singlets at 178.2 and 177.2 ppm occur for each isomer respectively. The remainder of the $^{13}\text{C}\{^1\text{H}\}$ spectrum is not of diagnostic value other than to confirm the presence of the two isomers.

The analogous platinum complex $[\text{Pt}(\kappa^3\text{-P,C,P'-1})\text{Cl}]\text{PF}_6$, **9**, was prepared by a similar procedure to that for **8** and isolated as a cream solid. The ^1H NMR spectrum of **9** mimics closely that of **8** as does the $^{31}\text{P}\{^1\text{H}\}$ NMR spectrum although this does have the usual Pt-195 satellites with $^1J_{\text{P-Pt}}$ coupling constants of 2552 and 2570 Hz. These compare with those reported for **3a/b** above confirming the retention of the *trans*-P arrangement in **9**. The NCN carbons for each isomer are observed as triplets at 166.2 and 164.9 ppm with $^2J_{\text{C-P}}$ of 9.2 Hz in the $^{13}\text{C}\{^1\text{H}\}$ NMR

spectrum of **9**. Similar values of 9.2 and 9.4 Hz have been observed in the complexes *trans*- $[\text{Pd}(\text{BenzNHC})(\text{PPh}_3)_2\text{Cl}]\text{BF}_4$ ¹⁷ where BenzNHC is a benzimidazole derived NHC and $[\text{Pt}(\text{[16]-ane-P}_2\text{C}^{\text{NHC}}_2)](\text{PF}_6)_2$ containing a macrocyclic ligand.¹⁸ A larger value of 25 Hz has been reported for a related P,C,P complex containing a central NHC donor.^{8h}

As a chelating complex was not obtained from the reaction of **1**·HPF₆ with $[\text{Rh}(1,5\text{-COD})\text{Cl}]_2$, access to a κ^3 complex through the addition of base to a preformed $[\text{Rh}(\kappa^2\text{-P,C,P'-1})(\text{L})_2]^{n+/-}$ species was precluded and a different means of synthesis was required. $[\text{Rh}(\text{acac})(\text{CO})_2]$ was considered an ideal starting material as it contains an internal base (acac⁻) for *in situ* deprotonation after coordination of one or both phosphines. The reaction of **1**·HPF₆ with 1 equivalent of $[\text{Rh}(\text{acac})(\text{CO})_2]$ in THF at reflux gave, after removal of all volatiles, a yellow solid that was recrystallised from MeOH. The ^1H NMR spectrum confirmed the loss of the amidinium proton and the coordinated acac ligand suggesting that the desired reaction had occurred to give $[\text{Rh}(\kappa^3\text{-P,C,P'-1})(\text{CO})]\text{PF}_6$, **10**. The structure of the complex was confirmed upon closer inspection of the ^1H NMR spectrum which showed the characteristic features for κ^3 -**1** coordination, *i.e.* the presence of two isomers with two methyl resonances upfield of TMS and eight doublets for the benzylic hydrogens with each geminal pair being separated by >2 ppm. The $^{31}\text{P}\{^1\text{H}\}$ NMR spectrum consisted of two almost overlapping ABX multiplets with $^2J_{\text{P-P}}$ having a magnitude of 268 and 274 Hz and $^1J_{\text{P-Rh}} = 134$ Hz. This value is typical of a *trans*-phosphine complex¹⁹ being only slightly smaller than the value of 142 Hz for the *trans*-PPh₃ groups in $\text{Rh}(\text{PPh}_3)_3\text{Cl}$ ²⁰ and is exactly the same as the values quoted for complexes of the type *trans*- $[\text{Rh}(\text{L})(\text{PPh}_3)_2(\text{CO})]\text{X}^{21}$ where L is a saturated five-membered NHC. The infrared spectrum of **10** showed the CO stretch for the coordinated carbonyl ligand at 2008 cm^{-1} which is directly comparable to the values observed by Lappert and co-workers for *trans*- $[\text{Rh}(5\text{-NHC})(\text{PPh}_3)_2(\text{CO})]\text{X}^{21}$.

The examples above are systems in which deprotonation of the amidinium group succeeds coordination of the phosphine donors (this is known for the preformed Pd and Pt complexes **2** and **3a/b** and implied for the Rh). The common approach of *in situ* deprotonation of **1**·HPF₆ followed by addition of the resultant free carbene to an appropriate metal precursor had, to this point, not been necessary. However, it was of interest to know if such an approach could be used with **1**·HPF₆ as the *N,N'*-dibenzyl analogue of **1**·HPF₆ does furnish complexes *via* this approach while the *N,N'*-di(2-pyridyl) derivative does not.^{11a} Part of the problem with benzyl derived NHCs is the facility by which they undergo a 1,2-migration of the benzylic substituent in their unprotected form.²² Fortunately when this approach was attempted in THF with AgOTf the complex $[\text{Ag}(\kappa^3\text{-P,C,P'-1})(\text{OTf})]$, **11**, was isolated as a grey solid. This was encouraging as the common method of preparing Ag-NHC complexes directly from the amidinium salt and Ag₂O did not work with **1**·HPF₆. The $^{31}\text{P}\{^1\text{H}\}$ NMR spectrum of the isolated solid appears relatively complex because of the presence of P-Ag(107/109) coupling but is essentially two superimposed ABX patterns, one for each of the two isotopomers. The value of the $^1J_{\text{P-Ag(107)}}$ and $^1J_{\text{P-Ag(109)}}$ coupling constants are 378 and 327 Hz respectively, reflecting the inherent difference in the gyromagnetic ratio for the two isotopes. The relatively small




Fig. 5 The two conformers for $\kappa^3\text{-P,C,P'-1}$ at a square planar metal centre.



value of 65.5 Hz for $^2J_{P-P}$ is not consistent with the large values of 300–600 Hz expected for *trans*-orientated P-donors and it would appear that the silver complex does not contain a PCPM plane. This is largely to be expected as d^{10} Ag(I) does not tend to form octahedral or square planar complexes and, if **11** is monomeric (although bridging forms of $PC_{NHC}P$ ligands are known²³ only monomeric species were observed in the mass spectrum of **11**), then the likely geometry would approximate to tetrahedral as noted for the related $[Cu(\kappa^3-P,C,P'-1)X]$ species.²⁴ The absolute magnitude of the $^2J_{P-P}$ coupling constant compares well with known tetrahedral phosphine containing complexes of silver(I) such as $[AgBr(\kappa^2-P,P'-dppp)(\kappa^1-P-dpppO)]$ where a value of 58 Hz is observed.²⁵ The $^1J_{P-Ag}$ coupling constants are somewhat larger than those of 253 and 220 Hz observed in $[AgBr(\kappa^2-P,P'-dppp)(\kappa^1-P-dpppO)]$.²⁵ Even though the coordination in the Ag(I) complex does not contain a planar PCPAg unit, there are still two possible isomers as shown in Fig. 6. The related copper(I) systems $[Cu(\kappa^3-P,C,P'-1)X]$ show a halide-dependant diastereomerism with the R_{Cu} configuration being favoured when $X = I^-$ and the S_{Cu} configuration when $X = Cl^-$.²⁴ The 1H NMR spectrum of **11** shows no evidence of the presence of more than one isomer although some of the signals in the 1H NMR spectrum are broadened slightly, most notably for one set of benzylic CH_2 resonances (AB pattern) and one of the methyl signals. The absence of a CH_3 resonance upfield of TMS in the 1H NMR spectrum would suggest that the preferred isomer of **11** has the *S* configuration at the metal and that this isomer is formed preferentially. Some very slight broadening is also evident for the benzyl methylenes in the $^{13}C\{^1H\}$ NMR spectrum of **11** although the spectrum is otherwise well resolved and there is some evidence of a second species in solution as each of the major peaks is associated with a minor peak. The NCN resonance however was not clearly resolved in the spectrum and was seen as a broad unstructured peak at 211 ppm. This is not surprising as this carbon is expected to be coupled to both phosphorus atoms and the silver for each of the isotopomers of the complex. The chemical shift of this carbon is downfield that of the dibenzyl analogue in a complex of the type $[Ag(L)_2]OTf^{11a}$ but compares to other silver complexes of 6- and 7-membered NHCs.²⁶

There have been very few studies of PCP ligands where the central donor is an NHC carbon.^{8b,c,e,22} Of these, none contain a

ring-expanded NHC and only one consists of two 7-membered chelates.^{8e} The Cu(I) complex of the latter ligand does not show a normal coordination mode but rather the NCN carbon acts as a bridging ligand in a dimeric Cu(I)–Cu(I) complex.^{8e} Such a motif is more common for monovalent silver²⁷ but is not considered to apply to complex **11** as both phosphines are clearly attached to a single Ag(I) centre as evidenced by the presence of $^2J_{P-P}$ coupling. It is also noteworthy that the related Cu(I) complex is a tetrahedral monomer.²⁴

Catalysis

Complexes of staunchly *trans*- P_2 chelates are not anticipated to catalyse transformations which involve reductive elimination. This is especially so when the catalytic cycle requires a transition from a square planar d^8 complex to a tetrahedral or lower coordinate d^{10} system upon reduction, *e.g.* $Pd(II) \rightarrow Pd(0)$. Such inactivity is associated with the relative position of the two potential coupling partners: being mutually *trans* they cannot eliminate. Less rigid P_2 donors may show a preference for *trans*-coordination but their increased flexibility enables *cis*-bidentate binding or monodentate coordination (either through bridging or transient loss of one P-donor) and thus facilitate catalysis by cycles that do involve reductive elimination. Both $[1-H]^+$ and **1** have been shown to be relatively promiscuous in their mode of metal binding and selected palladium and/or platinum complexes were anticipated to be active for catalysis.

In order to test this assumption some preliminary C–C coupling and hydrosilylation catalytic tests were performed with **2**, **8** and **5**. The results of the Suzuki coupling of various aryl halides and phenylboronic acid are shown in Table 1. Good conversions were seen for the aryl bromides after 18 h at 80 °C but no activity was observed with the aryl chlorides. The palladium complexes **2** and **8** show little difference in reactivity which may suggest a common catalytic cycle. The fact that $\kappa^3-P,C,P'-1$ has been shown to support d^8 and d^{10} metal systems coupled with




Fig. 6 Diastereomeric possibilities for κ^3-1 at pseudo tetrahedral silver(I) in complex **11**.

Table 1 Summary of the Suzuki coupling for complexes **2** and **8**^a

Catalyst	Substrate	Conversion ^b (%)
2	Bromobenzene	84
2	4-Bromotoluene	100
2	4-Bromoanisole	72
2	4-Bromoacetophenone	100
2	2-Bromotoluene	59
2	2-Bromopyridine	54
2	4-Chlorotoluene	0
2	4-Chloroanisole	0
2	2-Chlorotoluene	0
8	4-Bromotoluene	100
8	4-Bromoanisole	60
8	4-Bromoacetophenone	100
8	2-Bromotoluene	84
8	4-Chlorotoluene	0
8	4-Chloroanisole	0
8	2-Chlorotoluene	0

^a Catalytic conditions: 1×10^{-3} mol substrate, 1.2×10^{-3} mol $PhB(OH)_2$, 2×10^{-3} mol K_2CO_3 , 0.5 mol% catalyst, 3 ml 1,4-dioxane, 80 °C, 18 h. ^b Total GC-MS yield based upon consumption of substrate using *n*-decane as internal standard.



Table 2 Summary of the hydrosilylation results for complex **5**^a

Substrate	Conversion ^b (%)	Selectivity ^c
1-Octyne	100	98.5 : 1 : 0.5 ^d
2-Hexyne	100	63 : 37 ^e
Phenylacetylene	100	>99 : 1 ^f
Diphenylacetylene	100	—
Trimethylsilylacetylene	100	75 : 25 ^g
Styrene	29	89 : 11 ^h
Cyclohexenone	100	—
Octene	0	—

^a Catalytic conditions: 1×10^{-3} mol substrate, 3 ml toluene, 70 °C, substrate:Et₃SiH ratio 1 : 1, 0.01 mol% catalyst. ^b Total GC-MS yield based upon consumption of substrate using *n*-decane as internal standard. ^c Determined by ¹H NMR of the distilled products. ^d 1-(*E*)-Triethylsilyloct-1-ene: 1-(*Z*)-triethylsilyloct-1-ene: 2-triethylsilyloct-1-ene. ^e 2-(*E*)-Triethylsilylhex-2-ene: 3-(*E*)-triethylsilylhex-2-ene. ^f 2-(*E*)-Triethylsilylstyrene: 2-triethylsilylstyrene. ^g 1-(*E*)-Triethylsilyl-2-trimethylsilyl-ethene: 1-triethylsilyl-1-trimethylsilyl-ethene. ^h 2-Triethylsilyl-1-ethylbenzene: 2-(*E*)-triethylsilylstyrene.

the basic conditions of the catalysis tends to suggest the presence of $\kappa^3\text{-P,C,P'}$ complexes in the catalytic cycle for both **2** and **8**.

Platinum complexes have long been known as catalysts for the hydrosilylation of unsaturated hydrocarbons. Zerovalent platinum complexes containing divinyl-disiloxane feature prominently in the literature²⁸ usually with a secondary controlling ligand as Karstedt's complex Pt₂(dvdms)₃, although active, is poorly selective.²⁹ Given this history it is not surprising that **5** acts as a catalyst for the hydrosilylation of alkynes with Et₃SiH (Table 2). Complete conversion is observed for all the alkynes examined after 18 h at 70 °C with a catalyst loading of 0.01 mol%. Regio-control is an important consideration in hydrosilylation as alkynes can give a number of products upon addition, e.g. with Et₃SiH and terminal alkynes as substrates, *E* and *Z* 1-triethylsilyl and 2-triethylsilyl products can be formed. Although the thermodynamically more stable 1-substituted *E*-alkene is usually the major product, the observed selectivity for this isomer is very high for phenylacetylene and oct-1-yne with **5** compared with the best of the reported highly selective Pt(0) catalysts.^{28g,h,30} This selectivity is reduced for trimethylsilylacetylene where the 2-substituted isomer is produced as 25% of the product: this is not surprising as the alkyne substrate is electronically distinct and tends to form more 2-substituted product in other catalyst systems.^{28e} The internal hex-2-yne gives two products with the 2-isomer predominating (Table 2). Although the selectivity here is modest it is better than the essentially 1 : 1 mixtures observed with complexes of the type Pt(ER–NHC)(dvdms)₂.^{28h} The reactivity towards alkenes is poor as exemplified by the two entries in the table. Oct-1-ene was chosen as a typical terminal alkene and styrene as a somewhat activated alkene; only the latter showed any conversion under the conditions employed for the reaction.

Experimental

Methods and materials

All synthetic procedures and manipulations were performed under dry argon or nitrogen using standard Schlenk line

techniques. All solvents were freshly distilled from sodium (toluene), sodium/benzophenone (THF) or calcium hydride (acetonitrile, methanol and dichloromethane) under nitrogen before use. All other chemicals were obtained commercially and used as received. The ¹H and ¹³C NMR spectra were recorded on a Jeol Eclipse 300 MHz or Bruker 400 MHz spectrometer and referenced to tetramethylsilane (δ = 0 ppm). All ³¹P NMR spectra were recorded on a Jeol Eclipse 300 spectrometer operating in the proton-decoupled mode at 121.7 MHz. Mass spectra were obtained using a Waters LCT Premier XE mass spectrometer. Elemental analyses were performed by Stephen Boyer at London Metropolitan University, UK.³¹

Syntheses

1,2,2-Trimethyl-*N,N'*-cyclopentane-1,3-diylbis[1-{2-(diphenylphosphanyl)phenyl}methanimine]. A solution of 2-(diphenylphosphanyl)benzaldehyde (3 g, 10.3 mmol) and 1,2,2-trimethylcyclopentane-1,3-diamine (0.71 g, 5 mmol) were refluxed in degassed EtOH (70 ml) for 2 h under N₂. After cooling, the solution was concentrated to small volume *in vacuo* whereupon a white solid proliferated. This was filtered under N₂, washed sparingly with cold EtOH and dried at the pump. Yield = 3.1 g (90%). ¹H NMR (CDCl₃, 400 MHz) δ 8.71 (1H, d, *J* 4.7 Hz), 8.65 (1H, d, *J* 5.2 Hz), 7.91 (2H, m), 7.20 (24H, m), 6.74 (2H, m), 3.25 (1H, t, *J* 8.5 Hz), 1.87 (1H, m), 1.62 (2H, m), 1.37 (1H, m), 0.74 (3H, s), 0.57 (3H, s), 0.56 (3H, s) ppm. ³¹P (CDCl₃, 121.6 MHz): δ –12.7, –13.7 ppm.

***N,N'*-Bis{2-(diphenylphosphanyl)benzyl}-1,2,2-trimethylcyclopentane-1,3-diamine**. To a solution of 1,2,2-trimethyl-*N,N'*-cyclopentane-1,3-diylbis[1-{2-(diphenylphosphanyl)phenyl}methanimine] (2 g, 2.91 mmol) in MeOH was added portionwise NaBH₄ (0.3 g, 7.90 mmol). The mixture was stirred overnight at RT and then hydrolysed by the careful addition of conc. HCl (2 ml). The methanol was removed under reduced pressure and the residue dissolved in water (50 ml). The aqueous solution was made basic by the addition of 6 M NaOH solution and the desired phosphine extracted into diethyl ether (3 $\times$ 50 ml). After drying over MgSO₄ and removal of volatiles the diphosphine was obtained as an off-white solid. Yield = 1.8 g (90%). ¹H NMR (CDCl₃, 400 MHz) δ 7.60 (1H, m), 7.45 (3H, m), 7.25 (20H, m), 7.15 (2H, m), 6.81 (2H, m), 3.83 (4H, m), 2.65 (1H, t, *J* 6.6 Hz), 1.80 (1H, m), 1.60 (1H, m), 1.33 (3H, m br), 0.91 (3H, s), 0.71 (3H, s), 0.58 (3H, s) ppm. ¹³C{¹H} DEPT NMR (CDCl₃, 75.6 MHz) δ 146.1 (d, *J* 24.2 Hz), 145.4 (d, *J* 24.2 Hz), 137.0 (d obs), 135.2 (d, *J* 10.0 Hz), 134–128 (aromatics), 127.8 (d, *J* 5.5 Hz), 68.1 (s), 64.7 (s), 51.0 (CH₂, d, *J* 21.9 Hz), 47.2 (s), 45.8 (CH₂, d, *J* 23.1 Hz), 34.3 (s), 28.2 (s), 23.8 (s), 20.7 (s) ppm. ³¹P (CDCl₃, 121.6 MHz): δ –12.7, –13.7 ppm.

1-HPF₆. *N,N'*-Bis{2-(diphenylphosphanyl)benzyl}-1,2,2-trimethylcyclopentane-1,3-diamine (2 g, 2.89 mmol) and NH₄PF₆ (0.52 g, 3.2 mmol) were heated in triethylorthoformate (15 ml) at 100 °C under N₂ for 2 h. After cooling, the white precipitate was filtered, washed with cold EtOH (2 $\times$ 3 ml), then diethyl ether and air-dried. Yield = 2.3 g (94%). ¹H NMR (CD₃CN, 400 MHz) δ 7.79 (1H, br), 7.52–7.04 (25H, m), 6.95 (1H, m), 4.84 (1H, dd, *J* 14.5, 3.1 Hz), 4.73 (1H, dd, *J* 14.5, 1.7 Hz), 4.42 (1H, dd, *J* 14.6, 1.3 Hz), 4.28 (1H, d, *J* 14.6 Hz), 3.20 (1H, d, *J* 4.0 Hz), 2.36 (1H, m), 1.89 (2H, m), 1.68 (1H, m), 1.31



(3H, s), 0.95 (3H, s), 0.84 (3H, s) ppm. $^{13}\text{C}\{^1\text{H}\}$ DEPT NMR (CDCl_3 , 75.6 MHz) δ 151.7 (C, d, J 10.4 Hz), 137–127 (aromatics), 71.3 (CH, s), 65.1 (C, s), 54.7 (CH_2 , d, J 17.3 Hz), 50.7 (CH_2 , d, J 25.4 Hz), 40.2 (C, s), 38.5 (CH_2 , s), 30.5 (CH_2 , s), 20.6 (CH_3 , s), 15.9 (CH_3 , s), 13.5 (CH_3 , s) ppm. ^{31}P (CD_3CN , 121.6 MHz): δ -16.6, -17.5 ppm. MS: 701 ($[\text{1-H}]^+$, 100%).

trans-[Pd(κ^2 -P,C,P'-1)Cl₂]PF₆, 2. A mixture of K_2PdCl_4 (100 mg, 3.04×10^{-4} mol) and **1** (258 mg, 3.04×10^{-4} mol) in ethanol (20 ml) was refluxed for 24 h under N_2 . After cooling, the yellow precipitate was filtered in air, washed with EtOH (2×3 ml) and air-dried. Yield = 270 mg (87%). The complex can be recrystallised from CHCl_3 as golden yellow crystals. ^1H NMR (CD_3CN , 250 MHz) δ 9.58 (1H, s br), 8.05 (5H, m br), 7.7–7.4 (21H, m), 7.36 (1H, m), 7.11 (1H, m), 5.02 (1H, d, J 15.2 Hz), 4.77 (1H, d, J 14.5 Hz), 4.69 (1H, d, J 14.5 Hz), 4.61 (1H, d, J 15.2 Hz), 3.94 (1H, d, J 4.8 Hz), 1.76 (2H, m), 1.54 (3H, s), 1.25 (3H, s), 1.13 (3H, s) ppm. $^{13}\text{C}\{^1\text{H}\}$ DEPT NMR (d_6 -DMSO, 75.6 MHz) δ 154.5 (s), 140–127 (aromatics), 71.4 (s), 66.2 (s), 53.1 (s), 48.1 (s), 42.0 (s), 38.5 (s), 31.8 (s), 21.7 (s), 16.4 (s), 15.0 (s) ppm. ^{31}P (CD_3CN , 121.6 MHz): δ 24.7 (AB, $^2J_{\text{P-P}}$ 542 Hz), 22.3 (AB, $^2J_{\text{P-P}}$ 542 Hz) ppm. MS: 879 ($[\text{M}^+]$, 20%). *Anal.*: Calc. for $\text{C}_{47}\text{H}_{47}\text{N}_2\text{P}_3\text{Cl}_2\text{F}_6\text{Pd}$: C, 55.11; H, 4.63; N, 2.74%. Found: C, 55.2; H, 4.5; N, 2.6%.

trans-[Pt(κ^2 -1-H)Cl₂]PF₆, 3a. A mixture of K_2PtCl_4 (100 mg, 2.40×10^{-4} mol) and **1** (204 mg, 2.40×10^{-4} mol) in ethanol (15 ml) was refluxed for 24 h under N_2 . After cooling, the cream precipitate was filtered in air, washed with EtOH (2×3 ml) and air-dried. Yield = 217 mg (81%). The complex was treated with CHCl_3 (2×15 ml) to give a soluble fraction and an insoluble white solid (yield = 95 mg) **3b** (see below). The soluble *trans*-complex **3a** was recovered as a pale yellow solid upon removal of the CHCl_3 (yield = 122 mg). The complex can be recrystallised from MeCN as colourless crystals. ^1H NMR (CD_3CN , 250 MHz) δ 9.90 (1H, s br), 8.1–7.1 (28H, m), 5.12 (1H, d, J 15.5 Hz), 4.70 (3H, m), 4.00 (1H, br), 1.95 (2H, m), 1.62 (3H, s), 1.34 (3H, s), 1.19 (3H, s br) ppm. $^{13}\text{C}\{^1\text{H}\}$ DEPT NMR (d_6 -DMSO, 75.6 MHz) δ 154.5 (s), 140–126 (aromatics), 71.9 (s), 67.0 (s), 56.2 (s), 48.2 (s), 42.1 (s), 39.0 (s), 32.6 (s), 22.4 (s), 17.1 (s), 15.7 (s) ppm. ^{31}P (CD_3CN , 121.6 MHz): δ 20.0 ($^1J_{\text{P-Pt}}$ 2523 Hz), 19.7 ($^1J_{\text{P-Pt}}$ 2526 Hz) ppm. MS: 967 ($[\text{M}^+]$, 100%). *Anal.*: Calc. for $\text{C}_{47}\text{H}_{47}\text{N}_2\text{P}_3\text{Cl}_2\text{F}_6\text{Pt}$: C, 50.72; H, 4.27; N, 2.52%. Found: C, 50.8; H, 4.2; N, 2.6%.

trans-[Pt₂(μ -1-H)₂Cl₄](PF₆)₂, 3b. Compound **3b** was isolated during the preparation of **3a** (see above). ^1H NMR (d_6 -DMSO, 80 °C, 250 MHz) δ 8.31 (1H, s), 8.1–6.9 (28H, m), 5.80 (1H, d, J 16.3 Hz), 5.46 (2H, br), 4.77 (1H, d, J 16.3 Hz), 4.07 (1H, br), 1.58 (2H, m), 1.26 (1H, obs), 1.25 (3H, s), 0.83 (3H, s), 0.17 (3H, s br) ppm. ^{31}P (d_6 -DMSO, 80 °C, 121.6 MHz): δ 13.5 ($^1J_{\text{P-Pt}}$ 2556 Hz) ppm. MS: 967 ($[\text{M}^+]$, 100%). *Anal.*: Calc. for $\text{C}_{94}\text{H}_{94}\text{N}_4\text{P}_4\text{Cl}_4\text{F}_{12}\text{Pt}_2$: C, 50.72; H, 4.27; N, 2.52%. Found: C, 50.1; H, 4.0; N, 2.5%.

[Rh₂(COD)₂(μ -1-H)Cl₂]PF₆, 4. A mixture of $[\text{Rh}(\text{1,5-COD})\text{Cl}]_2$ (43 mg, 8.9×10^{-5} mol) and **1** (150 mg, 1.77×10^{-4} mol) in MeOH (12 ml) was heated close to reflux for 18 h under N_2 . After cooling, the yellow precipitate was filtered in air, washed with MeOH (2×3 ml) and air-dried. Yield = 130 mg (67%).

^1H NMR (CDCl_3 , 250 MHz) δ 8.18 (1H, s), 7.8–7.0 (28H, m), 6.42 (1H, d, J 16.5 Hz), 6.04 (1H, d, J 15.6 Hz), 5.78 (1H, d, J 14.9 Hz), 5.68 (1H, d, J 16.1 Hz), 5.42 (4H, m br), 4.04 (1H, d, J 4.2 Hz), 3.7–3.3 (4H, m br), 2.97 (1H, m), 2.55–1.85 (20H, m), 1.52 (3H, s), 1.47 (3H, s), 1.16 (3H, s br) ppm. ^{31}P (CDCl_3 , 121.6 MHz): δ 21.7 (d, $^1J_{\text{P-Rh}}$ 146 Hz), 21.4 (d, $^1J_{\text{P-Rh}}$ 146 Hz) ppm. MS: 874 ($[\text{Rh}(\text{1-H})\text{Cl}_2]^+$, 30%). *Anal.*: Calc. for $\text{C}_{63}\text{H}_{71}\text{N}_2\text{P}_3\text{Cl}_2\text{F}_6\text{Rh}_2$: C, 56.47; H, 5.35; N, 2.09%. Found: C, 56.4; H, 5.5; N, 2.2%.

[Pt₂(dvdms)₂(μ -1-H)]PF₆, 5. To a solution of Pt(dvdms) (0.18 ml of a 20% solution, 1.85×10^{-4} mol Pt) in THF (5 ml) was added a solution of **1**·HPF₆ (160 mg, 1.85×10^{-4} mol) in THF (5 ml) and the solution stirred for 24 h. The solvent was removed *in vacuo*, and the cream solid dissolved in toluene (5 ml) filtered and taken to dryness at the pump. The solid was washed with pentane to remove traces of dvdms and the desired compound isolated as a cream solid which was crystallised from toluene/Et₂O by vapour diffusion. Yield = 176 mg (59%). ^1H NMR (CDCl_3 , 300 MHz) δ 8.07 (1H, s br), 7.75–7.05 (28H, m), 4.98 (1H, d, J 16.7 Hz), 4.83 (1H, d, J 15.6 Hz), 4.64 (1H, d, J 15.6 Hz), 4.54 (1H, d, J 16.7 Hz), 2.65 (6H, m), 2.20 (6H, m), 1.85 (2H, m), 1.56 (2H, m), 0.97 (3H, s), 0.80 (3H, s), 0.60 (3H, s), 0.39 (3H, s), 0.38 (3H, s), 0.37 (6H, s), -0.05 (3H, s), -0.09 (6H, s), -0.11 (3H, s) ppm. $^{13}\text{C}\{^1\text{H}\}$ DEPT NMR (CDCl_3 , 75.6 MHz) δ 156.0 (s), 138–125 (aromatics), 71.5 (s), 61.2 (s), 54.3 (d, J 12.7 Hz), 53.8 (s), 52.4 (s), 51.4 (s), 51.1 (s), 50.4 (s), 48.4 (d, J 10.4 Hz), 47.6 (s), 46.7 (d, J 12.8 Hz), 41.0 (s), 38.3 (s), 31.3 (s), 21.5 (s), 17.7 (s), 13.6 (s), 1.4 (s), 0.99 (s), 0.98 (s), -1.72 (s), -1.75 (s), -1.79 (s) ppm. ^{31}P NMR (C_7D_8 , 121.6 MHz): δ 20.7 (s, $^1J_{\text{P-Pt}}$ 3494 Hz), 18.7 (s, $^1J_{\text{P-Pt}}$ 3506 Hz) ppm. ^{195}Pt NMR (CDCl_3 , 106.9 MHz): δ -5553 (dm, $^1J_{\text{P-Pt}}$ 3506 Hz), -5563 (dm, $^1J_{\text{P-Pt}}$ 3494 Hz) ppm. MS: 879 ($[\text{M}^+]$, 20%). *Anal.*: Calc. for $\text{C}_{63}\text{H}_{83}\text{N}_2\text{Si}_4\text{O}_2\text{P}_3\text{F}_6\text{Pt}_2$ (C_7H_8)_{0.5}: C, 48.23; H, 5.31; N, 1.69%. Found: C, 48.4; H, 5.7; N, 1.5%.

trans-[Pd₂(μ -1-H)₂I₄](PF₆)₂, 6. A solution of **2** (30 mg, 2.9×10^{-5} mol) and MeI (21 mg, 1.5×10^{-4} mol) in CHCl_3 (1 ml) was heated at 50 °C overnight. On return, the orange precipitate was filtered off and recrystallised from MeCN to give red crystals suitable for single-crystal X-ray crystallography. Yield = 20 mg (57%). Dissolution of the solid in CD_3CN gave complex $^{31}\text{P}\{^1\text{H}\}$ and ^1H NMR spectra indicative of mixtures of species in solution. MS: 1062 ($[\text{Pd}(\text{1-H})\text{I}_2]^+$, 15%). *Anal.*: Calc. for $\text{C}_{94}\text{H}_{94}\text{N}_4\text{P}_4\text{I}_4\text{Pd}_2\text{I}_{0.27}(\text{PF}_6)_{1.73}$: C, 46.86; H, 3.94; N, 2.33%. Found: C, 46.7; H, 4.0; N, 2.4%.

trans-[Pt₂(μ -1-H)₂I₄](PF₆)₂, 7. Prepared as described for **6** to give an orange solid. Yield = 43%. Dissolution of the solid in CD_3CN gave complex $^{31}\text{P}\{^1\text{H}\}$ and ^1H NMR spectra indicative of mixtures of species in solution. MS: 1151 ($[\text{Pt}(\text{1-H})\text{I}_2]^+$, 100%). *Anal.*: Calc. for $\text{C}_{94}\text{H}_{94}\text{N}_4\text{P}_4\text{I}_4\text{Pt}_2$: C, 43.56; H, 3.66; N, 2.16%. Found: C, 44.1; H, 3.8; N, 2.2%.

[Pd(κ^3 -1)Cl]PF₆, 8. A mixture of *trans*-[Pd(κ^2 -1-H)Cl₂]PF₆, **2** (200 mg, 1.95×10^{-4} mol) and KO^tBu (26 mg, 2.34×10^{-4} mol) in THF (10 ml) was stirred for 24 h under N_2 to give a blood-red solution. This was filtered, taken to dryness and the red residue extracted into DCM (10 ml). The solubles were



filtered away from a small amount of insoluble material and the volatiles removed *in vacuo*. The residue was dissolved in the minimum amount of THF and left to precipitate at 4 °C. The pink solid was isolated and dried *in vacuo*. Yield = 50 mg (26%). A slightly less pure material was obtained upon removal of the THF. Yield = 134 mg (69%). ^1H NMR (CDCl_3 , 250 MHz) δ 8.0–7.0 (28H, m), 6.68 (0.5H, d, J 14.5 Hz), 6.48 (0.5H, d, J 14.5 Hz), 6.36 (0.5H, d, J 15.6 Hz), 6.30 (0.5H, d, J 15.6 Hz), 4.22 (0.5H, d, J 15.6 Hz), 4.19 (0.5H, d, J 15.6 Hz), 4.15 (0.5H, d, J 14.5 Hz), 4.02 (0.5H, d, J 14.7 Hz), 3.11 (0.5H, d, J 4.2 Hz), 2.98 (0.5H, d, J 4.4 Hz), 1.51 (1H, m), 1.29 (1H, m), 1.15 (1.5H, s), 0.98 (1.5H, s), 0.75 (1H, m), 0.67 (1.5H, s), 0.65 (1.5H, s), 0.10 (1H, m), –0.10 (1.5H, s), –0.12 (1.5H, s) ppm. $^{13}\text{C}\{^1\text{H}\}$ NMR (CDCl_3 , 75.6 MHz) δ 178.2 (s), 177.2 (s), 144–125 (aromatics), 72.1 (s), 71.8 (s), 69.5 (s), 60.4 (d, J 9.6 Hz), 59.2 (d, J 9.7 Hz), 55.3 (d, J 9.7 Hz), 54.7 (d, J 7.0 Hz), 40.6 (s), 40.3 (s), 38.5 (s), 31.3 (s), 31.2 (s), 21.8 (s), 21.3 (s), 17.3 (s), 16.8 (s), 16.1 (s), 15.9 (s) ppm. ^{31}P (CDCl_3 , 121.6 MHz): δ 6.7 (AB, $^2J_{\text{P-P}}$ 436 Hz), 3.2 (AB, $^2J_{\text{P-P}}$ 436 Hz), 6.1 (AB, $^2J_{\text{P-P}}$ 435 Hz), 2.7 (AB, $^2J_{\text{P-P}}$ 435 Hz) ppm. MS: 832 ($[\text{M}^+] - \text{Cl}^- + \text{CN}^-$, 100%). *Anal.*: Calc. for $\text{C}_{47}\text{H}_{46}\text{N}_2\text{P}_3\text{-ClF}_6\text{Pd}$: C, 57.15; H, 4.70; N, 2.84%. Found: C, 57.3; H, 4.8; N, 2.9%.

[Pt(κ^3 -1)Cl]PF₆, 9. A mixture of *cis/trans*-[Pt(κ^2 -1-H)Cl₂]-PF₆, **3a/b** (200 mg, 1.80×10^{-4} mol) and KO^tBu (23 mg, 2.07×10^{-4} mol) in THF (10 ml) was stirred for 24 h under N₂. The mixture was filtered, and the filtrate left in air to slowly evaporate to small volume whereupon a colourless solid precipitated. The solid was isolated and dried in air. Yield = 98 mg (51%). A slightly less pure material was obtained upon removal of the remaining THF. Yield = 70 mg (36%). ^1H NMR (CDCl_3 , 300 MHz) δ 7.9–7.0 (28H, m), 6.88 (0.5H, d, J 14.5 Hz), 6.77 (0.5H, d, J 15.8 Hz), 6.67 (0.5H, d, J 15.8 Hz), 4.15 (1.5H, m), 3.98 (0.5H, d, J 15.0 Hz), 3.16 (0.5H, d, J 4.6 Hz), 3.03 (0.5H, d, J 4.4 Hz), 2.00 (0.5H, m), 1.54 (1H, m), 1.32 (1H, m), 1.17 (1.5H, s), 1.10 (0.5H, m), 0.97 (1.5H, s), 0.83 (0.5H, m), 0.70 (1.5H, s), 0.67 (1.5H, s), 0.21 (0.5H, m), –0.08 (1.5H, s), –0.12 (1.5H, s) ppm. $^{13}\text{C}\{^1\text{H}\}$ NMR (CD_3CN , 75.6 MHz) δ 75.6 MHz) δ 166.2 (t, J 9.2 Hz), 164.9 (t, J 9.2 Hz), 145–125 (aromatics), 73.8 (s), 71.3 (s), 68.3 (s), 68.2 (s), 60.2 (d, J 6.9 Hz), 58.8 (d, J 8.1 Hz), 55.3 (d, J 6.9 Hz), 54.3 (d, J 8.1 Hz), 41.0 (s), 40.8 (s), 38.8 (s), 32.9 (s), 31.2 (s), 22.1 (s), 21.4 (s), 17.5 (s), 16.6 (s), 16.3 (s) ppm. $^{31}\text{P}\{^1\text{H}\}$ NMR (CDCl_3 , 121.6 MHz): δ 5.27 (AB, $^2J_{\text{P-P}}$ 410 Hz, $^1J_{\text{P-Pt}}$ 2552 Hz), 2.97 (AB, $^2J_{\text{P-P}}$ 410 Hz, $^1J_{\text{P-Pt}}$ 2560 Hz), 4.98 (AB, $^2J_{\text{P-P}}$ 414 Hz, $^1J_{\text{P-Pt}}$ 2552), 1.02 (AB, $^2J_{\text{P-P}}$ 414 Hz, $^1J_{\text{P-Pt}}$ 2570 Hz) ppm. MS: 832 ($[\text{M}^+] - \text{Cl}^- + \text{CN}^-$, 100%). *Anal.*: Calc. for $\text{C}_{47}\text{H}_{46}\text{N}_2\text{P}_3\text{ClF}_6\text{Pt}$: C, 52.44; H, 4.32; N, 2.60%. Found: C, 51.9; H, 4.1; N, 2.6%.

[Rh(κ^3 -1)(CO)]PF₆, 10. A mixture of Rh(acac)(CO)₂ (60 mg, 2.36×10^{-4} mol) and **1-HPF₆** (200 mg, 2.36×10^{-4} mol) in THF (10 ml) was heated near reflux for 2 h. After cooling, the yellow solution was taken to dryness and the residue crystallised from hot MeOH in air to give an orange-yellow solid which was filtered, washed sparingly with cold MeOH (2 $\times$ 1 ml) and air-dried. Yield = 189 mg (82%). ^1H NMR (CD_3CN , 500 MHz) δ 8.1–7.1 (28H, m), 6.77 (0.5H, d, J 14.6 Hz), 6.65 (0.5H, d, J 14.4 Hz), 6.42 (0.5H, d, J 15.6 Hz), 6.35 (0.5H, d, J 15.4 Hz), 4.13 (0.5H, d, J 15.6 Hz), 4.06 (0.5H, d, J 15.4 Hz), 3.84 (0.5H,

d, J 14.6 Hz), 3.64 (0.5H, d, J 14.4 Hz), 2.88 (0.5H, d, J 4.6 Hz), 2.74 (0.5H, d, J 4.0 Hz), 1.43 (1H, m), 1.20 (1H, m), 1.06 (1.5H, s), 0.90 (1.5H, s), 0.77 (1H, m), 0.62 (1.5H, s), 0.60 (1.5H, s), 0.49 (1H, m), –0.18 (1.5H, s), –0.21 (1.5H, s) ppm. $^{13}\text{C}\{^1\text{H}\}$ NMR (CD_3CN , 75.6 MHz) δ 196.6 (br), 196.2 (br), 195.2 (br), 194.4 (br), 145–128 (aromatics), 72.4 (s), 70.1 (s), 68.2 (s), 61.4 (d, J 9.7 Hz), 59.6 (d, J 9.7 Hz), 56.8 (d, J 9.7 Hz), 55.6 (d, J 6.9 Hz), 41.2 (s), 39.2 (s), 33.3 (s), 31.2 (s), 22.1 (s), 21.5 (s), 17.8 (s), 16.9 (s), 16.2 (s), 15.9 (s) ppm. ^{31}P (CD_3CN , 121.6 MHz): δ 16.1 (ABX, $^2J_{\text{P-P}}$ 268 Hz, $^1J_{\text{P-Rh}}$ 134 Hz), 13.0 (ABX, $^2J_{\text{P-P}}$ 268 Hz, $^1J_{\text{P-Rh}}$ 134 Hz), 16.0 (ABX, $^2J_{\text{P-P}}$ 274 Hz, $^1J_{\text{P-Rh}}$ 134 Hz), 13.1 (ABX, $^2J_{\text{P-P}}$ 274 Hz, $^1J_{\text{P-Rh}}$ 134 Hz) ppm. MS: 831 ($[\text{M}^+]$, 100%). IR (KBr): 2009 cm^{-1} (C $\equiv$ O). *Anal.*: Calc. for $\text{C}_{48}\text{H}_{46}\text{N}_2\text{P}_3\text{OF}_6\text{Rh}$: C, 59.02; H, 4.76; N, 2.87%. Found: C, 58.9; H, 4.8; N, 3.0%.

S_{Ag}-[Ag(κ^3 -1)(OTf)], 11. To a suspension of **1-HPF₆** (150 mg, 1.77×10^{-4} mol) and AgOTf (45 mg, 1.77×10^{-4} mol) in THF (10 ml) was added KO^tBu (26 mg, 2.30×10^{-4} mol) whereupon the mixture darkened immediately. The mixture was stirred for 2 h in the absence of light, filtered to remove some fine solid and the volatiles removed *in vacuo*. The grey residue was dissolved in a small volume of CH₂Cl₂ (4 ml) and again filtered to remove a slight solid. The volatiles were removed at the pump and the residue triturated with toluene to give the desired product as a grey solid which was isolated by filtration under nitrogen. Yield = 131 mg (78%). ^1H NMR (CDCl_3 , 300 MHz) δ 7.7–6.7 (28H, m), 5.15 (1H, d, J 12.8 Hz), 4.84 (1H, d, J 15.8 Hz), 4.38 (1H, d, J 15.8 Hz), 4.10 (1H, d, J 12.8 Hz), 3.48 (1H, d, J 4.0 Hz), 2.28 (1H, m), 1.98 (2H, m), 1.47 (3H, s), 1.02 (3H, s), 0.83 (1H, m), 0.44 (3H, s br) ppm. $^{13}\text{C}\{^1\text{H}\}$ DEPT NMR (CDCl_3 , 75.6 MHz) δ 211.0 (v br), 142.8 (d, J 17.3 Hz), 140.8 (d, J 18.5 Hz), 135–127 (aromatics), 69.3, 68.2, 58.2, 53.0, 41.3, 40.3, 31.9, 22.2, 18.4, 15.4 ppm. ^{31}P (CDCl_3 , 121.6 MHz): δ 7.0 (ABX, $^2J_{\text{P-P}}$ 65.5 Hz, $^1J_{\text{P-Ag(107)}}$ 327 Hz, $^1J_{\text{P-Ag(109)}}$ 378 Hz), 4.1 (ABX, $^2J_{\text{P-P}}$ 65.5 Hz, $^1J_{\text{P-Ag(107)}}$ 327 Hz, $^1J_{\text{P-Ag(109)}}$ 378 Hz) ppm. MS: 845 ($[\text{M}^+] + \text{HCl}$, 80%), 809 ($[\text{M}^+]$, 10%). *Anal.*: Calc. for $\text{C}_{47}\text{H}_{46}\text{N}_2\text{P}_2\text{ClAg}$: C, 66.86; H, 5.50; N, 3.32%. Found: C, 66.0; H, 5.3; N, 3.1%.

Catalysis

Suzuki coupling

A mixture of substrate (1 mmol), PhB(OH)₂ (1.2 mmol), K₂CO₃ (2 mmol), 0.5 mol% catalyst in 1,4-dioxane (3 ml) with decane as internal GC standard (1 mmol) was heated at 80 °C for 18 h. A sample of the solution was diluted with CH₂Cl₂ and passed through a short column of silica. Total GC-MS yields based upon consumption of substrate using *n*-decane as internal standard were determined *via* an internal standard method of quantification on an Agilent Technologies 6890 N GC system equipped with an Agilent Technologies 5973 inert MS detector with MSD. Column: Agilent 190915-433 capillary, 0.25 mm $\times$ 30 m $\times$ 0.25 mm. Program for analysis: initial temperature of 40 °C, held for 2.5 min, ramp 5 °C min^{–1} next 150 °C, ramp 10 °C min^{–1} next 220 °C, held for 10 min. The temperature of both the injector and detector were maintained at 240 °C.



Table 3 Details of X-ray crystallographic data collection for the compounds **1**·HPF₆, **2**, **3a** and **6**

	1 ·HPF ₆	2	3a	6
Empirical formula	C ₄₇ H ₄₇ F ₆ N ₂ P ₃	C ₄₈ H ₄₈ Cl ₅ N ₂ P ₃ F ₆ Pd	C ₅₁ H ₅₃ Cl ₂ N ₄ P ₃ F ₆ Pt	C ₉₄ H ₉₄ I ₄ N ₄ P ₄ Pd ₂ , I _{0.27} (PF ₆) _{1.73}
Formula weight	846.78	1143.44	1194.87	2409.12
Crystal system	Triclinic	Monoclinic	Monoclinic	Orthorhombic
Space group	<i>P</i> 1	<i>C</i> 2	<i>P</i> 21	<i>P</i> 2(1)2(1)2
<i>a</i> /Å	11.9727(5)	26.7720(9)	11.9538(5)	27.8377(9)
<i>b</i> /Å	13.6776(6)	14.6590(6)	14.4075(4)	19.8686(12)
<i>c</i> /Å	13.8349(5)	28.2434(13)	29.2856(14)	21.6459(13)
β /°	75.759(2)	117.031(2)	93.809(2)	
<i>U</i> /Å ³	2112.19(15)	9873.3(7)	5032.6(3)	11 972.3(11)
<i>Z</i>	2	8	4	4
<i>D_c</i> /Mg m ⁻³	1.331	1.498	—	1.337
<i>F</i> (000)	884	4524	2392	4735
θ range/°	3.04 to 27.50	2.25 to 24.71	1.80 to 27.54	1.19 to 20.82
Index ranges	−14 ≤ <i>h</i> ≤ 15, −17 ≤ <i>k</i> ≤ 17, −17 ≤ <i>l</i> ≤ 17	−31 ≤ <i>h</i> ≤ 31, −17 ≤ <i>k</i> ≤ 15, −33 ≤ <i>l</i> ≤ 33	−15 ≤ <i>h</i> ≤ 15, −18 ≤ <i>k</i> ≤ 18, −38 ≤ <i>l</i> ≤ 37	−25 ≤ <i>h</i> ≤ 24, −16 ≤ <i>k</i> ≤ 19, −21 ≤ <i>l</i> ≤ 14
Reflections collected	14 999	15 228	19 515	12 167
Independent reflections	10 801	9398	11 357	8498
Data/restraints/parameters	14 999/606/1179	15 228/265/1218	19 515/173/1217	12 167/40/1145
Goodness of fit on <i>F</i> ²	1.047	1.016	1.026	1.061
Final <i>R</i> ₁ , <i>wR</i> ₂ [<i>I</i> > 2σ(<i>I</i>)] (all data)	0.0631, 0.1297 0.1009, 0.1500	0.0853, 0.1835 0.1530, 0.2224	0.0976, 0.1597 0.1889, 0.1947	0.1023, 0.2721 0.1281, 0.2961
Largest difference peak And hole/e Å ⁻³	0.284 and −0.321	1.346, −0.911	1.514, −1.057	3.294, −1.076
Flack parameter	0.03(12)	0.09(6)	0.049(14)	0.20(5)

Hydrosilylation

A mixture of substrate (1 mmol), Et₃SiH (1 mmol), and 0.05 mol% catalyst in toluene (3 ml) with decane as internal GC standard (1 mmol) was heated at 70 °C for 12 h. A sample of the solution was diluted with CH₂Cl₂ and passed through a short column of silica. Total GC-MS yields based upon consumption of substrate using *n*-decane as internal standard were determined via an internal standard method of quantification on an Agilent Technologies 6890 N GC system equipped with an Agilent Technologies 5973 inert MS detector with MSD. Column: Agilent 190915-433 capillary, 0.25 mm × 30 m × 0.25 mm. Program for analysis: initial temperature of 40 °C, held for 2.5 min, ramp 5 °C min⁻¹ next 150 °C, ramp 10 °C min⁻¹ next 220 °C, held for 10 min. The temperature of both the injector and detector were maintained at 240 °C. Selectivities were determined by ¹H NMR on the isolated materials.

Crystallography

Data collection was carried out on a Bruker–Nonius Kappa CCD diffractometer using graphite monochromated Mo Kα radiation (λ = 0.71073 Å). The instrument was equipped with an Oxford Cryosystems cooling apparatus. Data collection and cell refinement were carried out using COLLECT³² and HKL SCALEPACK.³³ Data reduction was applied using HKL DENZO and SCALEPACK.³³ The structures were solved and refined with SHELX-97.³⁴ Absorption corrections were performed using SORTAV.³⁵ Generally, all non-hydrogen atoms were generally refined anisotropically, while the hydrogen atoms were inserted in idealised positions with *U*_{iso} set at 1.2 or 1.5 times the *U*_{eq} of the parent atom. For **2**, the large asymmetric unit is partially accountable for the weak high angle data and two anion sites

show two-fold disorder. Geometric restraints were applied to the PF₆[−] during refinement to stop distortion of the anions and restraints were also applied in order to maintain similar atomic displacement parameters for atoms in close proximity. For **3a**, geometric restraints were applied to acetonitrile solvent molecules as well as to the atomic displacement parameters to stop distortion. For **6**, difference Fourier synthesis showed large spaces occupied by diffuse electron density, interpreted as disordered solvent, and hence SQUEEZE was used.³⁶ The large asymmetric unit combined with the disordered solvent are accountable for the weak high angle data which were omitted from the refinement. Geometric restraints were applied to the PF₆[−] during refinement to stop distortion of the anions. Initial examination of the data for **1**·HPF₆ and **2** suggested higher crystal symmetry but lower symmetry was necessary for the final ordered complex structures. Both anion sites show two-fold disorder. Geometric restraints were applied to the PF₆[−] during refinement to stop distortion of the anions. Restraints were also applied in order to maintain similar atomic displacement parameters for atoms in close proximity. The details of the data collection and structure solution are collected in Table 3.

Conclusions

A diphosphino(pro-NHC) ligand has been prepared containing a central bicyclic core derived from camphor and two closely similar but formally unique RPPH₂ arms. In its pro-ligand form (protonated NHC) the ligand binds solely through the phosphine donors and, depending on the metal ion, forms *trans*-chelating monomeric or bridging dimeric complexes. Addition of base to these preformed complexes liberates the NHC for coordination to give κ^3 -*P,C,P'* complexes which, for square planar metals, show conformational isomerism. Prior deprotonation of the pro-



ligand and subsequent addition of AgOTf gives the pseudo-tetrahedral complex $S_{Ag}-[Ag(\kappa^3-P,C,P'-I)(OTf)]$ which is isolated as a single isomer. Selected platinum and palladium complexes have been shown to be effective for the regioselective hydrosilylation of alkynes and for Suzuki coupling with aryl bromides. Studies of this and related ligands are continuing with a view to extending their application in catalysis.

Acknowledgements

The authors would like to thank Cardiff University for providing support for this work.

Notes and references

- (a) P. W. N. M. van Leeuwen and Z. Freixa, in *Modern Carbonylation Methods*, ed. L. Kollár, Wiley-VCH, 2008, p. 1; (b) P. C. J. Kamer, P. W. N. M. Van Leeuwen and J. N. H. Reek, *Acc. Chem. Res.*, 2001, **34**, 895; (c) Z. Freixa and P. W. N. M. Van Leeuwen, *J. Chem. Soc., Dalton Trans.*, 2003, 1890; (d) P. C. J. Kamer, P. W. N. M. van Leeuwen, J. F. Kees Goubitz, H. Schenk and C. Bo, *J. Am. Chem. Soc.*, 1998, **120**, 11616; (e) B. P. Morgan and R. C. Smith, *J. Organomet. Chem.*, 2008, **693**, 11; (f) E. E. Marlier, S. J. Terenaki, K. Ding, J. E. Mulliken and C. C. Lu, *Inorg. Chem.*, 2011, **50**, 9290.
- (a) Z. Freixa and P. W. N. M. van Leeuwen, *Coord. Chem. Rev.*, 2008, **252**, 1755; (b) N. Naser, D. J. Eisler and R. J. Puddephat, *Chem. Commun.*, 2010, **46**, 1953; (c) C. P. Butts, E. Filali, G. C. Lloyd-Jones, P.-O. Norrby, D. A. Sale and Y. Schramm, *J. Am. Chem. Soc.*, 2009, **131**, 9945; (d) C. P. Butts, J. Crosby, G. C. Lloyd-Jones and S. C. Stephen, *Chem. Commun.*, 1999, 1707; (e) I. J. S. Fairlamb and G. C. Lloyd-Jones, *Chem. Commun.*, 2000, 2447; (f) B. M. Trost and F. D. Toste, *J. Am. Chem. Soc.*, 1999, **121**, 4545; (g) L. Kaganovsky, D. Gelman and K. Rueck-Braun, *J. Organomet. Chem.*, 2010, **695**, 260; (h) A. Velian, S. Lin, A. J. M. Miller, M. W. Day and T. Agapie, *J. Am. Chem. Soc.*, 2010, **132**, 6296; (i) C. Azerraf, O. Grossman and D. Gelman, *J. Organomet. Chem.*, 2007, **692**, 761; (j) C. Azerraf and D. Gelman, *Organometallics*, 2009, **28**, 6578; (k) I. S. Mikhel, M. Garland, J. Hopewell, S. Mastroianni, C. L. McMullin, A. G. Orpen and P. G. Pringle, *Organometallics*, 2011, **30**, 974.
- (a) N. J. Destefano, D. K. Johnson, R. M. Lane and L. M. Venanzi, *Helv. Chim. Acta*, 1976, **59**, 2674; (b) Z. Freixa, M. S. Beentjes, G. D. Batema, C. B. Dieleman, G. P. F. van Strijdonck, J. N. H. Reek, P. C. J. Kamer, J. Fraanje, K. Goubitz and P. W. N. M. van Leeuwen, *Angew. Chem., Int. Ed.*, 2003, **42**, 1284; (c) F. B. Xu, Q. S. Li, X. S. Zeng, X. B. Leng and Z. Z. Zhang, *Organometallics*, 2002, **21**, 4894; (d) M. Sawamura, H. Hamashima and Y. Ito, *Tetrahedron: Asymmetry*, 1991, **2**, 593.
- (a) L. Chahen, B. Therrien and G. Süß-Fink, *J. Organomet. Chem.*, 2006, **691**, 4257; (b) G. Lamb, M. Clarke, A. M. Z. Slawin, B. Williams and L. Key, *Dalton Trans.*, 2007, 5582; (c) E. K. van den Beuken, A. Meetsma, H. Kooijman, A. L. Spek and B. L. Feringa, *Inorg. Chim. Acta*, 1997, **264**, 171; B. M. Trost, D. L. van Vranken and C. Brigel, *J. Am. Chem. Soc.*, 1992, **114**, 9327.
- (a) A. S. Goldman, A. H. Roy, Z. Huang, R. Ahuja, W. Schinski and M. Brookhart, *Science*, 2006, **312**, 257; (b) E. M. Schuster, M. Botoshansky and M. Gandelman, *Angew. Chem., Int. Ed.*, 2008, **47**, 4555; (c) E. Poverenov, I. Efremenko, A. I. Frenkel, Y. Ben-David, L. J. W. Shimon, G. Leitun, L. Konstantinovski, J. M. L. Martin and D. Milstein, *Nature*, 2008, **455**, 1093; (d) R. Ahuja, B. Punji, M. Findlater, C. Supplee, W. Schinski, M. Brookhart and A. S. Goldman, *Nat. Chem.*, 2011, **3**, 167; (e) M. van der Boom and D. Milstein, *Chem. Rev.*, 2003, **103**, 1759; (f) S. Musa, R. Romm, C. Azerraf, S. Kozhuz and D. Gelman, *Dalton Trans.*, 2011, **40**, 8760.
- (a) M. Albrecht and G. van Koten, *Angew. Chem., Int. Ed.*, 2001, **40**, 3750; (b) W. Leis, H. A. Mayer and W. C. Kaska, *Coord. Chem. Rev.*, 2008, **252**, 1787; (c) R. H. Crabtree, *J. Organomet. Chem.*, 2004, **689**, 4083; (d) D. F. Maclean, R. McDonald, M. J. Ferguson, A. J. Caddell and L. Turculet, *Chem. Commun.*, 2008, 5146; (e) M. Lersch and M. Tilset, *Chem. Rev.*, 2005, **105**, 2471; (f) M. Findlater, A. S. Goldman and M. S. Brookhart, *Alkane C-H activation by single-site metal catalysis*, in *Catalysis by Metal Complexes*, ed. P. J. Perez, Springer, New York, 2011, ch. 4, vol. 38; (g) J. Choi, A. MacArthur, M. S. Brookhart and A. S. Goldman, *Chem. Rev.*, 2011, **111**, 1761; (h) B. Punji, T. J. Emge and A. S. Goldman, *Organometallics*, 2010, **29**, 2702; (i) C. M. Jensen, *Chem. Commun.*, 1999, 2443.
- (a) M. D. Fryzuk and C. D. Montgomery, *Coord. Chem. Rev.*, 1989, **95**, 1; (b) G. van Koten, *Pure Appl. Chem.*, 1989, **61**, 1681; (c) A. Choualeb, A. J. Lough and D. G. Gusev, *Organometallics*, 2007, **26**, 5224; (d) C. Gunanathan, B. Gnanaprakasam, M. A. Iron, L. J. W. Shimon and D. Milstein, *J. Am. Chem. Soc.*, 2010, **132**, 14763; (e) R. Langer, G. Leitun, Y. Ben-David and D. Milstein, *Angew. Chem., Int. Ed.*, 2011, **50**, 1947; (f) M. J. Sgro and D. W. Stephan, *Dalton Trans.*, 2011, **40**, 2419; (g) A. A. Koridze, A. V. Polezhaev, S. V. Safronov, A. M. Sheloumov, F. M. Dolgushin, M. G. Ezemitskaya, B. V. Lokshin, P. V. Petrovskii and A. S. Peregudov, *Organometallics*, 2010, **29**, 4360.
- (a) A. M. Magill, D. S. McGuinness, K. J. Cavell, G. J. P. Britovsek, V. C. Gibson, A. J. P. White, D. J. Williams, A. H. White and B. W. Skelton, *J. Organomet. Chem.*, 2001, **617–618**, 546; (b) H. M. Lee, J. Y. Zeng, C. H. Hu and M. T. Lee, *Inorg. Chem.*, 2004, **43**, 6822; (c) F. E. Hahn, M. C. Jahnke and T. Pape, *Organometallics*, 2006, **25**, 5927; (d) T. Yagyu, S. Oya, M. Maeda and K. Jitsukawa, *Chem. Lett.*, 2006, **35**, 154; (e) S. Gischig and A. Togni, *Eur. J. Inorg. Chem.*, 2005, 4745; (f) J. C. C. Chen and I. J. B. Lin, *Organometallics*, 2000, **19**, 5113; (g) J. Y. Zeng, M.-H. Hsieh and H. M. Lee, *J. Organomet. Chem.*, 2005, **690**, 5662; (h) T. Steinke, B. K. Shaw, H. Jong, B. O. Patrick and M. D. Fryzuk, *Organometallics*, 2009, **28**, 2830; (i) L. P. Spencer, C. Beddie, M. B. Hall and M. D. Fryzuk, *J. Am. Chem. Soc.*, 2006, **128**, 12531; (j) L. P. Spencer and M. D. Fryzuk, *J. Organomet. Chem.*, 2005, **690**, 5788; (k) M. C. Jahnke, T. Pape and F. E. Hahn, *Eur. J. Inorg. Chem.*, 2009, 1960; (l) C. Michon, A. Ellern and R. J. Angelici, *Inorg. Chim. Acta*, 2006, **359**, 4549; (m) S. J. Roseblade, A. Ros, D. Monge, M. Alcarazo, E. Alvarez, J. M. Lassaletta and R. Fernandez, *Organometallics*, 2007, **26**, 2570.
- (a) D. Pugh, H. J. Wells, D. J. Evans and A. A. Danopoulos, *Dalton Trans.*, 2009, 7189; (b) A. A. Danopoulos, D. Pugh and J. A. Wright, *Angew. Chem., Int. Ed.*, 2008, **47**, 9765; (c) D. Pugh, A. Boyle and A. A. Danopoulos, *Dalton Trans.*, 2008, 1087; (d) A. A. D. Tulloch, A. A. Danopoulos, G. J. Tizzard, S. J. Coles, M. B. Hursthouse, R. S. Hay-Motherwell and W. B. Motherwell, *Chem. Commun.*, 2001, 1270; (e) E. Peris, J. Mata, J. A. Loch and R. H. Crabtree, *Chem. Commun.*, 2001, 201; (f) S. Grundemann, M. Albrecht, J. A. Loch, J. W. Fallner and R. H. Crabtree, *Organometallics*, 2001, **20**, 5485; (g) J. A. Loch, M. Albrecht, E. Peris, J. Mata, J. W. Fallner and R. H. Crabtree, *Organometallics*, 2002, **21**, 700; (h) D. J. Nielsen, K. J. Cavell, B. W. Skelton and A. H. White, *Inorg. Chim. Acta*, 2002, **327**, 116; (i) F. E. Hahn, M. C. Jahnke, V. Gomez-Benitez, D. Morales-Morales and T. Pape, *Organometallics*, 2005, **24**, 6458; (j) E. Mas-Marza, M. Poyatos, M. Sanau and E. Peris, *Organometallics*, 2004, **23**, 323; (k) G. T. S. Andavan, E. B. Bauer, C. S. Letko, T. K. Hollis and F. S. Tham, *J. Organomet. Chem.*, 2005, **690**, 5938; (l) H. V. Huynh, D. Yuan and Y. Han, *Dalton Trans.*, 2009, 7262; (m) D. S. McGuinness, J. A. Suttill, M. G. Gardiner and N. W. Davies, *Organometallics*, 2008, **27**, 4238; (n) D. J. Nielsen, K. J. Cavell, M. S. Viciu, S. P. Nolan, B. W. Skelton and A. H. White, *J. Organomet. Chem.*, 2005, **690**, 6133; (o) M. Raynal, C. S. J. Cazin, C. Vallée, H. Olivier-Bourbigou and P. Braunstein, *Chem. Commun.*, 2008, 3983; (p) A. R. Chianese, A. Mo, N. L. Lampland, R. L. Swartz and P. T. Bremer, *Organometallics*, 2010, **29**, 3019; (q) K. M. Schultz, K. I. Goldberg, D. G. Gusev and D. M. Heinekey, *Organometallics*, 2011, **30**, 1429.
- (a) R. Fränkel, U. Kernbach, M. Bakola-Christianopoulou, U. Plaia, M. Suter, W. Ponikvar, H. Nöth, C. Moinet and W. P. Fehlhammer, *J. Organomet. Chem.*, 2001, **617–618**, 530; (b) H. V. Rasika Dias and W. Jin, *Tetrahedron Lett.*, 1994, **35**, 1365; (c) H. Nakai, Y. Tang, P. Gantzel and K. Meyer, *Chem. Commun.*, 2003, 24; (d) X. Hu, I. Castro-Rodriguez and K. Meyer, *Organometallics*, 2003, **22**, 3016; (e) X. Hu and K. Meyer, *J. Organomet. Chem.*, 2005, **690**, 5474.
- (a) P. D. Newman, K. J. Cavell and B. M. Kariuki, *Organometallics*, 2010, **29**, 2724; (b) P. D. Newman, K. J. Cavell, A. J. Hallett and B. M. Kariuki, *Dalton Trans.*, 2011, **40**, 8807.
- R. C. Smith and J. D. Protasiewicz, *Organometallics*, 2004, **23**, 4215.
- L. Rigamonti, A. Forni, M. Manassero, C. Manassero and A. Pasini, *Inorg. Chem.*, 2010, **49**, 123.
- (a) C. J. Elsevier, B. Kowall and H. Kragten, *Inorg. Chem.*, 1995, **34**, 4836; (b) O. Niyomura, T. Iwasawa, N. Sawada, M. Tokunaga, Y. Obora and Y. Tsuji, *Organometallics*, 2005, **24**, 3468.



- 15 P. B. Hitchcock, M. F. Lappert, C. MacBeath, F. P. E. Scott and N. J. W. Warhurst, *J. Organomet. Chem.*, 1997, **534**, 139.
- 16 G. Berthon-Gelloz, O. Buisine, J.-F. Brière, G. Michaud, S. Stérin, G. Mignani, B. Tinant, J.-P. Declercq, D. Chapon and I. E. Markó, *J. Organomet. Chem.*, 2005, **690**, 6156.
- 17 T. Kösterke, T. Pape and F. E. Hahn, *J. Am. Chem. Soc.*, 2011, **133**, 2112.
- 18 A. Flores-Figueroa, T. Pape, K.-O. Feldmann and F. E. Hahn, *Chem. Commun.*, 2010, **46**, 324.
- 19 P. E. Garrou, *Chem. Rev.*, 1981, **81**, 229.
- 20 T. H. Brown and P. J. Green, *J. Am. Chem. Soc.*, 1970, **92**, 2359.
- 21 M. J. Doyle, M. F. Lappert, P. L. Pye and P. Terreros, *J. Chem. Soc., Dalton Trans.*, 1984, 2355.
- 22 (a) P. L. Arnold, M. Rodden and C. Wilson, *Chem. Commun.*, 2005, 1743; (b) C. Boehler, D. Stein, N. Donati and H. Grützmacher, *New J. Chem.*, 2002, **26**, 1291; (c) B. Cetinkaya, E. Cetinkaya, J. A. Chamizo, P. B. Hitchcock, H. A. Jasim, H. Küçükbay and M. E. Lappert, *J. Chem. Soc., Perkin Trans. 1*, 1998, 2047; (d) S. Solé, H. Gornitzka, O. Guerret and G. Bertrand, *J. Am. Chem. Soc.*, 1998, **120**, 9100.
- 23 P. L. Chiu and H. M. Lee, *Organometallics*, 2005, **24**, 1692.
- 24 P. D. Newman, K. J. Cavell and B. M. Kariuki, *Chem. Commun.*, 2012, **48**, 6511.
- 25 D. Affandi, S. J. Berners-Price, Effendy, P. J. Harvey, P. C. Healy, B. E. Ruch and A. H. White, *J. Chem. Soc., Dalton Trans.*, 1991, 1411.
- 26 (a) M. Iglesias, D. J. Beetstra, J. C. Knight, L.-L. Ooi, A. Stasch, S. J. Coles, L. Male, M. B. Hursthouse, K. J. Cavell, A. Dervisi and I. A. Fallis, *Organometallics*, 2008, **27**, 3279; (b) E. L. Kolychev, I. A. Portnyagin, V. V. Shuntikov, V. N. Khrustalev and M. S. Nechaev, *J. Organomet. Chem.*, 2009, **694**, 2454; (c) W. A. Herrmann, S. K. Schneider, K. Öfele, M. Sakamoto and E. Herdtweck, *J. Organomet. Chem.*, 2004, **689**, 2441.
- 27 (a) J. C. Garrison, R. S. Simons, W. G. Kofron, C. A. Tessier and W. J. Youngs, *Chem. Commun.*, 2001, 1780; (b) J. C. Garrison, R. S. Simons, W. G. Kofron, C. A. Tessier and W. J. Youngs, *J. Organomet. Chem.*, 2003, **673**, 1; (c) V. J. Catalano and M. A. Malwitz, *Inorg. Chem.*, 2003, **42**, 5483; (d) V. J. Catalano, M. A. Malwitz and A. O. Etogo, *Inorg. Chem.*, 2004, **43**, 5714.
- 28 (a) I. E. Markó, S. Stérin, O. Buisine, G. Mignani, P. Branlard, B. Tinant and J. P. Declercq, *Science*, 2002, **298**, 204; (b) I. E. Markó, S. Stérin, O. Buisine, G. Berthon, G. Michaud, B. Tinant and J. P. Declercq, *Adv. Synth. Catal.*, 2004, **346**, 1429; (c) O. Buisine, G. Berthon-Gelloz, J. F. Brière, S. Stérin, G. Mignani, P. Branlard, B. Tinant, J. P. Declercq and I. E. Markó, *Chem. Commun.*, 2005, 3856; (d) J. J. Hu, F. Li and T. S. A. Hor, *Organometallics*, 2009, **28**, 1212; (e) G. De Bo, G. Berthon-Gelloz, B. Tinant and I. E. Markó, *Organometallics*, 2006, **25**, 1881; (f) M. G. McLaughlin and M. J. Cook, *Chem. Commun.*, 2011, **47**, 11104; (g) M. Blug, X.-F. Le Goff, N. Mézailles and P. Le Floch, *Organometallics*, 2009, **28**, 2360; (h) J. J. Dunsford, B. M. Kariuki and K. J. Cavell, *J. Organomet. Chem.*, 2011, **696**, 188.
- 29 B. D. Karstedt, *U.S. Patent*, 3,775,452, 1973.
- 30 G. Berthon-Gelloz, J.-M. Schumers, G. De Bo and I. E. Markó, *J. Org. Chem.*, 2008, **73**, 4190.
- 31 S. Boyer, *Elemental Analysis Service*, London Metropolitan University, London, N7 7DD.
- 32 COLLECT, Nonius BV, 1998, Delft, The Netherlands.
- 33 Z. Otwinowski and W. Minor, *Methods Enzymol.*, 1997, **276**, 307.
- 34 G. M. Sheldrick, *Acta Crystallogr., Sect. A: Found. Crystallogr.*, 2008, **A64**, 112.
- 35 R. H. Blessing, *Acta Crystallogr., Sect. A: Found. Crystallogr.*, 1995, **51**, 33.
- 36 P. v. d. Sluis and A. L. Spek, *Acta Crystallogr., Sect. A: Found. Crystallogr.*, 1990, **46**, 194.

