

Cite this: *Chem. Sci.*, 2016, **7**, 5815Received 16th March 2016
Accepted 19th May 2016DOI: 10.1039/c6sc01191h
www.rsc.org/chemicalscience

Nickel-catalyzed cyclization of alkyne-nitriles with organoboronic acids involving *anti*-carbometalation of alkynes†


Xingjie Zhang, Xin Xie and Yuanhong Liu*

A nickel-catalyzed regioselective addition/cyclization of *o*-(cyano)phenyl propargyl ethers with arylboronic acids has been developed, which provides an efficient protocol for the synthesis of highly functionalized 1-naphthylamines with wide structural diversity. The reaction is characterized by a regioselective and *anti*-addition of the arylboronic acids to the alkyne and subsequent facile nucleophilic addition of the resulting alkenylmetal to the tethered cyano group. Mechanistic studies reveal that a Ni(i) species might be involved in the catalytic process.

Introduction

Transition-metal-catalyzed cascade reactions consisting of multiple carbometalation steps have attracted considerable attention in organic synthesis since these processes enable the rapid assembly of complex structures in an efficient, atom-economical and green manner.¹ Among these reactions, organoboron compounds are one of the most widely used reagents, not only due to their chemical stability and ready availability, but also because they can undergo a series of addition reactions to unsaturated compounds such as alkynes, dienes, enones, aldehydes/ketones, nitriles and isocyanates *etc.* in the presence of a transition metal catalyst, especially Rh, Pd or Ni complexes.² The development of cascade reactions by combining different types of these elemental reactions is undoubtedly important and attractive. In this regard, cascade reactions involving the addition of organoboron compounds to alkynes as the initial step have been realized mainly through Rh-^{1b,c,3} or Pd-catalysis,⁴ as reported by Murakami, Hayashi, Lu and other groups. These catalytic reactions generally proceed by *syn*-1,2-addition of the organometal species generated through transmetalation between the organoboron and the metal complex across the carbon–carbon triple bond, followed by nucleophilic attack of the resulting alkenylmetal on the remaining electrophile (Scheme 1, eqn (1)). So far most of the reported reactions proceed *via* formation of regioisomer A in which the R² group of R²B(OH)₂ locates on a carbon adjacent to the alkyne terminus R¹, leading

to an *exo*-alkene upon cyclization^{3,4} (Scheme 1, eqn (1)). Cyclizations involving the regioselective formation of the alkenylmetal with a metal α -to the R¹ substituent such as *syn*-B are quite rare⁵ (Scheme 1, eqn (2)), possibly because the subsequent cyclization process will involve a highly strained transition state. Thus, the development of new cyclization systems with controlled regiochemistry towards B is highly challenging. During our studies on nickel-catalyzed reactions, we found that such a transformation could be achieved by the addition of organoboron compounds to benzene-tethered alkyne-nitriles utilizing nickel as the catalyst, possibly through the isomerization of *syn*-B to *anti*-B. Herein, we report the first example of a nickel-catalyzed carboarylate cyclization of alkyne-nitriles with organoboronic acids involving regioselective and *anti*-carbonickelation of alkynes, which provides an efficient protocol for the synthesis of highly functionalized 1-naphthylamines. In addition, mechanistic studies revealed that Ni(i) species⁶ rather than Ni(ii) species were involved as the key intermediates, which has not been reported in Ni-catalyzed boron addition reactions.



State Key Laboratory of Organometallic Chemistry, Shanghai Institute of Organic Chemistry, Chinese Academy of Sciences, 345 Lingling Road, Shanghai 200032, China. E-mail: yhliu@sioc.ac.cn


† Electronic supplementary information (ESI) available. CCDC 1440646 and 1440647. For ESI and crystallographic data in CIF or other electronic format see DOI: 10.1039/c6sc01191h

Results and discussion

We chose the nickel-catalyzed reaction of *o*-(cyano)phenyl propargyl ether⁷ **1a** and phenylboronic acid as a model reaction for the optimization of the reaction conditions. Initially, we examined the reactions in the presence of $\text{Ni}(\text{COD})_2$ and various phosphine ligands such as PPh_3 in 1,4-dioxane at 90 °C. However, only a trace of the desired cyclization product was observed, along with some byproducts (Table 1, entry 1). Replacing $\text{Ni}(\text{COD})_2$ with a Ni(II) complex, $\text{Ni}(\text{acac})_2 \cdot 2\text{H}_2\text{O}$ (acac = acetylacetone), afforded the cyclized product 4-OTBS-substituted 1-naphthylamine **2a**, albeit in only 17–18% yields (entries 2–3). To our delight, the addition of 10 mol% of $^t\text{BuOK}$ as a base improved the yield of **2a** dramatically to 66% within a short reaction time (entry 4). The results suggest that a base is necessary for this reaction, possibly for promoting the transmetalation step by formation of a borate⁸ with the organoboronic acid. The structure of **2a** also revealed that arylation in the initial step occurred regioselectively on the alkyne carbon that is closer to the OTBS group. Subsequently, the effects of bases, phosphine ligands and solvents were evaluated. Of the various bases, Cs_2CO_3 gave the best result (73%, entry 5). Increasing the catalytic loading of Cs_2CO_3 to 20 mol% had little

effect on the yield of **2a** (entry 8). However, when a stoichiometric amount of Cs_2CO_3 was used, the yield was reduced rapidly (entry 9). It was remarkable that, unlike the use of more than one equivalent of base in most of the transition metal-catalyzed reactions involving organoborons, here only catalytic amounts of base were needed. Increasing the catalyst loading did not improve the yield of the product (entry 10). Triarylphosphine ligands and the N-heterocyclic carbene ligand IPr (IPr = 1,3-bis(2,6-diisopropylphenyl)imidazole-2-ylidene) were also effective, while ligands such as PPh_2Me and PCy_3 were less efficient (entries 11–17). Changing the solvent to THF or toluene afforded **2a** in satisfactory yields of 64–68% (entries 18–19). Addition of one equivalent of H_2O as a promoter or proton source did not afford a better result (entry 20). $\text{Ni}(\text{acac})_2$ also catalyzed the reaction efficiently (entry 21). When $\text{Ni}(\text{COD})_2$ was used as the catalyst, only trace amounts of **2a** were obtained (entry 22). Without the phosphine ligand, the reaction also proceeded to afford **2a** in 65% yield, albeit with a longer reaction time (entry 23). Without a nickel catalyst, no reaction occurred (entry 24). On the basis of the above optimization studies, the reaction conditions shown in Table 1, entry 5 were chosen as the best conditions.

Table 1 Optimization studies for the formation of 1-naphthylamine **2a**



Entry	Catalyst	Ligand	Base	Solvent	Time [h]	Yield ^a [%]
1	$\text{Ni}(\text{COD})_2$	PPh_3^b	—	1,4-Dioxane	3	Trace
2	$\text{Ni}(\text{acac})_2 \cdot 2\text{H}_2\text{O}$	PPh_3	—	1,4-Dioxane	24	18
3	$\text{Ni}(\text{acac})_2 \cdot 2\text{H}_2\text{O}$	$\text{P}(p\text{-CF}_3\text{C}_6\text{H}_4)_3$	—	1,4-Dioxane	24	17
4	$\text{Ni}(\text{acac})_2 \cdot 2\text{H}_2\text{O}$	$\text{P}(p\text{-CF}_3\text{C}_6\text{H}_4)_3$	$^t\text{BuOK}$	1,4-Dioxane	4	66
5	$\text{Ni}(\text{acac})_2 \cdot 2\text{H}_2\text{O}$	$\text{P}(p\text{-CF}_3\text{C}_6\text{H}_4)_3$	Cs_2CO_3	1,4-Dioxane	3	73
6	$\text{Ni}(\text{acac})_2 \cdot 2\text{H}_2\text{O}$	$\text{P}(p\text{-CF}_3\text{C}_6\text{H}_4)_3$	CsF	1,4-Dioxane	6	67
7	$\text{Ni}(\text{acac})_2 \cdot 2\text{H}_2\text{O}$	$\text{P}(p\text{-CF}_3\text{C}_6\text{H}_4)_3$	K_2CO_3	1,4-Dioxane	10	31
8	$\text{Ni}(\text{acac})_2 \cdot 2\text{H}_2\text{O}$	$\text{P}(p\text{-CF}_3\text{C}_6\text{H}_4)_3$	Cs_2CO_3^c	1,4-Dioxane	3	68
9	$\text{Ni}(\text{acac})_2 \cdot 2\text{H}_2\text{O}$	$\text{P}(p\text{-CF}_3\text{C}_6\text{H}_4)_3$	Cs_2CO_3^d	1,4-Dioxane	10	8 (59)
10 ^e	$\text{Ni}(\text{acac})_2 \cdot 2\text{H}_2\text{O}$	$\text{P}(p\text{-CF}_3\text{C}_6\text{H}_4)_3$	Cs_2CO_3	1,4-Dioxane	3	74
11	$\text{Ni}(\text{acac})_2 \cdot 2\text{H}_2\text{O}$	PPh_3	Cs_2CO_3	1,4-Dioxane	5	69
12	$\text{Ni}(\text{acac})_2 \cdot 2\text{H}_2\text{O}$	$\text{P}(p\text{-MeC}_6\text{H}_4)_3$	Cs_2CO_3	1,4-Dioxane	4	68
13	$\text{Ni}(\text{acac})_2 \cdot 2\text{H}_2\text{O}$	$\text{P}(\text{C}_6\text{F}_5)_3$	Cs_2CO_3	1,4-Dioxane	5	63
14	$\text{Ni}(\text{acac})_2 \cdot 2\text{H}_2\text{O}$	PPh_2Me	Cs_2CO_3	1,4-Dioxane	7	45
15	$\text{Ni}(\text{acac})_2 \cdot 2\text{H}_2\text{O}$	PCy_3	Cs_2CO_3	1,4-Dioxane	17	55
16	$\text{Ni}(\text{acac})_2 \cdot 2\text{H}_2\text{O}$	IPr	Cs_2CO_3	1,4-Dioxane	6	64
17	$\text{Ni}(\text{acac})_2 \cdot 2\text{H}_2\text{O}$	IPr	$^t\text{BuOK}$	1,4-Dioxane	7	62
18	$\text{Ni}(\text{acac})_2 \cdot 2\text{H}_2\text{O}$	$\text{P}(p\text{-CF}_3\text{C}_6\text{H}_4)_3$	Cs_2CO_3	THF	8	64
19	$\text{Ni}(\text{acac})_2 \cdot 2\text{H}_2\text{O}$	$\text{P}(p\text{-CF}_3\text{C}_6\text{H}_4)_3$	Cs_2CO_3	Toluene	3	68
20 ^f	$\text{Ni}(\text{acac})_2 \cdot 2\text{H}_2\text{O}$	$\text{P}(p\text{-CF}_3\text{C}_6\text{H}_4)_3$	Cs_2CO_3	1,4-Dioxane	6	57
21	$\text{Ni}(\text{acac})_2$	$\text{P}(p\text{-CF}_3\text{C}_6\text{H}_4)_3$	Cs_2CO_3	1,4-Dioxane	3	72
22	$\text{Ni}(\text{COD})_2$	$\text{P}(p\text{-CF}_3\text{C}_6\text{H}_4)_3^b$	Cs_2CO_3	1,4-Dioxane	9	Trace
23	$\text{Ni}(\text{acac})_2 \cdot 2\text{H}_2\text{O}$	—	Cs_2CO_3	1,4-Dioxane	5	65
24	—	$\text{P}(p\text{-CF}_3\text{C}_6\text{H}_4)_3$	Cs_2CO_3	1,4-Dioxane	10	(99)

^a Isolated yields. The yields of the recovered **1a** were shown in parentheses. ^b 10 mol% of the ligand was used. ^c 20 mol% of Cs_2CO_3 was used. ^d 1.0 equiv. of Cs_2CO_3 was used. ^e 10 mol% $\text{Ni}(\text{acac})_2 \cdot 2\text{H}_2\text{O}$, 10 mol% $\text{P}(p\text{-CF}_3\text{C}_6\text{H}_4)_3$ and 20 mol% Cs_2CO_3 were used. ^f One equiv. of H_2O was added.



Next, we proceeded to investigate the scope of this new cascade addition/cyclization reaction catalyzed by $\text{Ni}(\text{acac})_2 \cdot 2\text{H}_2\text{O}$. The reactivity of various organoboronic acids was first examined using **1a** as a reaction partner (Table 2). During this process, we found that the 5 mol% catalyst loading was not effective in some cases and thus 10 mol% $\text{Ni}(\text{acac})_2 \cdot 2\text{H}_2\text{O}$, 10 mol% $\text{P}(p\text{-CF}_3\text{C}_6\text{H}_4)_3$ and 20 mol% Cs_2CO_3 were used in most of the cases to achieve better product yields. As shown in Table 2, a wide range of diversely substituted aryl- or heteroaryl-boronic acids were suitable for this reaction, leading to the desired 1-naphthylamines **2a–2s** in generally good to high yields. Arylboronic acids bearing electron-donating groups such as *p*-Me, *p*-*t*Bu and *p*-MeO or electron-withdrawing groups such as *p*-F, *p*-Cl, *p*-CF₃, *p*-CN, *p*-CO₂Et and *p*-Ac on the aryl ring underwent the cyclization smoothly to provide the corresponding 1-naphthylamines **2b–2j** in 64–77% yields, and these functional groups were well tolerated during the reaction. Of note is that the CN and Ac groups remained intact under the reaction conditions, and no nickel-catalyzed boron additions to these groups were observed. The results indicated that electron-poor or -rich aryl substituents on the arylboronic acid had little influence on the

yields of products **2**. The sterically demanding *o*-MeO substituted arylboronic acid afforded **2k** with a longer reaction time and a lower yield of 47%, indicating that the reaction is markedly influenced by steric effects. Arylboronic acids with *–MeO* or *–Cl* substituents at the 3-, 3,4- or 3,5-positions of the phenyl ring, or with a biphenyl or 2-naphthyl ring transformed into products **2l–2m** and **2o–2r** efficiently in good yields. The use of 2-fluorophenylboronic acid gave **2n** in 41% yield. 2-Thienylboronic acid also participated in this cascade reaction, albeit with a lower yield of **2s**. However, when alkylboronic acids such as *n*-butylboronic acid were employed, no desired product was obtained.

The scope of *o*-(cyano)phenyl propargyl ethers was then examined (Table 3). A variety of electron-donating and -withdrawing groups on the aryl rings at the alkyne terminus were found to be compatible, such as *p*-Me, *p*-OMe, *p*-F, *p*-Cl, *p*-CF₃ and *p*-CO₂Me substituents, and the corresponding products **2t–2y** were formed in 61–76% yields. Interestingly, in contrast to the results of the reaction of **1a** with 2-thienylboronic acid, the presence of a 2-thienyl or 3-benzothienyl ring as the alkyne terminal did not have much influence on the reaction, and the corresponding products **2z** and **2za** were obtained in high yields of 80% and 74%, respectively. A 9H-fluorene substituent, which is a very useful unit in organofunctional materials, was also successfully incorporated into the product **2zb** in 70% yield. Alkenyl or alkyl-substituted alkynes, such as cyclohexenyl, cyclopropyl and propyl-substituted ones, however, afforded the desired products **2zc–2ze** in low yields of 23–35%. The structure of the 1-naphthylamine product was unambiguously confirmed by X-ray crystallographic analysis of **2o**.⁹ 1-Naphthylamines are important structural motifs found in a variety of biologically active substances. They also act as useful building blocks in

Table 2 Scope of the reaction with respect to arylboronic acids^a

1a	2.0 equiv	10 mol% $\text{Ni}(\text{acac})_2 \cdot 2\text{H}_2\text{O}$ 10 mol% $\text{P}(p\text{-CF}_3\text{C}_6\text{H}_4)_3$ 20 mol% Cs_2CO_3 1,4-dioxane, 90 °C	2
2a , 3 h, 73% ^b	2b , 8 h, 73% ^b	2c , 4 h, 77%	2d , 6 h, 68%
2e , 5 h, 68%	2f , 3 h, 69%	2g , 4 h, 76%	2h , 3 h, 74% ^c
2i , 5 h, 64%	2j , 6 h, 70% ^c	2k , 10 h, 47%	2l , 7.5 h, 75%
2m , 10.5 h, 65%	2n , 11 h, 41%	2o , 3 h, 68% ^b	2p , 2 h, 65%
2q , 6 h, 73%	2r , 6 h, 70%	2s , 10 h, 35%	

^a The yields given are for the isolated products. ^b 5 mol% $\text{Ni}(\text{acac})_2 \cdot 2\text{H}_2\text{O}$, 5 mol% $\text{P}(p\text{-CF}_3\text{C}_6\text{H}_4)_3$ and 10 mol% Cs_2CO_3 were used. ^c THF was used as the solvent.

Table 3 Scope of the *o*-(cyano)phenyl propargyl ethers^a

1	2.0 equiv	10 mol% $\text{Ni}(\text{acac})_2 \cdot 2\text{H}_2\text{O}$ 10 mol% $\text{P}(p\text{-CF}_3\text{C}_6\text{H}_4)_3$ 20 mol% Cs_2CO_3 1,4-dioxane, 90 °C	2
2t , 5 h, 76%	2u , 4 h, 69%	2v , 4 h, 76%	2w , 4 h, 61%
2x , 5.5 h, 74%	2y , 4.5 h, 62%	2z , 4 h, 80%	2za , 4 h, 74%
2zb , 4 h, 70%	2zc , 8 h, 35%	2zd , 6 h, 23%	2ze , 3 h, 29%

^a The yields given are for the isolated products.



synthetic chemistry and the dyestuffs industry. However, efficient methods for their synthesis are quite limited.¹⁰ Our reaction provides a convenient route to 1-naphthylamines.


To understand the reaction mechanism, we tried to isolate the possible reaction intermediates. First, a stoichiometric reaction of Ni(acac)₂ (1 equiv.), IPr (1 equiv.), PhB(OH)₂ (2 equiv.) and ⁴BuOK (2 equiv.) was carried out (Scheme 2, eqn (1)). It was found that in addition to a biphenyl product, a red crystalline compound IPrNi(acac) 3 was isolated in 62% yield. Complex 3 is paramagnetic as indicated by the appearance of broad signals in the ¹H NMR spectrum. The X-ray crystal analysis of 3 (ref. 9) clearly shows a rare, three-coordinate distorted T-shaped Ni(i) structure.^{6e,f} When Cs₂CO₃ was used instead of ⁴BuOK, the same Ni(i) complex was also observed, albeit in a low yield of 14%. In this case, a large amount of precipitate could be observed. The precipitate was assumed to be the PhB(OH)₂-Cs₂CO₃ adduct. Thus, the formation of this low-soluble adduct might prevent further reaction with the nickel complex. In fact, stirring a 1 : 1 mixture of PhB(OH)₂ and Cs₂CO₃ in 1,4-dioxane at 90 °C resulted in the formation of a large amount of white precipitate. This might also explain why the use of 1 equivalent of Cs₂CO₃ provided the cyclized product 2a in a low yield (Table 1, entry 9). Complex 3 might be formed by the comproportionation of Ni(0) and Ni(i) species.^{6c,d} To confirm this point, the stoichiometric reaction of Ni(COD)₂ with Ni(acac)₂ in the presence of IPr was carried out. To our delight, the same complex 3 was formed in moderate yield (Scheme 2, eqn (2)). This Ni(i) complex 3 was found to catalyze the cyclization of 1a

with PhB(OH)₂ to afford 2a (Scheme 2, eqn (3)), implying that the Ni(i) species was likely involved in the catalytic process. Recently, a Ni(i) species was proposed to be relevant in Ni-catalyzed cross-coupling reactions.⁶ Our results demonstrate that Ni(i) may also be involved in the Ni-catalyzed addition reactions of organoboron reagents to alkynes. In addition, the reaction of allene 4 with PhB(OH)₂ under Ni-catalyzed conditions afforded indeno[1,2-*b*]quinoline 5 *via* base-promoted cyano-Schmittel cyclization,⁷ while the desired 2a was not observed (Scheme 2, eqn (4)), indicating that the reaction does not involve the allene intermediate.


The origin of the regioselectivity in this reaction is not clear yet; it is possibly controlled by electronic factors in π -complex 12 (see Scheme 3).¹¹ To understand the role of the OTBS group, alkyne-nitrile 6 without the OTBS group was synthesized. The Ni-catalyzed reaction of 6 with PhB(OH)₂ afforded only 10% of the desired naphthylamine 7, along with small amounts of an unidentified byproduct (Scheme 2, eqn (5)). The results indicate that the presence of the OTBS group is crucial for this reaction.

In addition, it was found that in the presence of a radical scavenger such as TEMPO, the desired 2a and an *N*-arylated product 8 were formed in 74% combined yield (Scheme 2, eqn (6)). 8 was possibly formed by the Ni-catalyzed oxidative amidation of the arylboronic acid with amine 2a.^{12,13} The results indicate that the reaction was not inhibited by TEMPO, and this suggests that a radical species is not involved in this system.

Based on the above results, we propose the following reaction mechanism (Scheme 3). Initially, transmetalation of the arylboronic acid with the Ni(ii) complex, promoted by a base, provides diarylnickel(ii) species 9, together with HOBO, HC₂CO₃ and Cs(acac).¹⁴ 9 undergoes reductive elimination to form a Ni(0) species. The observation of biphenyl¹⁵ in the catalytic reaction of 1a with PhB(OH)₂ also indicates that Ni(ii) was



Scheme 2 Mechanistic studies.



Scheme 3 Possible reaction mechanism.



reduced in the reaction process. This Ni(0) species coproportionates with Ni(II) to afford Ni(I) complex **10**, which undergoes transmetalation with the arylboronic acid to give arylnickel(I) species **11**. Regioselective 1,2-addition of arylnickel(I) species **11** to the alkyne moiety in a *syn*-fashion takes place to give an alkenylnickel(I) intermediate *syn*-**13**. *cis*-to-*trans* isomerization of **13**,¹⁶ possibly through a carbene-like zwitterionic resonance species¹⁷ yields alkenylnickel(I) intermediate *anti*-**13** with a metal *trans*-to the Ar substituent. It was noted that most of the metal-catalyzed reactions of organoborons to alkynes gave the *syn*-addition product while few reactions produced the *anti*-addition product.^{2r,17} The regio- and stereochemistry for the addition process here are consistent with those observed for the cobalt(II)-catalyzed hydroarylation of propargyl-alcohols or -carbamates with arylboronic acids.¹⁸ The cyano group may play a role in facilitating the *cis*-*trans* isomerization by stabilizing the metal species and directing the subsequent addition reaction. Nucleophilic attack of the alkenylnickel in *anti*-**13** to the cyano group forms a cyclized intermediate **14**. Subsequent protonation of **14** produces the N–H imine **15** and a nickel(I) species **16**. Tautomerization of **15** affords the observed product **2**. **16** undergoes transmetalation with ArB(OH)₂ to regenerate the arylnickel(I) catalyst **11**.

Conclusions

In summary, we have developed a nickel-catalyzed regioselective addition/cyclization of *o*-(cyano)phenyl propargyl ethers with arylboronic acids, which provides an efficient protocol for the synthesis of highly functionalized 1-naphthylamines with wide structural diversity. The reaction is characterized by a regioselective and *anti*-addition of the arylboronic acids to the alkyne and subsequent facile nucleophilic addition of the resulting alkenylnickel to the tethered cyano group. Mechanistic studies reveal that a Ni(I) species might be involved in the catalytic process. Further mechanistic studies and the extension to alkynes tethered with a wide variety of electrophiles are currently ongoing in our laboratory.

Acknowledgements

We thank the National Natural Science Foundation of China (Grant No. 21125210, 21421091) for financial support.

Notes and references

- For reviews, see: (a) E. Negishi, C. Copéret, S. Ma, S.-Y. Liou and F. Liu, *Chem. Rev.*, 1996, **96**, 365–393; (b) T. Miura and M. Murakami, *Chem. Commun.*, 2007, 217–224; (c) S. W. Youn, *Eur. J. Org. Chem.*, 2009, 2597–2605.
- For selected papers, see: for Rh: (a) M. Sakai, M. Ueda and N. Miyaura, *Angew. Chem., Int. Ed.*, 1998, **37**, 3279–3281; (b) T. Hayashi, K. Inoue, N. Taniguchi and M. Ogasawara, *J. Am. Chem. Soc.*, 2001, **123**, 9918–9919; (c) C. G. Frost and K. J. Wadsworth, *Chem. Commun.*, 2001, 2316–2317; (d) M. Murakami and H. Igawa, *Chem. Commun.*, 2002, 390–391; (e) T. Matsuda, M. Makino and M. Murakami, *Org. Lett.*, 2004, **6**, 1257–1259; (f) H. Shimizu and M. Murakami, *Chem. Commun.*, 2007, 2855–2857; (g) T. Miura, Y. Takahashi and M. Murakami, *Chem. Commun.*, 2007, 3577–3579; (h) T. Matsuda, Y. Suda and A. Takahashi, *Chem. Commun.*, 2012, **48**, 2988–2990. For Pd: (i) T. Nishikata, Y. Yamamoto and N. Miyaura, *Angew. Chem., Int. Ed.*, 2003, **42**, 2768–2770; (j) C. H. Oh, T. W. Ahn and R. Reddy, *Chem. Commun.*, 2003, 2622–2623; (k) Y. Bai, J. Yin, W. Kong, M. Mao and G. Zhu, *Chem. Commun.*, 2013, **49**, 7650–7652. For Ni: (l) E. Shirakawa, G. Takahashi, T. Tsuchimoto and Y. Kawakami, *Chem. Commun.*, 2001, 2688–2689; (m) E. Shirakawa, G. Takahashi, T. Tsuchimoto and Y. Kawakami, *Chem. Commun.*, 2002, 2210–2211; (n) K. Hirano, H. Yorimitsu and K. Oshima, *Adv. Synth. Catal.*, 2006, **348**, 1543–1546; (o) J. D. Sieber, S. Liu and J. P. Morken, *J. Am. Chem. Soc.*, 2007, **129**, 2214–2215; (p) K. Hirano, H. Yorimitsu and K. Oshima, *Org. Lett.*, 2007, **9**, 5031–5033; (q) Y.-C. Wong, K. Parthasarathy and C.-H. Cheng, *Org. Lett.*, 2010, **12**, 1736–1739; (r) D. W. Robbins and J. F. Hartwig, *Science*, 2011, **333**, 1423–1427. So far only two reports involving Ni-catalyzed hydroarylation of alkynes with arylboronic acids have been published, see ref. 2l and r.

3 (a) R. Shintani, K. Okamoto, Y. Otomaru, K. Ueyama and T. Hayashi, *J. Am. Chem. Soc.*, 2005, **127**, 54–55; (b) T. Miura, M. Shimada and M. Murakami, *J. Am. Chem. Soc.*, 2005, **127**, 1094–1095; (c) T. Matsuda, M. Makino and M. Murakami, *Angew. Chem., Int. Ed.*, 2005, **44**, 4608–4611; (d) T. Miura, M. Shimada and M. Murakami, *Synlett*, 2005, 667–669; (e) T. Miura, H. Nakazawa and M. Murakami, *Chem. Commun.*, 2005, 2855–2856; (f) T. Miura, T. Sasaki, T. Harumashi and M. Murakami, *J. Am. Chem. Soc.*, 2006, **128**, 2516–2517; (g) T. Miura, Y. Takahashi and M. Murakami, *Org. Lett.*, 2007, **9**, 5075–5077; for Rh-catalyzed reaction of arylboronic acids with alkyne-nitriles to *exo*-alkenes, see ref. 3e.

- (a) J. Song, Q. Shen, F. Xu and X. Lu, *Org. Lett.*, 2007, **9**, 2947–2950; (b) T. Miura, T. Toyoshima, Y. Takahashi and M. Murakami, *Org. Lett.*, 2008, **10**, 4887–4889; (c) H. Tsukamoto, T. Suzuki, T. Uchiyama and Y. Kondo, *Tetrahedron Lett.*, 2008, **49**, 4174–4177; (d) X. Han and X. Lu, *Org. Lett.*, 2010, **12**, 108–111; (e) K. Shen, X. Han and X. Lu, *Org. Lett.*, 2012, **14**, 1756–1759.

5 For a Pd(0)-catalyzed cyclization of alkynals or alkynones involving formal anti-addition of organoboronic reagents *via* a different reaction mechanism, see: H. Tsukamoto, T. Ueno and Y. Kondo, *J. Am. Chem. Soc.*, 2006, **128**, 1406–1407.

- (a) T. J. Anderson, G. D. Jones and D. A. Vicic, *J. Am. Chem. Soc.*, 2004, **126**, 8100–8101; (b) G. D. Jones, C. McFarland, T. J. Anderson and D. A. Vicic, *Chem. Commun.*, 2005, 4211–4213; (c) G. Yin, I. Kalvet, U. Englert and F. Schoenebeck, *J. Am. Chem. Soc.*, 2015, **137**, 4164–4172; (d) L. M. Guard, M. Mohadjer Beromi, G. W. Brudvig, N. Hazari and D. J. Vinyard, *Angew. Chem., Int. Ed.*, 2015, **54**, 13352–13356. For Ni(I) complexes containing NHC ligands, see: Ni(Pr₂)Cl: (e) S. Miyazaki, Y. Koga,



T. Matsumoto and K. Matsubara, *Chem. Commun.*, 2010, **46**, 1932–1934; Ni(IMes)₂X: (f) K. Zhang, M. Conda-Sheridan, S. R. Cooke and J. Louie, *Organometallics*, 2011, **30**, 2546–2552.

7 For our recent results on cyano-Schmittel cyclization of (*o*-cyano)phenylpropargyl ethers, see: X. You, X. Xie, H. Chen, Y. Li and Y. Liu, *Chem.–Eur. J.*, 2015, **21**, 18699–18705.

8 Y. Makida, E. Marelli, A. M. Z. Slawin and S. P. Nolan, *Chem. Commun.*, 2014, **50**, 8010–8013.

9 CCDC 1440646 (**20**) and 1440647(3) contain the supplementary crystallographic data for this paper.†

10 (a) P. Sagar, R. Fröhlich and E.-U. Würthwein, *Angew. Chem., Int. Ed.*, 2004, **43**, 5694–5697; (b) Q. Shen and J. F. Hartwig, *J. Am. Chem. Soc.*, 2006, **128**, 10028–10029; (c) S.-C. Zhao, X.-Z. Shu, K.-G. Ji, A.-X. Zhou, T. He, X.-Y. Liu and Y.-M. Liang, *J. Org. Chem.*, 2011, **76**, 1941–1944; (d) R. S. Reddy, P. K. Prasad, B. B. Ahuja and A. Sudalai, *J. Org. Chem.*, 2013, **78**, 5045–5050; (e) P. Gao, J. Liu and Y. Wei, *Org. Lett.*, 2013, **15**, 2872–2875; (f) W. P. Hong, A. V. Iosub and S. S. Stahl, *J. Am. Chem. Soc.*, 2013, **135**, 13664–13667.

11 A. Arcadi, M. Aschi, M. Chiarini, G. Ferrara and F. Marinelli, *Adv. Synth. Catal.*, 2010, **352**, 493–498.

12 D. S. Raghuvanshi, A. K. Gupta and K. N. Singh, *Org. Lett.*, 2012, **14**, 4326–4329.

13 The reaction of amine **2a** with phenylboronic acid under the reaction conditions shown in Scheme 2, eqn (6) gave **8** in 43% yield.

14 P.-S. Lin, M. Jeganmohan and C.-H. Cheng, *Chem.–Asian J.*, 2007, **2**, 1409–1416.

15 Without ligand, only biphenyl was observed. With 10 mol% of P(*p*-CF₃C₆H₄)₃ as the ligand, in addition to biphenyl, 4,4'-bis(trifluoromethyl)biphenyl and 4-(trifluoromethyl) biphenyl were also observed, possibly due to aryl transfer from the phosphine ligand. See: I. Colon and D. R. Kelsey, *J. Org. Chem.*, 1986, **51**, 2627–2637.

16 A. Yamamoto and M. Suginome, *J. Am. Chem. Soc.*, 2005, **127**, 15706–15707.

17 M. Murakami, T. Yoshida, S. Kawanami and Y. Ito, *J. Am. Chem. Soc.*, 1995, **117**, 6408–6409.

18 P.-S. Lin, M. Jeganmohan and C.-H. Cheng, *Chem.–Eur. J.*, 2008, **14**, 11296–11299.

