



Cite this: *Org. Biomol. Chem.*, 2017, **15**, 3575

Received 3rd March 2017,  
Accepted 6th April 2017

DOI: 10.1039/c7ob00550d

rsc.li/obc

## Imidazolium-labeled glycosides as probes to harness glycosyltransferase activity in human breast milk†

I. Sittel and M. C. Galan  \*

**Imidazolium-labeled (ITag-) glycosides are used to harness the glycosyltransferase activity directly from human breast milk. The covalently attached ionic labels provide a bifunctional chemical handle that is used to monitor reaction progress by MS, as well as aid in product purification from complex mixtures. The technology is exemplified in the synthesis of biologically relevant oligosaccharide analogs, LacNAc-ITag, ITag-Lewis<sup>x</sup> and ITag-Lewis<sup>a</sup>, in a matter of days from human breast milk without having to isolate specific enzymes.**

The growing appreciation of oligosaccharides and glycoconjugates as key mediators in many important biological processes from cell signaling, embryogenesis, neural development to pathogen recognition, inflammation, innate immune responses and cancer, has led to their use as probes for biological research and as key compounds for drug and vaccine development.<sup>1</sup> Despite the many recent advances in the field, progress in this area is still impeded by a lack of general and efficient methods for the routine preparation of complex oligosaccharides. Traditional chemical approaches rely on lengthy and time consuming chemical steps to achieve good yields and stereocontrol.<sup>2</sup> On the other hand, the use of carbohydrate processing enzymes such as glycosyltransferases, to construct oligosaccharides offers several advantages over chemical approaches, since these enzymes are able to form glycosidic bonds with exquisite regio- and stereocontrol by catalyzing the transfer of an activated sugar donor to growing oligosaccharide chains without the need for protecting groups.<sup>2d,3</sup> However, access to these enzymes is limited as the proteins can be challenging to express, expensive to employ on scale and only a handful are currently commercially available.<sup>4</sup>

Human breast milk (HBM) is an extremely complex and highly variable biofluid comprised of carbohydrates, proteins,


lipids and enzymes, which has evolved over millennia to nourish infants and protect them from disease, whilst their own immune system matures.<sup>5</sup> HBM contains more than 200 different unbound human milk oligosaccharides (HMOs),<sup>5b,6</sup> consisting of five monosaccharide building blocks: galactose (Gal), glucose (Glc), *N*-acetylglucosamine (GlcNAc), fucose (Fuc) and sialic acid (Neu5Ac). All HMOs carry lactose (Gal-β-1,4-Glc) at the reducing end, which can be elongated through a β-1,3- or β-1,6-linkage by two different disaccharides, either Gal-β-1,3-GlcNAc (lacto-*N*-biose, type 1 chain) or Gal-β-1,4-GlcNAc *N*-acetylglucosamine (LacNAc, type 2 chain). Further chain elongation can take place by the incorporation of Fuc and Neu5Ac units.<sup>7</sup> Our understanding of the biosynthesis of these oligosaccharides is limited, since most of the specific glycosyltransferases that are responsible for the formation of HMO structures with specific glycosidic linkages have not been identified.<sup>7b,8</sup> In addition, HMO-specific glycosyltransferase expression levels are determined by the blood group, diet and environmental and genetic factors as well as lactational period of the donor.<sup>7a,9</sup>

Ionic liquid-based labels have emerged as a new class of chemical labels that can be employed for the immobilization and mass spectrometry (MS) detection of reagents from complex reaction mixtures, due to their unique physical and chemical properties.<sup>10</sup> Our group<sup>10c,11</sup> and others<sup>12</sup> have reported the synthesis of inexpensive and versatile imidazolium-based tags (ITags) and their application to chromatography-free chemical oligosaccharide synthesis. Furthermore, in our work, we demonstrated that the cationic labels are also compatible with enzymatic transformations and that ITag-glycoside substrates could be employed for the qualitative and quantitative reaction monitoring, using MS techniques, of bovine milk β-1,4-galactosyltransferase (β-1,4-GalT),<sup>13</sup> human α-1,3-fucosyltransferase (α-1,3-FucT VI)<sup>14</sup> and α-2,3-sialyltransferase (α-2,3-ST) from *pasteurella multocida*, as well as α-2,6-sialyltransferase (α-2,6-ST)<sup>15</sup> from *photobacterium damsela*. More recently, ITag-labeled mannosamine was also employed for the non-covalent labeling of live cells,<sup>16</sup> demonstrating the versatility of the probes.

School of Chemistry, University of Bristol, Cantock's Close, Bristol BS8 1TS, UK.  
E-mail: m.c.galan@bristol.ac.uk; Fax: (+44) (0)117 925 1295

† Electronic supplementary information (ESI) available: Full experimental and characterization data for all compounds, including NMR spectra and LC-MS traces. See DOI: 10.1039/c7ob00550d





**Fig. 1** General strategy to harness glycosyltransferase activity in human breast milk.

Encouraged by the simplicity and practicality of the approach, we hypothesized that ITag-glycan probes would be ideally suited to harness the biosynthetic machinery (*e.g.* glycosyltransferases) present in HBM, and thus gain expedient access to biologically important oligosaccharides, without the need to isolate or express specific enzymes (Fig. 1).


Initial experiments began by assessing our ability to monitor mixtures of ITag-glycoside products that had been generated *in vitro* in a multi-enzyme environment. To that end, methylimidazolium *N*-benzenesulfonyl-linked GlcNAc (GlcNAc-ITag) **1**, which was prepared in 4 steps and 59% overall yield from azidopropyl GlcNAc **6**<sup>15</sup> (see ESI† for details), was incubated in 3-morpholinopropane-1-sulfonic acid (MOPS) buffer (pH 7.8) at 37 °C with three different recombinant enzymes:  $\beta$ -1,4-GalT, which catalyzes the transfer of galactose from uridine 5'-diphosphogalactose (UDP-Gal) to C4 of GlcNAc to yield ITag-LacNAc **3**;  $\alpha$ -2,3-ST, which in the presence of cytidine 5'-monophospho-*N*-acetylneuraminic acid (CMP-Neu5Ac) will add Neu5Ac at C3' of the LacNAc disaccharide to yield 3'-Sialyl-LacNAc-ITag **9** and  $\alpha$ -1,3-FucT VI, which can fucosylate **3** at C3 of the GlcNAc unit in the presence of guanosine 5'-diphospho- $\beta$ -L-fucose (GDP-Fuc) to form the ITag-Lewis<sup>X</sup> trisaccharide [Fuc- $\alpha$ -1,3-(Gal- $\beta$ -1,4)-GlcNAc] **4** or generate Sialyl-Lewis<sup>X</sup>-ITag **10** from **9**<sup>17</sup> (Scheme 1). Incubations were carried out in the presence of an excess of an equimolar mixture of UDP-Gal, CMP-Neu5Ac and GDP-Fuc, which was added portion-wise over 3 days. LC-MS was used to analyze and monitor reaction progress. Four different ITagged species were identified in the mixture: starting material **1** ( $[M^+]$  513), LacNAc-ITag **3** ( $[M^+]$  675) and trisaccharides **4** ( $[M^+]$  821) and **9** ( $[M^+]$  966) in a ratio of 12:42:2:44 for **1**:**3**:**4**:**9** as determined by LC-MS. Comparison of retention times and MS data to that of isolated ITag-materials,<sup>13–15</sup> confirmed the structure identity of the individual ITag-glycosides.

Having demonstrated that we are able to detect ITag-glycosylation products from complex enzymatic mixtures, even at low concentrations ( $\mu$ M), GlcNAc-ITag **1**, which is a common motif in both type 1 and type 2 HMO chains,<sup>7a</sup> was used as a decoy to hijack glycosyltransferase activity in HBM.  $\beta$ -Linked GlcNAc units are further elongated in HMOs with a  $\beta$ -1,3- or  $\beta$ -1,4-linked galactose unit. Thus **1** was incubated at 37 °C with



**Scheme 1** One-pot *in vitro* chemo-enzymatic synthesis of ITagged human milk oligosaccharides **3**, **4** and **9**. See ESI† Fig. S1† for traces.

unpasteurized HBM ( $1 \text{ mg mL}^{-1}$ ) in the presence of excess UDP-Gal (2.4 equiv.), which was added portion-wise over 5 days, in order to bias the production of galactosylated species (Scheme 2). It is important to highlight that no reaction was observed without addition of the sugar nucleotide (Fig. S3 in ESI†). To our delight, LC-MS analysis of the milk sample showed >86% conversion of **1** to a galactose-containing ITag-disaccharide ( $M^+$  675). Importantly, ITag-oligosaccharide products display significantly different retention times in reverse-phase HPLC from other components in HBM (including unlabeled HMOs) to allow their efficient isolation. NMR



**Scheme 2** Incubation of GlcNAc-ITag **1** and LacNAc-ITag **3** with UDP-Gal and GDP-Fuc, respectively, in HBM and LC-MS traces of (A) total ion count of HBM incubation of **1** without addition of UDP-Gal; (B) extracted LC-MS trace for **1** ( $M^+$  513) before addition of UDP-Gal; (C) extracted LC-MS trace for **3** ( $M^+$  675) after 5 days of incubation with **1** and UDP-Gal; (D) extracted LC-MS trace for **4** ( $M^+$  821) after 17 days of incubation with **3** and GDP-Fuc.





no further conversion was observed. Donor nucleotide and fresh milk were added throughout the incubation period and in the case of fucosylated targets, the sample was freeze dried when no further conversion was observed and the reaction restarted by addition of fresh milk and GDP-Fucose to the dried sample (see specific details in ESI†). The reaction was then quenched by cooling on ice and the mixture filtered with ISO-Disc Filters (N 4-2, Nylon 4 mm × 0.2 μm) prior to analysis.

In conclusion, although metabolic oligosaccharide pathways present a number of significant challenges, we have shown that imidazolium-labeled (ITag-) glycosides can be used to harness glycosyltransferase activity directly from human breast milk, without having to isolate the specific enzymes. We demonstrate that ITags covalently attached to glycoside substrates provide a bifunctional chemical handle that can be used to monitor reaction progress by MS, as well as aid in the purification of the products from complex mixtures. Using this technology, we were able to access a series of biologically important oligosaccharides, LacNAc-ITag, ITag-Lewis<sup>x</sup> and ITag-Lewis<sup>a</sup>, in a matter of days from a readily available source, human breast milk. We believe that these cationic probes offer a practical and novel technology to harness enzyme activity in complex biosynthetic pathways. We believe that the strategy showcased here will find applications in and beyond the field of carbohydrates.

This research was supported by EPSRC CAF EP/L001926/1 and ERC-COG: 648239. We thank Anthony P. Corfield, Matthew J. Watt and Chloe C. Medina for fruitful discussions and Mrs Marion Copeland from The Precious Milk Bank (Southmead Hospital Bristol) for donation of milk samples.

## Notes and references

- (a) E. K. McCranie and B. O. Bachmann, *Nat. Prod. Rep.*, 2014, **31**, 1026; (b) J. Bauer and H. M. I. Osborn, *Future Med. Chem.*, 2015, **7**, 2285; (c) A. Varki, *Glycobiology*, 2017, **27**, 3.
- (a) M. C. Galan, D. Benito-Alifonso and G. M. Watt, *Org. Biomol. Chem.*, 2011, **9**, 3598; (b) J. P. Yasomanee and A. V. Demchenko, *Trends Glycosci. Glycotechnol.*, 2013, **25**, 13; (c) A. Fernandez-Tejada, F. J. Canada and J. Jimenez-Barbero, *ChemMedChem*, 2015, **10**, 1291; (d) L. Krasnova and C. H. Wong, *Annu. Rev. Biochem.*, 2016, **85**, 599; (e) A. P. Demchenko, *Handbook of Chemical Glycosylation: Advances in Stereoselectivity and Therapeutic Relevance*, Wiley, 2008.
- (a) T. Murata and T. Usui, *Trends Glycosci. Glycotechnol.*, 2000, **12**, 161; (b) A. Trincone and A. Giordano, *Curr. Org. Chem.*, 2006, **10**, 1163; (c) M. M. Palcic, *Curr. Opin. Chem. Biol.*, 2011, **15**, 226.
- (a) R. M. Schmaltz, S. R. Hanson and C. H. Wong, *Chem. Rev.*, 2011, **111**, 4259; (b) P. M. Danby and S. G. Withers, *ACS Chem. Biol.*, 2016, **11**, 1784; (c) M. C. Galan, A. P. Venot, J. Glushka, A. Imberty and G. J. Boons, *J. Am. Chem. Soc.*, 2002, **124**, 5964; (d) M. C. Galan, A. P. Venot and G. J. Boons, *Biochemistry*, 2003, **42**, 8522; (e) M. C. Galan, A. P. Venot, J. Glushka, A. Imberty and G. J. Boons, *Org. Biomol. Chem.*, 2003, **1**, 3891; (f) M. C. Galan, C. S. Dodson, A. P. Venot and G. J. Boons, *Bioorg. Med. Chem. Lett.*, 2004, **14**, 2205; (g) M. C. Galan, A. P. Venot, R. S. Phillips and G. J. Boons, *Org. Biomol. Chem.*, 2004, **2**, 1376; (h) C. A. G. M. Weijers, M. C. R. Franssen and G. M. Visser, *Biotechnol. Adv.*, 2008, **26**, 436.
- (a) N. J. Andreas, B. Kampmann and K. M. Le-Doare, *Early Hum. Dev.*, 2015, **91**, 629; (b) L. Bode, *Early Hum. Dev.*, 2015, **91**, 619; (c) T. W. Hale and P. Hartmann, *Textbook of Human Lactation*, Hale Publishing, 2007.
- Y. Yu, S. Mishra, X. Song, Y. Lasanajak, K. Bradley, M. Tappert, G. Air, D. Steinhauer, S. Halder, S. Cotmore, P. Tattersall, M. Agbandje-McKenna, R. Cummings and D. Smith, *J. Biol. Chem.*, 2012, **287**, 44784.
- (a) L. Bode, *Glycobiology*, 2012, **22**, 1147; (b) X. Chen, *Adv. Carbohydr. Chem. Biochem.*, 2015, **72**, 113.
- D. S. Newburg, *Curr. Med. Chem.*, 1999, **6**, 117.
- (a) G. Y. Wiederschain and D. S. Newburg, *J. Nutr. Biochem.*, 2001, **12**, 559; (b) S. Thurl, M. Munzert, J. Henker, G. Boehm, B. Muller-Werner, J. Jelinek and B. Stahl, *Br. J. Nutr.*, 2010, **104**, 1261.
- (a) J. P. Hallett and T. Welton, *Chem. Rev.*, 2011, **111**, 3508; (b) M. M. A. Pereira, *Curr. Org. Chem.*, 2012, **16**, 1680; (c) M. C. Galan, R. A. Jones and A. T. Tran, *Carbohydr. Res.*, 2013, **375**, 35; (d) T. Nokami, *J. Synth. Org. Chem., Jpn.*, 2014, **72**, 797.
- A. T. Tran, R. Burden, D. T. Racys and M. C. Galan, *Chem. Commun.*, 2011, **47**, 4526.
- (a) L. Gillard, A. T. Tran, F. D. Boyer and J. M. Beau, *Eur. J. Org. Chem.*, 2016, 1103; (b) L. Ma, S. Sun, X. B. Meng, Q. Li, S. C. Li and Z. J. Li, *J. Org. Chem.*, 2011, **76**, 5652; (c) J. Y. Huang, M. Lei and Y. G. Wang, *Tetrahedron Lett.*, 2006, **47**, 3047; (d) X. He and T. H. Chan, *Tetrahedron*, 2006, **62**, 3389.
- M. C. Galan, A. T. Tran and C. Bernard, *Chem. Commun.*, 2010, **46**, 8968.
- M. C. Galan, A. T. Tran, K. Bromfield, S. Rabbani and B. Ernst, *Org. Biomol. Chem.*, 2012, **10**, 7091.
- I. Sittel and M. C. Galan, *Bioorg. Med. Chem. Lett.*, 2015, **25**, 4329.
- D. Benito-Alifonso, S. Tremel, J. C. Sadler, M. Berry and M. C. Galan, *Chem. Commun.*, 2016, **52**, 4906.
- M. M. Palcic and O. Hindsgaul, *Trends Glycosci. Glycotechnol.*, 1996, **8**, 37.
- Although no significant galactosidase activity was observed, conversion levels seemed to stagnate without portion-wise addition of UDP-Galactose, likely due to donor hydrolysis.
- P. Chaturvedi, C. Warren, M. Altaye, A. Morrow, G. Ruiz-Palacios, L. Pickering and D. Newburg, *Glycobiology*, 2001, **11**, 365.
- (a) G. Y. Wiederschain and D. S. Newburg, *Bioactive Components of Human Milk*, 2001, vol. 501, p. 573; (b) Note that fresh milk was added during the incubation to keep enzyme levels up.



- 21 S. Thurl, J. Henker, M. Siegel, K. Tovar and G. Sawatzki, *Glycoconjugate J.*, 1997, **14**, 795.
- 22 (a) L. Shen, E. F. Grollman and V. Ginsburg, *Proc. Natl. Acad. Sci. U. S. A.*, 1968, **59**, 224; (b) T. Kumazaki and A. Yoshida, *Proc. Natl. Acad. Sci. U. S. A.*, 1984, **81**, 4193.
- 23 E. F. Grollman, A. Kobata and V. Ginsburg, *J. Clin. Invest.*, 1969, **48**, 1489.
- 24 S. Thurl, J. Henker, M. Siegel, K. Tovar and G. Sawatzki, *Glycoconjugate J.*, 1997, **14**, 795.
- 25 (a) S. Thurl, M. Munzert, J. Henker, G. Boehm, B. Mueller-Werner, J. Jelinek and B. Stahl, *Br. J. Nutr.*, 2010, **104**, 1261; (b) S. S. van Leeuwen, R. J. W. Schoemaker, G. J. Gerwig, E. J. M. van Leusen-van Kan, L. Dijkhuizen and J. P. Kamerling, *Glycobiology*, 2014, **24**, 728.
- 26 K. Stangier, M. M. Palcic, D. R. Bundle, O. Hindsgaul and J. Thiem, *Carbohydr. Res.*, 1997, **305**, 511.
- 27 G. Y. Wiederschain and D. S. Newburg, *Biochem. Mol. Med.*, 1996, **58**, 211.
- 28 P. H. Johnson, A. S. R. Donald, J. Feeney and W. M. Watkins, *Glycoconjugate J.*, 1992, **9**, 251.
- 29 A. Tran, R. Jones, J. Pastor, J. Boisson, N. Smith and M. Galan, *Adv. Synth. Catal.*, 2011, **353**, 2593.
- 30 M. M. Palcic, A. P. Venot, R. M. Ratcliffe and O. Hindsgaul, *Carbohydr. Res.*, 1989, **190**, 1.

